

A test of new empirical formulas for the prediction of hyperfine component frequencies in $^{127}\text{I}_2$

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1999 Metrologia 36 159

(<http://iopscience.iop.org/0026-1394/36/2/12>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 38.107.179.212

The article was downloaded on 20/02/2012 at 07:59

Please note that [terms and conditions apply](#).

Erratum

A test of new empirical formulas for the prediction of hyperfine component frequencies in $^{127}\text{I}_2$

A. Razet and S. Picard

Metrologia, 1997, **34**, 181-186

Page 183:

Please note that Figure 1b in this paper is incorrect; Figure 1 should be replaced by the following.




Figure 1. (a) ΔeQq shown as a function of the upper vibrational energy term $G(v')$. A fairly smooth curve is obtained. (b) ΔC as a function of $G(v')$. A large increase is observed close to the dissociation limit. (c) Δd as a function of $G(v')$. The curve shows behaviour similar to ΔC but there are values of different sign. (d) $\Delta \delta$ as a function of $G(v')$: $\Delta \delta$ is, apart from a large increase close to the dissociation limit, more or less constant.