

HYDRODYNAMIC MODELS OF TYPE I X-RAY BURSTS: METALLICITY EFFECTS

JORDI JOSÉ^{1,2}, FERMÍN MORENO¹, ANUJ PARIKH³, AND CHRISTIAN ILIADIS^{4,5}

¹ Departament de Física i Enginyeria Nuclear, EUETIB, Universitat Politècnica de Catalunya, C./ Comte d'Urgell 187, E-08036 Barcelona, Spain;
jordi.jose@upc.edu, moreno@ieec.fcr.es

² Institut d'Estudis Espacials de Catalunya (IEEC), Ed. Nexus-201, C/ Gran Capità 2-4, E-08034 Barcelona, Spain

³ Physik Department E12, Technische Universität München, James-Frank-Strasse, D-85748 Garching, Germany; anuj.pariikh@ph.tum.de

⁴ Department of Physics and Astronomy, University of North Carolina, Chapel Hill, NC 27599-3255, USA; iliadis@unc.edu

⁵ Triangle Universities Nuclear Laboratory, Durham, NC 27708-0308, USA
Received 2009 December 16; accepted 2010 May 24; published 2010 June 30

ABSTRACT

Type I X-ray bursts (XRBs) are thermonuclear stellar explosions driven by charged-particle reactions. In the regime for combined H/He-ignition, the main nuclear flow is dominated by the rp -process (rapid proton-captures and β^+ -decays), the 3α -reaction, and the αp -process (a suite of (α, p) and (p, γ) reactions). The main flow is expected to proceed away from the valley of stability, eventually reaching the proton drip line beyond $A = 38$. Detailed analysis of the relevant reactions along the main path has only been scarcely addressed, mainly in the context of parameterized one-zone models. In this paper, we present a detailed study of the nucleosynthesis and nuclear processes powering type I XRBs. The reported 11 bursts have been computed by means of a spherically symmetric (one-dimensional), Lagrangian, hydrodynamic code, linked to a nuclear reaction network that contains 325 isotopes (from ^1H to ^{107}Te), and 1392 nuclear processes. These evolutionary sequences, followed from the onset of accretion up to the explosion and expansion stages, have been performed for two different metallicities to explore the dependence between the extension of the main nuclear flow and the initial metal content. We carefully analyze the dominant reactions and the products of nucleosynthesis, together with the physical parameters that determine the light curve (including recurrence times, ratios between persistent and burst luminosities, or the extent of the envelope expansion). Results are in qualitative agreement with the observed properties of some well-studied bursting sources. Leakage from the predicted SbSnTe cycle cannot be discarded in some of our models. Production of ^{12}C (and implications for the mechanism that powers superbursts), light p -nuclei, and the amount of H left over after the bursting episodes will also be discussed.

Key words: hydrodynamics – nuclear reactions, nucleosynthesis, abundances – stars: neutron – X-rays: bursts

Online-only material: color figures

1. INTRODUCTION

Type I X-ray bursts (hereafter, XRBs) are cataclysmic stellar events. They are powered by thermonuclear runaways (TNRs) in the H/He-rich envelopes accreted onto neutron stars in close binary systems (see reviews by Bildsten 1998; Lewin et al. 1993, 1995; Psaltis 2006; Schatz & Rehm 2006; Strohmayer & Bildsten 2006). These events constitute the most frequent type of thermonuclear stellar explosion in the Galaxy (the third, in terms of total energy output after supernovae and classical novae), in part because of their short recurrence period (hours to days). About 90 Galactic low-mass X-ray binaries exhibiting such bursting behavior (with burst durations of $\tau_{\text{burst}} \sim 10\text{--}100$ s) have been found since the discovery of XRBs by Grindlay et al. (1976), and independently, by Belian et al. (1976). Type I XRBs and their associated nucleosynthesis have been extensively modeled by different groups (see pioneering work by Woosley & Taam 1976; Maraschi & Cavaliere 1977; Joss 1977), reflecting the astrophysical interest in determining the nuclear processes that power the explosion, the light curve, as well as in providing reliable estimates for the chemical composition of the neutron star surface (see Schatz et al. 1999; Parikh et al. 2008, and references therein).

With a neutron star hosting the explosion, temperatures, and densities in the accreted envelope reach high values: $T_{\text{peak}} > 10^9$ K, and $\rho \sim 10^6$ g cm $^{-3}$. As a result, detailed nucleosynthesis studies require the use of hundreds of isotopes, linked by thousands of nuclear interactions, extending all the way up to the SnSbTe-mass region (Schatz et al. 2001) or beyond (the extent of

the nuclear activity⁶ in the XRB nucleosynthesis study of Koike et al. 2004 reaches ^{126}Xe). Indeed, the extent of the rp -process in XRBs is still not clear: recent experimental work now shows that it will be more difficult to reach the SnSbTe-mass region (Elomaa et al. 2009). Because of computational constraints, XRB nucleosynthesis studies have been traditionally performed using limited nuclear reaction networks, truncated near Ni (Woosley & Weaver 1984; Taam et al. 1993, 1996; all using a 19-isotope network), Kr (Hanawa et al. 1983—274-isotope network; Koike et al. 1999—463 nuclides), Cd (Wallace & Woosley 1984; 16-isotope network), or Y (Wallace & Woosley 1981—250-isotope network). On the other hand, Schatz et al. (1999, 2001) have carried out very detailed nucleosynthesis calculations with a network containing more than 600 isotopes (up to Xe in Schatz et al. 2001), but using a one-zone approach. Koike et al. (2004) have also performed detailed one-zone nucleosynthesis calculations, with temperature and density profiles obtained from a spherically symmetric evolutionary code, linked to a 1270-isotope network extending up to ^{198}Bi .

Until recently, it has not been possible to couple hydrodynamic stellar calculations (in one dimension) and detailed networks. Recent efforts include Fisker et al. (2004, 2006, 2007, 2008), and Tan et al. (2007; ~ 300 isotopes, up to ^{107}Te), José & Moreno (2006; 2640 reactions and 478 isotopes, up to Te),

⁶ The nuclear activity reflects the changes in composition driven by different nuclear processes (i.e., p - and α -capture reactions, β -decays,...) that take place in the envelope at different stages of the burst. In this work, the extent of the nuclear activity is arbitrarily defined by the heaviest nucleus that achieves a mass fraction $> 10^{-9}$.

or Woosley et al. (2004) and Heger et al. (2007; up to 1300 isotopes with an adaptive network). This has prompted a detailed analysis of the nuclear activity powering the bursts. The most detailed work to date is that of Fisker et al. (2008), in the context of the one-dimensional general relativistic hydrodynamic code AGILE (Liebendörfer et al. 2002), linked to a nuclear reaction network containing 304 isotopes: a thorough analysis of the main nuclear activity in one characteristic burst is reported (although details for a sequence of five consecutive, “representative” bursts are also outlined). However, because of the specific choice of metallicity ($Z = 10^{-3}$, for the accreted matter) and mass-accretion rate ($\dot{M} \sim 10^{17} \text{ g s}^{-1}$) adopted, the nuclear activity does not extend much beyond mass $A \sim 65$, as a result of compositional inertia effects, that quench further extension of the nuclear path. Hence, the flow does not reach the SnSbTe-mass region, which was suggested as a natural endpoint in XRB nucleosynthesis studies (see Schatz et al. 1999, 2001).

Clearly, the identification of the most relevant reactions in the $A \sim 65$ –100 mass region remains to be addressed in detail in the framework of hydrodynamic simulations. This is particularly relevant since, as first pointed out by Hanawa et al. (1983), proton captures on heavy nuclei (i.e., the rp -process) have a dramatic effect on the shape of XRB light curves. To this end, a new set of type I XRBs have been computed with SHIVA, a one-dimensional, spherically symmetric, hydrodynamic, implicit, Lagrangian code, used extensively in the modeling of classical nova outbursts (see José & Hernanz 1998). The code has been linked to a fully updated nuclear reaction network containing 324 nuclides and 1392 nuclear processes, a subset of that used in Parikh et al. (2008), and includes the most relevant charged-particle-induced reactions occurring between ^1H and ^{107}Te , as well as their corresponding reverse processes. It is worth noting that the size of this network is similar (though slightly larger) to that adopted by Fisker et al. (2008). In order to set up the reaction rate library for our study, we started by adopting the proton drip line predicted by Audi et al. (2003a, 2003b). Experimental rates are available for a small subset of reactions (adopted from Angulo et al. 1999; Iliadis et al. 2001, and some recent updates for selected reactions). For all other reactions for which experimental rates are not available, we used the rates from the Hauser–Feshbach codes MOST (Goriely 1998; Arnould & Goriely 2006) and NON-SMOKER (Rauscher & Thielemann 2000; for details see Parikh et al. 2008). Neutron captures are disregarded since our early test calculations revealed that they play a minor role in XRB nucleosynthesis. All reaction rates incorporate the effects of thermal excitations in the target nuclei (Rauscher & Thielemann 2000). Screening factors are taken from Graboske et al. (1973) and DeWitt et al. (1973). For the weak interactions, β -delayed nucleon emission and laboratory decay rates (Audi et al. 2003a) have been adopted. For a discussion of employing stellar versus laboratory decay rates, see Woosley et al. (2004). It is worth noting, however, that many computed stellar decay rates (Fuller et al. 1982a, 1982b; Langanke & Martinez-Pinedo 2000) do not converge to their laboratory values at lower temperatures and densities, calling into question the model used for these calculations. Studies employing properly converging stellar decay rates for all isotopes relevant to XRB nucleosynthesis have not been performed by any group yet, and would certainly be interesting, although the results presented in this work would not be dramatically affected by their inclusion.

SHIVA uses a time-dependent formalism for convective transport whenever the characteristic convective timescale becomes

larger than the integration time step. Partial mixing between adjacent convective shells is treated by means of a diffusion equation (Prialnik et al. 1979). No additional semiconvection or thermohaline mixing is considered. Models make use of Iben’s (1975) opacity fits, better suited than the OPAL opacities for astrophysical environments that exhibit strong variations in metallicity, as in XRB nucleosynthesis. However, plans to incorporate these more realistic opacities are currently underway. The adopted equation of state includes contributions from the electron gas (with different degrees of degeneracy; Blinnikov et al. 1996), a multicomponent ion plasma, and radiation; Coulomb corrections to the electronic pressure are also taken into account.

Accretion is computed by redistributing material through a constant number of envelope shells (see Kutter & Sparks 1980 for details). To handle this, a tiny envelope, containing $1.1 \times 10^{18} \text{ g}$ of material (less than 1 permil of the total envelope mass accreted during the first bursting episode), distributed through all the envelope shells, is put initially in place (the influence of the number of envelope shells on burst properties will be discussed in Section 3). The model is then relaxed using a few, very large time steps, to guarantee hydrostatic equilibrium. The temperature at the bottom of the envelope barely reaches $2.7 \times 10^7 \text{ K}$, whereas the density is just $1.4 \times 10^3 \text{ g cm}^{-3}$ (corresponding to a pressure of $5.7 \times 10^{18} \text{ dyn cm}^{-2}$). Mass accretion and nuclear reactions are then initiated.

Special emphasis is placed on the effect of the initial metallicity of the accreted matter on the main nuclear path, which in turn, will affect the final post-burst envelope composition and the shape of the light curves.

The structure of the manuscript is as follows. In Section 2, we analyze the main features (nuclear path, nucleosynthesis, light curves, etc.) of a series of four bursts computed in a model with solar-like accreted material. The effect of the resolution adopted in this model is discussed in Section 3. A detailed analysis of the impact of the metallicity of the accreted material on burst properties is given in Section 4. Finally, a comparison with previous work, together with a thorough analysis of the corrections posed by general relativity, are discussed in Section 5.

2. MODEL 1

We summarize the gross properties of a series of thermonuclear bursts driven by mass accretion onto a $1.4 M_{\odot}$ neutron star ($L_{\text{ini}} = 1.6 \times 10^{34} \text{ erg s}^{-1} = 4.14 L_{\odot}$), at a rate $\dot{M}_{\text{acc}} = 1.75 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$ (corresponding to $0.08 \dot{M}_{\text{Edd}}$). The composition of the accreted material (see Table 1) is assumed to be solar-like ($X = 0.7048$, $Y = 0.2752$, $Z = 0.02$). All metals are initially assumed to be in the form of ^{14}N , following the rapid rearrangement of CNO isotopes that naturally occurs early in the burst (see Woosley et al. 2004). This model is qualitatively similar to model ZM, computed by Woosley et al. (2004) in the framework of the one-dimensional, hydrodynamic, implicit code *KEPLER*. This choice is made intentionally to compare with previous hydrodynamic studies. Note, however, that Woosley et al. assume a value of 10 km for the neutron star radius. In contrast, our model yields a value of 13.1 km, following the integration of the neutron star structure⁷ from the

⁷ The Harrison–Wheeler equation of state (see Shapiro & Teukolsky 1983) has been adopted to build up the initial neutron star configuration in hydrostatic equilibrium conditions. Although this equation of state is a rather crude approximation to the physical conditions in the very deep interior of a neutron star, the radius obtained is in good agreement with the values derived with more accurate equations of state (J. Lattimer 2009, private communication).




Figure 1. Main nuclear activity at the innermost envelope shell for model 1 ($M_{\text{NS}} = 1.4 M_{\odot}$, $\dot{M}_{\text{acc}} = 1.75 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$, $Z = 0.02$), at the early stages of accretion ($T_{\text{base}} = 9.9 \times 10^7 \text{ K}$). Upper panel: mass fractions of the most abundant species ($X > 10^{-5}$). Lower panel: main reaction fluxes ($F \geq 10^{-9} \text{ reactions s}^{-1} \text{ cm}^{-3}$). (A color version of this figure is available in the online journal.)

Table 1
Summary of the Models Computed in this Work^a

Model	$M_{\text{NS}} (M_{\odot})$	Metallicity	Envelope Shells	Bursts Computed
1	1.4	0.02	60	4
2	1.4	0.02	200	2
3	1.4	1×10^{-3}	60	5

Note. ^a A mass-accretion rate of $1.75 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$ has been adopted for all models.

core to its surface, in hydrostatic equilibrium. Differences in the neutron star size (and in turn, in surface gravity) may effect the strength of the explosion (mass accreted, peak temperature, nucleosynthesis, etc.).

2.1. First Burst

The piling-up of solar-like material on top of the neutron star during the accretion stage progressively compresses and

heats the envelope (consisting of 60 shells). Indeed, only 145 s since the beginning of accretion, the temperature at the base of the envelope reaches $T_{\text{base}} = 5 \times 10^7 \text{ K}$ (with ρ_{base} exceeding 10^4 g cm^{-3}).

The early nuclear activity is fully dominated by H-burning through hot CNO-cycle reactions, initiated by proton captures on ^{14}N nuclei. At this stage ($t = 2327 \text{ s}$), the envelope achieves $T_{\text{base}} \sim 10^8 \text{ K}$ ($\rho_{\text{base}} \sim 6.5 \times 10^4 \text{ g cm}^{-3}$), with an energy generation rate of $\epsilon_{\text{nuc}} \sim 1.2 \times 10^{14} \text{ erg g}^{-1} \text{ s}^{-1}$. The main reaction path (see Figure 1) is led by $^{15}\text{N}(p, \alpha)^{12}\text{C}$, which powers $^{12}\text{C}(p, \gamma)^{13}\text{N}(p, \gamma)^{14}\text{O}(\beta^+)^{14}\text{N}(p, \gamma)^{15}\text{O}(\beta^+)^{15}\text{N}$. This suite of nuclear processes competes with $^{13}\text{N}(\beta^+)^{13}\text{C}(p, \gamma)^{14}\text{N}$, and to a lesser extent, with $^{15}\text{N}(p, \gamma)^{16}\text{O}(p, \gamma)^{17}\text{F}(\beta^+)^{17}\text{O}(p, \alpha)^{14}\text{N}$. Besides H ($X = 0.689$) and ^4He ($Y = 0.290$), the next most abundant species in the envelope is now ^{15}O (1.2×10^{-2}). The amount of unburned ^{14}N has dropped to 2.4×10^{-3} (in the following, when discussing the nucleosynthesis, we will refer to abundances by mass, i.e., mass fractions). Other CNO-group nuclei, such as ^{12}C (5.4×10^{-5}), ^{13}N (4.2×10^{-4}),




Figure 2. Same as Figure 1, but for $T_{\text{base}} = 2.1 \times 10^8$ K.

(A color version of this figure is available in the online journal.)

^{14}O (6.1×10^{-3}), or ^{16}O (1.1×10^{-5}), have already achieved an abundance $\geq 10^{-5}$.

T_{base} reaches 2.1×10^8 K 4.49 hr (16,163 s) after the beginning of accretion. Mass accretion in highly degenerate conditions has compressed the envelope base to a density of $\rho_{\text{base}} = 2.7 \times 10^5$ g cm $^{-3}$ ($P_{\text{base}} = 9.1 \times 10^{21}$ dyn cm $^{-2}$). The total luminosity of the star has now increased to a value of 2.5×10^{35} erg s $^{-1}$. The main nuclear activity⁸ (Figure 2) is still dominated by proton captures and β^+ -decays, characteristics of the hot CNO cycle, now supplemented by $^{15}\text{N}(p, \gamma)^{16}\text{O}(p, \gamma)^{17}\text{F}(p, \gamma)^{18}\text{Ne}(\beta^+)^{18}\text{F}(p, \alpha)^{15}\text{O}$, and by the 3α reaction. The numerous p -captures have reduced the hydrogen content to 0.408. In turn, ^4He has increased to 0.570, becoming now the most abundant species at the base of the envelope (followed by the short-lived species ^{14}O [7.7×10^{-3}] and ^{15}O [1.4×10^{-2}]),

⁸ Equilibrium (p, γ) – (γ, p) pairs have been removed from the accompanying plots to highlight those reactions of lower flux that directly lead to the production of heavier species during the burst.

while most of the CNO nuclei have been reduced to 10^{-7} – 10^{-9} , by mass). The extension of the main nuclear activity reaches ^{40}Ca . Indeed, ^{32}S and ^{40}Ca are the only species in the Ne–Ca mass region with abundances exceeding 10^{-9} .

Convection sets in erratically, at ~ 1 m above the core–envelope interface (the overall envelope size, Δz , is ~ 14 m), when T_{base} reaches 3.9×10^8 K, and progressively extends throughout the whole envelope.

Shortly after, at $t = 5.88$ hr (21,181 s), T_{base} reaches 4×10^8 K (with $\rho_{\text{base}} = 2.9 \times 10^5$ g cm $^{-3}$ and $P_{\text{base}} = 1.2 \times 10^{22}$ dyn cm $^{-2}$). The hydrogen content has dropped to 0.209, whereas ^4He achieves 0.625. In turn, the rate of nuclear energy generation has increased to a value of 2.8×10^{16} erg g $^{-1}$ s $^{-1}$. The metallicity of this innermost envelope shell has increased from an initial value of 0.02 to 0.17, due to leakage from the CNO cycle (mainly powered by $^{15}\text{O}(\alpha, \gamma)$). As before, the next most abundant species are ^{14}O (6.9×10^{-2}) and ^{15}O (6.5×10^{-2}), but the number of isotopes with moderately large abundances




Figure 3. Same as Figure 1, but for $T_{\text{base}} = 4 \times 10^8$ K.
(A color version of this figure is available in the online journal.)

has now increased. Indeed, ^{40}Ca , ^{22}Mg , ^{18}Ne , ^{34}Ar , ^{48}Cr , and $^{42,44}\text{Ti}$ have achieved mass fractions of the order of 10^{-3} . The nuclear activity extends as far as ^{53}Co . The largest reaction fluxes (defined as the number of reactions per unit time and volume) correspond to the equilibrium processes $^{21}\text{Mg}(p, \gamma)^{22}\text{Al}(\gamma, p)^{21}\text{Mg}$, $^{30}\text{S}(p, \gamma)^{31}\text{Cl}(\gamma, p)^{30}\text{S}$, and $^{25}\text{Si}(p, \gamma)^{26}\text{P}(\gamma, p)^{25}\text{Si}$. Additional activity is powered by $3\alpha \rightarrow ^{12}\text{C}(p, \gamma)^{13}\text{N}(p, \gamma)^{14}\text{O}$, followed by $^{14}\text{O}(\alpha, p)^{17}\text{F}(p, \gamma)^{18}\text{Ne}$. The suite of secondary nuclear paths is rich and complex (see Figure 3), and is mainly dominated by p -capture reactions and β^+ -decays, as well as by the CNO-breakout reaction $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$. It is worth noting that the main nuclear path above Ca begins to move away from the valley of stability, toward the proton-drip line (see Figure 3, lower panel).

Just 2.3 s later ($t = 21,183$ s), T_{base} achieves 5×10^8 K. ρ_{base} has slightly decreased to 2.3×10^5 g cm $^{-3}$ because of a mild envelope expansion ($\Delta z \sim 15.5$ m). Note, however, that $P_{\text{base}} = 1.2 \times 10^{22}$ dyn cm $^{-2}$. Hence, the TNR is taking place nearly

at constant pressure. A time-dependent, convective mixing with adjacent shells, with a characteristic timescale of $\tau_{\text{conv}} \sim 10^{-4}$ s ($v_{\text{conv}} \sim 10^3\text{--}10^5$ cm s $^{-1}$), causes a slight increase in the H abundance at the base of the envelope. Indeed, the H abundance is now 0.288, by mass, whereas the ^4He content has slightly decreased to 0.563 (due to the high temperatures, favoring α -captures). The next most abundant species is now ^{18}Ne (4.4×10^{-2}), together with $^{14,15}\text{O}$ (4.2×10^{-2} and 1.8×10^{-2} , respectively). Several isotopes, such as $^{21,22}\text{Mg}$, $^{29,30}\text{S}$, $^{50,52}\text{Fe}$, ^{27}P , $^{24,25}\text{Si}$, $^{49,50,51}\text{Mn}$, and ^{34}Ar , have achieved abundances an order of magnitude lower ($\sim 10^{-3}$). The nuclear activity extends up to ^{57}Cu now, powering an energy generation rate of 1.2×10^{17} erg g $^{-1}$ s $^{-1}$, and an overall luminosity of 1.3×10^{36} erg s $^{-1}$. The largest fluxes still correspond to the forward and reverse reactions $^{30}\text{S}(p, \gamma)^{31}\text{Cl}(\gamma, p)^{30}\text{S}$, $^{21}\text{Mg}(p, \gamma)^{22}\text{Al}(\gamma, p)^{21}\text{Mg}$, and $^{25}\text{Si}(p, \gamma)^{26}\text{P}(\gamma, p)^{25}\text{Si}$, together with $^{14}\text{O}(\alpha, p)^{17}\text{F}(p, \gamma)^{18}\text{Ne}$. $^{14}\text{O} + \alpha$ becomes the most important α -capture reaction, overcoming $^{15}\text{O}(\alpha, \gamma)$, or the 3α . Additional




Figure 4. Same as Figure 1, but for $T_{\text{base}} = 7 \times 10^8$ K.

(A color version of this figure is available in the online journal.)

activity is driven by $^{18}\text{Ne}(\beta^+)^{18}\text{F}(p, \alpha)^{15}\text{O}$, $^{19}\text{Ne}(p, \gamma)^{20}\text{Na}(p, \gamma)^{21}\text{Mg}$, $^{21}\text{Na}(p, \gamma)^{22}\text{Mg}(p, \gamma)^{23}\text{Al}$, $^{12}\text{C}(p, \gamma)^{13}\text{N}(p, \gamma)^{14}\text{O}$, and $^{26}\text{Si}(p, \gamma)^{27}\text{P}$.

A qualitatively similar picture is found when T_{base} achieves 7×10^8 K ($t = 21,185$ s), with the most abundant species at the envelope base being H (0.308), ^4He (0.507), ^{18}Ne (4.8×10^{-2}), ^{22}Mg (2.5×10^{-2}), $^{29,30}\text{S}$ (1.1×10^{-2} , and 2.2×10^{-2} , respectively), and $^{24,25}\text{Si}$ (1.1×10^{-2} and 1.7×10^{-2} , respectively). The number of species with abundances of the order of 10^{-3} includes now $^{54,55,56}\text{Ni}$, ^{15}O , ^{28}S , ^{52}Fe , ^{27}P , ^{38}Ca , and $^{33,34}\text{Ar}$, with the main nuclear activity (see Figure 4) extending all the way up to ^{60}Zn . The largest reaction fluxes are achieved by the equilibrium processes described before, supplemented now by $^{26}\text{Si}(p, \gamma)^{27}\text{P}(p, \gamma)^{26}\text{Si}$, $^{22}\text{Mg}(p, \gamma)^{23}\text{Al}(p, \gamma)^{22}\text{Mg}$, $^{29}\text{S}(p, \gamma)^{30}\text{Cl}(p, \gamma)^{29}\text{S}$, and $^{16}\text{O}(p, \gamma)^{17}\text{F}(p, \gamma)^{16}\text{O}$, followed by p -capture reactions (and β^+ -decays) on Ne–Mg nuclei, such as $^{19}\text{Ne}(p, \gamma)^{20}\text{Na}(p, \gamma)^{21}\text{Mg}(\beta^+)^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$, or $^{17}\text{F}(p, \gamma)^{18}\text{Ne}(\beta^+)^{18}\text{F}(p, \alpha)^{15}\text{O}$, and by the α -capture reactions $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$, $^{14}\text{O}(\alpha, p)^{17}\text{F}$, and the 3α .

One second later ($t = 21,186$ s), T_{base} achieves 9×10^8 K. The hectic nuclear activity, which at this stage releases $\epsilon_{\text{nuc}} \sim 3.7 \times 10^{17} \text{ erg g}^{-1} \text{ s}^{-1}$, has reduced the H and ^4He abundances down to 0.262 and 0.457, respectively. The next most abundant species are now ^{22}Mg , ^{25}Si , $^{28,29,30}\text{S}$, $^{33,34}\text{Ar}$, and ^{60}Zn , all with mass fractions $\sim 10^{-2}$. The main nuclear activity has extended up to ^{68}Se . Aside from equilibrium (p, γ) and (γ, p) pairs (that involve ^{16}O – ^{17}F and a handful of species in the mass range Mg–Zn), the largest reaction fluxes correspond to a suite of p -captures and β^+ -decays (see Figure 5, lower panel), mainly $^{25}\text{Al}(p, \gamma)^{26}\text{Si}$ and $^{27,28,29}\text{P}(p, \gamma)^{28,29,30}\text{S}$. Moreover, the most important α -capture reactions, $^{22}\text{Mg}(\alpha, p)^{25}\text{Al}$, the 3α , $^{18}\text{Ne}(\alpha, p)^{21}\text{Na}$, and $^{14}\text{O}(\alpha, p)^{17}\text{F}$, have fluxes of the order of $\log F \sim -3$ (note the moderate extension of α -captures toward heavier species as a result of the higher temperatures). A very limited nuclear activity in the $A = 65$ – 100 mass region is, at this stage, driven by $^{65}\text{Ge}(p, \gamma)^{66}\text{As}(p, \gamma)^{67}\text{Se}(\beta^+)^{67}\text{As}(p, \gamma)^{68}\text{Se}$ (with $\log F \sim -8$), and $^{66}\text{As}(\beta^+)^{66}\text{Ge}(p, \gamma)^{67}\text{As}$ (as $\log F \sim -9$).




Figure 5. Same as Figure 1, but for $T_{\text{base}} = 9 \times 10^8$ K.
(A color version of this figure is available in the online journal.)

At $t = 21,188$ s, when T_{base} achieves 1×10^9 K, the energy generation rate by nuclear reactions reaches its maximum value: $\epsilon_{\text{nuc,max}} \sim 4.1 \times 10^{17} \text{ erg g}^{-1} \text{ s}^{-1}$. Two seconds later, the envelope will attain maximum expansion, with a size $\Delta z_{\text{max}} \sim 44$ m.

Proton and α -captures continue to reduce the overall H and He abundances at the envelope base (0.191 and 0.400, respectively). The next most abundant species is now ^{30}S (0.103)—a waiting point for the main nuclear path—followed by $^{33,34}\text{Ar}$, $^{37,38}\text{Ca}$, ^{42}Ti , ^{46}Cr , ^{50}Fe , ^{56}Ni , and ^{60}Zn (all with $X_i \sim 10^{-2}$). The nuclear activity has reached ^{76}Sr (see Figure 6). Time evolution of density, temperature, pressure, and rate of nuclear energy generation, at the innermost envelope shell, as well as of the overall neutron star luminosity and envelope size, are shown in Figures 7 and 8. In the $A = 65$ –100 mass region, in particular, the nuclear activity is now dominated by the chains $^{65}\text{Ge}(p, \gamma)^{66}\text{As}(p, \gamma)^{67}\text{Se}(\beta^+)^{67}\text{As}(p, \gamma)^{68}\text{Se}$ (with $\log F \sim -7$), $^{66}\text{As}(\beta^+)^{66}\text{Ge}(p, \gamma)^{67}\text{As}$ ($\log F \sim -8$), $^{65}\text{As}(p, \gamma)^{66}\text{Se}(\beta^+)^{66}\text{As}$, and $^{68}\text{Se}(\beta^+)^{68}\text{As}(p, \gamma)^{69}\text{Se}(p, \gamma)^{70}\text{Br}(p, \gamma)^{71}\text{Kr}(\beta^+)^{71}\text{Br}(p,$

$\gamma)^{72}\text{Kr}$ ($\log F \sim -9$). In terms of energy production, the most important contributions come from $^{39}\text{Ca}(p, \gamma)^{40}\text{Sc}$ ($\log F \sim -2$), multiple (p, γ) -reactions involving species in the mass range $A \sim 20$ –60, and a handful of α -capture reactions, such as $^{22}\text{Mg}(\alpha, p)^{25}\text{Al}$, $^{18}\text{Ne}(\alpha, p)^{21}\text{Na}$, the 3α , and $^{14}\text{O}(\alpha, p)^{17}\text{F}$ ($\log F \sim -3$).

Four seconds later, the envelope base achieves a maximum temperature of $T_{\text{peak}} \sim 1.06 \times 10^9$ K (similar values are reported in the simulations by Fisker et al. 2008). Besides H (0.220) and ^4He (0.370), the next most abundant isotope is now ^{60}Zn (0.159)—another waiting point for the nuclear flow—followed by ^{30}S (3.3×10^{-2}), ^{34}Ar (3.2×10^{-2}), ^{38}Ca (2.4×10^{-2}), ^{46}Cr (1.8×10^{-2}), ^{50}Fe (1.3×10^{-2}), $^{55,56}\text{Ni}$ (1.8×10^{-2} and 2.3×10^{-2} , respectively), and ^{59}Zn (1.8×10^{-2}). As shown in Figure 9, the extension of the main nuclear path reaches ^{80}Zr . The nuclear activity in the $A = 65$ –100 mass region is now dominated by $^{65}\text{Ge}(p, \gamma)^{66}\text{As}(p, \gamma)^{67}\text{Se}(\beta^+)^{67}\text{As}(p, \gamma)^{68}\text{Se}$ ($\log F \sim -6$), $^{65}\text{As}(p, \gamma)^{66}\text{Se}(\beta^+)^{66}\text{As}$ ($\log F \sim -7$), and




Figure 6. Same as Figure 1, but for $T_{\text{base}} = 10^9$ K.

(A color version of this figure is available in the online journal.)

$^{66}\text{As}(\beta^+)^{66}\text{Ge}(p, \gamma)^{67}\text{As}$, $^{68}\text{Se}(\beta^+)^{68}\text{As}(p, \gamma)^{69}\text{Se}(p, \gamma)^{70}\text{Br}(p, \gamma)^{71}\text{Kr}(\beta^+)^{71}\text{Br}(p, \gamma)^{72}\text{Kr}$ ($\log F \sim -8$). Energy production is due to dozens of (p, γ) -reactions involving species in the mass range $A \sim 20$ –60, plus some α -capture reactions, such as the 3α , $^{14}\text{O}(\alpha, p)^{17}\text{F}$, $^{22}\text{Mg}(\alpha, p)^{25}\text{Al}$, and $^{18}\text{Ne}(\alpha, p)^{21}\text{Na}$ ($\log F \sim -3$). Less than a second later ($t = 21,192.3$ s), the neutron star reaches maximum luminosity, $L_{\text{max}} = 3.8 \times 10^{38} \text{ erg s}^{-1}$ ($9.8 \times 10^4 L_{\odot}$).

The numerous proton captures on many species during the decline from T_{peak} reduce dramatically the H content in the innermost shell. Indeed, when T_{base} achieves 9.3×10^8 K ($t = 21,200$ s), the H abundance drops below 0.1, while $X(^4\text{He}) = 0.283$. Actually, the most abundant species in this shell is now ^{60}Zn (0.43 by mass), followed by ^{30}S (2.7×10^{-2}), ^{34}Ar (2.1×10^{-2}), ^{38}Ca (1.3×10^{-2}), ^{56}Ni (1.8×10^{-2}), and ^{64}Ge (3.3×10^{-2}). The nuclear activity reaches ^{90}Ru .

Five seconds later ($t = 21,205$ s), when T_{base} drops to 9.0×10^8 K, ^{60}Zn achieves a maximum abundance of 0.519,

by mass. H has been reduced to 1.3×10^{-2} ($X(^4\text{He}) = 0.226$). The next most abundant species are now ^{26}Si , ^{30}S , ^{34}Ar , ^{38}Ca , ^{56}Ni , and ^{64}Ge (all with mass fractions $\sim 10^{-2}$). The nuclear activity has not progressed beyond ^{90}Ru . The largest reaction fluxes correspond to proton captures and reverse photodisintegration reactions at equilibrium. Many other nuclear processes (β^+ -decays and α -induced reactions like the triple- α , $^{14}\text{O}(\alpha, p)^{17}\text{F}$, or $^{22}\text{Mg}(\alpha, p)^{25}\text{Al}$) contribute to the overall nuclear activity (see Figure 10). In the $A = 65$ –100 mass region, this is driven by $^{65}\text{Ge}(p, \gamma)^{66}\text{As}$ (followed either by $^{66}\text{As}(p, \gamma)^{67}\text{Se}(\beta^+)^{67}\text{As}$ or by $^{66}\text{As}(\beta^+)^{66}\text{Ge}(p, \gamma)^{67}\text{As}$), $^{67}\text{As}(p, \gamma)^{68}\text{Se}(\beta^+)^{68}\text{As}(p, \gamma)^{69}\text{Se}$ ($\log F \sim -5$), $^{69}\text{Se}(p, \gamma)^{70}\text{Br}(\beta^+)^{70}\text{Se}(p, \gamma)^{71}\text{Br}(\beta^+)^{71}\text{Se}$, $^{74}\text{Rb}(\beta^+)^{74}\text{Kr}$ ($\log F \sim -6$), and $^{70}\text{Br}(p, \gamma)^{71}\text{Kr}(\beta^+)^{71}\text{Br}$, $^{72}\text{Kr}(\beta^+)^{72}\text{Br}(p, \gamma)^{73}\text{Kr}(p, \gamma)^{74}\text{Rb}$, $^{74}\text{Kr}(p, \gamma)^{75}\text{Rb}(p, \gamma)^{76}\text{Sr}$ ($\log F \sim -7$). Energy production is due to (p, γ) -reactions involving species in the mass range $A \sim 20$ –55, and also due to two α -capture reactions, $^{14}\text{O}(\alpha, p)^{17}\text{F}$ and $^{22}\text{Mg}(\alpha, p)^{25}\text{Al}$ ($\log F \sim -3$).




Figure 7. Time evolution of density (upper left panel), temperature (upper right), pressure (lower left), and nuclear energy generation rate (lower right), at the innermost envelope shell for model 1 ($M_{\text{NS}} = 1.4 M_{\odot}$, $\dot{M}_{\text{acc}} = 1.75 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$, $Z = 0.02$), along the first bursting episode. The origin of the time coordinate is arbitrarily chosen as 21,150 s, for which $T_{\text{base}} \sim 3 \times 10^8 \text{ K}$.




Figure 8. Same as Figure 7, but for the overall neutron star luminosity (left panel), and envelope size (right panel), as measured from the core–envelope interface.




Figure 9. Same as Figure 1, but for the time when temperature at the envelope base reaches a peak value of $T_{\text{peak}} = 1.06 \times 10^9$ K. (A color version of this figure is available in the online journal.)

Following the fast decline in temperature, when $T_{\text{base}} = 8.0 \times 10^8$ K ($t = 21,212$ s), the ^{60}Zn abundance has dropped to 0.509, due to β^+ -decays. H has been heavily depleted (5×10^{-11}), whereas ^4He has been slightly reduced to an abundance of 0.190. The next most abundant species are ^{12}C , ^{30}P , ^{39}K , ^{56}Ni , ^{60}Cu , and ^{64}Ge . The extent of the nuclear activity (Figure 11) is still restricted to ^{90}Ru ; it will not proceed beyond this endpoint, first because of the heavy H depletion, and second, because the temperature is already too low to allow proton- or α -captures on heavier species due to their large Coulomb barriers. At this stage, the single, most important reaction, in terms of reaction fluxes, is the triple- α , followed by a suite of β^+ -decay reactions, such as $^{26}\text{Si}(\beta^+)^{26m}\text{Al}(\beta^+)^{26}\text{Mg}$, $^{34}\text{Cl}(\beta^+)^{34}\text{S}$, $^{60}\text{Zn}(\beta^+)^{60}\text{Cu}$, or $^{27}\text{Si}(\beta^+)^{27}\text{Al}$. Several α -captures follow the triple- α reaction as a chain: $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}(\alpha, \gamma)^{20}\text{Ne}(\alpha, \gamma)^{24}\text{Mg}(\alpha, \gamma)^{28}\text{Si}(\alpha, \gamma)^{32}\text{S}$, or through alternative paths, proceeding close to the valley of stability, up to $\sim \text{Ar}$, such as $^{13}\text{N}(\alpha, p)^{16}\text{O}$, $^{25,27}\text{Al}(\alpha, p)^{28,30}\text{Si}$, $^{22}\text{Mg}(\alpha, p)^{25}\text{Al}$, $^{22}\text{Na}(\alpha,$

$p)^{25}\text{Mg}$, or $^{26,27}\text{Si}(\alpha, p)^{29,30}\text{P}$, to quote some representative cases (see Figure 11). ϵ_{nuc} has already declined to a value of $\sim 3.8 \times 10^{15}$ erg $\text{g}^{-1} \text{s}^{-1}$.

When T_{base} drops to 4.3×10^8 K ($t = 21,254$ s), the nuclear energy generation rate has declined to a value of $\epsilon_{\text{nuc}} \sim 3 \times 10^{14}$ erg $\text{g}^{-1} \text{s}^{-1}$. As shown in Figure 12, the nuclear activity is dominated by $^{60}\text{Zn}(\beta^+)^{60}\text{Cu}$, because of its very large abundance ($X(^{60}\text{Zn}) = 0.416$), and is followed by the triple- α reaction ($X(^4\text{He}) = 0.137$), and by a suite of β^+ -decays of very abundant isotopes, such as ^{64}Ge , ^{30}P , ^{64}Ga , or ^{60}Cu (all with mass fractions $\sim 10^{-2}$, except ^{60}Cu [0.104]), followed by those of ^{38}K , ^{26m}Al , ^{68}Se , ^{25}Al , ^{68}As , ^{63}Ga , ^{59}Cu , and ^{61}Zn . Other species, such as ^{12}C , ^{26}Mg , ^{34}S , ^{39}K , or ^{56}Ni , have achieved an abundance of 10^{-2} by mass at this stage. The envelope has already shrunk to a size $\Delta z \sim 13$ m, whereas the overall luminosity of the star has decreased to $L_{\text{NS}} = 7.7 \times 10^{36}$ erg s^{-1} .

When T_{base} reaches 2×10^8 K ($t = 21,618$ s), ^{60}Zn has remarkably decayed into ^{60}Cu , which now constitutes the most




Figure 10. Same as Figure 1, but for $T_{\text{base}} = 9 \times 10^8$ K.
(A color version of this figure is available in the online journal.)

abundant species (with 0.393) at the base of the envelope. Because of the relatively low temperatures, the ^4He abundance is kept constant, at about 0.136. The next most abundant species are now ^{12}C , ^{26}Mg , ^{30}Si , ^{34}S , ^{39}K , ^{56}Ni , ^{60}Ni , ^{64}Ga , and ^{64}Zn . With the exception of the limited contribution of the triple- α reaction, the main nuclear path (Figure 13) is fully dominated by a suite of β^+ -decays on numerous species, all the way up to ^{72}Br .

When $t = 28,250$ s, T_{base} reaches a minimum value of 1.67×10^8 K, which we consider to mark the end of the first burst in our simulations. ^{60}Cu has decayed already into ^{60}Ni , now the most abundant species at the envelope base with a mass fraction of 0.504, followed by ^4He (0.136, not fully consumed during the TNR), and by ^{12}C , ^{26}Mg , ^{30}Si , ^{34}S , ^{39}K , ^{56}Ni , ^{60}Cu , and ^{64}Zn (see Figure 14). The marginal nuclear activity played by a suite of β -decays powers a rate of nuclear energy generation of $\epsilon_{\text{nuc}} \sim 8.8 \times 10^{11} \text{ erg g}^{-1} \text{ s}^{-1}$. At this stage, the size of the envelope has shrunk to $\Delta z \sim 8$ m, whereas

the overall luminosity of the star has decreased to $L_{\text{NS}} = 8.3 \times 10^{34} \text{ erg s}^{-1}$.

Profiles of density, temperature, rate of energy generation, pressure, and size, along the accreted envelope, for different snapshots during the first bursting episode computed in this model, are shown in Figures 15 and 16.

It is worth noting that the main nucleosynthetic activity takes place at the innermost, hottest envelope shell. Because of their lower temperatures and densities, all layers above the ignition shell show a similar nucleosynthetic pattern but somewhat diluted, limiting the extent of the nuclear activity to lower masses. Even though the specific reaction sequences have a clear dependence with depth (see Fisker et al. 2008, for the corresponding analysis in one bursting episode), it is clear that the identification of the main nuclear processes responsible for the nucleosynthesis in XRBs can rely on an accurate account of the activity at the ignition shell, at different stages of the TNR, as performed in this paper. However, we would like to




Figure 11. Same as Figure 1, but for $T_{\text{base}} = 8 \times 10^8$ K.
(A color version of this figure is available in the online journal.)

outline schematically how depth influences the extent of the nuclear activity throughout the envelope: while in the innermost shells of our computational domain (encompassing 6.8×10^{20} g) the nuclear activity reaches ^{90}Ru , at 1.7×10^{21} g above the core–envelope interface, the activity stops around ^{78}Sr (the final mass fraction of ^{90}Ru barely reaches 10^{-15} , by mass), while close to the surface (2.3×10^{21} g), the nuclear activity does not extend beyond ^{72}Se ($X(^{90}\text{Ru}) \sim 10^{-18}$). Moreover, the purely nucleosynthetic imprint in these shells is difficult to assess since it is partially poisoned by changes in the chemical composition driven by convective mixing.

All in all, the mean, mass-averaged chemical composition of the envelope at the end of this first burst is mainly dominated by the presence of intermediate-mass elements (far below the SnSbTe-mass region). This includes ^{60}Ni (0.32), ^4He (0.31), ^1H (0.17), ^{64}Zn (0.03), ^{12}C (0.02), ^{52}Fe (0.02), or ^{56}Fe (0.02) (see Table 2, for the mean composition of all species—stable

or with a half-life > 1 hr—which achieve $X_i > 10^{-9}$), with a nucleosynthesis endpoint (defined by the heaviest isotope with $X_i > 10^{-9}$) around ^{89}Nb (in agreement with the results reported by Fisker et al. 2008). In terms of overproduction factors, f (ratio of the mass-averaged composition of a given isotope over its solar abundance; see also Figures 17 and 19), ^{43}Ca , ^{45}Sc , ^{49}Ti , ^{51}V , $^{60,61}\text{Ni}$, $^{63,65}\text{Cu}$, $^{64,67,68}\text{Zn}$, ^{69}Ga , ^{74}Se , and ^{78}Kr achieve a value of $f \sim 10^4$.

It is important to stress that the presence of unburned H and ^4He in the envelope, at the end of the first burst, will have consequences for the subsequent eruptions (see Section 2.2). Note, however (Figure 17), that since the innermost envelope is devoid of H, the next burst will likely initiate well above the core–envelope interface. Moreover, the presence of unburned ^{12}C , particularly in the inner envelope layers, has important implications for studies of the physical mechanism that powers superbursts (see Section 2.2).




Figure 12. Same as Figure 1, but for $T_{\text{base}} = 4.3 \times 10^8$ K.
(A color version of this figure is available in the online journal.)

2.2. Second, Third, and Fourth Bursts

For conciseness, we will focus here on the main differences between the first and successive bursts computed for model 1. A first, remarkable difference is due to the so-called *compositional inertia* (Taam 1980; Woosley et al. 2004), which accounts for differences in the gross properties of the bursts driven by changes in the chemical content of the envelope. Indeed, after the first burst, the accreted matter will pile up on top of a metal-enriched envelope (the initial metallicity, $Z_{\text{ini}} \sim 0.02$, has risen to a mass-averaged value of $Z \sim 0.52$, at the end of the first burst) that is devoid of H at its innermost layers. This will cause a shift in the location of the ignition region, progressively moving away from the core–envelope interface in successive flashes (see Figure 18, right panel).

This is schematically shown as well in Figure 18 (left panel), which depicts the mass above the neutron star core, as well as the extension of the convective regions throughout the envelope,

for the four bursting episodes computed in this model: the peaks of the explosions (~ 10 – 100 s) correspond to the flat regions of the diagram, whereas the stages of steady accretion (~ 5 hr) are indicated by the steep slopes. Note that, in agreement with previous work (Woosley et al. 2004; Fisker et al. 2008), convection mainly develops around the peak of the bursts (during most of the explosion, energy transport is carried by radiation only). Note also that because of fuel consumption (H, He), the location of the ignition shell (and the extent of the convective regions) moves progressively away in mass from the neutron star core.

From the nucleosynthesis viewpoint (Table 2 and Figures 17 and 19), the nuclear activity extends progressively toward heavier species, reaching endpoints ($X_i > 10^{-9}$) around ^{89}Nb (1st burst), ^{97}Ru (2nd burst), ^{99}Rh (3rd burst), and ^{100}Pd (4th burst). The overall mean metallicity of the envelope at the end of each burst is 0.52 (1st burst), 0.71 (2nd burst), 0.80 (3rd burst), and 0.86 (4th burst). This increase in Z reflects both the nuclear




Figure 13. Same as Figure 1, but for $T_{\text{base}} = 2 \times 10^8$ K.
(A color version of this figure is available in the online journal.)

activity during each individual burst and the accumulated ashes from previous bursts. A similar mass-averaged ^{12}C yield of ~ 0.02 is systematically obtained at the end of each of the four bursts computed. This is not enough to power a superburst, which requires $X(^{12}\text{C})_{\text{min}} \geq 0.1$, at the envelope base (see Cumming & Bildsten 2001; Strohmayer & Brown 2002; Brown 2004; Cooper & Narayan 2004, 2005; Cumming 2005, or Cooper et al. 2006). With respect to overproduction factors, the increase in nuclear activity reported for successive bursts translates also into larger values, as high as $f \sim 10^6$, for ^{76}Se , $^{78,80}\text{Kr}$, and ^{84}Sr , or $f \sim 10^5$, for species such as $^{64,68}\text{Zn}$, $^{72,73}\text{Ge}$, $^{74,77}\text{Se}$, ^{82}Kr , $^{86,87}\text{Sr}$, ^{89}Y , and ^{94}Mo , for the 4th bursting episode (see Figures 17 and 19).

A summary of the gross properties of the four bursts computed in this model is given in Table 3. Peak temperatures and luminosities amount to $T_{\text{peak}} \sim (1.1\text{--}1.3) \times 10^9$ K and $L_{\text{peak}} \sim (1\text{--}2) \times 10^5 L_{\odot}$, respectively. Recurrence times between bursts

of $\tau_{\text{rec}} \sim 5\text{--}6.5$ hr and ratios between persistent⁹ and burst luminosities of $\alpha \sim 35\text{--}40$ (except for the first burst) have been obtained. These values are in agreement with those inferred from some observed XRB sources (see Galloway et al. 2008) such as the *textbook burster* GS 1826-24 [$\tau_{\text{rec}} = 5.74 \pm 0.13$ hr, $\alpha = 41.7 \pm 1.6$], 4U 1323-62 [$\tau_{\text{rec}} = 5.3$ hr, $\alpha = 38 \pm 4$], or 4U 1608-52 [$\tau_{\text{rec}} = 4.14\text{--}7.5$ hr, $\alpha = 41\text{--}54$]. As reported by Woosley et al. (2004), there is also some trend toward stabilization of these values with increasing burst number.

Figure 20 depicts the corresponding light curves from the second to the fourth burst. A quite interesting feature, observed in

⁹ We define $\alpha = \int_t^{t+\tau_{\text{rec}}} L(t) dt / \int_t^{t'+\tau_{0.01}} L(t) dt$, with the latter term integrated over the time during which the burst exceeds 1% of its peak luminosity, $\tau_{0.01}$. Note that during the interburst period, the accretion luminosity, $L_{\text{acc}} = GMM/R \sim 1.5 \times 10^{37}$ erg s⁻¹, will hide the thermal emission from the cooling ashes, as shown in Figures 8 and 20.




Figure 14. Same as Figure 1, but for the time when temperature at the envelope base achieves a minimum value of $T_{\min} = 1.67 \times 10^8$ K. (A color version of this figure is available in the online journal.)

some XRBs such as 4U 1608-52 (Penninx et al. 1989), 4U 17+2 (Kuulkers et al. 2002), or 4U 1709-267 (Jonker et al. 2004), is the appearance of a double-peaked burst in Figure 20 (lower left panel). Double- (or triple-) peaked bursts can be classified in two categories (Watts & Maurer 2007): the first one corresponds to the so-called *photospheric radius expansion* bursts, which exhibit multi-peaked bursts in the X-ray band but not in the bolometric luminosity. The second type of multi-peaked events is also visible in the bolometric light curves, and has been attributed to different causes, such as a stepped release of thermonuclear energy caused either by mixing induced by hydrodynamic instabilities (Fujimoto et al. 1988) or driven by a nuclear waiting-point impedance in the thermonuclear reaction flow (Fisker et al. 2004). A preliminary analysis of the 4th burst reported for model 1 suggests a likely nuclear physics origin (waiting-point impedance) for this double-peaked feature (see J. José & F. Moreno 2010, in preparation). The main nuclear activity and the dominant reaction fluxes at peak temperature, for the 4th burst computed in this model, are shown in Figure 21.

3. MODEL 2

In the previous section, we have reported results from a sequence of type I XRBs computed with a coarse resolution, in which the accreted envelope was discretized in 60 shells. We have checked the influence of the adopted number of envelope shells on the gross properties of the bursts by performing another simulation, identical to model 1, but computed with a finer resolution: 200 shells (hereafter, model 2).

A summary of the main properties of the two bursts computed for model 2 is given in Table 4: the recurrence times obtained are in the same range as those reported for model 1 (4–6 hr). The same applies to the ratios between persistent and burst luminosities, as well as to peak temperatures and luminosities. Similar light curves have also been obtained.

There is also good agreement from the nucleosynthesis viewpoint, with only minor differences in the final, mass-averaged abundances, as shown in Table 5 (particularly, for the heavier species of the network, since their low abundances are very sensitive to the specific thermal history of the explosion; see




Figure 15. Profiles of density (upper left panel), temperature (upper right), nuclear energy generation rate (lower left), and envelope size (measured from the core–envelope interface; lower right panel), for model 1 ($M_{\text{NS}} = 1.4 M_{\odot}$, $\dot{M}_{\text{acc}} = 1.75 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$, $Z = 0.02$), along the first bursting episode. Labels indicate different moments during the TNR, for which T_{base} reaches a value of (1) 9.9×10^7 K, (2) 2.1×10^8 K, (3) 4×10^8 K, (4) 6×10^8 K, (5) 8×10^8 K, (6) 1.06×10^9 K (T_{peak}), (7) 9.3×10^8 K, (8) 7×10^8 K, (9) 4.3×10^8 K, and (10) 1.7×10^8 K (T_{min}). Note the dramatic decrease in ϵ_{nuc} in the innermost layers of the envelope (profiles 8 to 10) as H is depleted.

(A color version of this figure is available in the online journal.)

also Figure 19). It is worth noting that both models reach almost identical nucleosynthesis endpoints ($X_i > 10^{-9}$): ^{89}Nb , for the first burst computed in both models, and ^{97}Ru (model 1) and ^{99}Rh (model 2), for the second burst. Furthermore, the amounts of unburned H, ^4He , and ^{12}C are very similar in both models. As expected from the above-mentioned similarities, there is also good agreement in terms of overproduction factors, dominated by ^{64}Zn and ^{60}Ni (with $f \sim 10^4$) in the first burst and by ^{64}Zn , ^{72}Ge , $^{74,76}\text{Se}$, $^{78,80}\text{Kr}$, and ^{84}Sr ($f \sim 10^5$), in the second, for both models.

All in all, we conclude that the resolution adopted in model 1 is appropriate for XRB simulations. This is in agreement with the studies performed by Fisker et al. (2004), who concluded that the minimum discretization of the envelope, in one-dimensional hydrodynamic simulations of XRBs, is about 25 shells.

4. MODEL 3

To test the impact of the metallicity of the accreted material (which reflects the surface composition of the companion star) on the overall properties of the bursts, we have computed another series of bursts (hereafter, model 3), driven by accretion of metal-deficient material ($Z \sim Z_{\odot}/20$) onto a $1.4 M_{\odot}$ neutron star ($L_{\text{ini}} = 1.6 \times 10^{34} \text{ erg s}^{-1} = 4.14 L_{\odot}$), at a rate $\dot{M}_{\text{acc}} = 1.75 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$. The composition of the accreted material is assumed to be $X = 0.759$, $Y = 0.240$, and $Z = 10^{-3}$, and as for model 1, all metals are initially added up in the form of ^{14}N . This model is indeed qualitatively similar to model zM, from Woosley et al. (2004; see also Fisker et al. 2008; Schatz et al. 2001). Both the envelope zoning and the initial relaxation phase are identical to those described for model 1.




Figure 16. Left panel: same as Figure 15, but for pressure. Right panel: time evolution of the total pressure at the outermost envelope shell. (A color version of this figure is available in the online journal.)




Figure 17. Left panels: mass fractions of the 10 most abundant, stable (or $\tau > 1$ hr) isotopes for model 1, at the end of the first (upper panel) and fourth bursts (lower panel), respectively. Right panels: same as left panels, but for overproduction factors relative to solar (for $f > 10^{-5}$).

(A color version of this figure is available in the online journal.)




Figure 18. Left panel: schematic representation of the convective regions in the accreted envelope for the four bursts computed in model 1. To simultaneously display the convective stages of the TNRs as compared with the overall duration of the bursts, a discontinuous time axis is used. Right panel: temperature profiles, showing the location of the ignition point (~ 5.6 m above the core-envelope interface), along the fourth bursting episode, for model 1 ($M_{\text{NS}} = 1.4 M_{\odot}$, $M_{\text{acc}} = 1.75 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$, $Z = 0.02$). Labels indicate different moments during the TNR, for which the temperature at the ignition shell reaches a value of (1) 3.4×10^8 K, (2) 4.5×10^8 K, (3) 7×10^8 K, (4) 9.2×10^8 K, (5) 9.7×10^8 K, (6) 1.04×10^9 K, (7) 1.06×10^9 K, (8) 8.9×10^8 K, (9) 7.4×10^8 K, and (10) 5.3×10^8 K. (A color version of this figure is available in the online journal.)




Figure 19. Mean post-burst composition in the envelope for each of the bursting episodes computed for model 1 (left) and model 2 (right). (A color version of this figure is available in the online journal.)

4.1. First Burst

The piling-up of matter on top of the neutron star during the accretion stage progressively compresses and heats the envelope.

At $t = 4337$ s, the envelope achieves $T_{\text{base}} \sim 10^8$ K ($\rho_{\text{base}} \sim 9.8 \times 10^4 \text{ g cm}^{-3}$). The nuclear activity is fully dominated by the CNO cycle. In contrast to model 1, the smaller metallicity of this model limits substantially the role of proton captures. Indeed, at this stage, H has only been reduced to 0.757 at the envelope base. The main reaction fluxes are actually an order of magnitude lower than those reported from model 1, for the same temperature, powering an energy generation rate of $\varepsilon_{\text{nuc}} \sim 6.2 \times 10^{12} \text{ erg g}^{-1} \text{ s}^{-1}$. Because of the lower metallicity

of this model, the time required to achieve $T_{\text{base}} \sim 10^8$ K is about twice the value reported for model 1, resulting in a thicker, more massive envelope which will affect the forthcoming explosion. Besides H and He (0.242), by far the most abundant nuclei in the envelope, the nuclear activity in the CNO region increases the chemical abundances of many species in this mass range, with ^{15}O (6.3×10^{-4}) being the most abundant CNO-group nucleus at the envelope base.

16.8 hr (60,347 s) after the beginning of accretion, T_{base} has reached 2×10^8 K (with $\rho_{\text{base}} = 5.7 \times 10^5 \text{ g cm}^{-3}$ and $P_{\text{base}} = 3.4 \times 10^{22} \text{ dyn cm}^{-2}$). The total luminosity of the star is only $6.9 \times 10^{34} \text{ erg s}^{-1}$. The main nuclear activity is governed by $^{15}\text{N}(p, \alpha)^{12}\text{C}$ and other reactions of the CNO cycle. $^7\text{Be}(p, \gamma)^8\text{B}$ is at equilibrium with its reverse photodisintegration reaction




Figure 20. Light curves corresponding to the second (upper left panel), third (upper right), and fourth bursts (lower left), and for the overall computed time (lower right), for model 1.

${}^8\text{B}(\gamma, p){}^7\text{Be}$. Because of the limited number of CNO catalysts in this low-metallicity model, some proton–proton chain reactions, such as the *pep*, ${}^3\text{He}(\alpha, \gamma){}^7\text{Be}$, or ${}^8\text{Be} \rightarrow 2{}^4\text{He}$, are relatively frequent. In terms of chemical abundances, the now frequent *p*-captures have reduced the hydrogen content down to a value of 0.695 (while $X({}^4\text{He}) = 0.303$). The next most abundant isotopes in the network are the short-lived species ${}^{15}\text{O}$ (1.4×10^{-3}) and ${}^{14}\text{O}$ (8×10^{-4}). The nuclear activity ($X_i > 10^{-9}$) reaches ${}^{40}\text{Ca}$ at this stage.

18.1 hr (65,081 s) from the onset of accretion, T_{base} reaches 4×10^8 K. Hydrogen continues to decrease smoothly ($X(\text{H}) = 0.689$), whereas the ${}^4\text{He}$ abundance reaches 0.280. The next most abundant nuclei are ${}^{14}\text{O}$ (1.4×10^{-2}) and ${}^{15}\text{O}$, now followed by ${}^{52}\text{Fe}$ and ${}^{18}\text{Ne}$, with mass fractions of the order of 10^{-3} . The nuclear activity extends up to ${}^{58}\text{Cu}$. The largest reaction fluxes correspond to different processes that operate almost at equilibrium with their inverse photodisintegration reactions, such as ${}^{21}\text{Mg}(p, \gamma){}^{22}\text{Al}(\gamma, p){}^{21}\text{Mg}$, ${}^{30}\text{S}(p, \gamma){}^{31}\text{Cl}(\gamma, p){}^{30}\text{S}$, ${}^{25}\text{Si}(p, \gamma){}^{26}\text{P}(\gamma, p){}^{25}\text{Si}$, and ${}^7\text{Be}(p, \gamma){}^8\text{B}(\gamma, p){}^7\text{Be}$.

Six seconds later ($t = 65,087$ s), T_{base} achieves 5×10^8 K. ρ_{base} has slightly decreased to 4.3×10^5 g cm $^{-3}$ because of a mild

envelope expansion ($\Delta z \sim 19.7$ m). ${}^4\text{He}$ has slightly decreased to 0.267, as a result of the frequent α -captures driven by the high temperatures achieved. The next most abundant species are now ${}^{18}\text{Ne}$ (10^{-2}), together with ${}^{14,15}\text{O}$ (3.5×10^{-3} and 7.5×10^{-3} , respectively), ${}^{52}\text{Fe}$ (2.8×10^{-3}), and ${}^{34}\text{Ar}$ (1.1×10^{-3}). The nuclear activity reaches ${}^{61}\text{Ga}$, powering an energy generation rate of 2.8×10^{16} erg g $^{-1}$ s $^{-1}$. The overall stellar luminosity is now 2.5×10^{35} erg s $^{-1}$.

A qualitatively similar picture is found when T_{base} achieves 7×10^8 K (at $t = 65,090$ s), with the nuclear activity extending all the way up to ${}^{68}\text{Se}$.

One second later ($t = 65,091$ s), T_{base} achieves 10^9 K. The nuclear activity (with $\epsilon_{\text{nuc}} \sim 1.5 \times 10^{17}$ erg g $^{-1}$ s $^{-1}$; see Figure 22) continues to reduce the H and ${}^4\text{He}$ abundances down to 0.680 and 0.224, respectively. The next most abundant species are now ${}^{28,29,30}\text{S}$, ${}^{33,34}\text{Ar}$, ${}^{25}\text{Si}$, ${}^{60}\text{Zn}$, and ${}^{38}\text{Ca}$ (all with mass fractions $\sim 10^{-2}$), with the main nuclear path reaching ${}^{72}\text{Kr}$. The largest absolute fluxes are achieved by nuclear interactions between equilibrium (*p*, γ)-(γ , *p*) pairs, which do not contribute to the net energy balance. Instead, the most important contributors to the energy production at this stage are




Figure 21. Upper panel: main nuclear activity at the ignition shell (~ 5.6 m above the core–envelope interface), when temperature reaches a peak value of $T_{\text{peak}} = 1.16 \times 10^9$ K, during the 4th burst computed for model 1. Lower panel: main reaction fluxes ($F \geq 10^{-9}$ reactions $\text{s}^{-1} \text{ cm}^{-3}$).

(A color version of this figure is available in the online journal.)

$^{25}\text{Al}(p, \gamma)^{26}\text{Si}$, $^{27,29,30}\text{P}(p, \gamma)^{28,30,31}\text{S}$, $^{28}\text{Si}(p, \gamma)^{29}\text{P}$, $^{32,33}\text{Cl}(p, \gamma)^{33,34}\text{Ar}$, $^{32,33}\text{Cl}(p, \gamma)^{33,34}\text{Ar}$, $^{35}\text{Ar}(p, \gamma)^{36}\text{K}$, $^{35,36,37}\text{K}(p, \gamma)^{36,37,38}\text{Ca}$, the chain $3\alpha \rightarrow ^{12}\text{C}(p, \gamma)^{13}\text{N}(p, \gamma)^{14}\text{O}(\alpha, p)^{17}\text{F}(p, \gamma)^{18}\text{Ne}(\alpha, p)^{21}\text{Na}(p, \gamma)^{22}\text{Mg}$, and a suite of β^+ -decays, such as $^{25}\text{Si}(\beta^+)^{25}\text{Al}$, $^{28,29,30}\text{S}(\beta^+)^{28,29,30}\text{P}$, and $^{33}\text{Ar}(\beta^+)^{33}\text{Cl}$. The activity in the $A = 65\text{--}100$ mass region is dominated by the suite of reactions depicted in Figure 22 (lower panel), mainly $^{65}\text{Ge}(p, \gamma)^{66}\text{As}$, $^{67}\text{Se}(\beta^+)^{67}\text{As}(p, \gamma)^{68}\text{Se}$, $^{65}\text{As}(p, \gamma)^{66}\text{Se}(\beta^+)^{66}\text{As}$ (with $\log F \sim -7$), and $^{65}\text{Se}(\beta^+)^{65}\text{As}$ ($\log F \sim -8$).

In contrast to model 1, which achieved a peak temperature of 1.06×10^9 K, model 3 reaches relatively higher values. Hence, at $t = 65,093$ s, T_{base} achieves 1.2×10^9 K. The H and ^4He abundances have been reduced to 0.648 and 0.205, respectively. The next most abundant nucleus is still ^{30}S (3.1×10^{-2}), followed by ^{38}Ca (2×10^{-2}), and by a large number of species with abundances of the order of 10^{-3} , such as $^{36,37}\text{Ca}$, $^{28,29}\text{S}$, $^{32,33,34}\text{Ar}$, $^{58,59,60}\text{Zn}$, $^{62,63,64}\text{Ge}$, $^{48,49,50}\text{Fe}$,

$^{53,54,55}\text{Ni}$, ^{41}Ti , and $^{44,45,46}\text{Cr}$. At this stage, the main nuclear path reaches ^{76}Sr .

One second later, at $t = 65,094$ s, the rate of nuclear energy generation achieves a maximum value of $\epsilon_{\text{nuc,max}} \sim 2.1 \times 10^{17} \text{ erg g}^{-1} \text{ s}^{-1}$.

At $t = 65,095$ s, while $T_{\text{base}} = 1.3 \times 10^9$ K, the main nuclear path reaches ^{80}Zr . Because of the large temperature achieved, the number of proton- and α -captures increases, which in turn efficiently reduces the H (0.621) and ^4He (0.197) abundances. The next most abundant nucleus is ^{60}Zn (2.8×10^{-2}), followed by ^{64}Ge , ^{38}Ca , ^{30}S , ^{55}Ni , and ^{59}Zn (with $X_i \sim 10^{-2}$, see Figure 23). The nuclear activity in the $A = 65\text{--}100$ mass region is now dominated by $^{65}\text{As}(p, \gamma)^{66}\text{Se}(\beta^+)^{66}\text{As}(p, \gamma)^{67}\text{Se}$ ($\log F \sim -4$), $^{65}\text{Ge}(p, \gamma)^{66}\text{As}$, and $^{67}\text{Se}(\beta^+)^{67}\text{As}(p, \gamma)^{68}\text{Se}$ ($\log F \sim -5$; see Figure 23). Energy production is mainly due to suite of (p, γ) reactions and β^+ -decays involving nuclear species in the mass range $A = 30\text{--}62$.

Table 2Mean Composition of the Envelope ($X_i > 10^{-9}$) at the End of Each Burst, for Model 1

Nucleus	Burst 1	Burst 2	Burst 3	Burst 4
¹ H	1.7×10^{-1}	8.8×10^{-2}	6.3×10^{-2}	3.9×10^{-2}
⁴ He	3.1×10^{-1}	2.0×10^{-1}	1.4×10^{-1}	1.0×10^{-1}
¹² C	1.7×10^{-2}	2.5×10^{-2}	2.0×10^{-2}	1.9×10^{-2}
¹³ C	8.4×10^{-5}	1.4×10^{-4}	1.3×10^{-4}	1.1×10^{-4}
¹⁴ N	2.1×10^{-3}	1.5×10^{-3}	1.1×10^{-3}	7.6×10^{-4}
¹⁵ N	2.6×10^{-3}	2.0×10^{-3}	1.4×10^{-3}	9.6×10^{-4}
¹⁶ O	5.7×10^{-4}	5.2×10^{-4}	3.3×10^{-4}	2.9×10^{-4}
¹⁷ O	3.7×10^{-6}	6.3×10^{-6}	6.8×10^{-6}	1.2×10^{-5}
¹⁸ O	6.7×10^{-5}	2.3×10^{-5}	1.7×10^{-5}	6.1×10^{-6}
¹⁸ F	6.2×10^{-5}	5.4×10^{-5}	3.1×10^{-5}	2.5×10^{-5}
¹⁹ F	2.1×10^{-4}	1.8×10^{-4}	1.1×10^{-4}	7.3×10^{-5}
²⁰ Ne	6.3×10^{-4}	5.4×10^{-4}	3.5×10^{-4}	4.0×10^{-4}
²¹ Ne	2.0×10^{-5}	9.5×10^{-6}	6.7×10^{-6}	6.2×10^{-6}
²² Ne	9.9×10^{-5}	4.2×10^{-5}	3.2×10^{-5}	2.9×10^{-5}
²² Na	3.3×10^{-3}	1.9×10^{-3}	1.2×10^{-3}	9.2×10^{-4}
²³ Na	3.7×10^{-4}	2.1×10^{-4}	1.2×10^{-4}	1.3×10^{-4}
²⁴ Mg	1.7×10^{-3}	1.4×10^{-3}	9.1×10^{-4}	1.3×10^{-3}
²⁵ Mg	2.6×10^{-3}	1.7×10^{-3}	1.2×10^{-3}	1.5×10^{-3}
²⁶ Mg	1.8×10^{-3}	2.5×10^{-3}	1.7×10^{-3}	1.1×10^{-3}
²⁶ Al ^g	2.6×10^{-4}	9.8×10^{-5}	5.7×10^{-5}	1.1×10^{-4}
²⁷ Al	1.8×10^{-3}	1.5×10^{-3}	8.8×10^{-4}	6.7×10^{-4}
²⁸ Si	1.2×10^{-3}	5.3×10^{-3}	7.0×10^{-3}	1.3×10^{-2}
²⁹ Si	3.1×10^{-4}	8.6×10^{-4}	4.6×10^{-4}	6.9×10^{-4}
³⁰ Si	3.5×10^{-3}	5.6×10^{-3}	3.8×10^{-3}	3.0×10^{-3}
³¹ P	5.1×10^{-4}	1.4×10^{-3}	1.7×10^{-3}	1.8×10^{-3}
³² S	3.8×10^{-4}	2.9×10^{-2}	5.8×10^{-2}	9.0×10^{-2}
³³ S	2.9×10^{-4}	3.1×10^{-3}	4.3×10^{-3}	5.1×10^{-3}
³⁴ S	3.2×10^{-3}	1.1×10^{-2}	1.6×10^{-2}	1.8×10^{-2}
³⁵ Cl	1.0×10^{-3}	4.8×10^{-3}	1.0×10^{-2}	1.0×10^{-2}
³⁶ Ar	4.2×10^{-4}	4.0×10^{-3}	9.2×10^{-3}	1.0×10^{-2}
³⁷ Cl	3.6×10^{-7}	6.4×10^{-7}	1.8×10^{-6}	4.1×10^{-6}
³⁷ Ar	2.3×10^{-4}	5.7×10^{-4}	1.0×10^{-3}	1.1×10^{-3}
³⁸ Ar	1.9×10^{-3}	4.9×10^{-3}	8.5×10^{-3}	8.9×10^{-3}
³⁹ K	3.8×10^{-3}	7.8×10^{-3}	1.2×10^{-2}	1.3×10^{-2}
⁴⁰ Ca	3.2×10^{-3}	4.3×10^{-3}	5.6×10^{-3}	5.3×10^{-3}
⁴¹ K	...	9.3×10^{-9}	4.8×10^{-8}	7.6×10^{-8}
⁴¹ Ca	6.5×10^{-5}	6.9×10^{-5}	6.0×10^{-5}	7.2×10^{-5}
⁴² Ca	5.5×10^{-4}	1.2×10^{-3}	1.7×10^{-3}	1.9×10^{-3}
⁴³ Ca	1.8×10^{-4}	3.6×10^{-4}	7.7×10^{-4}	1.2×10^{-3}
⁴³ Sc	4.3×10^{-4}	9.5×10^{-4}	1.2×10^{-3}	9.7×10^{-4}
⁴⁴ Ca	4.3×10^{-8}	3.7×10^{-6}	1.8×10^{-5}	3.2×10^{-5}
⁴⁴ Sc	2.5×10^{-7}	1.9×10^{-5}	3.6×10^{-5}	2.9×10^{-5}
⁴⁴ Ti	5.8×10^{-4}	7.2×10^{-4}	7.3×10^{-4}	7.5×10^{-4}
⁴⁵ Sc	8.2×10^{-5}	7.9×10^{-5}	9.3×10^{-5}	1.2×10^{-4}
⁴⁵ Ti	1.5×10^{-4}	1.7×10^{-4}	1.1×10^{-4}	1.0×10^{-4}
⁴⁶ Ti	9.5×10^{-4}	1.9×10^{-3}	2.4×10^{-3}	2.7×10^{-3}
⁴⁷ Ti	6.6×10^{-4}	8.7×10^{-4}	9.2×10^{-4}	9.5×10^{-4}
⁴⁸ Ti	2.1×10^{-7}	5.6×10^{-7}	1.6×10^{-6}	3.4×10^{-6}
⁴⁸ V	1.2×10^{-4}	1.4×10^{-4}	2.4×10^{-4}	3.7×10^{-4}
⁴⁸ Cr	1.8×10^{-3}	2.1×10^{-3}	2.0×10^{-3}	2.0×10^{-3}
⁴⁹ Ti	1.4×10^{-7}	4.4×10^{-7}	6.9×10^{-7}	8.9×10^{-7}
⁴⁹ V	1.5×10^{-3}	1.5×10^{-3}	1.2×10^{-3}	1.4×10^{-3}
⁵⁰ Cr	1.5×10^{-3}	2.3×10^{-3}	2.7×10^{-3}	3.0×10^{-3}
⁵¹ V	3.4×10^{-6}	6.4×10^{-6}	1.6×10^{-5}	3.0×10^{-5}
⁵¹ Cr	3.2×10^{-3}	4.5×10^{-3}	5.1×10^{-3}	5.8×10^{-3}
⁵² Cr	1.5×10^{-5}	2.2×10^{-5}	4.9×10^{-5}	9.4×10^{-5}
⁵² Mn	3.4×10^{-3}	2.3×10^{-3}	2.7×10^{-3}	4.0×10^{-3}
⁵² Fe	2.0×10^{-2}	1.5×10^{-2}	9.9×10^{-3}	1.2×10^{-2}
⁵³ Mn	1.2×10^{-3}	1.1×10^{-3}	8.6×10^{-4}	1.1×10^{-3}
⁵⁴ Fe	1.0×10^{-3}	1.4×10^{-3}	1.6×10^{-3}	1.8×10^{-3}
⁵⁵ Mn	6.6×10^{-9}	2.3×10^{-8}	7.1×10^{-8}	1.6×10^{-7}
⁵⁵ Fe	2.3×10^{-4}	3.1×10^{-4}	6.4×10^{-4}	1.0×10^{-3}
⁵⁵ Co	2.9×10^{-3}	3.8×10^{-3}	4.0×10^{-3}	4.3×10^{-3}
⁵⁶ Fe	8.0×10^{-8}	5.4×10^{-7}	1.7×10^{-6}	3.5×10^{-6}
⁵⁶ Co	2.2×10^{-4}	5.3×10^{-4}	1.1×10^{-3}	1.8×10^{-3}

Table 2

(Continued)

Nucleus	Burst 1	Burst 2	Burst 3	Burst 4
⁵⁶ Ni	2.4×10^{-2}	4.5×10^{-2}	5.2×10^{-2}	5.9×10^{-2}
⁵⁷ Fe	2.6×10^{-8}	1.6×10^{-7}	2.0×10^{-7}	2.5×10^{-7}
⁵⁷ Co	2.5×10^{-4}	1.2×10^{-4}	1.4×10^{-4}	2.0×10^{-4}
⁵⁷ Ni	6.5×10^{-3}	3.9×10^{-3}	2.7×10^{-3}	2.9×10^{-3}
⁵⁸ Ni	4.7×10^{-3}	2.7×10^{-3}	1.9×10^{-3}	2.2×10^{-3}
⁵⁹ Ni	7.3×10^{-3}	4.3×10^{-3}	3.3×10^{-3}	3.8×10^{-3}
⁶⁰ Ni	3.2×10^{-1}	3.3×10^{-1}	3.1×10^{-1}	3.1×10^{-1}
⁶¹ Ni	3.9×10^{-3}	1.7×10^{-3}	1.9×10^{-3}	2.4×10^{-3}
⁶¹ Cu	8.2×10^{-3}	6.0×10^{-3}	3.5×10^{-3}	3.8×10^{-3}
⁶² Ni	5.0×10^{-4}	3.1×10^{-4}	4.4×10^{-4}	5.8×10^{-4}
⁶² Zn	3.2×10^{-3}	2.5×10^{-3}	1.4×10^{-3}	1.5×10^{-3}
⁶³ Cu	4.1×10^{-3}	4.4×10^{-3}	3.1×10^{-3}	3.3×10^{-3}
⁶⁴ Zn	3.4×10^{-2}	1.0×10^{-1}	1.3×10^{-1}	1.3×10^{-1}
⁶⁵ Cu	2.5×10^{-7}	2.9×10^{-7}	5.1×10^{-7}	9.2×10^{-7}
⁶⁵ Zn	1.4×10^{-3}	2.5×10^{-3}	1.9×10^{-3}	2.2×10^{-3}
⁶⁶ Zn	3.2×10^{-5}	5.0×10^{-5}	1.3×10^{-4}	2.4×10^{-4}
⁶⁶ Ga	4.0×10^{-4}	5.1×10^{-4}	5.5×10^{-4}	6.3×10^{-4}
⁶⁶ Ge	5.4×10^{-4}	1.3×10^{-3}	7.5×10^{-4}	8.7×10^{-4}
⁶⁷ Ga	4.7×10^{-4}	1.0×10^{-3}	9.7×10^{-4}	1.1×10^{-3}
⁶⁸ Ge	2.9×10^{-3}	2.6×10^{-2}	4.4×10^{-2}	4.5×10^{-2}
⁶⁹ Ge	2.2×10^{-4}	1.3×10^{-3}	1.5×10^{-3}	1.8×10^{-3}
⁷⁰ Ge	7.3×10^{-5}	4.1×10^{-4}	4.6×10^{-4}	6.0×10^{-4}
⁷¹ As	7.7×10^{-5}	4.7×10^{-4}	6.0×10^{-4}	7.7×10^{-4}
⁷² Se	2.5×10^{-4}	5.7×10^{-3}	1.4×10^{-2}	1.6×10^{-2}
⁷³ Se	3.0×10^{-5}	4.2×10^{-4}	7.6×10^{-4}	1.0×10^{-3}
⁷⁴ Se	1.3×10^{-5}	1.8×10^{-4}	3.3×10^{-4}	4.8×10^{-4}
⁷⁵ Br	1.2×10^{-5}	1.8×10^{-4}	3.7×10^{-4}	5.3×10^{-4}
⁷⁶ Kr	3.0×10^{-5}	1.4×10^{-3}	4.5×10^{-3}	5.6×10^{-3}
⁷⁷ Kr	4.5×10^{-6}	1.5×10^{-4}	4.1×10^{-4}	6.1×10^{-4}
⁷⁸ Kr	2.7×10^{-6}	9.2×10^{-5}	2.5×10^{-4}	4.1×10^{-4}
⁷⁹ Kr	1.6×10^{-6}	5.9×10^{-5}	1.7×10^{-4}	2.7×10^{-4}
⁸⁰ Sr	3.9×10^{-6}	4.0×10^{-4}	1.6×10^{-3}	2.2×10^{-3}
⁸¹ Rb	8.4×10^{-7}	6.8×10^{-5}	2.7×10^{-4}	4.3×10^{-4}
⁸² Sr	7.9×10^{-7}	8.2×10^{-5}	3.2×10^{-4}	5.6×10^{-4}
⁸³ Sr	3.0×10^{-7}	4.3×10^{-5}	1.9×10^{-4}	3.2×10^{-4}
⁸⁴ Sr	1.9×10^{-7}	3.8×10^{-5}	1.4×10^{-4}	2.6×10^{-4}
⁸⁵ Y	1.1×10^{-7}	4.0×10^{-5}	2.1×10^{-4}	3.5×10^{-4}
⁸⁶ Zr	7.1×10^{-8}	4.4×10^{-5}	2.3×10^{-4}	4.6×10^{-4}
⁸⁷ Zr	2.7×10^{-8}	3.6×10^{-5}	1.8×10^{-4}	3.9×10^{-4}
⁸⁸ Zr	5.6×10^{-9}	1.8×10^{-5}	1.4×10^{-4}	2.6×10^{-4}
⁸⁹ Nb	4.0×10^{-9}	3.1×10^{-5}	2.9×10^{-4}	5.8×10^{-4}
⁹⁰ Mo	...	1.1×10^{-5}	1.1×10^{-4}	2.2×10^{-4}
⁹¹ Nb	...	5.4×10^{-6}	5.3×10^{-5}	1.1×10^{-4}
⁹² Mo	...	2.1×10^{-6}	1.9×10^{-5}	4.5×10^{-5}
⁹³ Tc	...	1.3×10^{-6}	1.3×10^{-5}	4.5×10^{-5}
⁹⁴ Tc	...	6.9×10^{-7}	7.6×10^{-6}	3.8×10^{-5}
⁹⁵ Ru	...	1.1×10^{-7}	1.1×10^{-6}	4.7×10^{-6}
⁹⁶ Ru	...	9.4×10^{-9}	1.1×10^{-7}	7.6×10^{-7}
⁹⁷ Ru	...	3.9×10^{-9}	5.1×10^{-8}	3.9×10^{-7}
⁹⁸ Ru	1.3×10^{-8}	9.2×10^{-8}
⁹⁹ Rh	2.4×10^{-9}	1.5×10^{-8}
¹⁰⁰ Pd	2.2×10^{-9}

Table 3

Summary of Burst Properties for Model 1

Burst	T_{peak} (K)	$t(T_{\text{peak}})$ (s)	τ_{rec} (hr)	L_{peak} (L_{\odot})	$\tau_{0.01}$ (s)	α
1	1.06×10^9	21192	5.9	9.7×10^4	75.8	60
2	1.15×10^9	44342	6.4	1.7×10^5	62.3	40
3	1.26×10^9	62137	4.9	2.1×10^5	55.4	34
4	1.12×10^9	80568	5.1	1.2×10^5	75.7	36

Table 4
Summary of Burst Properties for Model 2

Burst	T_{peak} (K)	$t(T_{\text{peak}})$ (s)	τ_{rec} (hr)	L_{peak} (L_{\odot})	$\tau_{0.01}$ (s)	α
1	1.05×10^9	21189	5.9	9.0×10^4	59.2	62
2	1.20×10^9	37783	4.6	1.5×10^5	73.9	31

Shortly after, at $t = 65,098$ s, a peak temperature of $T_{\text{peak}} = 1.4 \times 10^9$ K is achieved at the envelope base. This is followed, less than a second later, by a maximum expansion of the envelope, $\Delta z_{\text{max}} \sim 73.9$ m, and by a maximum luminosity, $L_{\text{max}} = 4.0 \times 10^{38}$ erg s $^{-1}$ ($10^5 L_{\odot}$). The main nuclear path reaches ^{93}Pd (already beyond the nucleosynthesis endpoint achieved in model 1). With respect to the chemical abundances, the envelope base is still dominated by H (0.560) and ^4He (0.175), with ^{60}Zn reaching a mass fraction of 0.111. The next most abundant species are ^{64}Ge (6.3×10^{-2}) and ^{68}Se (2×10^{-2}). Regarding the activity in the $A > 65$ mass region, at this stage is dominated by $^{66}\text{Se}(\beta^+)^{66}\text{As}(p, \gamma)^{67}\text{Se}(\beta^+)^{67}\text{As}(p, \gamma)^{68}\text{Se}$ ($\log F \sim -4$), $^{65}\text{Ge}(p, \gamma)^{66}\text{As}$, and $^{68}\text{Se}(\beta^+)^{68}\text{As}(p, \gamma)^{69}\text{Se}(p, \gamma)^{70}\text{Br}(p, \gamma)^{71}\text{Kr}(\beta^+)^{71}\text{Br}(p, \gamma)^{72}\text{Kr}$ ($\log F \sim -5$; see Figure 24, for additional processes down to $\log F \sim -8$). The most important contributors to the energy production are at this stage $^{44,45}\text{V}(p, \gamma)^{45,46}\text{Cr}$, $^{49}\text{Mn}(p, \gamma)^{50}\text{Fe}(\beta^+)^{50}\text{Mn}(p, \gamma)^{51}\text{Fe}$, $^{52}\text{Co}(p, \gamma)^{53}\text{Ni}$, $^{53}\text{Co}(p, \gamma)^{54}\text{Ni}(\beta^+)^{54}\text{Co}(p, \gamma)^{55}\text{Ni}(\beta^+)^{55}\text{Co}(p, \gamma)^{56}\text{Ni}$, and $^{58}\text{Zn}(\beta^+)^{58}\text{Cu}(p, \gamma)^{59}\text{Zn}(\beta^+)^{59}\text{Cu}(p, \gamma)^{60}\text{Zn}$ ($\log F \sim -3$).

At $t = 65,110$ s, following the decline from peak temperature, the envelope base achieves $T_{\text{base}} = 1.3 \times 10^9$ K (Figure 25). At this stage, the main nuclear activity has already reached the SnSbTe-mass region (^{104}Sn , in particular). The chemical abundances at the envelope base are still dominated by H (0.471), now followed by ^{64}Ge (0.162) and ^{68}Se (0.161), while ^4He has dropped to 0.131. The next most abundant species shift to ^{60}Zn (2.1×10^{-2}) and ^{72}Kr (1.9×10^{-2}), with a suite of nuclei reaching $\sim 10^{-3}$ (^{30}S , ^{67}Se , $^{37,38}\text{Ca}$, ^{76}Sr , $^{62,63}\text{Ge}$, ^{59}Zn , ^{55}Ni , ^{34}Ar , or ^{50}Fe). The nuclear activity in the $A > 65$ mass region is now powered by $^{65}\text{As}(p, \gamma)^{66}\text{Se}(\beta^+)^{66}\text{As}(p, \gamma)^{67}\text{Se}(\beta^+)^{67}\text{As}(p, \gamma)^{68}\text{Se}(\beta^+)^{68}\text{As}(p, \gamma)^{69}\text{Se}(p, \gamma)^{70}\text{Br}(p, \gamma)^{71}\text{Kr}(\beta^+)^{71}\text{Br}(p, \gamma)^{72}\text{Kr}$ ($\log F \sim -4$), $^{65}\text{Ge}(p, \gamma)^{66}\text{As}$, $^{69}\text{Br}(p, \gamma)^{70}\text{Kr}(\beta^+)^{70}\text{Br}$, $^{72}\text{Kr}(\beta^+)^{72}\text{Br}(p, \gamma)^{73}\text{Kr}(p, \gamma)^{74}\text{Rb}(p, \gamma)^{75}\text{Sr}(\beta^+)^{75}\text{Rb}(p, \gamma)^{76}\text{Sr}$, $^{78}\text{Y}(p, \gamma)^{79}\text{Zr}$, and $^{86}\text{Tc}(p, \gamma)^{87}\text{Ru}$ ($\log F \sim -5$; see Figure 25). Energy production is not due to a handful of nuclear processes but to dozens of different reactions (from $3\alpha \rightarrow ^{12}\text{C}$ all the way to $^{71}\text{Br}(p, \gamma)^{72}\text{Kr}$).

Twenty-two seconds later ($t = 65,132$ s), the temperature at the envelope base has decreased to $T_{\text{base}} = 1.2 \times 10^9$ K (Figure 26). The H content has been slightly reduced to 0.370, whereas ^4He reaches 8.74×10^{-2} . Indeed, after H, the most abundant species at the envelope base are now ^{68}Se (0.270) and ^{64}Ge (0.104), followed by ^{72}Kr (8.7×10^{-2}), ^{76}Sr (2.8×10^{-2}), and ^{80}Zr (1.2×10^{-2}). At this stage, some of the heaviest species of the network have already achieved an abundance of $\sim 10^{-3}$ (such as $^{88,89}\text{Ru}$, $^{92,93}\text{Pd}$, ^{96}Cd , ^{99}In , or $^{101,102}\text{Sn}$, together with the lighter nuclei ^{30}S , ^{38}Ca , $^{59,60}\text{Zn}$, and ^{67}Se), with the nuclear activity extending all the way up to ^{107}Te . The nuclear activity in the $A = 65$ –100 mass region is similar to that described above for $T_{\text{base}} = 1.3 \times 10^9$ K, and is depicted in Figure 26 (lower panel).

At $t = 65,264$ s, the envelope base achieves $T_{\text{base}} = 10^9$ K (Figure 27). Now, the most abundant element at the envelope base is ^{105}Sn (0.228). This is followed by a large number of species with abundances $\sim 10^{-2}$, such as ^{104}Sn , ^{68}Se , ^{72}Kr , ^{104}In ,

Table 5
Mean Composition of the Envelope ($X_i > 10^{-9}$) at the End of Each Burst, for Model 2

Nucleus	Burst 1	Burst 2
^1H	2.3×10^{-1}	1.0×10^{-1}
^4He	3.0×10^{-1}	1.9×10^{-1}
^{12}C	1.8×10^{-2}	2.2×10^{-2}
^{13}C	6.8×10^{-5}	6.2×10^{-5}
^{14}N	2.0×10^{-3}	1.4×10^{-3}
^{15}N	2.6×10^{-3}	2.0×10^{-3}
^{16}O	5.3×10^{-4}	3.9×10^{-4}
^{17}O	4.8×10^{-6}	3.0×10^{-6}
^{18}O	6.3×10^{-5}	2.8×10^{-5}
^{18}F	2.1×10^{-4}	5.7×10^{-5}
^{19}F	3.6×10^{-4}	1.7×10^{-4}
^{20}Ne	5.4×10^{-4}	4.1×10^{-4}
^{21}Ne	2.2×10^{-5}	7.3×10^{-6}
^{22}Ne	7.7×10^{-5}	4.4×10^{-5}
^{22}Na	3.6×10^{-3}	1.6×10^{-3}
^{23}Na	2.3×10^{-4}	1.7×10^{-4}
^{24}Mg	1.2×10^{-3}	1.3×10^{-3}
^{25}Mg	1.8×10^{-3}	1.4×10^{-3}
^{26}Mg	3.1×10^{-3}	1.6×10^{-3}
$^{26}\text{Al}^{\text{g}}$	1.0×10^{-4}	1.1×10^{-4}
^{27}Al	1.6×10^{-3}	1.1×10^{-3}
^{28}Si	1.2×10^{-3}	7.5×10^{-3}
^{29}Si	4.6×10^{-4}	6.7×10^{-4}
^{30}Si	4.7×10^{-3}	4.6×10^{-3}
^{31}P	6.1×10^{-4}	1.4×10^{-3}
^{32}S	4.7×10^{-4}	2.7×10^{-2}
^{33}S	4.0×10^{-4}	2.4×10^{-3}
^{34}S	4.3×10^{-3}	9.6×10^{-3}
^{35}Cl	1.1×10^{-3}	3.8×10^{-3}
^{36}Ar	5.8×10^{-4}	2.9×10^{-3}
^{37}Cl	1.6×10^{-7}	5.3×10^{-7}
^{37}Ar	2.9×10^{-4}	4.0×10^{-4}
^{38}Ar	2.6×10^{-3}	5.2×10^{-3}
^{39}K	4.9×10^{-3}	8.7×10^{-3}
^{40}Ca	3.1×10^{-3}	3.4×10^{-3}
^{41}K	...	8.7×10^{-9}
^{41}Ca	6.8×10^{-5}	4.5×10^{-5}
^{42}Ca	7.3×10^{-4}	1.2×10^{-3}
^{43}Ca	9.6×10^{-5}	4.3×10^{-4}
^{43}Sc	7.2×10^{-4}	1.2×10^{-3}
^{44}Ca	5.8×10^{-9}	2.3×10^{-6}
^{44}Sc	9.6×10^{-8}	1.1×10^{-5}
^{44}Ti	5.6×10^{-4}	4.8×10^{-4}
^{45}Sc	3.5×10^{-5}	4.9×10^{-5}
^{45}Ti	2.1×10^{-4}	1.2×10^{-4}
^{46}Ti	1.2×10^{-3}	1.9×10^{-3}
^{47}Ti	6.8×10^{-4}	6.3×10^{-4}
^{48}Ti	2.8×10^{-8}	2.5×10^{-7}
^{48}V	4.4×10^{-5}	9.4×10^{-5}
^{48}Cr	1.9×10^{-3}	1.7×10^{-3}
^{49}Ti	2.4×10^{-8}	2.9×10^{-7}
^{49}V	1.4×10^{-3}	1.2×10^{-3}
^{50}Cr	1.6×10^{-3}	2.2×10^{-3}
^{51}V	6.4×10^{-7}	5.5×10^{-6}
^{51}Cr	3.4×10^{-3}	4.9×10^{-3}
^{52}Cr	1.6×10^{-6}	8.1×10^{-6}
^{52}Mn	1.0×10^{-3}	1.5×10^{-3}
^{52}Fe	1.7×10^{-2}	1.3×10^{-2}
^{53}Mn	9.9×10^{-4}	8.4×10^{-4}
^{54}Fe	9.2×10^{-4}	1.2×10^{-3}
^{55}Mn	...	1.3×10^{-8}
^{55}Fe	7.9×10^{-5}	2.6×10^{-4}
^{55}Co	2.8×10^{-3}	3.7×10^{-3}
^{56}Fe	1.4×10^{-8}	3.6×10^{-7}
^{56}Co	1.1×10^{-4}	5.0×10^{-4}

Table 5
(Continued)

Nucleus	Burst 1	Burst 2
⁵⁶ Ni	3.2×10^{-2}	5.0×10^{-2}
⁵⁷ Fe	2.6×10^{-9}	6.6×10^{-8}
⁵⁷ Co	6.9×10^{-5}	7.4×10^{-5}
⁵⁷ Ni	5.1×10^{-3}	3.4×10^{-3}
⁵⁸ Ni	3.4×10^{-3}	2.3×10^{-3}
⁵⁹ Ni	4.8×10^{-3}	3.7×10^{-3}
⁶⁰ Ni	2.9×10^{-1}	3.2×10^{-1}
⁶¹ Ni	9.1×10^{-4}	1.4×10^{-3}
⁶¹ Cu	6.3×10^{-3}	5.8×10^{-3}
⁶² Ni	9.9×10^{-5}	1.9×10^{-4}
⁶² Zn	1.8×10^{-3}	2.1×10^{-3}
⁶³ Cu	2.2×10^{-3}	3.2×10^{-3}
⁶⁴ Zn	2.3×10^{-2}	1.2×10^{-1}
⁶⁵ Cu	2.4×10^{-8}	2.7×10^{-7}
⁶⁵ Zn	5.4×10^{-4}	3.1×10^{-3}
⁶⁶ Zn	1.7×10^{-6}	2.9×10^{-5}
⁶⁶ Ga	6.4×10^{-5}	6.2×10^{-4}
⁶⁶ Ge	2.8×10^{-4}	1.7×10^{-3}
⁶⁷ Ga	1.8×10^{-4}	1.2×10^{-3}
⁶⁸ Ge	1.6×10^{-3}	2.8×10^{-2}
⁶⁹ Ge	7.3×10^{-5}	1.8×10^{-3}
⁷⁰ Ge	2.0×10^{-5}	5.9×10^{-4}
⁷¹ As	2.4×10^{-5}	6.2×10^{-4}
⁷² Se	1.2×10^{-4}	5.8×10^{-3}
⁷³ Se	8.0×10^{-6}	6.1×10^{-4}
⁷⁴ Se	3.2×10^{-6}	2.7×10^{-4}
⁷⁵ Br	3.3×10^{-6}	2.7×10^{-4}
⁷⁶ Kr	1.2×10^{-5}	1.6×10^{-3}
⁷⁷ Kr	1.2×10^{-6}	2.3×10^{-4}
⁷⁸ Kr	6.7×10^{-7}	1.5×10^{-4}
⁷⁹ Kr	4.0×10^{-7}	9.2×10^{-5}
⁸⁰ Sr	1.5×10^{-6}	4.9×10^{-4}
⁸¹ Rb	2.3×10^{-7}	1.1×10^{-4}
⁸² Sr	2.2×10^{-7}	1.4×10^{-4}
⁸³ Sr	9.4×10^{-8}	7.0×10^{-5}
⁸⁴ Sr	6.4×10^{-8}	6.3×10^{-5}
⁸⁵ Y	4.5×10^{-8}	6.4×10^{-5}
⁸⁶ Zr	3.1×10^{-8}	7.8×10^{-5}
⁸⁷ Zr	1.4×10^{-8}	6.9×10^{-5}
⁸⁸ Zr	3.5×10^{-9}	3.3×10^{-5}
⁸⁹ Nb	2.5×10^{-9}	5.7×10^{-5}
⁹⁰ Mo	...	2.4×10^{-5}
⁹¹ Nb	...	1.3×10^{-5}
⁹² Mo	...	7.3×10^{-6}
⁹³ Tc	...	6.9×10^{-6}
⁹⁴ Tc	...	5.0×10^{-6}
⁹⁵ Ru	...	9.2×10^{-7}
⁹⁶ Ru	...	9.1×10^{-8}
⁹⁷ Ru	...	3.7×10^{-8}
⁹⁸ Ru	...	9.1×10^{-9}
⁹⁹ Rh	...	1.8×10^{-9}

⁹⁴Pd, ⁶⁴Ge, ¹⁰³In, ⁷⁶Sr—all more abundant than H (1.8×10^{-2}) and ⁴He (2.6×10^{-2}), at this stage—together with ^{102,103}Sn, ⁹⁵Ag, ¹⁰⁷Te, ^{100,101,102}In, ⁸⁰Zr, ⁶⁰Zn, and ^{98,99}Cd. Note that, since the heaviest element included in our network, ¹⁰⁷Te, achieved already an abundance of 2.2×10^{-2} , leakage from the SnSbTe-mass region cannot be discarded. Note, however, that α -emission for ¹⁰⁷Te was not included, which may account for the high abundances reported here. Further studies to explore possible nucleosynthesis beyond the SnSbTe-mass region are underway with a larger network, so detailed abundances in this region should be taken with caution. The set of equilibrium

(p, γ)-(γ, p) pairs is now accompanied by a handful of β^+ -decays (such as ⁸⁰Zr(β^+)⁸⁰Y, ⁷⁶Sr(β^+)⁷⁶Rb, ⁸⁴Mo(β^+)⁸⁴Nb, and ⁸²Nb(β^+)⁸²Zr) as the nuclear processes with largest absolute fluxes, since H depletion and the low temperature limit the extent of charged-particle reactions. Indeed, these weak interactions will become progressively more important during the last stages of the burst. At this stage, the activity in the $A = 65$ –100 mass region is dominated by ⁷⁶Sr(β^+)⁷⁶Rb(p, γ)⁷⁷Sr(p, γ)⁷⁸Y, ⁷⁹Y(p, γ)⁸⁰Zr(β^+)⁸⁰Y(p, γ)⁸¹Zr(p, γ)⁸²Nb(β^+)⁸²Zr(p, γ)⁸³Nb(p, γ)⁸⁴Mo(β^+)⁸⁴Nb(p, γ)⁸⁵Mo, ⁸⁹Ru(p, γ)⁹⁰Rh (log $F \sim -4$), and more than 60 different nuclear processes with log $F \sim -5$ (see Figure 27, lower panel), involving nuclei in the mass range $A = 65$ –104. Indeed, energy production is driven by (p, γ) and β^+ processes involving species in this mass range.

At $t = 65,362$ s, the temperature at the envelope base has already declined to $T_{\text{base}} = 7.6 \times 10^8$ K (Figure 28). H is now fully depleted (7.6×10^{-12}), while ⁴He barely reaches 1.8×10^{-2} . As before, the most abundant element at the envelope base is ¹⁰⁵Sn (0.251), followed by ¹⁰⁴In (0.142), and by a large number of species with abundances $\sim 10^{-2}$. The depletion of H dramatically reduces the fluxes of most of the (p, γ) reactions, which are now overcome by many β^+ -decays (such as those affecting ^{68,69}Se, ⁶⁸As, ⁶⁴Ge, ^{71,72}Br, ⁶⁰Zn, ⁶⁰Cu, ^{72,73}Kr, ¹⁰⁴Sn, ⁶⁴Ga, ⁸²Zr, or ⁷⁶Sr), and by a suite of α -capture reactions, such as $3\alpha \rightarrow ^{12}\text{C}(\alpha, \gamma)^{16}\text{O}(\alpha, \gamma)^{20}\text{Ne}(\alpha, \gamma)^{24}\text{Mg}(\alpha, \gamma)^{28}\text{Si}(\alpha, \gamma)^{32}\text{S}$, or $^{13}\text{N}(\alpha, p)^{16}\text{O}$.

At $t = 69,715$ s, after a long decline, a minimum temperature is achieved at the envelope base, $T_{\text{base}} = 2 \times 10^8$ K (Figure 29), which we consider to mark the end of the first bursting episode for this model. At this stage, H is fully depleted (3.2×10^{-23}) at the envelope base, while ⁴He has a mass fraction of 1.5×10^{-2} only. The distribution of the most abundant elements almost follows the one described for $T_{\text{base}} = 7.6 \times 10^8$ K (Figure 28), and is dominated by ¹⁰⁵Sn (0.251), followed by ¹⁰⁴In (0.147), and by a large number of species with abundances $\sim 10^{-2}$, such as ⁹⁴Pd, ^{100,101,102,103}In, ⁶⁸Ge, ⁶⁴Zn, ⁷²Se, ⁷⁶Rb, ¹⁰⁷Te, ⁶⁰Ni, ⁹⁹Cd, ^{97,98}Ag, ^{89,90}Ru, and ⁸⁰Y. At this stage, the dominant interactions are all β^+ -decays (^{60,61}Cu(β^+)^{60,61}Ni, ^{66,67}Ge(β^+)^{66,67}Ga, ⁶⁵Ga(β^+)⁶⁵Zn, ⁵¹Mn(β^+)⁵¹Cr, ⁵²Fe(β^+)⁵²Mn, ⁶³Zn(β^+)⁶³Cu, ⁵⁶Ni(β^+)⁵⁶Co, or ⁴³Sc(β^+)⁴³Ca), except for the triple- α reaction.

Depth also influences the extent of the nuclear activity throughout the envelope, but in contrast to model 1, the nuclear activity in all shells of our computational domain essentially reaches the SnSbTe-mass region. Indeed, the inner part of the envelope (encompassing 3.4×10^{21} g) is, at the end of the burst, dominated by large amounts of ¹⁰⁵Sn and ¹⁰⁴In, the most abundant nuclei with mass fractions ~ 0.1 –0.2; at 5.6×10^{21} g above the core-envelope interface, the most abundant isotopes are H (0.26) and ⁴He (0.12), while the most abundant species in the SnSbTe-mass region achieve a mass fraction $\sim 10^{-3}$; and close to the surface (7.7×10^{21} g), shells are largely dominated by the presence of unburned H (0.75) and ⁴He (0.24), with $X(\text{SnSbTe}) \sim 10^{-7}$.

The mean, mass-averaged chemical composition of the whole envelope, at the end of the first bursting episode, is dominated by the presence of unburned H (0.18) and ⁴He (0.084), followed by ¹⁰⁵Ag (0.075), ¹⁰⁴Pd (0.053), ⁶⁴Zn (0.042), ⁹⁵Ru (0.031), ⁶⁸Ge (0.028), ⁹⁴Tc (0.026), and ¹⁰³Ag (0.026), with a nucleosynthesis endpoint around ¹⁰⁷Cd. In contrast, the first burst computed in model 1 yielded, in general, lighter nuclei, ⁶⁰Ni, ⁴He, ¹H, ⁶⁴Zn, ¹²C, and ^{52,56}Fe, with a more modest nucleosynthesis endpoint around ⁸⁹Nb (Table 2).




Figure 22. Main nuclear activity at the innermost envelope shell for model 3 ($M_{\text{NS}} = 1.4 M_{\odot}$, $\dot{M}_{\text{acc}} = 1.75 \times 10^{-9} M_{\odot} \text{ yr}^{-1}$, $Z = 10^{-3}$), at the time when temperature at the envelope base reaches $T_{\text{base}} = 10^9$ K. Upper panel: mass fractions of the most abundant species ($X > 10^{-3}$). Lower panel: main reaction fluxes ($F \geq 10^{-11}$ reactions $\text{s}^{-1} \text{ cm}^{-3}$) responsible for the nuclear activity in the $A = 65$ – 100 mass region—except for equilibrium (p, γ) – (γ, p) pairs.

(A color version of this figure is available in the online journal.)

In terms of overproduction factors, f (Figure 30), while model 1 showed moderate values ($f \sim 10^4$) for a handful of intermediate-mass elements, such as ^{43}Ca , ^{45}Sc , ^{49}Ti , ^{51}V , $^{60,61}\text{Ni}$, $^{63,65}\text{Cu}$, $^{64,67,68}\text{Zn}$, ^{69}Ga , ^{74}Se , or ^{78}Kr , model 3 achieves moderate overproduction factors ($\geq 10^4$), for all stable species heavier than ^{64}Zn , and as high as $\sim 10^8$ for ^{98}Ru , $^{102,104}\text{Pd}$, and ^{106}Cd .

4.2. Second, Third, Fourth, and Fifth Bursts

Tables 6–8 summarize the most relevant properties that characterize the bursting episodes computed for model 3. Recurrence times between bursts of $\tau_{\text{rec}} \sim 9$ hr (except for the first one, for which $\tau_{\text{rec}} \sim 18$ hr), ratios between persistent and burst luminosities of $\alpha \sim 20$ – 30 , and peak luminosities around $L_{\text{peak}} \sim 10^5 L_{\odot}$ represent the basic observables associated

with this model. Indeed, the recurrence times obtained are in agreement with the values reported for the XRB sources (see Galloway et al. 2008) 1A 1905+00 [$\tau_{\text{rec}} = 8.9$ hr], 4U 1254-69 [$\tau_{\text{rec}} = 9.2$ hr], or XTE J1710-281 [$\tau_{\text{rec}} = 8.9$ hr, $\alpha = 22$ – 100]. A striking result is the quick stabilization of the recurrence times, that show a regular periodicity after the second burst.

It is worth noting that both the recurrence periods and the ratios between persistent and burst luminosities are larger than those reported for model 1 (see Table 6, for comparison), showing a clear dependence on the metallicity of the accreted material: the smaller the metal content, the larger the recurrence time (and the smaller the value of α).

The corresponding light curves (see Figure 31) exhibit, in turn, a clear pattern: as shown in Figure 32 (left panel), where light curves of the third bursting episode computed in models 1 and 3 are compared, explosions in metal-deficient envelopes




Figure 23. Same as Figure 22, but for $T_{\text{base}} = 1.3 \times 10^9$ K.
(A color version of this figure is available in the online journal.)

Table 6
Properties of the Last Burst Computed in Models 1 and 3

Property	Model 1	Model 3
$\rho_{\text{max,base}}$ ($g cm^{-3}$)	1.3×10^6	2.6×10^6
$P_{\text{max,base}}$ ($dyn cm^{-2}$)	4.2×10^{22}	1.1×10^{23}
$\rho_{\text{max,ign}}$ ($g cm^{-3}$)	5.4×10^5	1.0×10^6
$P_{\text{max,ign}}$ ($dyn cm^{-2}$)	1.3×10^{22}	2.9×10^{22}
τ_{acc} (hr)	5.1	8.8
Δm_{acc} (M_{\odot})	1.0×10^{-12}	1.8×10^{-12}
T_{peak} (K)	1.1×10^9	1.3×10^9
L_{peak} (L_{\odot})	1.2×10^5	1.0×10^5
Δz_{max} (m)	40	44
α	36	30

Note. $\rho_{\text{max,base}}$ and $P_{\text{max,base}}$ are the maximum density and pressure achieved at the base of the envelope, whereas $\rho_{\text{max,ign}}$ and $P_{\text{max,ign}}$ correspond to the maximum values attained at the ignition shell (defined as the first shell that reaches $T > 4 \times 10^8$ K, for this burst).

Table 7
Summary of Burst Properties for Model 3

Burst	T_{peak} (K)	$t(T_{\text{peak}})$ (s)	τ_{rec} (hr)	L_{peak} (L_{\odot})	$\tau_{0.01}$ (s)	α
1	1.40×10^9	65110	18.1	1.0×10^5	423	34
2	1.39×10^9	98879	9.4	1.1×10^5	296	24
3	1.32×10^9	130816	8.9	9.8×10^4	281	24
4	1.30×10^9	162777	8.9	1.0×10^5	252	27
5	1.26×10^9	194266	8.8	1.0×10^5	250	30

(such as model 3) are characterized by lower peak luminosities and longer decline times. A similar pattern has been reported by Heger et al. (2007), in the framework of one-dimensional, hydrodynamic models of XRBs performed with the KEPLER code. It is worth noting that no double-peaked bursts have been obtained in model 3.

Larger peak temperatures, around $T_{\text{peak}} \sim (1.3\text{--}1.4) \times 10^9$ K, have also been obtained in model 3. This, together with the




Figure 24. Same as Figure 22, but for the time when temperature at the envelope base reaches a peak value of $T_{\text{peak}} = 1.4 \times 10^9$ K. (A color version of this figure is available in the online journal.)

longer exposure times to high temperatures (driven by the slower decline phase) cause a dramatic extension of the main nuclear path toward the SnSbTe-mass region or beyond.

From the nucleosynthesis viewpoint, and as shown in Table 8 and Figures 31 and 32, the nuclear activity already reaches the end of the network (^{107}Te) at the late stages of the first bursting episode. The overall mean metallicity of the envelope at the end of each burst is now 0.74 (1st burst), 0.85 (2nd burst), 0.90 (3rd burst), 0.92 (4th burst), and 0.93 (5th burst). Note that, although the accreted material is more metal deficient in model 3 than in model 1, the post-burst mean metallicity of the envelope is larger in model 3. This results from the combination of higher temperatures and longer burst durations, which favors the extension of the nuclear activity: for short bursts (like those obtained in metal-rich envelopes), only the fastest p - and/or α -capture reactions can naturally occur (those that proceed with a

characteristic time shorter than the overall exposure time to high temperatures); in contrast, for long-duration bursts, the overall number of p - and/or α -capture reactions increases dramatically. This, in particular, affects CNO breakout through $^{15}\text{O}(\alpha, \gamma)^{19}\text{Ne}$ and $^{14}\text{O}(\alpha, p)^{17}\text{F}$, which are favored in the longer bursts obtained for model 3. It is also worth noting that the abundance pattern obtained after the different bursts is very similar. This is clearly shown in Figure 33, that depicts the main nuclear activity at peak and at the end of the fourth burst computed for model 3, when compared with that corresponding to the first burst—Figures 24 and 29. This fact justifies our emphasis on the reaction sequences that characterize the first burst (see discussion in Section 5.2).

A final ^{12}C yield of $\sim 2 \times 10^{-3}$ is obtained at the end of each bursting episode (except in the first one, for which $X(^{12}\text{C}) = 8 \times 10^{-4}$). As reported for model 1, the amount of unburned ^{12}C left over turns out to be too small to power a




Figure 25. Same as Figure 22, but for $T_{\text{base}} = 1.3 \times 10^9 \text{ K}$.
(A color version of this figure is available in the online journal.)

superburst. Finally, huge overproduction factors (see Figures 31 and 32), involving heavy species such as $^{102,104,105}\text{Pd}$, ^{98}Ru , or ^{94}Mo (with $f \sim 10^8$), have been obtained in model 3, in contrast with the somewhat more modest values achieved in model 1, where maximum overproduction factors are about $f \sim 10^6$, and involving lighter species, such as ^{76}Se , $^{78,80}\text{Kr}$, or ^{84}Sr .

5. DISCUSSION

5.1. General Relativity Corrections

The calculations reported here have been performed assuming Newtonian gravity. Since the envelope layers are very thin, it is easy to introduce general relativity corrections to this Newtonian framework (see Ayasli & Joss 1982; Lewin et al. 1993; Taam

et al. 1993; Cumming et al. 2002; Woosley et al. 2004). To this end, the surface gravity is rewritten as $g = GM_*/R_*^2(1+z)$, where M_* is the mass, R_* is the stellar radius (defined in such a way that the surface area is $4\pi R_*^2$), and z is the gravitational redshift given by $1+z = (1 - 2GM_*/R_*c^2)^{-1/2}$. Our models of $M_* = 1.4 M_\odot$ require $R_* = 14.3 \text{ km}$, and a gravitational redshift of $z = 0.19$.

Following Woosley et al. (2004), once the redshift and radius are determined, it is straightforward to derive the set of correcting factors to the physical magnitudes described above for a suitable observer at infinity. Hence, recurrence times and burst durations should be increased by a factor of $1+z$. The mass-accretion rate as well as the burst luminosity have to take into account both the difference in surface area (compared to the Newtonian framework) and the gravitational redshift term. The energy and rest mass-accretion rate scale as $R_*^2/(1+z)$,




Figure 26. Same as Figure 22, but for $T_{\text{base}} = 1.2 \times 10^9 \text{ K}$.
(A color version of this figure is available in the online journal.)

while the luminosity $\propto R_*^2/(1+z)^2$. However, when M_* is taken exactly as M_{NS} (Newtonian framework), the surface area and redshift corrections for energy and mass-accretion rate cancel out, since $g \propto (1+z)/R_*^2 = \text{const}$, and hence, no correction to the observed burst energy or mass-accretion rate is necessary, while the luminosity correction is simply given by $1/(1+z) = 0.84$. In addition, the accretion luminosity for an observer at infinity changes only by a factor of 1.012; that is, the ratio between gravitational energy released per unit mass in general relativity, $c^2 z/(1+z)$, and the Newtonian value, $GM_{\text{NS}}/R_{\text{NS}}$. Finally, the luminosity measured at infinity will be smaller by a factor of $(1+z) = 1.19$.

5.2. Comparison with Previous Work

For consistency, the results discussed in this paper have been compared with those reported in previous work (obtained with similar hydrodynamic codes or in the framework of one-zone models).

As emphasized in Section 2, model 1 is qualitatively similar to model ZM of Woosley et al. (2004). The 12 bursts computed by Woosley et al. (2004) in a Newtonian frame were characterized by recurrence times of about $\sim 2.7 \text{ hr}$, peak luminosities of $L_{\text{peak}} \sim (1.5-2) \times 10^{38} \text{ erg s}^{-1}$, and ratios between persistent and burst luminosities of $\alpha \sim 60-65$. Our calculations (model




Figure 27. Same as Figure 22, but for $T_{\text{base}} = 10^9$ K.
(A color version of this figure is available in the online journal.)

1, Newtonian frame) yield $\tau_{\text{rec}} \sim 5\text{--}6.5$ hr, $L_{\text{peak}} \sim (3\text{--}7) \times 10^{38}$ erg s $^{-1}$, and $\alpha \sim 35\text{--}40$.

The role played by the metallicity of the accreted material (model 3, with $Z = Z_{\odot}/20 = 0.001$) qualitatively agrees with the pattern reported by Woosley et al. (2004; see also, Heger et al. 2007). Longer recurrence times of ~ 9 hr, peak temperatures of about $(1.3\text{--}1.4) \times 10^9$ K, and ratios between persistent and burst luminosities of $\alpha \sim 20\text{--}30$ (with $L_{\text{peak}} \sim 10^{38}$ erg s $^{-1}$) have been obtained in the five bursts computed in model 3. In turn, the 15 bursts computed by Woosley et al. (2004)

for model zM are characterized by recurrence times of about 3–3.5 hr, peak luminosities of $L_{\text{peak}} \sim 10^{38}$ erg s $^{-1}$, and ratios between persistent and burst luminosities of $\alpha \sim 50\text{--}60$. Results reveal a dependence of burst properties on the metallicity of the accreted material: the smaller the metal content, the larger the recurrence time (and the smaller the α). In turn, explosions in metal-deficient envelopes (i.e., model 3) are characterized by lower peak luminosities and longer decline times, in agreement with the pattern described in Woosley et al. (2004) and Heger et al. (2007). Model 3 bears as well a clear resemblance with




Figure 28. Same as Figure 22, but for $T_{\text{base}} = 7.6 \times 10^8$ K.

(A color version of this figure is available in the online journal.)

the model computed by Fisker et al. (2008). In that work, five representative bursting sequences were analyzed, with $\tau_{\text{rec}} \sim 3.5\text{--}4$ hr, $L_{\text{peak}} \sim (7\text{--}8) \times 10^{37}$ erg s $^{-1}$, and $\alpha \sim 65\text{--}70$, as measured at infinity.

Despite the qualitative similarities in the gross properties of the bursts presented in this paper (as well as in the role played by the metallicity of the accreted material) and those reported in previous work, a quantitative comparison reveals some discrepancies that are worth analyzing. In model 1 (with $Z = Z_{\odot}$), our computations yield systematically larger (by a factor of ~ 2) recurrence times and peak luminosities (and hence, lower α) than model ZM of Woosley et al. (2004). Similar results are found in the low-metallicity case (model 3, with $Z = Z_{\odot}/20$) when compared with model zM of Woosley et al. (2004), except for the peak luminosities that turn out to be very similar. It is also worth noting that the values reported by Fisker et al. (2008) show discrepancies with respect to

Woosley et al. (2004), in particular, lower peak luminosities (and larger α). A major difference concerns the much larger effect played by the metallicity of the accreted material in this work as compared with Woosley et al. (2004), who explained the moderate effect found as due to compositional inertia washing out the influence of the initial metallicity. Another striking issue concerns the extremely large differences in the gross physical characteristics—nucleosynthesis, energies, or recurrence times—between the first and subsequent bursts, as reported by Woosley et al. (2004). In terms of nucleosynthesis or nuclear activity, Figures 30–32 reveal a similar behavior for the different bursts (although a somewhat lower production of intermediate-mass elements as well as of the heaviest elements is reported for the first burst computed in model 3).

Very limited information on the nucleosynthetic yields obtained in model ZM is given in Woosley et al. (2004). Thus, we will restrict the discussion on the extent of the nuclear activity




Figure 29. Same as Figure 22, but for the time when temperature at the envelope base achieves a minimum value of $T_{\min} = 2 \times 10^8 \text{ K}$. (A color version of this figure is available in the online journal.)

and on the resulting chemical abundance pattern to model 3, through a brief comparison with the work reported by Schatz et al. (2001), Fisker et al. (2008), and Woosley et al. (2004; for model zM). It is worth noting that both the nucleosynthetic endpoint (located in the SnSbTe-mass region) and the main nuclear path in the $A \sim 50$ – 100 mass region obtained in this work (passing through a suite of different nuclei, such as ^{55}Co , ^{60}Zn , ^{70}Br , ^{75}Rb , ^{85}Mo , ^{90}Rh , or $^{100,105}\text{Sn}$) are similar to those reported by Schatz et al. (2001) in the framework of one-zone calculations. Whereas the main nuclear path in the first burst of model zM (Woosley et al. 2004) is very similar to the one reported in this work, compositional inertia causes a more limited extension of the nuclear activity in the successive bursts of Woosley et al.: hence, while the three most abundant nuclei at the bottom of the envelope are ^{106}Sn and $^{104,106}\text{In}$ at the end of the first burst, this switches to ^{64}Zn , ^{68}Se , and ^{32}S (a similar trend is also reported by Fisker et al. 2008). In this work, the mass-averaged composition at the end of the first burst computed for model 3 (see Table 8) is dominated (aside from some residual H and ^4He) by the presence of ^{104}Pd (0.05, by mass) and ^{105}Ag (0.08), while $X(^{64}\text{Zn}) \sim 0.04$. But the peak at the end of the abundance distribution (see Figure 30) increases with subsequent bursts up to a plateau value, which indicates that these heavy nuclei are still

produced in similar quantities. This is very different to the results reported by Woosley et al. (2004). Indeed, at the end of the fifth burst, the abundance pattern shows still a significant presence of heavy species (i.e., $X(^{105}\text{Ag}) \sim 0.1$, $X(^{104}\text{Pd}) \sim 0.08$, and $X(^{94}\text{Tc}) \sim 0.05$), together with a simultaneous increase in the abundances of intermediate-mass elements, such as ^{60}Ni (0.06), ^{64}Zn (0.09), ^{68}Ge (0.07), or ^{72}Se (0.04). It is finally worth noting that, in agreement with all previous hydrodynamic studies, both models 1 and 3 yield very small post-burst abundances of ^{12}C , below the threshold amount required to power superbursts. Even though only a few bursts have been computed for these models, they already show a trend on the amount of ^{12}C that may be expected after many more bursts.

Finally, it is also worth mentioning that large differences exist between the hydrodynamic simulations reported here (see also Woosley et al. 2004; Fisker et al. 2008) and those based on one-zone models (i.e., Schatz et al. 1999, 2001) as regards the shape of the light curve accompanying the bursting episodes (the primary difference being the presence of a long-lasting plateau in the latter).

The origin of the discrepancies reported is not totally clear and would require additional hydrodynamic studies. Note, however, that the local surface gravity of our model is somewhat smaller

Table 8Mean Composition of the Envelope ($X_i > 10^{-9}$) at the End of Each Burst, for Model 3

Nucleus	Burst 1	Burst 2	Burst 3	Burst 4	Burst 5
¹ H	1.8×10^{-1}	8.6×10^{-2}	5.5×10^{-2}	4.5×10^{-2}	3.7×10^{-2}
⁴ He	8.4×10^{-2}	5.8×10^{-2}	4.4×10^{-2}	3.8×10^{-2}	3.3×10^{-2}
¹² C	7.7×10^{-4}	1.7×10^{-3}	1.8×10^{-3}	1.7×10^{-3}	1.5×10^{-3}
¹³ C	4.2×10^{-6}	8.9×10^{-6}	1.4×10^{-5}	2.3×10^{-5}	2.7×10^{-5}
¹⁴ N	2.8×10^{-4}	2.1×10^{-4}	1.7×10^{-4}	1.5×10^{-4}	1.4×10^{-4}
¹⁵ N	4.9×10^{-4}	3.6×10^{-4}	2.9×10^{-4}	2.4×10^{-4}	2.2×10^{-4}
¹⁶ O	1.4×10^{-5}	2.9×10^{-5}	4.1×10^{-5}	3.7×10^{-5}	3.8×10^{-5}
¹⁷ O	4.8×10^{-7}	6.8×10^{-7}	2.1×10^{-6}	1.3×10^{-6}	2.0×10^{-6}
¹⁸ O	...	3.7×10^{-9}	9.4×10^{-9}	9.5×10^{-9}	8.1×10^{-9}
¹⁸ F	6.4×10^{-9}	1.3×10^{-8}	2.8×10^{-8}	2.4×10^{-8}	2.3×10^{-8}
¹⁹ F	8.7×10^{-8}	1.6×10^{-7}	1.7×10^{-7}	2.0×10^{-7}	8.0×10^{-8}
²⁰ Ne	1.5×10^{-5}	2.9×10^{-5}	3.5×10^{-5}	3.0×10^{-5}	3.1×10^{-5}
²¹ Ne	1.9×10^{-8}	4.4×10^{-8}	4.9×10^{-8}	5.2×10^{-8}	4.4×10^{-8}
²² Ne	2.9×10^{-7}	6.6×10^{-7}	4.3×10^{-7}	1.1×10^{-6}	7.5×10^{-7}
²² Na	1.1×10^{-5}	2.4×10^{-5}	2.0×10^{-5}	2.3×10^{-5}	1.9×10^{-5}
²³ Na	2.6×10^{-6}	6.5×10^{-6}	6.5×10^{-6}	6.5×10^{-6}	5.7×10^{-6}
²⁴ Mg	8.2×10^{-5}	1.4×10^{-4}	1.4×10^{-4}	1.2×10^{-4}	1.1×10^{-4}
²⁵ Mg	4.9×10^{-5}	7.3×10^{-5}	7.4×10^{-5}	7.2×10^{-5}	6.6×10^{-5}
²⁶ Mg	1.1×10^{-5}	1.9×10^{-5}	1.8×10^{-5}	2.2×10^{-5}	2.0×10^{-5}
²⁶ Al ^g	1.2×10^{-5}	1.4×10^{-5}	1.3×10^{-5}	1.2×10^{-5}	1.1×10^{-5}
²⁷ Al	4.1×10^{-5}	6.3×10^{-5}	6.5×10^{-5}	6.8×10^{-5}	6.2×10^{-5}
²⁸ Si	2.6×10^{-4}	6.2×10^{-4}	5.2×10^{-4}	3.9×10^{-4}	3.6×10^{-4}
²⁹ Si	8.6×10^{-6}	1.1×10^{-5}	9.2×10^{-6}	1.3×10^{-5}	1.1×10^{-5}
³⁰ Si	5.5×10^{-5}	8.8×10^{-5}	8.0×10^{-5}	1.1×10^{-4}	9.9×10^{-5}
³¹ P	2.3×10^{-5}	3.6×10^{-5}	2.6×10^{-5}	3.2×10^{-5}	2.6×10^{-5}
³² S	3.7×10^{-5}	2.1×10^{-3}	4.9×10^{-3}	5.8×10^{-3}	7.7×10^{-3}
³³ S	5.7×10^{-6}	4.2×10^{-5}	6.1×10^{-5}	5.6×10^{-5}	9.2×10^{-5}
³⁴ S	2.6×10^{-5}	6.7×10^{-5}	7.2×10^{-5}	7.3×10^{-5}	7.5×10^{-5}
³⁵ Cl	2.9×10^{-5}	1.5×10^{-4}	4.2×10^{-4}	5.0×10^{-4}	5.9×10^{-4}
³⁶ Ar	8.2×10^{-6}	2.9×10^{-3}	6.8×10^{-3}	9.9×10^{-3}	1.2×10^{-2}
³⁷ Cl	3.5×10^{-9}	2.8×10^{-8}	5.4×10^{-8}	1.5×10^{-7}	2.2×10^{-7}
³⁷ Ar	3.7×10^{-6}	2.6×10^{-5}	5.1×10^{-5}	1.1×10^{-4}	1.6×10^{-4}
³⁸ Ar	1.5×10^{-5}	6.4×10^{-5}	1.1×10^{-4}	1.7×10^{-4}	2.0×10^{-4}
³⁹ K	4.7×10^{-5}	9.4×10^{-4}	1.4×10^{-3}	2.1×10^{-3}	2.5×10^{-3}
⁴⁰ Ca	1.3×10^{-4}	1.3×10^{-3}	1.8×10^{-3}	2.3×10^{-3}	2.6×10^{-3}
⁴¹ K	1.1×10^{-9}
⁴¹ Ca	2.5×10^{-6}	6.4×10^{-6}	7.7×10^{-6}	1.4×10^{-5}	1.7×10^{-5}
⁴² Ca	3.7×10^{-6}	1.6×10^{-5}	1.8×10^{-5}	3.3×10^{-5}	3.5×10^{-5}
⁴³ Ca	3.3×10^{-6}	1.1×10^{-5}	1.0×10^{-5}	2.0×10^{-5}	2.6×10^{-5}
⁴³ Sc	1.4×10^{-5}	4.0×10^{-5}	3.7×10^{-5}	5.3×10^{-5}	5.2×10^{-5}
⁴⁴ Ca	...	4.2×10^{-9}	5.7×10^{-7}	1.6×10^{-6}	5.1×10^{-6}
⁴⁴ Sc	6.0×10^{-9}	2.8×10^{-8}	2.2×10^{-6}	3.3×10^{-6}	9.2×10^{-6}
⁴⁴ Ti	1.5×10^{-5}	4.9×10^{-5}	6.3×10^{-5}	8.0×10^{-5}	9.0×10^{-5}
⁴⁵ Sc	1.2×10^{-6}	1.9×10^{-6}	6.0×10^{-6}	1.3×10^{-5}	2.3×10^{-5}
⁴⁵ Ti	4.0×10^{-6}	6.0×10^{-6}	1.8×10^{-5}	1.9×10^{-5}	2.3×10^{-5}
⁴⁶ Ti	7.0×10^{-6}	1.4×10^{-5}	1.8×10^{-5}	3.9×10^{-5}	5.4×10^{-5}
⁴⁷ Ti	1.4×10^{-5}	3.0×10^{-5}	3.9×10^{-5}	5.7×10^{-5}	6.7×10^{-5}
⁴⁸ Ti	2.6×10^{-9}	5.8×10^{-9}	4.9×10^{-8}	2.0×10^{-7}	5.4×10^{-7}
⁴⁸ V	2.4×10^{-6}	4.8×10^{-6}	1.3×10^{-5}	2.7×10^{-5}	4.7×10^{-5}
⁴⁸ Cr	5.2×10^{-5}	1.0×10^{-4}	1.2×10^{-4}	1.3×10^{-4}	1.3×10^{-4}
⁴⁹ Ti	1.9×10^{-9}	2.2×10^{-8}	4.8×10^{-8}	8.3×10^{-8}	1.2×10^{-7}
⁴⁹ V	4.7×10^{-5}	6.3×10^{-5}	7.9×10^{-5}	8.8×10^{-5}	9.5×10^{-5}
⁵⁰ Cr	3.9×10^{-5}	5.7×10^{-5}	6.9×10^{-5}	1.0×10^{-4}	1.3×10^{-4}
⁵¹ V	5.3×10^{-8}	1.1×10^{-7}	3.2×10^{-7}	7.9×10^{-7}	1.7×10^{-6}
⁵¹ Cr	1.2×10^{-4}	1.9×10^{-4}	2.2×10^{-4}	2.7×10^{-4}	3.1×10^{-4}
⁵² Cr	5.6×10^{-7}	7.4×10^{-7}	4.0×10^{-6}	1.3×10^{-5}	2.8×10^{-5}
⁵² Mn	2.1×10^{-4}	2.5×10^{-4}	4.0×10^{-4}	6.0×10^{-4}	7.7×10^{-4}
⁵² Fe	2.1×10^{-3}	2.2×10^{-3}	2.1×10^{-3}	1.8×10^{-3}	1.6×10^{-3}
⁵³ Mn	8.8×10^{-5}	2.1×10^{-4}	3.6×10^{-4}	4.0×10^{-4}	4.2×10^{-4}
⁵⁴ Fe	7.7×10^{-5}	1.2×10^{-4}	1.5×10^{-4}	1.8×10^{-4}	1.9×10^{-4}
⁵⁵ Mn	5.1×10^{-9}	1.6×10^{-8}	3.4×10^{-8}
⁵⁵ Fe	1.0×10^{-5}	1.7×10^{-5}	4.5×10^{-5}	7.7×10^{-5}	1.1×10^{-4}
⁵⁵ Co	2.2×10^{-4}	3.1×10^{-4}	3.0×10^{-4}	3.0×10^{-4}	2.9×10^{-4}
⁵⁶ Fe	1.0×10^{-9}	1.3×10^{-8}	1.4×10^{-7}	4.7×10^{-7}	9.9×10^{-7}
⁵⁶ Co	4.6×10^{-6}	2.2×10^{-5}	8.4×10^{-5}	1.6×10^{-4}	2.5×10^{-4}

Table 8

(Continued)

Nucleus	Burst 1	Burst 2	Burst 3	Burst 4	Burst 5
⁵⁶ Ni	7.8×10^{-4}	2.1×10^{-3}	2.7×10^{-3}	3.3×10^{-3}	3.6×10^{-3}
⁵⁷ Fe	...	3.3×10^{-8}	6.2×10^{-8}	1.0×10^{-7}	1.5×10^{-7}
⁵⁷ Co	1.3×10^{-5}	1.5×10^{-5}	2.9×10^{-5}	4.8×10^{-5}	6.6×10^{-5}
⁵⁷ Ni	5.6×10^{-4}	5.8×10^{-4}	5.7×10^{-4}	5.1×10^{-4}	4.8×10^{-4}
⁵⁸ Ni	4.1×10^{-4}	4.0×10^{-4}	4.0×10^{-4}	3.7×10^{-4}	3.6×10^{-4}
⁵⁹ Ni	6.9×10^{-4}	6.4×10^{-4}	6.3×10^{-4}	5.8×10^{-4}	5.7×10^{-4}
⁶⁰ Ni	2.4×10^{-2}	3.7×10^{-2}	4.6×10^{-2}	5.2×10^{-2}	5.8×10^{-2}
⁶¹ Ni	1.5×10^{-3}	2.1×10^{-3}	3.8×10^{-3}	5.5×10^{-3}	6.3×10^{-3}
⁶¹ Cu	5.7×10^{-3}	6.9×10^{-3}	6.5×10^{-3}	4.7×10^{-3}	3.7×10^{-3}
⁶² Ni	3.3×10^{-4}	5.8×10^{-4}	1.4×10^{-3}	1.8×10^{-3}	2.0×10^{-3}
⁶² Zn	3.6×10^{-3}	4.4×10^{-3}	3.3×10^{-3}	2.3×10^{-3}	1.9×10^{-3}
⁶³ Cu	4.1×10^{-3}	3.3×10^{-3}	2.9×10^{-3}	2.5×10^{-3}	2.5×10^{-3}
⁶⁴ Zn	4.2×10^{-2}	5.6×10^{-2}	7.0×10^{-2}	8.1×10^{-2}	9.1×10^{-2}
⁶⁵ Cu	8.1×10^{-7}	1.2×10^{-6}	4.9×10^{-6}	1.1×10^{-5}	1.7×10^{-5}
⁶⁵ Zn	8.3×10^{-3}	9.7×10^{-3}	1.2×10^{-2}	1.2×10^{-2}	1.2×10^{-2}
⁶⁶ Zn	1.3×10^{-4}	2.3×10^{-4}	9.9×10^{-4}	1.8×10^{-3}	2.3×10^{-3}
⁶⁶ Ga	2.8×10^{-3}	2.7×10^{-3}	3.1×10^{-3}	2.7×10^{-3}	2.2×10^{-3}
⁶⁶ Ge	6.8×10^{-3}	4.9×10^{-3}	3.8×10^{-3}	2.3×10^{-3}	1.9×10^{-3}
⁶⁷ Ga	4.1×10^{-3}	3.4×10^{-3}	3.6×10^{-3}	3.2×10^{-3}	3.0×10^{-3}
⁶⁸ Ge	2.8×10^{-2}	4.5×10^{-2}	5.6×10^{-2}	6.5×10^{-2}	7.1×10^{-2}
⁶⁹ Ge	1.0×10^{-2}	1.0×10^{-3}	1.2×10^{-2}	1.1×10^{-2}	1.1×10^{-2}
⁷⁰ Ge	6.5×10^{-3}	5.2×10^{-3}	5.3×10^{-3}	4.6×10^{-3}	4.2×10^{-3}
⁷¹ As	6.1×10^{-3}	4.8×10^{-3}	5.0×10^{-3}	4.4×10^{-3}	4.2×10^{-3}
⁷² Se	1.7×10^{-2}	2.8×10^{-2}	3.4×10^{-2}	3.8×10^{-2}	4.1×10^{-2}
⁷³ Se	9.1×10^{-3}	6.5×10^{-3}	7.0×10^{-3}	6.4×10^{-3}	6.3×10^{-3}
⁷⁴ Se	7.2×10^{-3}	6.5×10^{-3}	7.7×10^{-3}	7.4×10^{-3}	7.2×10^{-3}
⁷⁵ Br	8.4×10^{-3}	4.5×10^{-3}	3.3×10^{-3}	2.2×10^{-3}	2.0×10^{-3}
⁷⁶ Kr	1.3×10^{-2}	1.8×10^{-2}	2.0×10^{-2}	2.2×10^{-2}	2.3×10^{-2}
⁷⁷ Kr	7.2×10^{-3}	6.0×10^{-3}	6.2×10^{-3}	5.9×10^{-3}	5.7×10^{-3}
⁷⁸ Kr	8.7×10^{-3}	7.3×10^{-3}	7.5×10^{-3}	6.9×10^{-3}	6.5×10^{-3}
⁷⁹ Kr	7.0×10^{-3}	3.5×10^{-3}	3.0×10^{-3}	2.5×10^{-3}	2.4×10^{-3}
⁸⁰ Sr	1.3×10^{-2}	1.3×10^{-2}	1.4×10^{-2}	1.5×10^{-2}	1.5×10^{-2}
⁸¹ Rb	8.0×10^{-3}	6.4×10^{-3}	6.4×10^{-3}	6.0×10^{-3}	5.8×10^{-3}
⁸² Sr	1.7×10^{-2}	1.3×10^{-2}	1.2×10^{-2}	1.2×10^{-2}	1.1×10^{-2}
⁸³ Sr	1.3×10^{-2}	1.2×10^{-2}	1.4×10^{-2}	1.3×10^{-2}	1.3×10^{-2}
⁸⁴ Sr	1.3×10^{-2}	6.1×10^{-3}	3.8×10^{-3}	2.6×10^{-3}	2.2×10^{-3}
⁸⁵ Y	9.8×10^{-3}	7.9×10^{-3}	7.7×10^{-3}	7.2×10^{-3}	6.9×10^{-3}
⁸⁶ Zr	1.7×10^{-2}	1.9×10^{-2}	2.1×10^{-2}	2.1×10^{-2}	2.0×10^{-2}
⁸⁷ Zr	2.1×10^{-2}	9.1×10^{-3}	5.1×10^{-3}	3.8×10^{-3}	3.0×10^{-3}
⁸⁸ Zr	5.9×10^{-3}	5.5×10^{-3}	5.0×10^{-3}	5.1×10^{-3}	5.3×10^{-3}
⁸⁹ Nb	1.8×10^{-2}	2.0×10^{-2}	1.9×10^{-2}	1.8×10^{-2}	1.8×10^{-2}
⁹⁰ Mo	1.2×10^{-2}	1.1×10^{-2}	9.9×10^{-3}	9.6×10^{-3}	<




Figure 30. Upper left panel: mass fractions of the 10 most abundant stable (or $\tau > 1$ hr) isotopes, at the end of the first burst, for model 3. Lower left panel: mean post-burst composition in the envelope for each of the bursting episodes computed for model 3. Right panels: overproduction factors relative to solar ($f > 10^{-5}$), at the end of the first (upper panel) and third bursts (lower panel), for model 3.

(A color version of this figure is available in the online journal.)

relativity corrections are introduced; see Section 5.1), for our $1.4 M_\odot$ neutron star (corresponding to a surface gravity of $g = 1.08 \times 10^{14} \text{ cm s}^{-2}$); in turn, the calculations reported by Fisker et al. (2008), in a general relativity framework, relied on a 11 km ($1.4 M_\odot$) neutron star, for which $g = 1.53 \times 10^{14} \text{ cm s}^{-2}$. Although XRB properties depend weakly upon the neutron star mass (or surface gravity), part of the differences outlined between the three studies can be attributed to the combined effect of the adopted neutron star size (surface gravity) and to differences in the input physics (i.e., nuclear reaction network, opacities, treatment of convection). In particular, the use of Iben's opacities may have some effect on the peak luminosities achieved since the larger OPAL opacities will likely decrease the amount of energy radiated away from the star. Moreover, the inclusion of semiconvection and thermohaline mixing would have a minor effect in the properties of the explosions, likely affecting the appearance of marginal convective transport between bursts (see Woosley et al. 2004; Fisker et al. 2008). It is however worth noting that the convective pattern shown in Figures 18 and 32 is

similar to those reported in previous work: namely that convection sets in as soon as superadiabatic gradients are established in the envelope, following the early stages of the TNR and the corresponding rise in temperature; it reaches the surface and begins to recede before the observed burst properly commences, shutting off thereafter (Woosley et al. 2004; Fisker et al. 2008).

The potential impact of XRB nucleosynthesis on Galactic abundances is still a matter of debate. Matter accreted onto a neutron star of mass M and radius R releases $GMm_p/R \sim 200 \text{ MeV nucleon}^{-1}$, whereas only a few MeV nucleon^{-1} are released from thermonuclear fusion. Thus ejection from a neutron star is unlikely. However, it has been suggested that radiation-driven winds during photospheric radius expansion may lead to ejection of a tiny fraction of the envelope (containing nuclear processed material; see Weinberg et al. 2006; MacAlpine et al. 2007). Indeed, XRBs have been proposed as a possible source of the light p -nuclei $^{92,94}\text{Mo}$ and $^{96,98}\text{Ru}$ (Schatz et al. 1998, 2001). No matter is ejected in any of the models reported in this work, a result fully independent of the adopted resolution




Figure 31. Light curves corresponding to the first (upper left panel), second (upper right), fourth (lower left), and fifth bursts (lower right), computed for model 3.




Figure 32. Left panel: light curve comparison for the third burst computed in models 1 and 3. Right panel: development of convective regions in model 3, as a function of time. For illustrative purposes, the convective pattern obtained during the third burst is shown in the accompanying inset.

(A color version of this figure is available in the online journal.)




Figure 33. Upper panel: main nuclear activity at the ignition shell when temperature reaches a peak value of $T_{\text{peak}} = 1.23 \times 10^9$ K, during the 4th burst computed for model 3. Lower panel: same as in the upper panel, but at the end of the burst.

(A color version of this figure is available in the online journal.)




Figure 34. Distribution of the unstable species ^{92}Pd , ^{94}Ag , and $^{96,98}\text{In}$ (that power the light p -nuclei $^{92,94}\text{Mo}$ and $^{96,98}\text{Ru}$) throughout the accreted envelope, at the end of the fifth bursting episode of model 3.

(A color version of this figure is available in the online journal.)

and in agreement with all previous hydrodynamic simulations (Woosley et al. 2004; Fisker et al. 2008). Moreover, it is worth noting that, as shown in Figures 17 and 30, the abundances of many species synthesized during the bursts decrease remarkably toward the outer envelope layers, because of inefficient convective transport (see Figures 18 and 32). To assess the possible contribution to the Galactic abundances, one has to rely on the abundances of the outer envelope layers (the only ones that have a chance to be ejected by radiation-driven winds). This shows the limitations posed by one-zone nucleosynthesis calculations, in which the chemical species synthesized in the innermost layers are, by construction, assumed to represent the whole (chemically homogeneous) envelope. The mass fractions of these p -nuclei, obtained in model 3, drop by more than an order of magnitude in the outer envelope layers (as compared with the values achieved at the innermost envelope; see Figure 34); the resulting overproduction factors, $f \sim 10^6$, are several orders of magnitude smaller than those required to account for the origin of these problematic nuclei (see Weinberg et al. 2006; Bazin et al. 2008), in sharp contrast with the results obtained on the basis of one-zone calculations (Schatz et al. 1998, 2001).

The authors are grateful to the anonymous referee for a critical reading of the manuscript and for valuable comments that have

improved the presentation of this paper. This work has been partially supported by the Spanish MICINN grants AYA2007-66256 and EUI2009-04167, by the E.U. FEDER funds, by the U.S. Department of Energy under Contract No. DE-FG02-97ER41041, by the DFG cluster of excellence “Origin and Structure of the Universe,” and by the ESF EUROCORES Program EuroGENESIS.

REFERENCES

- Angulo, C., et al. 1999, *Nucl. Phys. A*, **656**, 3
- Arnould, M., & Goriely, S. 2006, *Nucl. Phys. A*, **777**, 157
- Audi, G., Bersillon, O., Blachot, J., & Wapstra, A. H. 2003a, *Nucl. Phys. A*, **729**, 3
- Audi, G., Wapstra, A. H., & Thibault, C. 2003b, *Nucl. Phys. A*, **729**, 337
- Ayasli, S., & Joss, P. C. 1982, *ApJ*, **256**, 637
- Bazin, D., et al. 2008, *Phys. Rev. Lett.*, **101**, 252501
- Belian, R. D., Conner, J. P., & Evans, W. D. 1976, *ApJ*, **206**, L135
- Bildsten, L. 1998, in *The Many Faces of Neutron Stars*, ed. R. Bucccheri, J. van Paradijs, & M. A. Alpar (Dordrecht: Kluwer), 419
- Blinnikov, S. I., Dunina-Barkovskaya, N. V., & Nadyozhin, D. K. 1996, *ApJS*, **106**, 171
- Brown, E. F. 2004, *ApJ*, **614**, L57
- Cooper, R. L., Mukhopadhyay, B., Steeghs, D., & Narayan, R. 2006, *ApJ*, **642**, 443
- Cooper, R. L., & Narayan, R. 2004, *BAAS*, **36**, 941
- Cooper, R. L., & Narayan, R. 2005, *ApJ*, **629**, 422
- Cumming, A. 2005, *Nucl. Phys. A*, **758**, 439c
- Cumming, A., & Bildsten, L. 2001, *ApJ*, **559**, L127
- Cumming, A., Morsink, S., Bildsten, L., Friedman, J. L., & Holz, D. E. 2002, *ApJ*, **564**, 343
- DeWitt, H. E., Graboske, H. C., & Cooper, M. S. 1973, *ApJ*, **181**, 439
- Elomaa, V.-V., et al. 2009, *Phys. Rev. Lett.*, **102**, 252501
- Fisker, J. L., Görres, J., Wiescher, M. C., & Davids, B. 2006, *ApJ*, **650**, 332
- Fisker, J. L., Schatz, H., & Thielemann, F.-K. 2008, *ApJS*, **174**, 261
- Fisker, J. L., Tan, W. P., Görres, J., Wiescher, M. C., & Cooper, R. L. 2007, *ApJ*, **665**, 637
- Fisker, J. L., Thielemann, F.-K., & Wiescher, M. C. 2004, *ApJ*, **608**, L61
- Fujimoto, M. Y., Sztajno, M., Lewin, W. H. G., & van Paradijs, J. 1988, *A&A*, **199**, L9
- Fuller, G. M., Fowler, W. A., & Newman, M. J. 1982a, *ApJ*, **252**, 715
- Fuller, G. M., Fowler, W. A., & Newman, M. J. 1982b, *ApJS*, **48**, 279
- Galloway, D. K., Munro, M. P., Hartman, J. M., Savov, P., Psaltis, D., & Chakrabarty, D. 2008, *ApJS*, **179**, 360
- Goriely, S. 1998, in *Nuclei in the Cosmos V*, ed. N. Prantzos & S. Harissopulos (Paris: Ed. Frontières), 314
- Graboske, H. C., DeWitt, H. E., Grossman, A. S., & Cooper, M. S. 1973, *ApJ*, **181**, 457
- Grindlay, J., Gursky, H., Schnopper, H., Parsignault, D. R., Heise, J., Brinkman, A. C., & Schrijver, J. 1976, *ApJ*, **205**, L127
- Hanawa, T., Sugimoto, D., & Hashimoto, M.-A. 1983, *PASJ*, **35**, 491
- Heger, A., Cumming, A., Galloway, D. K., & Woosley, S. E. 2007, *ApJ*, **671**, L141
- Iben, I. 1975, *ApJ*, **196**, 525
- Iliadis, C., D'Auria, J. M., Starrfield, S., Thompson, W. J., & Wiescher, M. N. 2001, *ApJS*, **134**, 151
- Jonker, P. G., Galloway, D. K., McClintock, J. E., Buxton, M., Garcia, M., & Murray, S. 2004, *MNRAS*, **354**, 666
- José, J., & Hernanz, M. 1998, *ApJ*, **494**, 680
- José, J., & Moreno, F. 2006, in *Proc. International Symp. on Nuclear Astrophysics—Nuclei in the Cosmos*, ed. A. Mengoni et al. (PoS NIC-IX; Geneva: CERN), 123
- Joss, P. C. 1977, *Nature*, **270**, 310
- Koike, O., Hashimoto, M., Arai, K., & Wanajo, S. 1999, *A&A*, **342**, 464
- Koike, O., Hashimoto, M., Kuromizu, R., & Fujimoto, S. 2004, *ApJ*, **603**, 242
- Kuulkers, E., et al. 2002, *A&A*, **382**, 503
- Kutter, G. S., & Sparks, W. M. 1980, *ApJ*, **239**, 988
- Langanke, K., & Martinez-Pinedo, G. 2000, *Nucl. Phys. A*, **673**, 481
- Lewin, W. H. G., van Paradijs, J., & Taam, R. E. 1993, *Space Sci. Rev.*, **62**, 233
- Lewin, W. H. G., van Paradijs, J., & Taam, R. E. 1995, in *X-Ray Binaries*, ed. W. H. G. Lewin, J. van Paradijs, & E. P. J. van den Heuvel (Cambridge: Cambridge Univ. Press), 175
- Liebendörfer, M., Rosswog, S., & Thielemann, F.-K. 2002, *ApJS*, **141**, 229
- MacAlpine, G. M., Ecklund, T. C., Lester, W. R., & Vanderveer, S. J. 2007, *AJ*, **133**, 81
- Maraschi, L., & Cavaliere, A. 1977, in *Highlights in Astronomy*, Vol. 4, Part I, ed. E. Muller (Dordrecht: Reidel), 127
- Parikh, A., José, J., Moreno, F., & Iliadis, C. 2008, *ApJS*, **178**, 110
- Penninx, W., Damen, E., van Paradijs, J., Tan, J., & Lewin, W. H. G. 1989, *A&A*, **208**, 146
- Prialnik, D., Shara, M. M., & Shaviv, G. 1979, *A&A*, **72**, 192
- Psaltis, D. 2006, in *Compact Stellar X-ray Sources*, ed. W. H. G. Lewin & M. van der Klis (Cambridge: Cambridge Univ. Press), 1
- Rauscher, T., & Thielemann, F.-K. 2000, *At. Data Nucl. Data Tables*, **75**, 1
- Schatz, H., Bildsten, L., Cumming, A., & Wiescher, M. 1999, *ApJ*, **524**, 1014
- Schatz, H., & Rehm, K. E. 2006, *Nucl. Phys. A*, **777**, 601
- Schatz, H., et al. 1998, *Phys. Rep.*, **294**, 167
- Schatz, H., et al. 2001, *Phys. Rev. Lett.*, **86**, 3471
- Shapiro, S. L., & Teukolsky, S. A. 1983, *Black Holes, White Dwarfs, and Neutron Stars: The Physics of Compact Objects* (New York: Wiley)
- Strohmayer, T., & Bildsten, L. 2006, in *Compact Stellar X-ray Sources*, ed. W. H. G. Lewin & M. van der Klis (Cambridge: Cambridge Univ. Press), 113
- Strohmayer, T., & Brown, E. F. 2002, *ApJ*, **566**, 1045
- Taam, R. E. 1980, *ApJ*, **241**, 358
- Taam, R. E., Woosley, S. E., & Lamb, D. Q. 1996, *ApJ*, **459**, 271
- Taam, R. E., Woosley, S. E., Weaver, T. A., & Lamb, D. Q. 1993, *ApJ*, **413**, 324
- Tan, W. P., Fisker, J. L., Görres, J., Couder, M., & Wiescher, M. C. 2007, *Phys. Rev. Lett.*, **98**, 242503
- Wallace, R. K., & Woosley, S. E. 1981, *ApJS*, **45**, 389
- Wallace, R. K., & Woosley, S. E. 1984, in *AIP Conf. Proc.*, Vol. 115, *High Energy Transients in Astrophysics*, ed. S. E. Woosley (Melville, NY: AIP), 319
- Watts, A. L., & Maurer, I. 2007, *A&A*, **467**, L33
- Weinberg, N., Bildsten, L., & Schatz, H. 2006, *ApJ*, **639**, 1018
- Woosley, S. E., & Taam, R. E. 1976, *Nature*, **263**, 101
- Woosley, S. E., & Weaver, T. A. 1984, in *AIP Conf. Proc.*, Vol. 115, *High Energy Transients in Astrophysics*, ed. S. E. Woosley (Melville, NY: AIP), 273
- Woosley, S. E., et al. 2004, *ApJS*, **151**, 75