

PAPER

Probing quantum coherence in qubit arrays

To cite this article: J Almeida *et al* 2013 *J. Phys. B: At. Mol. Opt. Phys.* **46** 104002

View the [article online](#) for updates and enhancements.

You may also like

- [Enhanced optical Anderson localization effects in modulated Bloch lattices](#)
Ramy El-Ganainy, Mohammad-Ali Miri and Demetrios Christodoulides
- [Electron localization in amorphous solids: numerical studies for the distorted diamond lattice](#)
Th Koslowski and W von Niessen
- [Analysis of localization phenomena in weakly interacting disordered lattice gases](#)
T Schulte, S Drenkelforth, J Kruse et al.



Easy-to-use and Helium-3 free
cryogenics solutions



LEARN MORE

Probing quantum coherence in qubit arrays

J Almeida^{1,2,6}, P C de Groot^{3,4,5}, S F Huelga^{1,2,6}, A M Liguori-Schremp^{1,2}
and M B Plenio^{1,2}

¹ Institut für Theoretische Physik, Universität Ulm, Albert-Einstein-Allee 11, D-89069, Ulm, Germany

² Institute for Integrated Quantum Science and Technology, University Ulm, Albert-Einstein-Allee 11, D-89069, Ulm, Germany

³ Kavli Institute of Nanoscience, Delft University of Technology, 2600 GA Delft, The Netherlands

⁴ Fakultät für Physik, Ludwig-Maximilians-Universität, Schellingstrasse 4, D-80799 München, Germany

⁵ Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-Strasse 1, D-85748 Garching, Germany

E-mail: javier.almeida@uni-ulm.de and susana.huelga@uni-ulm.de

Received 7 September 2012, in final form 23 October 2012

Published 9 May 2013

Online at stacks.iop.org/JPhysB/46/104002

Abstract

We discuss how the observation of population localization effects in periodically driven systems can be used to quantify the presence of quantum coherence in interacting qubit arrays. Essential for our proposal is the fact that these localization effects persist beyond tight-binding Hamiltonian models. This result is of special practical relevance in those situations where direct system probing using tomographic schemes becomes infeasible beyond a very small number of qubits. As a proof of principle, we study analytically a Hamiltonian system consisting of a chain of superconducting flux qubits under the effect of a periodic driving. We provide extensive numerical support of our results in the simple case of a two-qubit chain. For this system we also study the robustness of the scheme against different types of noise and disorder. We show that localization effects underpinned by quantum coherent interactions should be observable within realistic parameter regimes in chains with a larger number of qubits.

(Some figures may appear in colour only in the online journal)

1. Introduction

Transport processes are of fundamental importance in a wide variety of physical and biological systems, ranging from the actual motion of particles on a lattice [1, 2] to the transfer of classical and quantum information across spin or harmonic chains [3–5]. Relevant for our purposes, there exist specific features of the transport process that are intrinsically linked to the dynamics of the chain and in particular to whether or not the chain elements can interact coherently [6]. In the mid-80s the motion of a charged particle on a one-dimensional lattice under the influence of a time-dependent electric field was studied and shown to exhibit *dynamic localization* (DL) [1]. The canonical situation to illustrate this phenomenon is provided by an infinite linear chain of sites along which a charged particle

moves under the combined influence of a nearest-neighbour exchange interaction and a time-dependent external driving. In that setting, it was found that the mean-square displacement of the particle as a function of the field modulation E_1 , rather than exhibiting a diffusive behaviour, does not grow without bounds but oscillates sinusoidally. A related phenomenon is the so-called *coherent destruction of tunnelling* (CDT), initially formulated in dissipationless conditions for a symmetric, externally driven, double-well potential [7] and subsequently also studied in a dissipative environment [8] (and references therein). Both DL and CDT are genuine manifestations of coherent quantum effects resulting from the interference between different transition paths that leads to the selective inhibition of transport [9]. In contrast, in the classical case, and from an initially localized state, an equilibrium state would be attained in which neighbouring sites would be equally populated.

⁶ Authors to whom any correspondence should be addressed.

Ample experimental evidence supports the existence of both types of localization effects in a variety of systems. DL has been observed in Rydberg atoms, where the localization regime is characterized by ‘freezing’ the width of the wave packet with respect to the Rydberg levels [10], in driven quantum wells or semiconductor superlattices, where a suppression of the conductance was observed [11] and in ultracold atoms interacting with a standing wave of near-resonant light, where this phenomenon was found in the suppression of momentum [12, 13]. There also exist experimental proposals to use CDT as a means to control the dynamics of ultracold atoms in optical lattices [14] and to create long-distance entanglement between atoms with the possibility to use it in the implementation of quantum logical gates [14, 15]. CDT has also been recently observed in both noninteracting [16] and interacting systems [17]. Further, motivated by the desire to study the effects of quantum coherence and dephasing noise and the interplay of the two on transport processes in biological systems [18], recently the detection of DL was proposed as a way to demonstrate the possible existence of coherence effects in ion channels [19], i.e. protein complexes that regulate the flow of particular ions across the cell membrane and that are essential for a wide variety of cellular functions.

Here we extend this work and discuss the possibility of observing localization effects beyond the canonical setting, including deviations from a strict tight-binding Hamiltonian as well as the inclusion of non-Hamiltonian (noisy) effects. We will show that signatures of localization effects can still be observed in this case and apply these results to the problem of qualitatively witnessing quantum coherence in an interacting chain of superconducting qubits.

2. Renormalization of intra-qubit interactions by means of an external modulation

Motivated by specific qubit realizations in the solid state, we analyse an array of interacting qubits subject to a Hamiltonian of the form ($\hbar = 1$):

$$H_0 = \sum_{k=1}^N \frac{\omega_k}{2} \sigma_k^z + \sum_{k=1}^{N-1} J_{k,k+1} \sigma_k^x \sigma_{k+1}^x, \quad (1)$$

with N denoting the number of qubits in the chain, ω_k the site energies for each qubit and $J_{k,k+1}$ the coupling between neighbouring qubits k and $k + 1$. In the presence of a time-dependent external driving of the form

$$H_{ac}(t) = \frac{1}{2} \sum_{k=1}^N k \cdot E_{ac} \cos(\omega t) \sigma_k^z \quad (2)$$

the Hamiltonian $H(t) = H_0 + H_{ac}(t)$ of the chain reads

$$H = \frac{1}{2} \sum_{k=1}^N (\omega_k + k E_{ac} \cos(\omega t)) \sigma_k^z + \sum_{k=1}^{N-1} J_{k,k+1} \sigma_k^x \sigma_{k+1}^x. \quad (3)$$

With the substitution $\sigma_k^x = \sigma_k^+ + \sigma_k^-$, the Hamiltonian above can be rewritten as the sum of three contributions,

$$H(t) = H_z(t) + H_1 + H_2 \quad (4)$$

with,

$$H_z(t) \equiv \frac{1}{2} \sum_{k=1}^N (\omega_k + k E_{ac} \cos(\omega t)) \sigma_k^z \quad (5)$$

$$H_1 \equiv \sum_{k=1}^{N-1} J_{k,k+1} (\sigma_k^+ \sigma_{k+1}^- + \text{h.c.}) \quad (6)$$

$$H_2 \equiv \sum_{k=1}^{N-1} J_{k,k+1} (\sigma_k^+ \sigma_{k+1}^+ + \text{h.c.}). \quad (7)$$

Defining the total excitation number operator as

$$\hat{N} \equiv \sum_{k=1}^N \sigma_k^+ \sigma_k^-, \quad (8)$$

it is easy to see that $[H_z(t), \hat{N}] = [H_1, \hat{N}] = 0$ while $[H_2, \hat{N}] \neq 0$. For this reason, the term H_1 is usually referred to as an exchange interaction, in the sense that it allows for a hopping of the excitations within the chain, but it does not create or annihilate them. This is the canonical interaction in previous studies of dynamical localization in systems that can be modelled with a tight-binding Hamiltonian [1].

In the following lines we will show that the interactions described by the terms H_1 and H_2 can indeed be enhanced or inhibited separately by the proper tuning of frequency ω and amplitude E_{ac} of the external field.

To gain an insight into the problem, it is convenient to move to an interaction picture with respect to the time-dependent term $H_z(t)$. That is, we first define

$$U_0(t) \equiv \exp_+ \left\{ -i \int_0^t d\tau H_z(\tau) \right\} = \exp \left\{ -i \int_0^t d\tau H_z(\tau) \right\}, \quad (9)$$

where we have made use of the fact that $H_z(t)$ commutes with itself at different times to write the last equality above. Computing explicitly the integral above and taking into account that the operators acting on different sites commute, we have that

$$U_0 = \prod_{k=1}^N \exp \left\{ -i \frac{\omega_k t}{2} \sigma_k^z - i \frac{k E_{ac}}{2\omega} \sin(\omega t) \sigma_k^z \right\}. \quad (10)$$

Hence, the interaction picture Hamiltonian of our chain can be written as

$$H'(t) \equiv U_0(t)^\dagger (H_1 + H_2) U_0(t) \equiv H'_1(t) + H'_2(t), \quad (11)$$

where we have defined $H'_1(t) \equiv U_0(t)^\dagger H_1 U_0(t)$ and $H'_2(t) \equiv U_0(t)^\dagger H_2 U_0(t)$. Now, making use of equation (9) and the Jacobi–Anger expansion $e^{iz \sin(\phi)} = \sum_{n=-\infty}^{\infty} J_n(z) e^{in\phi}$, where $J_n(x)$ are the Bessel functions of the first kind, we can proceed further and evaluate the form of these terms explicitly as

$$\begin{aligned} H'_1(t) &= \sum_{k=1}^{N-1} J_{k,k+1} \sigma_k^+ \sigma_{k+1}^- \exp \left\{ i(\omega_k - \omega_{k+1})t - i \frac{E_{ac}}{\omega} \sin(\omega t) \right\} \\ &\quad + \text{h.c.} \\ &= \sum_{k=1}^{N-1} J_{k,k+1} \sigma_k^+ \sigma_{k+1}^- e^{i(\omega_k - \omega_{k+1})t} \sum_{n=-\infty}^{\infty} J_n \left(\frac{E_{ac}}{\omega} \right) e^{in\omega t} \\ &\quad + \text{h.c.} \end{aligned} \quad (12)$$

$$\begin{aligned}
 H'_2(t) &= \sum_{k=1}^{N-1} J_{k,k+1} \sigma_k^+ \sigma_{k+1}^+ \exp \left\{ i(\omega_k + \omega_{k+1})t \right. \\
 &\quad \left. + i \frac{E_{ac}(2k+1)}{\omega} \sin(\omega t) \right\} + \text{h.c.} \\
 &= \sum_{k=1}^{N-1} J_{k,k+1} \sigma_k^+ \sigma_{k+1}^+ e^{i(\omega_k + \omega_{k+1})t} \\
 &\quad \times \sum_{n=-\infty}^{\infty} J_n \left(\frac{E_{ac}(2k+1)}{\omega} \right) e^{in\omega t} + \text{h.c.} \quad (13)
 \end{aligned}$$

For the sake of clarity we will consider in the following the case of a homogeneous chain with $\omega_k = \omega_0$ and $J_{k,k+1} = J$ for all values of k . Under this assumption the Hamiltonian $H'(t)$ can be written as

$$H'(t) = \sum_{k=1}^{N-1} g(t) \sigma_k^+ \sigma_{k+1}^- + g'(t) \sigma_k^+ \sigma_{k+1}^+ + \text{h.c.} \quad (14)$$

with time-dependent renormalized couplings $g(t)$ and $g'(t)$ defined as

$$g(t) \equiv J \sum_{n=-\infty}^{\infty} J_n \left(\frac{E_{ac}}{\omega} \right) e^{in\omega t} \quad (15)$$

$$g'_k(t) \equiv J \sum_{n=-\infty}^{\infty} J_n \left(\frac{E_{ac}(2k+1)}{\omega} \right) e^{i(2\omega_0 + n\omega)t}. \quad (16)$$

In the regime where the tunnelling frequency of the qubits is much smaller than the frequency of the driving field, that is $J \ll \omega$, we can invoke the rotating wave approximation in the series above and neglect those terms that rotate faster than J . In particular, for $g(t)$ this means that only the non-rotating term with $n = 0$ survives and we can write

$$g(t) \equiv g = J \cdot J_0 \left(\frac{E_{ac}}{\omega} \right). \quad (17)$$

Applying the same reasoning to $g'(t)$ it follows that the only possibility to have surviving terms is that a resonance between ω_0 and ω occurs such that $2\omega_0 + n'\omega = 0$ for some integer value $n' \in \mathbb{Z}$. In this case, equation (16) can be further simplified, yielding

$$g'_k(t) \equiv g'_k = J \cdot J_{|n'|} \left(\frac{|n'|E_{ac}(2k+1)}{2\omega_0} \right). \quad (18)$$

We therefore see that the effect of the external modulation can be interpreted as renormalization of the coupling constants, imprinting a periodic dependence that will lead to a selective inhibition of transport. In the following sections we will provide numerical evidence of the accuracy of the expressions derived above. This type of localization effect will be later exploited to detect signatures of coherent interaction in arrays of coupled qubits.

3. System description

We will study the persistence of population localization effects induced by the renormalization of the hopping coupling, as explained in the previous section, in the dynamics of superconducting qubit arrays. Superconducting qubits are

effective two-level systems with a controllable transition frequency between their eigenstates, whose potential to be manufactured lithographically in a controlled manner and in a variety of geometries makes them a promising candidate for the implementation of quantum registers and information processors [20]. On the other hand, while the fabrication of structures involving many qubits is indeed feasible [21], its effective probing, and in particular the verification that the system does exhibit quantum coherence, is beyond the realm of current technology even for moderate system size. The current state of the art is provided by the tomographic analysis and the entanglement verification of three-qubit systems [22–24].

In general, the superconducting qubit Hamiltonian can be written as $H_s = (\epsilon/2) \sigma^z + (\Delta/2) \sigma^x$. Depending on the particular qubit realization, the parameters ϵ and Δ refer to different variables defining ‘charge’, ‘phase’ or ‘flux’ qubits. The latter, also called ‘persistent current qubit’, consists of a superconducting loop interrupted by three Josephson junctions, two with capacitance C_1 and the third with C_2 [25, 26, 20]. The values of the three Josephson junction coupling constants, $E_{J,1}$ corresponding to capacitance C_1 and $E_{J,2}$ to capacitance C_2 , respectively, are chosen so that the Josephson part of the Hamiltonian alone defines a bistable system. At a value of external magnetic flux $\Phi \approx 0.5\Phi_0$ ($\Phi_0 = h/2e$ is the superconducting flux quantum), the system carries either a clockwise or counterclockwise persistent current, each generating an equal but opposite magnetic flux and defining the two possible states of this qubit. With an appropriate choice of the parameters $E_{J,1,2}$ and $C_{1,2}$, the barrier in phase space separating the left and right current states can be made low enough so that tunnelling between the two classical states can occur. For the flux qubits, Δ represents the tunnelling amplitude, while the energy bias $\epsilon = -I_p(\Phi - 0.5\Phi_0)$ is proportional to the detuning $\Phi - 0.5\Phi_0$, with I_p the circulating current. The energies of the ground and first excited states are thus $E_{\mp} = \mp \sqrt{\epsilon^2 + \Delta^2}$.

For flux qubits, the most natural implementation for the interaction is through the mutual inductances between the loops. The flux generated by the one-qubit loop, which depends on its internal state, adds to the total flux picked up by the neighbouring qubits, thereby changing the energy biases of those qubits. The mutual inductance, and therefore the strength of the interaction (J), depends on the geometry of the qubit loops, specifically their size and their proximity to each other. The strength of the coupling can be enhanced in two steps. First by physically connecting the loops so that their persistent currents share a common line. In this case the kinetic inductance of the shared line adds to the geometrical inductance, where the former term can easily be the dominating part in the total interaction strength. To reach very strong coupling, for example to reach the regime $J > \Delta$, a fourth junction can be placed in the shared line. If the capacitance and Josephson junction coupling constant of this fourth junction are large compared to parameters of the qubit junctions, then the single-qubit properties are not significantly altered, while still the interaction strength can be enhanced dramatically. Coupling strengths of several GHz are easily achieved. For the sake of the suggested experiments in this paper it is




Figure 1. Schematic depiction of a superconducting qubit chain with controllable injection and readout as well as individual addressing for local modulation. By means of population measurements on a selected qubit, the presence of quantum coherence is inferred from dynamical localization effects, as described in the main text.

however not necessary to reach such high coupling values since the hopping inhibition between the qubits is enhanced in the regime where the hopping constant is smaller than the driving frequency, i.e. $J \ll \omega$ (see equations (15), (16) and the explanations below those expressions).

In the experiment described in [27], it is demonstrated how the standard flux qubit design, for which only ϵ is a tunable parameter, can be extended to also have a tunable Δ . The main change is that the smallest junction of the qubit is replaced by a small loop containing one junction in each arm, i.e. in a SQUID (superconducting quantum interference device) geometry. The SQUID acts as a single junction with tunable E_J , tuned by the flux penetrating the small loop. Local control lines can be used to change the flux through this loop, thereby controlling the $E_{J,2}$ of the qubit, and thus the Δ . Figure 1 shows a schematic of a possible implementation of a chain of strongly interacting flux qubits with tunable tunnelling splitting Δ . When the array of interacting superconducting flux qubits is operated at their corresponding degeneracy point, the chain is described by a Hamiltonian of the form (1), with $\omega_i = \Delta_i$. Typical values for the tunnelling amplitudes are in the range of 5–20 GHz while nearest-neighbour couplings $J \sim 200$ MHz. The residual next-to-nearest coupling is smaller than 10 MHz.

As far as the coupling to the environment is concerned, typically, noise sources that are located relatively far away from the chain couple to multiple or all qubits in the chain, for example magnetic coils, or some parts of the control and readout circuits. Noise sources that are located much closer, such as local control lines for the individual qubits, or microscopic noise sources in the materials surrounding the qubits, couple only, or mostly, to a single qubit. In this work we focus on the latter type: we suppose the array to be in contact with an external environment that acts locally on each qubit and can lead in principle to both pure dephasing and dissipation. In our model each qubit is coupled to its local environment via a spin-boson Hamiltonian of the form

$$H_{SB} = H_s + H_b + H_{s-b}, \quad (19)$$

where the bath is modelled as a collection of harmonic oscillators, $H_b = \sum_k \omega_k \hat{a}_k^\dagger \hat{a}_k$, and the system and bath couple linearly through the Hamiltonian [28]

$$H_{s-b} = \sigma_z^i \hat{X}_i. \quad (20)$$

Here $\hat{X}_i = \sum_k g_k (\hat{a}_k + \hat{a}_k^\dagger)$ denotes the bath's *force* operator. In the qubit eigenbasis and using the same symbols to denote the Pauli matrices in this new frame in order not to complicate




Figure 2. Coherent destruction of tunnelling. Simulations corresponding to a $N = 2$ chain with $\omega_1 = \omega_2 = 10$ GHz and hopping $J = 10$ MHz. The *black dots* have been computed with $\omega = 0.3$ GHz and $H(t) = H_z(t) + H_1$ (see definitions in the text). The *black line* has been computed with $\omega = 0.3$ GHz and $H(t) = H_z(t) + H_1 + H_2$. The *red line* has been computed with $\omega = 2.0$ GHz and $H(t) = H_z(t) + H_1 + H_2$. The dissipation rate is the same in all curves, $\gamma_{\text{diss}} = 1$ MHz. The dashed vertical lines represent the first $z_{0,1} \simeq 2.4048$ and second $z_{0,2} \simeq 5.5201$ zeros of the Bessel function $J_0(z)$.

the notation, we can derive, using the standard assumptions, a Markovian master equation $\dot{\rho} = -i[H, \rho] + L_{\text{deph}}(\rho) + L_{\text{diss}}(\rho)$ for the qubit array where the coupling to the local environment is described in terms of Lindblad terms of the form [29–32]

$$L_{\text{deph}}(\rho) = \gamma_{\text{deph}} \sum_{i=1}^N (2\sigma_i^+ \sigma_i^- \rho \sigma_i^+ \sigma_i^- - \{\sigma_i^+ \sigma_i^-, \rho\}) \quad (21)$$

$$L_{\text{diss}}(\rho) = \gamma_{\text{diss}} \sum_{i=1}^N (2\sigma_i^- \rho \sigma_i^+ - \{\sigma_i^+ \sigma_i^-, \rho\}), \quad (22)$$

with γ_{deph} and γ_{diss} denoting the dephasing and dissipation rate, respectively, $\sigma_i^\pm = (\sigma_i^x \pm i\sigma_i^y)/2$ acting on the i th qubit of the chain and $\{, \}$ the anticommutation operation. The value of the noise rates in these expressions depends strongly on the selected qubit operating point via the parameter $\theta = \arctan(\Delta/\epsilon)$. Measured values for $T_1 = \gamma_{\text{diss}}^{-1}$ range from 150 to 500 ns while pure dephasing times (T_2^*) are typically around 300 ns. In the specific case where each qubit in the chain is operated at the degeneracy point, so that $\epsilon = 0$ for every qubit, the pure dephasing term, which has a rate $\gamma_{\text{deph}} \sim \cos(\theta)$ [29], cancels out and the chain is subject to dissipative noise only. This is the first parameter regime that we are going to analyse in the following section with the aim of unveiling coherent dynamics through dynamical localization effects in a driven chain. Later, still operating each qubit at its degeneracy point, we will relax the constraint of having only dissipative noise and we will consider possible dephasing effects arising from terms with the form of equation (21).

4. Numerical results

The phenomenon described above concerning the renormalization of the coupling constants in the interaction Hamiltonian (4) under the effect of an external sinusoidal driving field can be clearly observed in figure 2. There we study the dynamics




Figure 3. Resonance conditions. Simulations for a $N = 2$ chain with $\omega_1 = \omega_2 = 10$ GHz, $J = 10$ MHz and a dissipation rate $\gamma_{\text{diss}} = 1$ MHz. For each value of ω , the electric field is fixed so that $E_{\text{ac}}/\omega = z_{0,1}$, where $z_{0,1} \simeq 2.4048$ is the first zero of the Bessel function $J_0(z)$. The axis on the top represents some integer values $n' = 2\omega_0/\omega$ for which the resonance condition in the coupling constant g' is fulfilled.

of a chain consisting in two inductively coupled superconducting qubits operated at their degeneracy points, so that the (undriven) system Hamiltonian is given by equation (1). In this case, as discussed before, the noise is purely dissipative when expressed in the rotated basis. The system is initialized so that only the first site is excited. The different curves correspond to the maximum value of the population that has been transferred from the first to the second qubit within a sufficiently long time interval (1 μ s). The black line corresponds to an off-resonance situation ($\omega_0 = 10$ GHz, $\omega = 0.3$ GHz, with no integer value n' such that $2\omega_0 + n'\omega = 0$). In this situation the coupling of the contribution given by H_2 effectively renormalizes to zero and the dynamics is governed by the tight-binding term H_1 . According to expression (15), it is the Bessel function $J_0(E_{\text{ac}}/\omega)$ which governs this behaviour and when its argument E_{ac}/ω coincides with one of its zeros, then the hopping between both qubits is suppressed. On the other hand, the red curve has been computed on a resonance situation ($\omega_0 = 10$ GHz, $\omega = 2.0$ GHz, such that $2\omega_0 + n'\omega = 0$ for $n' = -10$). In this situation both couplings g and g' in equations (15) and (16) are in general different from zero and the total dynamics is hence more convoluted. Note that the fact that both the red and black curves are indistinguishable to the eye for low values of E_{ac}/ω is due to the slow buildup of the Bessel function $J_{n'}(z)$ (with $n' = 10$) that governs the H_2 term. Finally, the black circles in figure 2 have been computed using the same parameters that we used for the off-resonance situation but this time we explicitly excluded the contribution given by the H_2 terms. The fact that the black line and the black circles superimpose each other is a clear indication that the term H_2 effectively renormalizes to zero when the field is out of resonance with this term.

The existence of resonance conditions for the terms contributed by H_2 in expression (4) is clearly illustrated in figure 3, which has been evaluated with the same parameters as in figure 2. We have however fixed the ratio $E_{\text{ac}}/\omega = z_{0,1}$ with $z_{0,1} \simeq 2.4048$, the first zero of the Bessel function of order zero. With this constraint the population transfer induced by the terms contributed by H_1 in equation (4) is completely




Figure 4. Effect of frequency inhomogeneities. Maximum value of the population transferred from qubit 1 to qubit 2 in a $N = 2$ chain after a fixed time $t_{\text{max}} = 500$ ns. The frequency of the external field is $\omega = 1.3$ GHz and its magnitude is fixed so that $E_{\text{ac}}/\omega = 1.2$, far from the closest zero of the $J_0(z)$ Bessel function. The parameters of the qubits are $\omega_0 = 10$ GHz and $J = 10$ MHz. The dephasing and dissipation rates have been set to zero for clarity.

suppressed. We have however the freedom to use different values of the driving field ω such that the term H_2 is either on- or off-resonance with the frequency ω_0 . We can clearly see in figure 3 those values (compatible with the discrete grid used to sample the frequency ω) where the resonance condition is fulfilled and a peak in the population transfer appears, induced entirely by the terms in H_2 .

As a result, the presence of a *correction* to the canonical Hamiltonian H_1 does not hinder the manifestation of localization effects. By appropriate tuning of the external driving we can select resonance conditions that lead to inhibition of transport and provide a fingerprint of the underlying coherent evolution. In figure 4 we illustrate the effect of an inhomogeneous distribution of the tunnelling amplitudes of the qubits within the chain. Not surprisingly, the presence of this sort of disorder in the array leads to a quick loss of contrast. However, the fact that the tunnelling amplitude of superconducting flux qubits is actually tunable can provide a mechanism to overcome this difficulty by minimizing or even suppressing the local disorder in the chain⁷. On the other hand, real implementations of interacting superconducting flux qubits typically result in interaction strength differences of the order of a few per cent between different nearest neighbours. It is important to stress at this point that these differences have little effect in our previous discussion since they do not affect the resonance conditions in equations (15) and (16) that are the key to the hopping inhibition effect presented in this work.

In real implementations of superconducting flux qubits we should also expect deviations from a purely dissipative model of noise. This motivates the study of the effect of pure dephasing processes in our system. To this end we will introduce in the master equation a Lindblad term of the form given in equation (22), with dephasing rate γ_{deph} . To single out the effect of pure dephasing, we will take a vanishingly small dissipation rate, that is $\gamma_{\text{diss}} = 0$.

In figure 5 we have studied how the transference of population in a chain is affected by pure dephasing noise terms. In figure 5 (top) we plot the dynamics of

⁷ Recent work on tunable schemes is for instance reported in [33].




Figure 5. Effect of pure dephasing noise. Top: population transfer from qubit 1 to qubit 2 in a $N = 2$ chain and the following parameters: $\omega_1 = \omega_2 = 10$ GHz, $J = 10$ MHz, $\omega = 1.3$ GHz. The dissipation rate has been set to zero ($\gamma_{diss} = 0.0$) for clarity and a new dephasing rate γ_{deph} has been included ($\mathcal{L}_{deph} \equiv \gamma_{deph} \sum \mathcal{L}_k(\sigma_z^k)$, with $\mathcal{L}_k(\sigma_z^k) = \sigma_k^z \rho \sigma_k^z - \rho$). Each panel corresponds to a fixed value $z = E_{ac}/\omega$ with $z_{0,1} \simeq 2.4048$, the first zero of the Bessel function $J_0(z)$. Bottom: maximum value of the population transferred to the second qubit in the time interval and with the same parameters as those in the figure on top and the following dephasing rates (in GHz): $\gamma_{deph} = 0.0$ (blue), $\gamma_{deph} = 0.001$ (green), $\gamma_{deph} = 0.005$ (red), $\gamma_{deph} = 0.01$ (turquoise), $\gamma_{deph} = 0.05$ (cyan), $\gamma_{deph} = 0.1$ (yellow).

the population for different values of the dephasing rate, $\gamma_{deph} = \{0.0, 0.001, 0.01, 0.1\}$ GHz. It is worth remarking that the inclusion of these terms in the master equation leads to an asymptotic stationary state that is maximally mixed [34]. However, the transient dynamics still provides valuable information. It is remarkable to note that even though the dephasing rapidly destroys the coherent oscillations of the population, still the renormalization of the hopping rate survives for surprisingly large values of γ_{deph} . This fact can be clearly appreciated in the panel with $E_{ac}/\omega = z_{0,1}$ where the hopping is strongly inhibited and still notable for dephasing rates as high as $\gamma_{deph} = 100$ MHz. In figure 5 (bottom) we have plotted the same information in a more compact format. In this graph the effect of dephasing can be clearly seen to reduce the visibility of the population transfer oscillations (as a function of E_{ac}/ω). Nonetheless, as stated above, the hopping modulation is quite visible also in this graph even for large values of the dephasing rate.




Figure 6. Quantum coherence in a $N = 2$ chain. In this graph we plot the magnitude $C \equiv \max \sum_{i,j \neq i} \text{abs}(\rho_{i,j}(t))$ as a function of the visibility of the transport pattern (population fringes in figure 5). We have considered 27 values of the pure dephasing rate in a logarithmic scale from $\gamma_{deph} = 0.1$ GHz ($C_{max} \simeq 0$) to $\gamma_{deph} = 0.0$ GHz ($C_{max} = 0.5$). For convenience we have fixed $\gamma_{diss} = 0$.




Figure 7. Hopping inhibition in a $N = 6$ chain. Maximum value of the population transferred from the first qubit to the last one in a $N = 6$ chain within a time interval of $2.8 \mu s$. The frequency of the external field is $\omega = 1.3$ GHz and its magnitude is fixed so that $E_{ac}/\omega = 1.2$, far from the closest zero of the $J_0(z)$ Bessel function. The parameters of the qubits are $\omega_1 = 10$ GHz and $J = 10$ MHz. For this computation we have considered dissipation noise only, with the following rates (in GHz): $\gamma_{diss} = 0.0001$ (blue), $\gamma_{diss} = 0.0005$ (green), $\gamma_{diss} = 0.001$ (red), $\gamma_{diss} = 0.005$ (turquoise, slightly visible).

The qualitative relation between the properties of the transport dynamics and the coherence in the system is illustrated in figure 6. In this figure we have represented the coherence of the chain, quantified by the sum of the off-diagonal elements of the density matrix $C \equiv \max \sum_{i,j \neq i} \text{abs}(\rho_{i,j}(t))$, as a function of the visibility of the population fringes depicted in figure 5. For the sake of clarity we have used a purely dephasing noise since it uniquely affects the coherence terms of the density matrix. The main conclusions are however unaffected by including dissipation terms. We can see in this graph that the degree of quantum coherence of the system can be properly quantified by the transport schemes proposed in this work. For small values of the dephasing rate, the relation between coherence and visibility is indeed essentially linear. As a result, population measurements alone would allow, in the presence of a tunable driving, signatures of quantum coherence in the system to be detected. The scalability of the procedure is illustrated in figure 7 for an array of $N = 6$ qubits, a system size that is currently untractable with tomographic schemes. As

opposed to the previous graphs computed for $N = 2$, now the one-excitation sector contains more than two eigenstates and hence the population dynamics is affected by more than one characteristic frequency. This results in a more convoluted dynamics and different possible protocols to measure the hopping inhibition. For this graph we have chosen to start with a chain where only the first qubit is initially on its excited configuration; we measure then the maximum population transferred to the last qubit within a time interval of $2.8 \mu\text{s}$, long enough to allow the initial excitation to reach the end of the chain. Note in this graph that the perfect hopping inhibition is again only achieved at the zeros of the corresponding Bessel functions. The fact that the hopping seems to vanish in an extended interval around this point is only a consequence of the interplay between the finite time interval used to perform the population measure and the arrival time required for a wave packet created at the beginning of the chain to reach its end (see [15] for a thorough study of this topic), which increases as the hopping rate decreases.

5. Conclusions

To summarize, we have analysed the persistence of localization effects beyond exact tight binding Hamiltonians and beyond a closed system description. Introducing an a.c. interaction term of the form of equation (2), we have seen that the original ZZ coupling, which provides the natural model for the actual coupling in superconducting architectures, can be effectively mapped into the Hamiltonian (14) where the excitation-preserving and non-preserving terms are affected by two different renormalized couplings g and g' . We have seen that these effective couplings are determined by certain resonance conditions (for the coupling g there is always a resonance for $n' = 0$, for g' the stronger condition $2\omega_0 + n'\omega = 0$ is required) and, given that the resonance conditions are fulfilled, by the arguments of the Bessel functions that define these couplings. In conclusion, we have proposed a method that allows us to tune independently two kinds of interaction of very different nature but with the common feature of leading to localization phenomena. These are shown to be useful for witnessing the coherent behaviour of coupled qubit arrays. As a proof of principle, using typical parameter regimes in chains of superconducting flux qubits, we have shown that transport inhibition can be qualitatively linked to the degree of coherence in the system. These types of experiments, which involve population measurements only, can provide a benchmark for quantum behaviour in systems whose complexity makes them unsuited for detailed tomographic analysis, ranging from arrays of self-assembled quantum dots [35] to coupled nanomagnets [36].

Acknowledgments

Financial support from the EU Integrated Project Q-Essence, STREP actions CORNER and PICC and the Humboldt Foundation is gratefully acknowledged. We thank H Mooij, A Rivas and B Röthlisberger for helpful discussions and their comments on the precursor of this manuscript and to A

Bermúdez and M Paternostro for their feedback on the current manuscript.

References

- [1] Dunlap D H and Kenkre V M 1986 Dynamic localization of a charged particle moving under the influence of an electric field *Phys. Rev. B* **34** 3625
- [2] Holthaus M and Hone D W 1996 Localization effects in ac-driven tight-binding lattices *Phil. Mag. B* **74** 104–37
- [3] Audenaert K, Eisert J, Plenio M B and Werner R F 2002 Entanglement properties of the harmonic chain *Phys. Rev. A* **66** 042327
- [4] Plenio M B, Hartley J and Eisert J 2004 Dynamics and manipulation of entanglement in coupled harmonic systems with many degrees of freedom *New J. Phys.* **6** 36
- [5] Bose S 2007 Quantum communication through spin chain dynamics: an introductory overview *Contemp. Phys.* **48** 13
- [6] For a recent review, see, Kohler S, Lehmann J and Hänggi P 2005 *Phys. Rep.* **406** 379
- [7] Grossmann F, Dittrich T, Jung P and Hänggi P 1991 Coherent destruction of tunneling *Phys. Rev. Lett.* **67** 516
- [8] Grifoni M and Hänggi P 1998 Driven quantum tunneling *Phys. Rep.* **304** 229
- [9] Kayanuma Y and Saito K 2008 Coherent destruction of tunneling, dynamical localization and the Landau–Zehner formula *Phys. Rev. A* **77** 010101
- [10] Blümel R, Buchleitner A, Graham R, Sirko L, Smilansky U and Walther H 1991 Dynamical localization in the microwave interaction of Rydberg atoms: the influence of noise *Phys. Rev. A* **44** 4521
- [11] Keay B J, Zeuner S, Allen S J Jr, Maranowski K D, Gossard A C, Bhattacharya U and Rodwell M J W 1995 Dynamic localization, absolute negative conductance, and stimulated, multiphoton emission in sequential resonant tunneling semiconductor superlattices *Phys. Rev. Lett.* **75** 4102
- [12] Moore F L, Robinson J C, Bharucha C F, Williams P E and Raizen M G 1994 Observation of dynamical localization in atomic momentum transfer: a new testing ground for quantum chaos *Phys. Rev. Lett.* **73** 2974
- [13] Bharucha C F, Robinson J C, Moore F L, Sundaram B, Niu Q and Raizen M G 1999 Dynamical localization of ultracold sodium atoms *Phys. Rev. E* **60** 3881
- [14] Creffield C E 2007 Quantum control and entanglement using periodic driving fields *Phys. Rev. Lett.* **99** 110501
- [15] Galve F, Zueco D, Kohler S, Lutz E and Hänggi P 2009 Entanglement resonance in driven spin chains *Phys. Rev. A* **79** 032332
- [16] Kierig E *et al* 2008 Single-particle tunneling in strongly driven double-well potentials *Phys. Rev. Lett.* **100** 190405
- [17] Eckardt A and Holthaus M 2008 Avoided-level-crossing spectroscopy with dressed matter waves *Phys. Rev. Lett.* **101** 245302
- [18] Fleming G R, Huelga S F and Plenio M B 2011 Focus on quantum effects and noise in biomolecules *New J. Phys.* **13** 115002
- [19] Vaziri A and Plenio M B 2010 Quantum coherence in ion channels: resonances, transport and verification *New J. Phys.* **12** 085001
- [20] Devoret M H, Wallraff A and Martinis J M 2004 Superconducting qubits: a short review arXiv:cond-mat/0411174
- [21] You J Q and Nori F 2011 Atomic physics and quantum optics using superconducting circuits *Nature* **474** 589
- [22] Johnson M W *et al* 2011 Quantum annealing with manufactured spins *Nature* **473** 194

- [22] Neeley M *et al* 2010 Generation of three-qubit entangled states using superconducting phase qubits *Nature* **467** 570
- [23] DiCarlo L *et al* 2010 Preparation and measurement of three-qubit entanglement in a superconducting circuit *Nature* **467** 574
- [24] Lucero E *et al* 2012 Computing prime factors with a Josephson phase qubit quantum processor *Nature Phys.* **8** 719–23
- [25] Mooij J E, Orlando T P, Tian L, van der Wal C H, Levitov L S, Lloyd S and Mazo J J 1999 A superconducting persistent current qubit *Science* **285** 1036
- [26] Orlando T P, Mooij J E, Tian L, van der Wal C H, Levitov L S, Lloyd S and Mazo J J 1999 Superconducting persistent-current qubit *Phys. Rev. B* **60** 15398–413
- [27] Paauw F G, Fedorov A, Harmans C J P M and Mooij J E 2009 Tuning the gap of a superconducting flux qubit *Phys. Rev. Lett.* **102** 090501
- [28] Shnirman A, Makhlin Y and Schön G 2002 Noise and decoherence in quantum two-level systems *Phys. Scr. T* **102** 147
- [29] Tsomokos D I, Hartmann M J, Huelga S F and Plenio M B 2007 Entanglement dynamics in chains of qubits with noise and disorder *New J. Phys.* **9** 79
- [30] Rivas A, Plato A D K, Huelga S F and Plenio M B 2010 Markovian master equations: a critical study *New J. Phys.* **12** 113032
- [31] Rivas A and Huelga S F 2011 *Open Quantum Systems. An Introduction* (Heidelberg: Springer)
- [32] Oxtoby N P, Rivas A, Huelga S F and Fazio R 2009 Probing a composite spin-boson environment *New J. Phys.* **11** 063028
- [33] Bialczak R C *et al* 2011 Fast tunable coupler for superconducting qubits *Phys. Rev. Lett.* **106** 060501
- [34] Rivas A, Oxtoby N P and Huelga S F 2009 Stochastic resonance phenomena in spin chains *Eur. Phys. J. B* **69** 51
- [35] Miller B T *et al* 1997 Few-electron ground states of charge-tunable self-assembled quantum dots *Phys. Rev. B* **56** 6764
- [36] Timco G A *et al* 2009 Engineering the coupling between molecular spin qubits by coordination chemistry *Nature Nanotechnol.* **4** 173