

Proper time flow equation for gravity

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JHEP02(2005)035

(<http://iopscience.iop.org/1126-6708/2005/02/035>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 38.107.179.213

The article was downloaded on 22/02/2012 at 13:21

Please note that [terms and conditions apply](#).

Proper time flow equation for gravity

Alfio Bonanno

*INAF - Osservatorio Astrofisico di Catania
Via S. Sofia 78, I-95123 Catania, Italy
INFN, Via S. Sofia 64, I-95123 Catania, Italy
E-mail: abo@ct.astro.it*

Martin Reuter

*Institute of Physics, University of Mainz
Staudingerweg 7, D-55099 Mainz, Germany
E-mail: reuter@zino.physik.uni-mainz.de*

ABSTRACT: We analyze a proper time renormalization group equation for Quantum Einstein Gravity in the Einstein-Hilbert truncation and compare its predictions to those of the conceptually different exact renormalization group equation of the effective average action. We employ a smooth infrared regulator of a special type which is known to give rise to extremely precise critical exponents in scalar theories. We find perfect consistency between the proper time and the average action renormalization group equations. In particular the proper time equation, too, predicts the existence of a non-gaussian fixed point as it is necessary for the conjectured nonperturbative renormalizability of Quantum Einstein Gravity.

KEYWORDS: Renormalization Group, Models of Quantum Gravity, Nonperturbative Effects.

Contents

1. Introduction	1
2. The proper time flow equation	2
3. The β-functions of g and λ	4
4. Universal quantities	6
5. Conclusions	7

1. Introduction

Recently a lot of work went into the exploration of the Wilsonian renormalization group (RG) flow of Quantum Einstein Gravity [1]–[9] and its possible impact on black holes and cosmology [10]–[16]. By definition, Quantum Einstein Gravity (QEG) is the conjectured quantum field theory of the spacetime metric whose bare action is (infinitesimally close to) an ultraviolet (UV) attractive non-gaussian fixed point of the RG flow. As outlined by S. Weinberg long ago [17] this theory would be nonperturbatively renormalizable or “asymptotically safe” and, most probably, predictive and internally consistent even at the shortest sub-Planckian distances. Most of the existing RG calculations were performed within the framework of the exact RG equation pertaining to the effective average action [18, 19, 20], applied to a truncated space of action functionals (theory space). It turned out that the UV fixed point necessary for asymptotic safety does indeed exist within the Einstein-Hilbert [6, 3, 4] and the R^2 -truncation [5], and detailed analyses of the reliability of those approximations [3, 5] revealed that the fixed point found is unlikely to be a truncation artifact. But clearly it is necessary to collect further evidence for its existence at the exact level, both within the effective average action formalism (by generalizing the truncation) and with conceptually independent methods.

In fact, a conceptually independent support of the asymptotic safety conjecture comes from the work of Forgács and Niedermaier [21] who were able to analyze the path integral over the metrics with two Killing vectors exactly and to show that at least this sector is asymptotically safe. In view of recent progress [22] on causal, i.e. lorentzian dynamical triangulations [23] one also might hope that it will soon be possible to compare the analytical results to Monte Carlo data.

In the present paper we are going to analyze the non-gaussian fixed point of QEG using the “proper time renormalization group” which has been extensively used for other theories recently [24]–[27]. While this formalism is less complete and systematic than the exact RG

equation of the average action [20], it has had quite spectacular successes in high precision calculations of critical exponents [25]. Even though, at the technical level, the proper time method is reminiscent of the average action approach, it is well known [26] that the proper time RG equation is *not* a special case of the exact RG equation of the average action; it is conceptually independent in this sense. The approach we are going to use implements a certain RG improvement of one-loop perturbation-theory, employing an infrared (IR)-regularized proper time representation of the corresponding one-loop determinants. We shall derive a flow equation for the scale dependent gravitational action from it whose structure is quite different from that of the exact RG equation of the average action, for every possible choice of its built-in cutoff operator \mathcal{R}_k [19]. Apart from the conceptual independence the main motivation for the present analysis are the remarkably precise critical exponents which have been obtained for the Wilson-Fisher fixed point with this method [25]. In fact, we shall be mostly interested in the universal properties of the non-gaussian fixed point of QEG, (g^*, λ^*) , in particular its critical exponents θ' and θ'' and the product $g^* \lambda^*$.

2. The proper time flow equation

The ultimate goal of the RG approach would be the calculation of the integral $Z = \int \mathcal{D}\gamma_{\mu\nu} \exp(-S[\gamma_{\mu\nu}])$ over all metrics $\gamma_{\mu\nu}$ where S is an arbitrary bare action, invariant under general coordinate transformations. We are going to employ the background field formalism and start by writing $\gamma_{\mu\nu}$ as a background $\bar{g}_{\mu\nu}$ plus a (not necessarily small) fluctuation: $\gamma_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}$. Furthermore, we gauge fix the path integral by adding gauge fixing and ghost terms to S . We use an a priori arbitrary gauge fixing term $S_{\text{gf}}[h; g]$ of the background-type, meaning that it is invariant under a *combined* transformation of $h_{\mu\nu}$ and $\bar{g}_{\mu\nu}$. This property will guarantee that the effective action becomes invariant under general coordinates transformation. Thus, denoting the Faddeev-Popov ghosts by C^μ and \bar{C}_μ , and their action by $S_{\text{gh}}[h, C, \bar{C}; \bar{g}]$, the path integral reads $Z = \int \mathcal{D}h \mathcal{D}C \mathcal{D}\bar{C} \exp(-S[\bar{g}+h] - S_{\text{gf}} - S_{\text{gh}})$. Next we couple $h_{\mu\nu}$ to a source, Legendre transform $\ln Z$ in the usual way, and thus obtain the effective action $\Gamma[\bar{h}; \bar{g}]$. Here $\bar{h}_{\mu\nu} \equiv \langle h_{\mu\nu} \rangle$ is the expectation value of the fluctuation. The corresponding expectation value of the complete metric will be obtained by $g_{\mu\nu} = \langle \gamma_{\mu\nu} \rangle \equiv \bar{g}_{\mu\nu} + \bar{h}_{\mu\nu}$. It is customary to consider Γ a functional of g and \bar{g} rather than \bar{h} and \bar{g} , and to define $\Gamma[g, \bar{g}] \equiv \Gamma[\bar{h} = g - \bar{g}; \bar{g}]$. As always in the background formalism [1, 28, 20] the “ordinary” effective action, the generating functional of one-particle irreducible Green’s functions, $\Gamma[g]$, is obtained by setting $\bar{h} = 0$ or $\bar{g} = g$: $\Gamma[g] = \Gamma[g, g] \equiv \Gamma[\bar{h} = 0; \bar{g}]$.

As in every field theory, the one-loop correction to the effective action is given by a set of determinants:

$$\Gamma_1[g, \bar{g}] = \frac{1}{2} \ln \det \widehat{S}^{(2)} - \ln \det S_{\text{gh}}^{(2)}. \tag{2.1}$$

Here we set $\widehat{S}[\bar{h}; g] \equiv S[\bar{g} + \bar{h}] + S_{\text{gf}}[\bar{h}; \bar{g}]$ and $\widehat{S}^{(2)}$ denotes the matrix of the second functional derivatives of \widehat{S} with respect to $\bar{h}_{\mu\nu}$, and likewise for the ghosts.

In order to evaluate $\ln \det \Omega = \text{Tr} \ln \Omega$ for $\Omega = \widehat{S}^{(2)}$ and $S_{\text{gh}}^{(2)}$ we use the proper time representation $\text{Tr} \ln \Omega = - \int_0^\infty (ds/s) \text{Tr} \exp(-s\Omega)$. We regularize the short-distance singularities of this expression by introducing a UV cutoff Λ which suppresses the contributions to the s -integral with $s \lesssim \Lambda^{-2}$. Furthermore, in order to set up a Wilsonian RG equation, we also introduce a IR cutoff k which suppresses the contributions due to large proper times $s \gtrsim k^{-2}$. In principle one could employ a sharp proper time cutoff which amounts to replacing $\int_0^\infty ds$ with $\int_{1/\Lambda^2}^{1/k^2} ds$. In this paper we use instead a more general class of proper time cutoffs. One writes $\text{Tr}[\ln \Omega]_{\text{reg}} = - \text{Tr} \int_0^\infty f_k(s) \exp(-s\Omega)$ where the shape of the function $f_k(s)$ is chosen such that it interpolates smoothly between $f_k(s) \approx 0$ for $s \gg k^{-2}$ and $f_k(s) \approx 1$ for $s \ll k^{-2}$, and similarly in the UV. Actually the UV cutoff will play no role in that follows since we shall concentrate on the k -derivative of the regularized determinant. Introducing the ‘‘RG-time’’ $t = \ln k$ we have

$$\partial_t \text{Tr}[\ln \Omega]_{\text{reg}} = - \text{Tr} \int_0^\infty \frac{ds}{s} \partial_t f_k(s) \exp(-s\Omega). \tag{2.2}$$

For an appropriately chosen $f_k(s)$ the scale derivative $\partial_t f_k(s)$ is independent of Λ . Hence, at the level of (2.2), we may safely perform the limit $\Lambda \rightarrow \infty$.

For actual calculations we shall use the following one-parameter family of smooth cutoffs $f_k \equiv f_k^m$ which has been extensively used in the literature [24], in particular in high precision calculation of the critical exponents¹ [25]:

$$f_k^m(s) = \frac{\Gamma(m+1, \mathcal{Z}sk^2) - \Gamma(m+1, \mathcal{Z}s\Lambda^2)}{\Gamma(m+1)}. \tag{2.3}$$

Here m is an arbitrary real, positive parameter which controls the shape of the f_k^m in the interpolating regions, and $\Gamma(\alpha, x) = \int_x^\infty dt t^{\alpha-1} e^{-t}$ denotes the incomplete Gamma-function. Furthermore, \mathcal{Z} is a constant (actually a matrix in field space) which has to be adjusted in the following way. If the field of type ‘‘ a ’’ has a kinetic term $-Z_a D^2$, we set $\mathcal{Z} = Z_a$ for this field to make sure that the eigenvalues of $-D^2$ are cut off precisely at k^2 , rather than k^2/Z_a [19]. The scale derivative of (2.3) is given by

$$\partial_t f_k^m(s) = -2 \frac{1}{m!} (\mathcal{Z}sk^2)^{m+1} \exp(-\mathcal{Z}sk^2) \tag{2.4}$$

which is indeed independent of Λ . Inserting (2.4) into (2.2) and performing the integral we obtain

$$\partial_t \text{Tr}[\ln \Omega]_{\text{reg}} = 2 \text{Tr} \left(\frac{\mathcal{Z}k^2}{\Omega + \mathcal{Z}k^2} \right)^{m+1}. \tag{2.5}$$

For m integer, in the limit $\Lambda \rightarrow \infty$, eq. (2.3) yields the following explicit representation:

$$f_k^m(s) = \exp(-\mathcal{Z}sk^2) \sum_{\mu=0}^m \frac{1}{\mu!} (\mathcal{Z}sk^2)^\mu. \tag{2.6}$$

¹We do not perform the rescaling $k^2 \rightarrow mk^2$ as in [27].

If we regularize the determinants in (2.1) with a proper time cutoff then Γ_1 and, as a consequence, the complete one loop effective action becomes k -dependent. We denote it by $\widehat{S}_k[g, \bar{g}] \equiv \widehat{S}[\bar{h}; \bar{g}] + \Gamma_1[g, \bar{g}]_{\text{reg}}$. Its scale derivative is given by

$$\partial_t \widehat{S}_k[g, \bar{g}] = -\frac{1}{2} \text{Tr} \int_0^\infty \frac{ds}{s} \partial_t f_k^m(s) [\exp(-s\widehat{S}^{(2)}) - 2 \exp(-sS_{\text{gh}}^{(2)})]. \quad (2.7)$$

Up to this point we performed a standard one-loop calculation, with the classical action \widehat{S} appearing on the r.h.s. of (2.7), and the quantum corrected \widehat{S}_k on its l.h.s. . Now we perform a RG-improvement of this one-loop calculation by feeding-back \widehat{S}_k into the r.h.s. of (2.7). This leads to the following “proper time RG equation” for the functional \widehat{S}_k :

$$\partial_t \widehat{S}_k[g, \bar{g}] = -\frac{1}{2} \text{Tr} \int_0^\infty \frac{ds}{s} \partial_t f_k^m(s) [\exp(-s\widehat{S}_k^{(2)}) - 2 \exp(-sS_{\text{gh}}^{(2)})]. \quad (2.8)$$

For a detailed discussion of flow equations of this type we refer to the literature [24, 25]. Note that only *after* the derivatives implicit in $\widehat{S}_k^{(2)}$ have been performed one can set $g = \bar{g}$ in order to obtain the “ordinary” running action $S_k[g] \equiv \widehat{S}_k[g, g]$ since these derivatives are to be performed at fixed \bar{g} .

The above flow equation could be generalized to allow for an RG evolution in the ghost sector. Starting from the standard one-loop effective action $\Gamma_1[g, \bar{g}, \xi, \bar{\xi}]_{\text{reg}}$ in presence of nonvanishing ghost expectation values $\xi = \langle C \rangle$ and $\bar{\xi} = \langle \bar{C} \rangle$, and following the same steps as above, we arrive at

$$\partial_t \widetilde{S}_k[g, \bar{g}, \xi, \bar{\xi}] = -\frac{1}{2} \text{STr} \int_0^\infty \frac{ds}{s} \partial_t f_k^m(s) \exp(-s\widetilde{S}_k^{(2)}). \quad (2.9)$$

The functional $\widetilde{S}_k[g, \bar{g}, \xi, \bar{\xi}]$ is a generalization of $\widehat{S} + S_{\text{gh}}$ at the tree-, and $\widehat{S} + S_{\text{gh}} + \Gamma_1[g, \bar{g}, \xi, \bar{\xi}]_{\text{reg}}$ at the one loop level, respectively. In (2.9), $\widetilde{S}_k^{(2)}$ is a supermatrix involving derivatives with respect to g , ξ and $\bar{\xi}$, and the supertrace takes care of the relative minus sign between the graviton and the ghost contributions. In the present paper we are not going to use the generalized flow equation (2.9).

3. The β -functions of g and λ

The RG equation (2.8) describes a flow on the infinite dimensional space of functionals $\widehat{S}[g, \bar{g}]$. We can try to obtain nonperturbative solutions (RG trajectories) by truncating this theory space. In the following we shall consider the Einstein-Hilbert approximation [1] whose truncated theory space is 2-dimensional. The corresponding ansatz for \widehat{S}_k is parametrized by a k -dependent Newton constant G_k and cosmological constant $\bar{\lambda}_k$:

$$\widehat{S}_k[g, \bar{g}] = 2\kappa Z_{Nk} \int d^d x \sqrt{\bar{g}} \{-R(g) + 2\bar{\lambda}_k\} + \kappa^2 Z_{Nk} \int d^d x \sqrt{\bar{g}} \bar{g}^{\mu\nu} (\mathcal{F}_\mu^{\alpha\beta} g_{\alpha\beta}) (\mathcal{F}_\nu^{\rho\sigma} g_{\rho\sigma}). \quad (3.1)$$

The first term on the r.h.s. of (3.1) is the familiar Einstein-Hilbert action in d -dimensions with the k -dependent couplings. We wrote $G_k \equiv \bar{G}/Z_{Nk}$ for the running Newton constant, with a fixed constant \bar{G} , and introduced the convenient abbreviation $\kappa = (32\bar{G})^{-1/2}$. The

second term of (3.1) is the gauge fixing term for the background version of the harmonic coordinate condition [28, 1], with gauge fixing parameter $\alpha = 1$. The operator $\mathcal{F}_\mu^{\alpha\beta}$ is given by $\mathcal{F}_\mu^{\alpha\beta} = \delta_\mu^\beta \bar{g}^{\alpha\gamma} \bar{D}_\gamma - \frac{1}{2} \bar{g}^{\alpha\beta} \bar{D}_\mu$ where the covariant derivative \bar{D}_μ is constructed from the background metric. Note that $\mathcal{F}_\mu^{\alpha\beta} \bar{g}_{\alpha\beta} = 0$ so that the gauge fixing term drops from (3.1) upon setting $g = \bar{g}$. The ghost lagrangian pertaining to this ansatz [1] is $\propto \bar{C}_\mu \mathcal{M}^\mu_\nu C^\nu$ with the kinetic operator $\mathcal{M} \propto S_{\text{gh}}^{(2)}$ given by

$$\mathcal{M}[g, \bar{g}]^\mu_\nu = \bar{g}^{\mu\rho} \bar{g}^{\sigma\lambda} \bar{D}_\lambda (g_{\rho\nu} D_\sigma + g_{\sigma\nu} D_\rho) - \bar{g}^{\rho\sigma} \bar{g}^{\mu\nu} \bar{D}_\lambda g_{\sigma\nu} D_\rho. \quad (3.2)$$

Next we insert the truncation ansatz into the flow equation (2.8). As a result, it boils down to a coupled system of ordinary differential equations for the two functions Z_{Nk} and $\bar{\lambda}_k$. They are obtained by performing a derivative expansion of the traces on the r.h.s. of (2.8) and then comparing the coefficients of the two invariants $\int \sqrt{g}$ and $\int \sqrt{g} R$ on both sides of the equation. After having performed the second functional derivatives in $\widehat{S}_k^{(2)}$ we may set $\bar{h} = 0$. This leads to $\widehat{S}_k^{(2)}[g, g]^{\mu\nu}{}_{\rho\sigma} = 2\kappa^2 Z_{Nk} [-K^{\mu\nu}{}_{\rho\sigma} D^2 + U^{\mu\nu}{}_{\rho\sigma}]$ where

$$K^{\mu\nu}{}_{\rho\sigma} = \frac{1}{4} [\delta_\rho^\mu \delta_\sigma^\nu + \delta_\sigma^\mu \delta_\rho^\nu - g^{\mu\nu} g_{\rho\sigma}] \quad (3.3)$$

and

$$U^{\mu\nu}{}_{\rho\sigma} = \frac{1}{4} [\delta_\rho^\mu \delta_\sigma^\nu + \delta_\sigma^\mu \delta_\rho^\nu - g^{\mu\nu} g_{\rho\sigma}] (R - 2\bar{\lambda}_k) + \frac{1}{2} [g^{\mu\nu} R_{\rho\sigma} + g_{\rho\sigma} R^{\mu\nu}] - \frac{1}{4} [\delta_\rho^\mu R^\nu{}_\sigma + \delta_\sigma^\mu R^\nu{}_\rho + \delta_\rho^\nu R^\mu{}_\sigma + \delta_\sigma^\nu R^\mu{}_\rho] - \frac{1}{2} [R^\nu{}_\rho{}^\mu{}_\sigma + R^\nu{}_\sigma{}^\mu{}_\rho]. \quad (3.4)$$

It is sufficient to retain the terms proportional to $\int \sqrt{g}$ and $\int \sqrt{g} R$ from the derivative expansion of the traces $\text{Tr exp}(\dots)$ appearing in (2.8). They can be obtained straightforwardly by means of standard heat kernel techniques. Since the calculation is similar to (but simpler than) the one in [1] we omit the details here. Also the subsequent s -integration can be performed easily in terms of Gamma functions. Equating the result to the l.h.s. of the flow equation,

$$\partial_t \widehat{S}_k[g, g] = 2\kappa^2 \int d^d x \sqrt{g} [-R(g) \partial_t Z_{Nk} + 2\partial_t (Z_{Nk} \bar{\lambda}_k)] \quad (3.5)$$

we obtain two differential equations of the form $\partial_k Z_{Nk} = \dots$ and $\partial_k (Z_{Nk} \bar{\lambda}_k) = \dots$. It is convenient to rewrite them in terms of the dimensionless Newton constant $g(k) \equiv k^{d-2} G_k \equiv k^{d-2} Z_{Nk}^{-1} \bar{G}$ and the dimensionless cosmological constant $\lambda(k) \equiv k^{-2} \bar{\lambda}_k$. This leads to the following system of equations:

$$\partial_t g = \beta_g(g, \lambda) \equiv [d - 2 + \eta_N] g \quad (3.6a)$$

$$\partial_t \lambda = \beta_\lambda(g, \lambda). \quad (3.6b)$$

The anomalous dimension $\eta_N \equiv -\partial_t \ln Z_{Nk}$ is given by

$$\eta_N = 8(4\pi)^{1-\frac{d}{2}} \left[\frac{d(7-5d)}{24} (1-2\lambda)^{\frac{d}{2}-m-2} - \frac{d+6}{6} \right] g \frac{\Gamma(m+2-\frac{d}{2})}{\Gamma(m+1)} \quad (3.7)$$

and the beta-function of λ reads

$$\beta_\lambda = -(2 - \eta_N)\lambda + 4(4\pi)^{1-\frac{d}{2}} \left[\frac{d(d+1)}{4} (1 - 2\lambda)^{\frac{d}{2}-m-1} - d \right] g \frac{\Gamma(m+1-\frac{d}{2})}{\Gamma(m+1)}. \quad (3.8)$$

The equations (15) with (3.7) and (3.8) are much simpler than their counterparts from the average action. In particular they do not contain terms (proportional to η_N) stemming from the differentiation of the \mathcal{Z} -factors (see ref. [1]), and they allow for an explicit evaluation of the threshold functions.

4. Universal quantities

We have calculated the non-trivial fixed point $(g^*, \lambda^*) \neq 0$ in $d = 4$, implied by the simultaneous vanishing of β_g and β_λ in (15), and the critical exponents θ', θ'' [3, 4] associated to the stability matrix for the non-trivial fixed point. The results are depicted in table 1 for various values of the cut-off parameter m . The most important result is that the proper time RG equation, too, predicts a non-gaussian fixed point. It exists for any value of m . The nonuniversal coordinates λ^* and g^* show a significant m -dependence, and it is impressive to see how this m -dependence cancels out in the product λ^*g^* which is universal in an exact calculation [4]. For every value of m , the stability matrix $(-\partial_i\beta_j)$, $i, j \in \{g, \lambda\}$, has a pair of complex conjugate eigenvalues $\theta' \pm i\theta''$. Remarkably, those critical exponents and the product λ^*g^* tend to constant values as m increases; they form a “plateau” as it is shown in figure 1. The universal quantities θ', θ'' , and λ^*g^* show excellent stability properties for this class of regulators, and these values are in complete agreement with the values found in the framework of the exact flow equation for the effective average action for gravity [3, 4]. In fact, the differences between the proper time and the average action RG equations are of the same order of magnitude as the residual scale dependence which is present in each of the two approaches and which is due to the truncated theory space.




Figure 1: The universal quantities θ', θ'' , and λ^*g^* as a function of the regulator parameter m .

m	g^*	λ^*	λ^*g^*	θ'	θ''
3/2	0.763	0.192	0.147	2.000	1.658
2	1.166	0.118	0.138	1.834	1.230
3	1.890	0.066	0.125	1.769	1.081
4	2.589	0.046	0.119	1.750	1.001
5	3.281	0.035	0.115	1.742	0.959
6	3.970	0.028	0.113	1.737	0.934
10	6.718	0.016	0.108	1.729	0.886
40	27.271	0.0038	0.103	1.722	0.840

Table 1: Fixed point values and critical exponents for various values of the cutoff parameter m . Note the plateau behavior of the universal quantities θ' , θ'' , and λ^*g^* as the value of m increases.

5. Conclusions

We have presented a nonperturbative flow equation for gravity which employs a smooth cutoff function in the proper time integration. The specific cutoff function employed in this paper is known to give rise to particularly accurate critical exponents in several scalar theories. The main result of our analysis is to have shown that the non-gaussian fixed point exists already in a simple improved 1-loop calculation, an approach less sophisticated than that of the exact RG equation.

We have then shown that the universal quantities determined by the non-gaussian fixed point are in very good agreement with the results from the average action [3, 4, 5], and in particular, that they show a very weak dependence on the regulator. Our result can be taken as a further, conceptually different indication that the fixed point is rather “robust” and should survive also in a more general truncation.

References

- [1] M. Reuter, *Nonperturbative evolution equation for quantum gravity*, *Phys. Rev. D* **57** (1998) 971 [[hep-th/9605030](#)].
- [2] D. Dou and R. Percacci, *The running gravitational couplings*, *Class. and Quant. Grav.* **15** (1998) 3449 [[hep-th/9707239](#)].
- [3] O. Lauscher and M. Reuter, *Ultraviolet fixed point and generalized flow equation of quantum gravity*, *Phys. Rev. D* **65** (2002) 025013 [[hep-th/0108040](#)].
- [4] M. Reuter and F. Saueressig, *Renormalization group flow of quantum gravity in the Einstein-Hilbert truncation*, *Phys. Rev. D* **65** (2002) 065016 [[hep-th/0110054](#)].
- [5] O. Lauscher and M. Reuter, *Is quantum Einstein gravity nonperturbatively renormalizable?*, *Class. and Quant. Grav.* **19** (2002) 483 [[hep-th/0110021](#)]; *Flow equation of quantum Einstein gravity in a higher-derivative truncation*, [hep-th/0205062](#); *Towards nonperturbative renormalizability of quantum Einstein gravity*, *Int. J. Mod. Phys. A* **17** (2002) 993 [[hep-th/0112089](#)].
- [6] W. Souma, *Non-trivial ultraviolet fixed point in quantum gravity*, *Prog. Theor. Phys.* **102** (1999) 181 [[hep-th/9907027](#)].

- [7] R. Percacci and D. Perini, *Asymptotic safety of gravity coupled to matter*, *Phys. Rev. D* **68** (2003) 044018 [[hep-th/0304222](#)]; *Constraints on matter from asymptotic safety*, *Phys. Rev. D* **67** (2003) 081503 [[hep-th/0207033](#)];
J. Polonyi and E. Regos, *Casimir effect and running Newton constant*, [hep-th/0404185](#).
- [8] M. Reuter and F. Saueressig, *A class of nonlocal truncations in quantum Einstein gravity and its renormalization group behavior*, *Phys. Rev. D* **66** (2002) 125001 [[hep-th/0206145](#)].
- [9] D.F. Litim, *Fixed points of quantum gravity*, *Phys. Rev. Lett.* **92** (2004) 201301 [[hep-th/0312114](#)].
- [10] A. Bonanno and M. Reuter, *Quantum gravity effects near the null black hole singularity*, *Phys. Rev. D* **60** (1999) 084011 [[gr-qc/9811026](#)]; *Renormalization group improved black hole spacetimes*, *Phys. Rev. D* **62** (2000) 043008 [[hep-th/0002196](#)].
- [11] A. Bonanno and M. Reuter, *Cosmology of the planck era from a renormalization group for quantum gravity*, *Phys. Rev. D* **65** (2002) 043508 [[hep-th/0106133](#)].
- [12] A. Bonanno and M. Reuter, *Cosmology with self-adjusting vacuum energy density from a renormalization group fixed point*, *Phys. Lett. B* **527** (2002) 9 [[astro-ph/0106468](#)];
Cosmological perturbations in renormalization group derived cosmologies, *Int. J. Mod. Phys. D* **13** (2004) 107–122 [[astro-ph/0210472](#)].
- [13] E. Bentivegna, A. Bonanno and M. Reuter, *Confronting the IR fixed point cosmology with high redshift supernova data*, *JCAP* **01** (2004) 001 [[astro-ph/0303150](#)].
- [14] A. Bonanno, G. Esposito and C. Rubano, *A class of renormalization group invariant scalar field cosmologies*, *Gen. Rel. Grav.* **35** (2003) 1899 [[hep-th/0303154](#)]; *Arnowitt-Deser-Misner gravity with variable g and λ and fixed point cosmologies from the renormalization group*, *Class. and Quant. Grav.* **21** (2004) 5005 [[gr-qc/0403115](#)].
- [15] M. Reuter and H. Weyer, *Renormalization group improved gravitational actions: a brans-dicke approach*, *Phys. Rev. D* **69** (2004) 104022 [[hep-th/0311196](#)].
- [16] M. Reuter and H. Weyer, *Running newton constant, improved gravitational actions and galaxy rotation curves*, *Phys. Rev. D* **70** (2004) 124028 [[hep-th/0410117](#)]; *Quantum gravity at astrophysical distances?*, *JCAP* **12** (2004) 001 [[hep-th/0410119](#)].
- [17] S. Weinberg, *General relativity, an Einstein centenary survey*, S.W. Hawking, W. Israel eds., Cambridge University Press, Cambridge 1979.
- [18] C. Wetterich, *Exact evolution equation for the effective potential*, *Phys. Lett. B* **301** (1993) 90.
- [19] J. Berges, N. Tetradis and C. Wetterich, *Non-perturbative renormalization flow in quantum field theory and statistical physics*, *Phys. Rept.* **363** (2002) 223 [[hep-ph/0005122](#)];
C. Wetterich, *Effective average action in statistical physics and quantum field theory*, *Int. J. Mod. Phys. A* **16** (2001) 1951 [[hep-ph/0101178](#)].
- [20] M. Reuter and C. Wetterich, *Effective average action for gauge theories and exact evolution equations*, *Nucl. Phys. B* **417** (1994) 181;
Exact evolution equation for scalar electrodynamics, *Nucl. Phys. B* **427** (1994) 291; *Average action for the Higgs model with abelian gauge symmetry*, *Nucl. Phys. B* **391** (1993) 147;
M. Reuter, *Effective average action of Chern-Simons field theory*, *Phys. Rev. D* **53** (1996) 4430 [[hep-th/9511128](#)]; *Renormalization of the topological charge in Yang-Mills theory*, *Mod. Phys. Lett. A* **12** (1997) 2777 [[hep-th/9604124](#)].

- [21] M. Niedermaier, *Dimensionally reduced gravity theories are asymptotically safe*, *Nucl. Phys.* **B 673** (2003) 131 [[hep-th/0304117](#)];
M. Niedermaier, *On the renormalization of truncated quantum Einstein gravity*, *J. High Energy Phys.* **12** (2002) 066 [[hep-th/0207143](#)];
P. Forgacs and M. Niedermaier, *A fixed point for truncated quantum Einstein gravity*, [hep-th/0207028](#).
- [22] J. Ambjørn, J. Jurkiewicz and R. Loll, *Emergence of a 4D world from causal quantum gravity*, *Phys. Rev. Lett.* **93** (2004) 131301 [[hep-th/0404156](#)].
- [23] A. Dasgupta and R. Loll, *A proper-time cure for the conformal sickness in quantum gravity*, *Nucl. Phys.* **B 606** (2001) 357 [[hep-th/0103186](#)];
J. Ambjørn, J. Jurkiewicz and R. Loll, *Dynamically triangulating lorentzian quantum gravity*, *Nucl. Phys.* **B 610** (2001) 347 [[hep-th/0105267](#)];
J. Ambjørn, J. Jurkiewicz and R. Loll, *Computer simulations of 3D lorentzian quantum gravity*, *Nucl. Phys.* **94** (Proc. Suppl.) (2001) 689 [[hep-lat/0011055](#)].
- [24] S.-B. Liao, *On connection between momentum cutoff and the proper time regularizations*, *Phys. Rev.* **D 53** (1996) 2020 [[hep-th/9501124](#)];
S.-B. Liao, *Operator cutoff regularization and renormalization group in Yang-Mills theory*, *Phys. Rev.* **D 56** (1997) 5008 [[hep-th/9511046](#)];
B.J. Schaefer, O. Bohr and J. Wambach, *Finite-temperature gluon condensate with renormalization group flow equations*, *Phys. Rev.* **D 65** (2002) 105008 [[hep-th/0112087](#)];
B.-J. Schaefer and H.-J. Pirner, *The equation of state of quarks and mesons in a renormalization group flow picture*, *Nucl. Phys.* **A 660** (1999) 439 [[nucl-th/9903003](#)];
Application of the heat-kernel method to the constituent quark model at finite temperature, *Nucl. Phys.* **A 627** (1997) 481 [[hep-ph/9706258](#)];
J. Meyer, G. Papp, H.J. Pirner and T. Kunihiro, *Renormalization group flow equation at finite density*, *Phys. Rev.* **D 61** (2000) 035202 [[nucl-th/9908019](#)];
G. Papp, B.J. Schaefer, H.J. Pirner and J. Wambach, *On the convergence of the expansion of renormalization group flow equation*, *Phys. Rev.* **D 61** (2000) 096002 [[hep-ph/9909246](#)].
- [25] A. Bonanno and D. Zappala, *Towards an accurate determination of the critical exponents with the renormalization group flow equations*, *Phys. Lett.* **B 504** (2001) 181 [[hep-th/0010095](#)].
- [26] D.F. Litim and J.M. Pawłowski, *Predictive power of renormalisation group flows: a comparison*, *Phys. Lett.* **B 516** (2001) 197 [[hep-th/0107020](#)]; *Perturbation theory and renormalisation group equations*, *Phys. Rev.* **D 65** (2002) 081701 [[hep-th/0111191](#)];
Completeness and consistency of renormalisation group flows, *Phys. Rev.* **D 66** (2002) 025030 [[hep-th/0202188](#)]; *Wilsonian flows and background fields*, *Phys. Lett.* **B 546** (2002) 279 [[hep-th/0208216](#)];
D.F. Litim and L. Vergara, *Subleading critical exponents from the renormalisation group*, *Phys. Lett.* **B 581** (2004) 263 [[hep-th/0310101](#)].
- [27] M. Mazza and D. Zappala, *Proper time regulator and renormalization group flow*, *Phys. Rev.* **D 64** (2001) 105013 [[hep-th/0106230](#)];
A. Bonanno and G. Lacagnina, *Spontaneous symmetry breaking and proper-time flow equations*, *Nucl. Phys.* **B 693** (2004) 36 [[hep-th/0403176](#)].
- [28] S.L. Adler, *Einstein gravity as a symmetry breaking effect in quantum field theory*, *Rev. Mod. Phys.* **54** (1982) 729.