

PERSPECTIVE • OPEN ACCESS


Baseline data on forest loss and associated uncertainty: advances in national forest monitoring

To cite this article: Johanne Pelletier and Scott J Goetz 2015 *Environ. Res. Lett.* **10** 021001

View the [article online](#) for updates and enhancements.

You may also like

- [Using spatial statistics to identify emerging hot spots of forest loss](#)
Nancy L Harris, Elizabeth Goldman, Christopher Gabris et al.
- [Using payment for ecosystem services to meet national reforestation commitments: impacts of 20+ years of forestry incentives in Guatemala](#)
Evan Patrick, Van Butsic and Matthew D Potts
- [Measurement and monitoring needs, capabilities and potential for addressing reduced emissions from deforestation and forest degradation under REDD+](#)
Scott J Goetz, Matthew Hansen, Richard A Houghton et al.



The Breath Biopsy® Guide
Fourth edition

FREE

DOWNLOAD THE FREE E-BOOK

BREATH BIOPSY

OWLSTONE MEDICAL

Environmental Research Letters



PERSPECTIVE

Baseline data on forest loss and associated uncertainty: advances in national forest monitoring

OPEN ACCESS

RECEIVED

16 December 2014

REVISED

7 January 2015

ACCEPTED FOR PUBLICATION

9 January 2015

PUBLISHED

4 February 2015

Johanne Pelletier and Scott J Goetz

The Woods Hole Research Center, 149 Woods Hole Rd, Falmouth, MA 02540-1644, USA

E-mail: johannepelletier@gmail.com

Keywords: REDD+, national forest monitoring, forest cover change

Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Abstract

Countries participating in climate change mitigation via the United Nations Framework Convention on Climate Change reducing emissions from deforestation and forest degradation mechanism are required to establish national forest monitoring systems. The design of national forest monitoring system includes provision of transparent, consistent and accurate estimates of emissions and removals from forests, while also taking into account national circumstances and capabilities. One key component of these systems lies in satellite remote sensing approaches and techniques to determine baseline data on forest loss against which future rates of change can be evaluated. Advances in approaches meeting these criteria for measuring, reporting and verification purposes are therefore of tremendous interest. A robust example advancing such approaches, focused on Peru, is provided in the recent paper of Potapov *et al* (2014 *Environ. Res. Lett.* **9** 124012).

Many developing countries are engaged in preparation for reducing emissions from deforestation and forest degradation (REDD+) under the United Nations Framework Convention on Climate Change (UNFCCC). An indication of the level of engagement is the fact that some 47 countries have now joined the readiness fund of the World Bank's forest carbon partnership facility (FCPF), a fund directed to preparing tropical and sub-tropical countries for REDD+. Readiness for REDD+ requires participating countries develop a robust national forest monitoring systems (NFM) that allow for transparent, consistent over time and accurate estimation of greenhouse gas (GHG) emissions and removals from forests (UNFCCC 2009, 2013). NFM require a combination of remote sensing and ground-based forest carbon inventory approaches (UNFCCC 2009, Goetz *et al* 2014) and following the guidance and guidelines provided by Intergovernmental Panel on Climate Change for estimating carbon fluxes from forests (UNFCCC 2013). The UNFCCC REDD+ decisions also specify a step-wise approach to monitoring improvements towards performance-based payments (UNFCCC 2010). Thus, the preparation of countries for measuring and reporting GHG emissions based on forest cover change (activity data) and carbon density (emission factors) is rapidly advancing. However, as

recently as 2010, only 4 of 99 developing countries were approaching a state of effective national forest monitoring, with many countries having large capacity gaps, particularly in tropical Africa (Romijn *et al* 2012).

For measuring activity data, there is general consensus that satellite remote sensing is the most practical way to establish baseline deforestation rates against which future rates of change can be monitored (Achard *et al* 2010, Baker *et al* 2010). Many developing countries lack the resources and expertise to use remote sensing to evaluate forest extent and change, and some do not have national land-cover map available, so widespread technical capacity building efforts are needed. Several countries have 'snapshot' land-cover maps for one or two points in time, typically years apart, but they fail to capture annual deforestation events and associated land-use/cover dynamics (Pelletier *et al* 2013). Moreover, the methodologies used to create such maps often lack technical details necessary for adequately assessing map accuracy. When good practices for estimating uncertainty of changes in forest area are applied, the variability of the estimates have a major impact on associated GHG emissions estimates (Olofsson *et al* 2013). Good practice change assessment should include estimates of the accuracy of change, change area adjusted for the classification error, confidence intervals associated with the

accuracy, and error statistics on the area of change (Olofsson *et al* 2014). These good practices allow identifying and targeting the sources of uncertainty, as well as prioritization of capacity advancement in a continuous manner to advance remote sensing approaches and techniques.

In this sense, the work of Potapov *et al* (2014) is important for advancing operational REDD+ measurement and reporting because it responds to the very criteria that are desirable for accurately estimating emissions from deforestation as part of NFM. Potapov *et al* establish a statistically robust baseline on the rates and extent of forest loss for the humid tropical biome of Peru. In doing so they provide a well-documented, repeatable change detection approach that is consistent over time, and can be extended forward and backward in time for forest change monitoring. They provide an example of operational national REDD+ forest monitoring that is well-documented and transparent, thereby establishing a reproducible approach that other countries might well emulate. Their methodology for assessing accuracy, based on samples of high spatial resolution imagery for estimating total forest and change area statistics, also provides a source of validation for their 'wall to wall' full coverage maps. Their work is a national-scale refinement and implementation of the global approach recently advanced by Hansen *et al* (2013).

The approach of Potapov *et al* (2014) also advances the attribution of forest cover change by separating natural from anthropogenic disturbances, an important criteria for addressing the drivers of change and establishing payment for performance under REDD+ (Goetz *et al* 2014). In the case of Peru, they find anthropogenic disturbance accounts for over 92% of gross forest loss. Their approach further attributes forest cover loss by year of change and forest type, which establishes a basis for potentially assessing carbon removals of GHG from the atmosphere via vegetation recovery and regrowth. The application of this multi-step approach establishes a firm basis for assessing activity data for REDD+, as well as potentially providing a means to maintain consistency with national GHG inventories in the broader land use, land use change and forestry sector. Establishing these baseline data sets for ongoing forest change can also serve as input for establishing forest reference emission levels and forest reference levels.

Broad-scale efforts are underway to advance NFM in developing countries, with significant technical and financial support dedicated to this in REDD+ participating countries (e.g. UN-REDD¹ and FCPF²). The design phase of these efforts includes the

selection of appropriate remote sensing approaches and techniques that are also cost-efficient and tailored to national circumstances and capabilities (De Sy *et al* 2012, GFOI 2013). The work of Potapov *et al* (2014) provides an advance in this regard as well since they established meaningful collaboration and real partnership between Peru's Ministry of Environment (MINAM) and the University of Maryland, thereby ensuring sustainable development of capabilities for national scale forest monitoring and response to the drivers of deforestation. The suitability of their approach is further aligned with several important criteria for REDD+ measuring, reporting and verification, including data continuity, cost and access, as well as data processing repeatability across space and time. This collaborative effort demonstrates a clear advance in the implementation of technological capabilities and the development of a robust national forest monitoring system. The work of Potapov *et al* (2014) clearly shows there is no current technological limit to robust measurement, monitoring, reporting and verification of deforestation activities. Their work thus paves the way to improved emissions estimates from forest loss and associated efforts to mitigate climate change under REDD+.

Acknowledgments

We acknowledge support from the NASA Applied Sciences Program (SERVIR), Grant Number: NNX12AL27G.

References

- Achard F, Stibig H-J, Eva H D, Lindquist E J, Bouvet A, Arino O and Mayaux P 2010 Estimating tropical deforestation from earth observation data *Carbon Manage.* **1** 271–87
- Baker D J *et al* 2010 Achieving forest carbon information with higher certainty: a five-part plan *Environ. Sci. Policy* **13** 249–60
- De Sy V, Herold M, Achard F, Asner G P, Held A, Kellndorfer J and Verbesselt J 2012 Synergies of multiple remote sensing data sources for REDD+ monitoring *Curr. Opin. Environ. Sustainability* **4** 1–11
- GFOI 2013 *Integrating Remote-Sensing and Ground-Based Observations for Estimation of Emissions and Removals of Greenhouse Gases in Forests: Methods and Guidance from the Global Forest Observations Initiative* (Geneva, Switzerland: Group on Earth Observations) 2014
- Goetz S J, Hansen M, Houghton R A, Walker W, Laporte N and Busch J 2014 *Measurement and Monitoring for REDD+: The Needs, Current Technological Capabilities and Future Potential, Why Forests? Why Now?* ed F Seymour and J Busch (Washington, DC: Center For Global Development)
- Hansen M C *et al* 2013 High-resolution global maps of 21st-century forest cover change *Science* **342** 850–3
- Olofsson P, Foody G M, Herold M, Stehman S V, Woodcock C E and Wulder M A 2014 Good practices for estimating area and assessing accuracy of land change *Remote Sens. Environ.* **148** 42–57
- Olofsson P, Foody G M, Stehman S V and Woodcock C E 2013 Making better use of accuracy data in land change studies: estimating accuracy and area and quantifying uncertainty using stratified estimation *Remote Sens. Environ.* **129** 122–31

¹ The UN-REDD programme supports national readiness in 56 partner countries in Africa, Asia and Latin America, with a financial contribution of totaling US\$195.7 million in June 2014. Available at: <http://www.un-redd.org/>

² The FCPF of the World Bank's support to countries via the readiness fund sums up to US\$360 million. Available at: <https://www.forestcarbonpartnership.org/>

- Pelletier J, Martin D and Potvin C 2013 REDD+ CO₂ flux estimation and reporting for early actions: dealing with uncertainty *Environ. Res. Lett.* **8** 034009
- Potapov P V *et al* 2014 National satellite-based humid tropical forest change assessment in Peru in support of REDD+ implementation *Environ. Res. Lett.* **9** 124012
- Romijn E, Herold M, Kooistra L, Murdiyarso D and Verchot L 2012 Assessing capacities of non-annex I countries for national forest monitoring in the context of REDD+ *Environ. Sci. Policy* **19–20** 33–48
- UNFCCC 2009 Methodological guidance for activities relating to reducing emissions from deforestation and forest degradation and the role of conservation, sustainable management of forests and enhancement of forest carbon stocks in developing countries Decision 4/CP.15 United Framework Convention on Climate Change FCCC/CP/2009/11/Add.1
- UNFCCC 2010 The Cancun agreements: Outcome of the work of the Ad Hoc working group on long-term cooperative action under the convention. United framework convention on climate change, Decision 1/CP.16 United Framework Convention on Climate Change FCCC/CP/2010/7/Add.1
- UNFCCC 2013 Modalities for national forest monitoring systems. Warsaw framework for REDD-plus Decision 11/CP.19 United Framework Convention on Climate Change FCCC/CP/2013/10/Add.1