

PAPER

Superweak momentum transfer near optical vortices

To cite this article: Stephen M Barnett and M V Berry 2013 *J. Opt.* **15** 125701

View the [article online](#) for updates and enhancements.

You may also like

- [Analysis of quality factor of quartz-crystal tuning forks using L-shaped bar model with torsion spring](#)
Hideaki Itoh and Keisuke Sugiura
- [Modeling of integrated shape memory alloy and Macro-Fiber Composite actuated trailing edge](#)
Aghna Mukherjee, Shaikh Faruque Ali and A Arockiarajan
- [Improving energy harvesting by nonlinear dual-beam energy harvester with an annular potential energy function](#)
Nan Shao, Hongxin Yang, Zhen Huang et al.

Superweak momentum transfer near optical vortices

Stephen M Barnett¹ and M V Berry²

¹ School of Physics and Astronomy, SUPA, University of Glasgow, Glasgow G12 8QQ, UK

² H H Wills Physics Laboratory, Tyndall Avenue, Bristol BS8 1TL, UK

E-mail: stephen.barnett@glasgow.ac.uk and asymptotico@bristol.ac.uk

Received 27 August 2013, accepted for publication 13 September 2013

Published 18 October 2013

Online at stacks.iop.org/JOpt/15/125701

Abstract

Near a vortex in a monochromatic light beam, the length of the local wavevector (phase gradient) can exceed the wavenumber in any of the plane waves in the superposition representing the beam. One way to detect these ‘superweak’ momenta could be by ‘superkicks’ imparted to a small particle located near the vortex, by absorbing individual large-momentum photons from the beam. A model for this process is a two-level atom with a transition resonant with the light beam. A semiclassical analysis shows that the momentum distribution of the atom is shifted by interaction with the vortex beam, by amounts that can almost reach the target superkicks and are usually greater than the momenta in the plane waves comprising the beam.

Keywords: weak measurement, quantum optics, singularities, semiclassical

1. Introduction

A fundamental feature of phase singularities [1–4] of monochromatic fields representing waves in three-dimensional space \mathbf{r} is that they are optical vortex lines, near which the phase varies on scales smaller than the wavelength λ . It follows that the length of the local wavevector $\mathbf{k}(\mathbf{r})$ —the phase gradient—exceeds the free-space wavenumber $k_0 = 2\pi/\lambda$. Such rapid variations are now recognized as ‘superoscillations’ [5, 6], where the ‘weak value’ [7] of momentum $\hbar\mathbf{k}(\mathbf{r})$ exceeds the free-space momentum $\hbar k_0$. The question we address here, in the spirit of quantum weak measurement theory [8–10], is whether such large local momenta could be imparted as ‘superkicks’ to test particles (‘atoms’) in the field. This situation appears paradoxical, because the momentum $\hbar\mathbf{k}(\mathbf{r})$ imparted to the absorbing particle exceeds that of any single photon in the field, raising concerns about global momentum conservation. Our resolution follows from considering the quantum mechanics of the motion of an absorbing atom. We need to account for the wave nature both of the electromagnetic field and the atom used to probe it.

A local model for the field strength near a vortex line of strength m on the z axis of a linearly polarized paraxial

wave, is

$$\begin{aligned} E_m(\mathbf{r}, t) &= E_m(\mathbf{r}) \exp(-i\omega t), & \omega &= ck_0, \\ E_m(\mathbf{r}) &= Nr^m \exp\{i(m\phi + k_0 z)\} \\ &= N(x + iy)^m \exp(ik_0 z), \end{aligned} \quad (1.1)$$

in which here and hereafter N denotes a generic multiplier or normalization constant. The corresponding local wavevector is

$$\mathbf{k}(\mathbf{r}) = \nabla \arg E_m(\mathbf{r}, t) = \frac{m}{r} \mathbf{e}_\phi + k_0 \mathbf{e}_z. \quad (1.2)$$

Its transverse part exceeds k_0 inside the cylinder $k_0 r < m$. The factor $(x + iy)^m$ describes any locally symmetric vortex on strength m ; it is also a small- r approximation to the exact Bessel beam solution [11] of the Helmholtz equation:

$$\begin{aligned} E_{m,\text{Bessel}}(\mathbf{r}) &= NJ_m(Kr) \exp \\ &\times \left\{ i \left(m\phi + z\sqrt{k_0^2 - K^2 - \omega t} \right) \right\}. \end{aligned} \quad (1.3)$$

Paraxiality corresponds to $K \ll k_0$.

In section 2 we derive the approximate Hamiltonian for a model quantum detector in the form of a two-level atom placed in the field, and show that when the atom gets excited its position wavefunction inherits the form of the

optical field. In section 3 we calculate the corresponding atomic momentum distribution, incorporating the momentum uncertainty implied by localizing the atom near the vortex. In particular, we calculate the average momentum, and elucidate the conditions under which this can exceed the momentum of free-space optical photons.

This study, developing an idea envisaged earlier [7, 12], complements and extends existing explorations [13, 14] of possible quantum effects associated with the cores of optical vortices.

2. Atom Hamiltonian

As a probe for the optical field, we consider a model two-level atom of mass M , with electronic ground and excited states $|g\rangle$ and $|e\rangle$ and corresponding energies ε_g and ε_e , coupled by electric dipole interaction. It suffices to work within the semiclassical approximation, in which the atom is treated quantum mechanically and the field is described classically [15, 16]. We can describe the coupled atom–field system by the state $|\Psi_0(t)\rangle$, driven by the Hamiltonian

$$\hat{H}_0 = \frac{\hat{p}^2}{2M} + \varepsilon_e |e\rangle\langle e| + \varepsilon_g |g\rangle\langle g| - \hat{\mu}[E_m(\mathbf{r}, t) + E_m^*(\mathbf{r}, t)], \quad (2.1)$$

where \hat{p}^2 is the square of the total atomic momentum and $\hat{\mu}$ is the electric dipole operator. Neither of the electronic states has an intrinsic dipole moment, so the dipole operator has only off-diagonal matrix elements in this basis and can be written in the form

$$\hat{\mu} = \mu(|g\rangle\langle e| + |e\rangle\langle g|). \quad (2.2)$$

We immediately eliminate some of the fast oscillations by transforming to a new state $|\Psi(t)\rangle$ (equivalent to the interaction picture):

$$|\Psi_0(t)\rangle = \exp\left\{-\frac{i}{\hbar}t(\varepsilon_g |g\rangle\langle g| + \varepsilon_e |e\rangle\langle e|)\right\} |\Psi(t)\rangle. \quad (2.3)$$

Thus the Schrödinger equation is

$$i\hbar\partial_t|\Psi(t)\rangle = \hat{H}|\Psi(t)\rangle \quad (2.4)$$

with the new Hamiltonian

$$\begin{aligned} \hat{H} = & \frac{\hat{p}^2}{2M} - \mu \left(|g\rangle\langle e| \exp\left\{-\frac{i}{\hbar}(\varepsilon_e - \varepsilon_g)t\right\} \right. \\ & + |e\rangle\langle g| \exp\left\{\frac{i}{\hbar}(\varepsilon_e - \varepsilon_g)t\right\} \left. \right) \\ & \times (E_m(\mathbf{r}, t) + E_m^*(\mathbf{r}, t)). \end{aligned} \quad (2.5)$$

(Here we have used the fact that $|g\rangle$ and $|e\rangle$ are orthogonal, and the consequence that the operators $|e\rangle\langle e|$ and $|g\rangle\langle g|$ commute.)

We can make three further simplifications. First, we can make the atom sufficiently massive and slowly-moving such that the first (kinetic) term has little effect during the short times will be of interest to us and can be neglected. Second, we choose exact resonance, so that $\varepsilon_e - \varepsilon_g = \hbar\omega$. Third, we note that from the form of the electric field amplitude (1.1),

\hat{H} contains time-independent terms and also terms rotating at 2ω : twice the optical frequency. The effect of these rapidly rotating terms will tend to average out over relevant time scales, and discarding them leads to the final time-independent ‘rotating wave’ Hamiltonian [16]:

$$\hat{H}_{\text{RW}} = -\mu(|g\rangle\langle e|E_m^*(\mathbf{r}) + |e\rangle\langle g|E_m(\mathbf{r})). \quad (2.6)$$

We can now analyse the effect on the atomic motion of absorbing a quantum of energy near to the vortex core, where the local optical momentum $|\mathbf{k}(\mathbf{r})| > k_0$. In doing so it is essential to include a quantum description of the motional state of the atom as well as its electronic state.

Let the initial state of the atom be

$$\langle \mathbf{r} | \Psi(0) \rangle = |g\rangle \psi_{\text{init}}(\mathbf{r}), \quad (2.7)$$

corresponding to the atom being in its electronic ground state and in the motional state $\psi_{\text{init}}(\mathbf{r})$. After a short time this state will evolve to a superposition of the ground and excited states. We can describe this using first-order perturbation theory, because we are interested only in the effects of a single absorption event, with the result

$$\begin{aligned} \langle \mathbf{r} | \Psi(t) \rangle & \approx \left(1 - i \frac{\hat{H}_{\text{RW}} t}{\hbar} \right) \langle \mathbf{r} | \Psi(0) \rangle \\ & = |g\rangle \psi_{\text{init}}(\mathbf{r}) + i \frac{\mu t}{\hbar} E_m(\mathbf{r}) |e\rangle \psi_{\text{init}}(\mathbf{r}). \end{aligned} \quad (2.8)$$

If after this short time the atom has made a transition to the excited state (by absorbing a single quantum from the field) then its associated motional wavefunction will be

$$\psi_m(\mathbf{r}) = N(x + iy)^m \exp(ik_0 z) \psi_{\text{init}}(\mathbf{r}). \quad (2.9)$$

Clearly, the phase of the electric field has been imprinted on the motional wavefunction and this encodes the ‘kick’ given to the atom. Our focus will be on the transverse momentum, but we note the immediate obvious consequence of (2.9) that for $\psi_m(\mathbf{r})$ the average momentum in the z -direction exceeds by $\hbar k_0$ that for $\psi_{\text{init}}(\mathbf{r})$. This is the familiar atomic recoil associated with the conservation of linear momentum.

3. Momentum distribution

It follows from (2.9) that the momentum state of the excited atom is given in terms of the initial momentum state $\bar{\psi}_0(\mathbf{k})$ (Fourier transform of (2.9)) by

$$\bar{\psi}_m(\mathbf{k}) = N(\partial_{k_x} + i\partial_{k_y})^m \bar{\psi}_{\text{init}}(k_x, k_y, k_z - k_0). \quad (3.1)$$

An immediate consequence is that the final momentum state can contain only momenta that were present in the initial state: the only effect of the interaction with the light is a redistribution of momenta that the atom already possessed as a result of its localization near the vortex. In particular, if the initial momentum state has bounded support in \mathbf{k} , the final state has the same bounded support.

Nevertheless, it is possible for the atom to acquire momenta larger than k_0 . We explore this for an initial (unexcited) atomic wavefunction located close to $\mathbf{r} = (x_0, 0, 0)$,




Figure 1. Momentum distribution (3.5) of excited atom in transverse planes, for (a) $x_0 = \sigma$, $m = 1$, (b) $x_0 = \sigma$, $m = 3$, (c) $x_0 = 2\sigma$, $m = 3$, (d) $x_0 = 5\sigma$, $m = 4$.

in the form of the Gaussian

$$\psi_{\text{init}}(\mathbf{r}) = N \exp\left(-\frac{(x-x_0)^2 + y^2 + z^2}{2\sigma^2}\right). \quad (3.2)$$

In order to localize the atom in the region $k_0 r < m$ where the light is superoscillatory, we restrict the position x_0 and the width σ to

$$x_0 < \frac{m}{k_0} = \frac{2\pi m}{\lambda}, \quad \sigma < m\lambda. \quad (3.3)$$

The corresponding initial momentum distribution is

$$\bar{\psi}_{\text{init}}(\mathbf{k}) = N \exp(-\frac{1}{2}\sigma^2(k_x^2 + k_y^2 + k_z^2) + ik_x x_0). \quad (3.4)$$

An easy calculation (iterating (3.1) for successive values of m) gives the final momentum probability distribution

$$|\bar{\psi}_m(\mathbf{k})|^2 = N^2 \exp(-(k_z - k_0)^2 \sigma^2) \times (k_x^2 + (k_y + x_0/\sigma^2)^2)^m \exp(-(k_x^2 + k_y^2)\sigma^2). \quad (3.5)$$

This shows that in addition to the obvious z momentum shift by k_0 , the distribution is skewed towards $+y$ (if $x_0 > 0$), as illustrated in figure 1 for several values of x_0/σ and m .

To quantify this y kick, we calculate the momentum expectation values for different vortex strengths m :

$$\langle \mathbf{k}_m \rangle = \frac{\int \int d\mathbf{k} \mathbf{k} |\bar{\psi}_m(\mathbf{k})|^2}{\int \int d\mathbf{k} |\bar{\psi}_m(\mathbf{k})|^2} \equiv k_m \mathbf{e}_y + k_0 \mathbf{e}_z. \quad (3.6)$$

The k_y shifts can be evaluated exactly in terms of Laguerre polynomials [17]:

$$k_m = \frac{x_0 L_{m-1}^{(1)}(-x_0^2/\sigma^2)}{\sigma^2 L_m(-x_0^2/\sigma^2)} = \frac{1}{2} \partial_{x_0} \log(L_m(-x_0^2/\sigma^2)). \quad (3.7)$$

The first few shifts are

$$\begin{aligned} k_1 &= \frac{x_0}{\sigma^2 + x_0^2}, & k_2 &= \frac{2x_0(2\sigma^2 + x_0^2)}{2\sigma^4 + 4x_0^2\sigma^2 + x_0^4}, \\ k_3 &= \frac{3x_0(6\sigma^4 + 6x_0^2\sigma^2 + x_0^4)}{6\sigma^6 + 18x_0^2\sigma^4 + 9x_0^4\sigma^2 + x_0^6}, & (3.8) \\ k_4 &= \frac{4x_0(24\sigma^6 + x_0^2(6\sigma^2 + x_0^2)^2)}{24\sigma^8 + 96x_0^2\sigma^6 + 72x_0^4\sigma^4 + 16x_0^6\sigma^2 + x_0^8}. \end{aligned}$$

These functions should be compared with the corresponding superkicks that we hope to detect in the optical field, given by (cf (1.2))

$$k_{m,\text{super}} = \frac{m}{x_0}. \quad (3.9)$$

Comparisons are shown in figure 2. The average momentum shifts acquired by the atoms reach their greatest value $k_{m\text{max}}$ when $x_0 = x_{0\text{max}} \sim \sigma$; the precise values are given in table 1.




Figure 2. Full curves: mean momentum shifts k_m in y direction (equations (3.6) and (3.7)); dashed curves: local optical momenta (target superkicks) $k_{m,\text{super}}$ (equation (3.9)); as functions of the atom position x_0 , for (a) $m = 1$, (b), $m = 2$, (c) $m = 3$, (d) $m = 4$.




Figure 3. Full curves: momentum distributions (3.4) of the excited atom along the symmetry line $k_x = 0$ for fixed k_z , for atom positions x_0 corresponding to the maximum average shifts in table 1, for (a) $m = 1$; (b) $m = 3$; (c) $m = 5$; (d) $m = 7$. Dashed curves: the corresponding initial atomic momentum distributions (3.4). (The curves are normalized to facilitate comparison.)

Table 1. Maximal momentum shifts for vortex strengths m .

m	1	2	3	4	5	6
$x_{0\text{max}}/\sigma$	1	0.939	0.900	0.874	0.854	0.840
$k_{m\text{max}}\sigma$	0.5	0.858	1.151	1.404	1.629	1.835

Figure 3 shows the momentum distributions along the symmetry line $k_x = 0$ for x_0 corresponding to these maximum shifts, illustrating how the shifts get bigger for larger m .

It is clear from figure 2 that the average momentum shifts tend to the superkick values (3.9) as x_0/σ increases but fall short for smaller x_0 . The shifts vanish as $x_0 \rightarrow 0$, because then the atomic wavefunction overlaps the other side of the vortex where $\mathbf{k}(\mathbf{r})$ is in the opposite direction. Notwithstanding the shortfall, in the superoscillatory region near the vortex where the atom shifts are maximal (table 1) these shifts exceed the momenta k_0 in the plane-wave superpositions comprising the field, by an amount

$$\frac{k_{m\text{max}}}{k_0} = \frac{(k_{m\text{max}}\sigma) \lambda}{2\pi \sigma} \sim \frac{(k_{m\text{max}}\sigma) \lambda}{2\pi x_0}, \quad (3.10)$$

which is greater than unity if the atom is close enough to the vortex.

We have calculated the momentum distribution of the atom on the assumption that a resonant transition has taken place. However, the probability that such a transition will occur is very small when the atom is near a vortex. From (2.9), the Fermi golden rule [16] gives the transition probability per unit time as

$$\begin{aligned} P &\propto \int d\mathbf{r} |\langle \mathbf{r}, e | \Psi(t) \rangle|^2 \\ &\propto N \int d\mathbf{r} |E_m(\mathbf{r})|^2 |\psi_{\text{init}}(\mathbf{r})|^2 \\ &\approx |E_m(x_0, 0, 0)|^2 = Nx_0^{2m} \end{aligned} \quad (3.11)$$

where we have assumed that ψ_{init} is localized near $(x_0, 0, 0)$ as in (3.2), and the field has the vortex form (2.1). This is consistent with the following picture [7, 12] for the time-averaged force on the atom (classical radiation pressure), resulting from the collective effect of many quantum transitions. The force, given by the probability per unit time multiplied by the momentum shift at each transition, is proportional to the large-momentum transfer (the superkick) weighted by the weak light intensity. Classically, this corresponds to the important distinction, emphasized earlier [7], between the local momentum $\hbar\mathbf{k}(\mathbf{r})$ and the local current

$$\text{Im}[E_m^*(\mathbf{r})\nabla E_m(\mathbf{r})] = |E_m(\mathbf{r})|^2 \mathbf{k}(\mathbf{r}). \quad (3.12)$$

4. Concluding remarks

It is important to consider how superkicks might be observed in an experiment. The natural way to proceed would be to trap a suitable atom and to cool it to its motional ground state. In this state the atomic motional wavefunction will be approximately Gaussian, the width of which will be determined by the experimentally controllable trapping potential. If the trap is switched off then the atomic wavepacket will spread but, for sufficiently short times, the atom will remain localized near to the trap position. Applying a short laser pulse carrying orbital angular momentum will induce a transition to the atomic excited state and with it the superkick of interest. This should be observable by examining the momentum probability distribution for the excited atom.

A number of improvements on this basic idea may make superweak momentum transfer more readily observable. The first is the use of a Bose–Einstein condensate rather than a single atom. Such condensates have the advantage that many atoms share the same motional state; a transition for a small proportion of these may be more readily observable than a low-probability transition for a single trapped atom. Moreover, the use of a two-photon Raman transition rather than a single-photon transition would make it possible to remove the large-momentum kick in the z direction, the existence of which might otherwise mask the azimuthal superkick we seek. Finally, using a magnetic trap may remove the requirement for turning off the trap before applying the electromagnetic field. If the field is tuned so as to flip the

electron spin in the trapped atom then the trapping potential will become repulsive and the trapped atom will be ejected from the trap. The superkick should then be visible as a preferred azimuthal direction of ejection from the trap. Assessing which of these ideas provides the greatest potential for observing superweak momentum transfer requires further details of the particular experimental arrangement and lies beyond the scope of this paper.

Finally, we note a curious feature of the classical mechanics of the test particles we have been considering, as represented by the time-averaged force on them, in circumstances where this is proportional to the Poynting vector [7, 18]. For the scalar model we have used here, the time-averaged force is

$$\mathbf{F}(\mathbf{r}) \propto |E_m(\mathbf{r})|^2 \mathbf{k}(\mathbf{r}), \quad (4.1)$$

in which the local wavevector momentum is multiplied by the wave intensity, in contrast with the bare wavevector as in the quantum kicks $\hbar\mathbf{k}(\mathbf{r})$. As is known [19, 20] and has been emphasized [21], this force has non-zero curl, so it is not derivable from a potential. In such ‘curl forces’ [22], the classical motion is Newtonian but not Hamiltonian or Lagrangian, so—for example—Noether’s theorem does not apply, and the link between symmetries and conservation laws is broken. It is hard to see how such non-Hamiltonian physics can be directly quantized, but our analysis reported here, in which classical radiation pressure (here associated with a curl force) is quantumly deconstructed into individual superkicks, points to a route where quantum effects might nevertheless be understood.

Acknowledgments

SMB thanks Erling Riis for advice on atom traps; his research is supported by the UK EPSRC and the Leverhulme Trust. MVB acknowledges generous hospitality of the University of Wrocław, where the first draft of this paper was written; his research is supported by the Leverhulme Trust.

Note added in proof. Professor K Bliokh has suggested that what we call superkicks might be related to experiments carried out on the transverse momenta of evanescent waves near an interface. See [23, 24], and discussion on p 14 of [25]. There are indeed similarities in that a momentum transfer in excess of the free-space momentum is indicated, but the proximity of a medium suggests that the apparent momentum imbalance might be made up by recoil of the medium. In contrast, the phenomenon we have treated could be demonstrated, at least in principle, arbitrarily far from a material interface, and a quantum treatment of the motion of the absorbing atom is necessary in order to account for momentum conservation.

References

- [1] Nye J F and Berry M V 1974 Dislocations in wave trains *Proc. R. Soc. A* **336** 165–90
- [2] Nye J F 1999 *Natural Focusing and Fine Structure of Light: Caustics and Wave Dislocations* (Bristol: Institute of Physics Publishing)
- [3] Dennis M R, O’Holleran K and Padgett M J 2009 Singular optics: optical vortices and polarization singularities *Prog. Opt.* **53** 293–363

- [4] Berry M V 1998 Much ado about nothing: optical dislocation lines (phase singularities, zeros, vortices...) *Proc. SPIE* **3487** 1–5
- [5] Berry M V 1994 *Faster than Fourier in Quantum Coherence and Reality; In Celebration of the 60th Birthday of Yakir Aharonov* ed J S Anandan and J L Safko (Singapore: World Scientific) pp 55–65
- [6] Berry M V 2013 A note on superoscillations associated with Bessel beams *J. Opt.* **15** 044006
- [7] Berry M V 2009 Optical currents *J. Opt. A* **11** 094001
- [8] Aharonov Y, Albert D Z and Vaidman L 1988 How the result of a measurement of a component of the spin of a spin 1/2 particle can turn out to be 100 *Phys. Rev. Lett.* **60** 1351–4
- [9] Aharonov Y and Rohrlich D 2005 *Quantum Paradoxes: Quantum Theory for the Perplexed* (Weinheim: Wiley–VCH)
- [10] Aharonov Y, Popescu S and Tollaksen J 2010 A time-symmetric formulation of quantum mechanics *Phys. Today* **63** 27–33
- [11] Durnin J 1987 Exact solutions for nondiffracting beams. I. The scalar theory *J. Opt. Soc. Am.* **4** 651–4
- [12] Berry M V 2013 Five momenta *Eur. J. Phys.* **44** 1337–48
- [13] Berry M V and Dennis M R 2004 Quantum cores of optical phase singularities *J. Opt. A: Pure Appl. Opt.* **6** S178–80
- [14] Barnett S M 2008 On the quantum core of an optical vortex *J. Mod. Opt.* **55** 2279–92
- [15] Loudon R 2000 *The Quantum Theory of Light* (Oxford: Oxford University Press)
- [16] Barnett S M and Radmore P M 1997 *Methods in Theoretical Quantum Optics* (Oxford: Oxford University Press)
- [17] DLMF 2010 *NIST Handbook of Mathematical Functions* (Cambridge: Cambridge University Press) <http://dlmf.nist.gov>
- [18] Bliokh K Y, Bekshaev A Y, Kofman A G and Nori F 2013 Photon trajectories, anomalous velocities and weak measurements: a classical interpretation *New J. Phys.* **15** 073022
- [19] Ashkin A and Gordon J P 1983 Stability of radiation-pressure particle traps: an optical Earnshaw theorem *Opt. Lett.* **8** 511–3
- [20] Gómez-Medina R, Nieto-Vesperinas M and Sáenz J J 2011 Nonconservative electric and magnetic optical forces on submicron dielectric particles *Phys. Rev. A* **83** 033825
- [21] Berry M V and Shukla P 2013 Physical curl forces: dipole dynamics near optical vortices *J. Phys. A: Math. Theor.* at press
- [22] Berry M V and Shukla P 2012 Classical dynamics with curl forces, and motion driven by time-dependent flux *J. Phys. A: Math. Theor.* **45** 305201
- [23] Huard S and Imbert C 1978 Mesure de l'impulsion échangée au cours de l'interaction onde évanescence-atome *Opt. Commun.* **24** 185–9
- [24] Huard S 1979 Exchanged momentum between moving atoms and a surface wave: theory and experiment *Can. J. Phys.* **57** 612–8
- [25] Bliokh K Y, Bekshaev A Y, Kofman A G and Nori F 2013 Photon trajectories, anomalous velocities and weak measurements: a classical interpretation *New J. Phys.* **15** 073022