

CHANDRA X-RAY SOURCES IN M101

W. D. PENCE, S. L. SNOWDEN,¹ AND K. MUKAI¹

NASA Goddard Space Flight Center, Laboratory for High Energy Astrophysics, Code 662, Greenbelt, MD 20771

AND

K. D. KUNTZ

Joint Center for Astrophysics, Department of Physics, University of Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, MD 21250

Received 2001 May 18; accepted 2001 July 5

ABSTRACT

A deep (98.2 ks) *Chandra* Cycle 1 observation has revealed a wealth of discrete X-ray sources as well as diffuse emission in the nearby face-on spiral galaxy M101. From this rich data set we have created a catalog of the 110 sources from the S3 chip detected with a significance of greater than 3σ . This detection threshold corresponds to a flux of $\sim 10^{-16}$ ergs cm $^{-2}$ s $^{-1}$ and a luminosity of $\sim 10^{36}$ ergs s $^{-1}$ for a distance to M101 of 7.2 Mpc. The sources display a distinct correlation with the spiral arms of M101 and include a variety of X-ray binaries, supersoft sources, supernova remnants, and other objects of which only ~ 27 are likely to be background sources. There are only a few sources in the interarm regions, and most of these have X-ray colors consistent with that of background active galactic nuclei. The derived $\log N - \log S$ relation for the sources in M101 (background subtracted) has a slope of -0.80 ± 0.05 over the range of 10^{36} – 10^{38} ergs s $^{-1}$. The nucleus is resolved into two nearly identical X-ray sources, each with a 0.5–2.0 keV flux of 4×10^{37} ergs s $^{-1}$. One of these sources coincides with the optical nucleus, and the other coincides with a cluster of stars 110 pc to the south. The field includes 54 optically identified supernova remnants (SNRs), of which 12 are detected by *Chandra*. Two of the SNR sources are variable and hence must be compact objects. In total, eight of the X-ray sources show evidence for short-term temporal variation during this observation. Two of these variable sources are now brighter than the *ROSAT* detection threshold, but they were not detected in the previous *ROSAT* observations taken in 1992 and 1996. There are also two variable sources previously seen with *ROSAT* that apparently have faded below the *Chandra* detection threshold. The brightest source in the field shows extreme long-term and short-term temporal variability. At its peak brightness it has a super-Eddington luminosity greater than 10^{39} ergs s $^{-1}$. There are 10 supersoft sources (SSSs) in the field, which can be divided into two distinct subclasses: the brighter class (three objects) has a luminosity of $\sim 10^{38}$ ergs s $^{-1}$ and a blackbody temperature of ~ 70 eV, whereas the other class (seven objects) is an order of magnitude fainter and has a blackbody temperature of only ~ 50 eV.

Subject headings: galaxies: individual (M101) — galaxies: spiral — X-rays: galaxies

On-line material: machine-readable table

1. INTRODUCTION

M101 is a nearby (7.2 Mpc; Stetson et al. 1998), face-on spiral galaxy that is ideal for studies of the X-ray source population in a galaxy similar to the Milky Way. Since M101 is viewed through a relatively small amount of obscuring foreground Galactic hydrogen gas ($N_{\text{H}} = 1.2 \times 10^{20}$ cm $^{-2}$; Stark et al. 1992), it is especially well suited for surveying soft X-ray sources, without the observational biases introduced by the thick absorption in the Galactic plane when studying X-ray sources in our own Galaxy. Although M101 has a slightly later morphological type (Scd vs. Sbc) and a significantly larger D_{25} isophotal diameter (60 kpc vs. 23 kpc) than the Milky Way (de Vaucouleurs & Pence 1978; de Vaucouleurs et al. 1991), one would expect that the properties of the X-ray source populations in these two galaxies should be quite similar.

M101 was previously observed with the *Einstein* X-ray observatory (McCammon & Sanders 1984; Trinchieri, Fabiano, & Romaine 1990) and later with *ROSAT* to study both the discrete sources (Wang, Immler, & Pietsch 1999) and the diffuse X-ray emission (Snowden & Pietsch 1995).

M101 was an obvious candidate for follow-up studies with the *Chandra* X-ray observatory, and this paper presents the basic data and analysis of the population of discrete X-ray sources detected in a Cycle 1 *Chandra* observation. This is one in a series of papers that analyze various aspects of this rich data set. A previous paper (Snowden et al. 2001) reinvestigated some of the hypernova remnant candidates in M101, and two other papers currently in preparation will study the brighter black hole X-ray binary candidates in more detail and analyze the diffuse X-ray emission in M101.

2. DATA PROCESSING

The present data set consists of a 98.2 ks exposure taken on 2000 March 26–27 with the *Chandra* observatory ACIS imaging spectrometer with the nucleus of M101 centered near the “aim point” of the S3 CCD chip. The analysis in this paper is limited to the sources detected within the 8.4×8.4 (17.6 \times 17.6 kpc) field of view of the S3 chip, which has much better angular resolution and extended sensitivity to lower X-ray energies than the other chips. This chip covers the central region of M101 out to a minimum distance of 6.3 kpc from the nucleus toward the northwest and a maximum distance of 14.9 kpc at the east corner of the chip. By comparison, the prominent spiral arms extend out

¹ Universities Space Research Association.

to about 14 kpc, and the $R_{2.5}$ isophotal radius (de Vaucouleurs et al. 1991) is 30.2 kpc, so the S3 chip covers about 50% and 11%, respectively, of these circular areas in M101.

The standard CIAO pipeline software system (“Rev 2” as of 2000 December) was used to clean and calibrate the data set. In addition to the standard data cleaning, we also excluded the data during a 64 s interval of unusually high background count rate, which reduced the exposure duration to 98,180 s. Finally, we excluded all photon events that had a nominal pulse-invariant (PI) channel energy greater than 8.0 keV where the S3 chip has virtually no sensitivity. The nominal lower energy cutoff in the data set is 0.125 keV.

3. SOURCE DETECTION

The CIAO *celldetect*² program was initially used with a threshold of 2.5σ to generate a preliminary candidate list of

² At the time that we received the *Chandra* data files the *celldetect* program appeared to produce the most reliable list of sources. Later, after the release CIAO v2.1, we repeated the source detection procedure using the *wavdetect* program but concluded that the final list of sources presented in Table 1 would not change significantly. The log N –log S analysis in § 4 was based on the results from *wavdetect*.

X-ray sources in the 0.125–8.0 keV band image. At this low threshold setting about 20% of the sources were clearly spurious, mainly located in the wings of brighter sources near the edges of the chip. These spurious sources were rejected by inspecting their positions on an adaptively smoothed image of the S3 chip field of view constructed from all the X-ray events in which sources with as few as 10 net counts are clearly visible. The final adopted set of sources has formal signal-to-noise ratios close to 3.0 or higher. Note that the detection threshold for faint point sources increases with distance from the “aim point” because of the decreased angular resolution, and as a result the faintest point sources (with $F_X \sim 10^{-16}$ ergs cm⁻² s⁻¹) are detectable only within a radius of about 2′ (4 kpc) from the nucleus of M101.

The positions of our final 110 *Chandra* X-ray sources in M101 are shown in Figures 1 and 2 and listed in Table 1. The source positions derived from the *Chandra* data are generally accurate to better than 1″. The mean difference between our X-ray positions and the astrometric optical positions of 12 previously identified supernova remnants (SNRs) catalogued by Matonick & Fesen (1997) is only $0'.47 \pm 0'.20$, about the size of one ACIS CCD pixel.

As can be seen in Figure 2, the X-ray sources are strongly correlated with the position of the optical spiral arms and




FIG. 1.—Adaptively smoothed *Chandra* image of the M101 field covered by the S3 chip showing the location of the 110 X-ray sources. The nuclear source is indicated by the square, to the right and slightly above center.




FIG. 2.—Location of the 110 *Chandra* X-ray sources plotted on top of the image of M101 from the digitized Palomar Sky Survey E plate. It can be seen that most of the sources are located within the spiral arm regions. The squares indicated the eight “interarm” sources that are discussed in the text. Some of the regions were shown in white simply to improve the contrast with the underlying image.

bright H II regions in M101. There are relatively few X-ray sources in the interarm regions, which indicates that most of the X-ray sources are associated with the young stellar population in M101. To quantify this association, we divided the area covered by the S3 chip into either “spiral arm” or “interarm” regions based on the appearance in the blue optical photograph of M101, with the result that 75% of the area of the chip was classified as covering spiral arm regions (also includes the nuclear region) and 25% as covering interarm regions. Using this spatial criteria, only eight of the X-ray sources (sources 5, 7, 8, 19, 29, 39, 64, and 97) were found to be located in the interarm regions and not associated with optical emission in M101. This gives a mean surface density of $1.9 \text{ sources arcmin}^{-2}$ in the spiral arms compared with only $0.45 \text{ sources arcmin}^{-2}$ in the interarm regions (both before background subtraction).

We calculated the expected number of background sources in our field based on the point-source luminosity function determined from a similar *Chandra* observation of a blank field by Mushotzky et al. (2000), the H I column density map of M101 by Kamphuis (1993), and the Hartmann & Burton (1997) measure of the Galactic H II column.

Given that the *Chandra* response is rather hard and that the AGN spectra are also hard, the diminution of the surface density of AGNs due to the H I absorption in M101 is rather small, only about 8%. From this we estimate that about 27 of the detected sources, only one-quarter of our total sample, are likely to be serendipitous background (or perhaps in a few cases, foreground) sources. Based on the above division of the chip area into “spiral arm” and “interarm” regions, about 20 of these background sources should appear to be located in the spiral arm and nuclear regions. This corresponds to a mean background surface density of $0.38 \text{ sources arcmin}^{-2}$, which is remarkably close to the measured density of sources in the “interarm” regions of $0.45 \text{ sources arcmin}^{-2}$. This suggests that most of these interarm sources are background AGNs, and indeed, six of the eight interarm sources do have relatively hard SR1 and HR1 colors (see § 6 and Fig. 6) that are consistent with the expected power-law spectrum of an AGN. Furthermore, the brightest of these sources (source 5) was previously identified as an AGN by Wang et al. (1999). The other two sources located in the interarm regions (sources 7 and 8) have very soft spectra and thus are more likely to be

TABLE 1
S3 CHIP X-RAY SOURCES IN M101

SOURCE	R.A. (J2000)	DECL. (J2000)	AREA (pixels)	NET COUNTS			NET COUNTS			$\log F_x$ (ergs cm ⁻² s ⁻¹)		
				S1	M1	H1	S2	M2	H2	S2	M2	H2
1	14 02 49.15	54 18 42.1	344	9	5	3	2	18	-4	-16.26	-15.27	...
2	14 02 49.61	54 20 00.2	261	6	12	6	-4	27	0	...	-15.13	...
3	14 02 51.63	54 20 53.1	199	4	3	9	2	8	5	-16.23	-15.57	-15.01
4	14 02 52.08	54 19 46.3	224	10	5	0	3	11	1	-16.16	-15.49	-15.57
5	14 02 52.82	54 21 11.8	185	160	164	355	52	441	185	-14.90	-13.85	-13.45
6	14 02 53.25	54 18 56.1	255	4	10	7	4	15	3	-16.00	-15.37	-15.30
7	14 02 55.64	54 20 29.0	144	12	1	1	3	9	2	-16.19	-15.55	-15.33
8	14 02 57.05	54 19 48.1	142	31	-1	-2	32	-3	-1	-15.12
9	14 02 57.49	54 18 34.7	223	-5	6	14	-4	8	11	...	-15.49	-14.64
10	14 02 59.43	54 20 43.0	90	19	12	1	7	25	1	-15.78	-15.17	...
11	14 03 00.47	54 20 02.9	94	24	10	2	9	25	3	-15.67	-15.16	-15.29
12	14 03 00.79	54 20 17.5	88	19	1	-1	12	8	-0	-15.54	-15.68	...
13	14 03 01.17	54 23 42.0	229	169	0	-4	124	44	-3	-14.52	-14.93	...
14	14 03 01.29	54 21 33.7	90	8	10	1	2	18	-0	-16.33	-15.33	...
15	14 03 02.12	54 23 25.5	192	-4	9	5	-1	8	3	...	-15.59	-15.21
16	14 03 02.57	54 24 17.1	280	43	-1	-3	46	-2	-4	-14.82
17	14 03 03.84	54 21 33.2	95	5	26	102	3	61	70	-16.17	-14.66	-13.87
18	14 03 04.73	54 19 24.9	88	7	9	2	4	14	1	-16.02	-15.42	...
19	14 03 05.22	54 17 52.4	221	40	31	55	16	91	19	-15.41	-14.57	-14.43
20	14 03 05.31	54 23 11.9	143	7	8	0	-1	16	0	...	-15.35	...
21	14 03 06.00	54 19 45.1	92	4	0	22	2	10	15	-16.34	-15.44	-14.55
22	14 03 06.16	54 20 12.3	94	1	1	12	2	2	11	-16.35	-16.15	-14.69
23	14 03 06.59	54 22 28.9	91	4	7	15	4	14	9	-16.02	-15.34	-14.77
24	14 03 06.99	54 22 09.9	95	1	1	25	-1	13	16	...	-15.33	-14.52
25	14 03 07.39	54 19 38.3	95	9	62	230	2	163	137	-16.36	-14.25	-13.58
26	14 03 07.89	54 18 36.9	137	12	9	17	7	24	8	-15.78	-15.14	-14.84
27	14 03 07.99	54 21 23.3	93	28	9	-1	15	23	-1	-15.42	-15.19	...
28	14 03 08.32	54 18 12.0	170	4	2	38	3	6	35	-16.13	-15.61	-14.17
29	14 03 08.42	54 24 59.7	359	18	51	100	9	119	41	-15.67	-14.43	-14.10
30	14 03 08.52	54 20 57.7	95	29	0	-0	27	2	1	-15.19	-16.29	...
31	14 03 08.65	54 23 36.8	168	18	3	1	12	10	-0	-15.53	-15.59	...
32	14 03 09.07	54 22 59.0	104	3	6	20	1	21	7	...	-15.17	-14.85
33	14 03 09.41	54 20 53.0	92	4	11	8	4	15	5	-16.03	-15.34	-15.04
34	14 03 09.76	54 21 15.2	95	11	11	35	5	38	15	-15.93	-14.92	-14.55
35	14 03 10.50	54 17 49.1	209	6	6	16	3	15	10	-16.09	-15.28	-14.70
36	14 03 10.70	54 21 26.7	94	15	3	-0	5	14	-0	-15.93	-15.44	...
37	14 03 12.44	54 17 54.3	198	82	31	16	27	96	5	-15.18	-14.58	-15.01
38	14 03 12.54	54 20 53.2	60	95	83	119	38	204	54	-15.03	-14.19	-13.97
39	14 03 12.55	54 24 18.4	250	8	18	14	4	31	6	-15.99	-15.04	-14.96
40	14 03 12.55	54 20 56.5	60	143	95	76	68	208	38	-14.77	-14.20	-14.12
41	14 03 12.75	54 17 43.7	222	7	3	6	2	15	-1	-16.34	-15.39	...
42	14 03 12.82	54 19 01.2	88	12	3	0	6	9	1	-15.84	-15.61	...
43	14 03 13.31	54 21 57.5	93	26	4	1	13	17	2	-15.51	-15.33	-15.49
44	14 03 13.61	54 20 54.0	88	7	3	8	2	12	5	-16.32	-15.43	-15.03
45	14 03 13.63	54 20 09.6	94	216	2	-2	170	48	-1	-14.39	-14.90	...
46	14 03 13.69	54 21 47.3	95	5	1	30	2	12	23	-16.36	-15.35	-14.36
47	14 03 13.69	54 19 09.1	93	32	62	22	3	112	2	-16.16	-14.52	-15.49
48	14 03 13.90	54 18 11.4	165	23	14	1	11	26	1	-15.57	-15.15	...
49	14 03 13.95	54 21 36.7	88	24	19	21	12	46	7	-15.54	-14.88	-14.88
50	14 03 14.25	54 20 52.0	93	8	5	2	1	12	3	...	-15.46	-15.29
51	14 03 14.37	54 18 05.7	180	63	47	58	21	126	22	-15.30	-14.44	-14.38
52	14 03 14.62	54 21 52.1	92	14	7	-1	8	13	-0	-15.72	-15.47	...
53	14 03 14.84	54 21 10.5	92	13	7	6	3	20	4	-16.16	-15.24	-15.15
54	14 03 14.87	54 21 23.2	94	2	7	5	-0	14	1	...	-15.43	...
55	14 03 15.08	54 20 18.1	94	19	1	1	7	14	1	-15.78	-15.43	...
56	14 03 15.47	54 17 03.7	320	39	26	4	9	60	0	-15.68	-14.80	...
57	14 03 15.82	54 17 48.4	219	6	30	146	-1	101	83	...	-14.46	-13.80
58	14 03 16.02	54 18 19.8	158	14	25	11	0	46	3	...	-14.89	-15.23
59	14 03 16.50	54 21 05.8	94	12	2	14	6	13	10	-15.85	-15.36	-14.73
60	14 03 16.71	54 18 34.8	133	17	9	7	8	23	3	-15.72	-15.19	-15.29
61	14 03 17.21	54 20 09.0	94	15	1	4	7	9	5	-15.78	-15.55	-15.05

TABLE 1—*Continued*

SOURCE	R.A. (J2000)	DECL. (J2000)	AREA (pixels)	NET COUNTS			NET COUNTS			$\log F_x$ (ergs cm ⁻² s ⁻¹)		
				S1	M1	H1	S2	M2	H2	S2	M2	H2
62.....	14 03 17.68	54 18 35.9	138	4	5	47	1	21	35	...	-15.11	-14.18
63.....	14 03 18.06	54 18 23.0	157	105	103	157	40	240	84	-15.01	-14.12	-13.79
64.....	14 03 18.66	54 24 30.2	293	15	41	53	1	81	27	-16.49	-14.58	-14.26
65.....	14 03 19.26	54 21 49.6	93	16	24	29	7	43	20	-15.78	-14.86	-14.42
66.....	14 03 20.44	54 16 31.9	427	0	11	88	2	48	49	-16.26	-14.74	-14.00
67.....	14 03 20.72	54 19 42.3	94	6	10	2	1	16	2	...	-15.35	-15.50
68.....	14 03 21.11	54 19 08.2	110	16	3	-1	2	17	-1	-16.22	-15.34	...
69.....	14 03 21.47	54 17 02.9	349	1	10	32	-1	22	21	...	-15.10	-14.39
70.....	14 03 21.54	54 19 45.9	88	174	275	423	40	640	193	-15.01	-13.71	-13.43
71.....	14 03 21.67	54 19 20.0	102	70	73	113	25	173	57	-15.22	-14.27	-13.96
72.....	14 03 21.68	54 20 24.0	94	21	-1	-2	20	1	-2	-15.32
73.....	14 03 22.69	54 20 38.9	93	29	8	2	11	28	1	-15.58	-15.13	...
74.....	14 03 23.76	54 18 18.5	207	-0	-1	20	-1	9	11	...	-15.49	-14.68
75.....	14 03 24.05	54 23 37.5	227	50	19	-1	16	54	-2	-15.42	-14.84	...
76.....	14 03 24.18	54 19 48.9	94	53	228	704	10	584	392	-15.62	-13.70	-13.12
77.....	14 03 25.18	54 20 50.2	94	8	11	21	3	27	11	-16.16	-15.07	-14.69
78.....	14 03 25.31	54 23 43.2	246	-1	5	21	-4	14	15	...	-15.30	-14.54
79.....	14 03 25.31	54 20 16.1	91	3	8	13	3	15	7	-16.15	-15.32	-14.89
80.....	14 03 25.33	54 21 13.2	92	29	0	3	28	1	4	-15.17	...	-15.15
81.....	14 03 25.85	54 21 25.2	98	15	4	-1	3	17	-1	-16.18	-15.36	...
82.....	14 03 26.28	54 19 51.6	119	14	5	-1	1	17	0	-16.52	-15.34	...
83.....	14 03 26.64	54 16 56.0	414	18	4	-1	8	16	-3	-15.69	-15.38	...
84.....	14 03 26.69	54 20 42.9	102	41	14	5	14	41	4	-15.48	-14.93	-15.07
85.....	14 03 27.16	54 18 31.4	223	51	36	24	26	76	9	-15.20	-14.66	-14.74
86.....	14 03 27.73	54 24 00.5	308	1	4	11	-1	5	11	...	-15.68	-14.66
87.....	14 03 27.90	54 20 04.5	130	2	2	14	-1	11	8	...	-15.43	-14.83
88.....	14 03 28.66	54 22 02.1	155	8	6	2	1	12	2	-16.46	-15.45	-15.39
89.....	14 03 28.77	54 21 13.5	130	15	6	5	5	17	4	-15.92	-15.30	-15.14
90.....	14 03 28.89	54 20 59.5	137	12	14	2	5	25	-1	-15.93	-15.19	...
91.....	14 03 29.01	54 21 49.4	152	8	11	2	-1	19	3	...	-15.25	-15.21
92.....	14 03 29.90	54 20 57.9	143	78	-0	0	46	32	0	-14.96	-15.06	...
93.....	14 03 29.98	54 22 29.0	193	38	170	189	3	340	54	-16.21	-14.01	-13.98
94.....	14 03 30.73	54 22 22.0	205	0	13	47	1	35	24	-16.51	-14.93	-14.34
95.....	14 03 31.51	54 20 52.9	169	6	4	3	5	5	3	-15.91	-15.80	-15.27
96.....	14 03 31.97	54 23 25.0	303	21	2	-7	19	2	-6	-15.29	-16.14	...
97.....	14 03 32.25	54 15 35.3	736	8	12	30	6	27	18	-15.83	-15.05	-14.47
98.....	14 03 32.40	54 21 03.1	187	6723	2210	375	2828	6417	62	-13.17	-12.77	-13.92
99.....	14 03 33.37	54 17 59.8	371	215	-1	-5	145	69	-4	-14.45	-14.74	...
100.....	14 03 33.58	54 23 07.4	301	3	-0	12	-2	3	13	...	-15.85	-14.59
101.....	14 03 34.00	54 20 11.4	227	20	-1	-1	22	-4	0	-15.28
102.....	14 03 35.35	54 21 24.1	244	4	5	20	2	20	7	-16.24	-15.21	-14.88
103.....	14 03 35.56	54 17 08.9	534	67	58	123	23	173	53	-15.26	-14.27	-13.99
104.....	14 03 36.05	54 19 24.8	300	303	227	174	117	528	58	-14.55	-13.82	-13.95
105.....	14 03 36.27	54 18 14.8	401	8	23	120	-0	87	65	...	-14.52	-13.90
106.....	14 03 39.29	54 18 26.7	446	14	16	39	7	44	18	-15.79	-14.86	-14.46
107.....	14 03 41.37	54 19 04.0	437	140	136	27	23	278	2	-15.26	-14.13	-15.40
108.....	14 03 47.36	54 22 29.6	581	23	24	12	8	45	6	-15.73	-14.88	-14.90
109.....	14 03 52.05	54 21 48.0	755	54	37	8	23	76	0	-15.26	-14.69	...
110.....	14 03 53.86	54 21 57.8	756	271	283	223	80	625	72	-14.72	-13.75	-13.86

NOTE.—Units of right ascension are hours, minutes, and seconds, and units of declination are degrees, arcminutes, and arcseconds. Table 1 is also available in machine-readable form in the electronic edition of the *Astrophysical Journal*.

physically associated with M101. Source 8, in fact, is one of the supersoft sources (SSSs) discussed in § 9.

4. THE $\log N$ – $\log S$ RELATION

To create a $\log N$ – $\log S$ relation for the point sources toward M101 that would be more easily comparable to relations derived for other galaxies, we applied somewhat more restrictive criteria than for the catalogue of sources. We chose all sources where the effective exposure was greater than 75% of the maximum and having a significance

greater than 4.5 in the 0.5–2.0 keV energy band (the M2 band defined in § 6), where the source counts were measured within the elliptical point-spread function (PSF) region containing 75% of the encircled energy, and the background was measured from an annular region 2–4 times the size of the 95% encircled energy ellipse. The source rates were corrected for vignetting. (This choice of energy range balances the need to detect the most numerous types of objects in M101, which are soft, with the desire to detect hard objects as well.) The object catalog was binned into $\Delta \log (S/\text{counts})$




FIG. 3.— $\log N$ – $\log S$ relation for M101. The points with error bars are the relation for all the point sources with $\sigma > 4.5$, the lower smooth line is the calculated relation for the background AGN absorbed by the M101 disk, and the squares are the difference between those two relations. The dotted line is the fit; the vertical lines show the region over which the fit was made. A flux conversion factor equivalent to 10^{-14} ergs cm $^{-2}$ s $^{-1}$ = 3.33×10^{-3} counts s $^{-1}$, which is appropriate for a background AGN with a $\Gamma = 1.42$ power-law X-ray spectrum, was used in these calculations.

s $^{-1}$) bins, and each bin was divided by the number of square arcminutes over which such a flux could have been detected.³ The $\log N$ – $\log S$ relation is shown in Figure 3 along with the relation due to the background AGN seen through the disk of M101 (as calculated from the relation of Mushotzky et al. 2000). A flux conversion factor equivalent to 10^{-14} ergs cm $^{-2}$ s $^{-1}$ = 3.33×10^{-3} counts s $^{-1}$, which is appropriate for a background AGN with a $\Gamma = 1.42$ power-law X-ray spectrum, was used in these calculations. The resulting $\log N$ – $\log S$ relation for the sources not due to the background AGN is nearly linear with $N(>S) \propto S^{-0.80 \pm 0.05}$. Excluding the eight interarm sources identified in the previous section from the analysis only makes a slight difference, resulting in a slope of -0.76 ± 0.05 . The uncertainty of the slope was calculated following the Lampton, Margon, & Bowyer (1976) criteria and is larger than the bootstrap uncertainty (± 0.02) or the variation produced by changing the flux range of the fit (± 0.02). The relatively flat slope derived here implies that the amount of diffuse emission due to unresolved faint point sources will be small.

Figure 4 shows the surface density of point sources as a function of radius from the nucleus. The optical bulge of M101 is visible out to ~ 0.75 (Okamura, Kanazawa, & Kodaira 1976), and we do indeed see an increase in the point-source surface density in the bulge region. Because the total number of sources within the bulge region is small




FIG. 4.—Surface density of point sources as a function of the distance from the nucleus, calculated for annuli with $\Delta R = 0.5$.

(~ 14), it is difficult to determine whether the “bulge sources” follow a different $\log N$ – $\log S$ relation than the “disk sources.” However, to within the uncertainties, we do not observe a difference in the shapes of the relations between bulge ($R < 1'$) and disk sources ($1' < R < 4.5'$).

The above analysis could be repeated for the 2.0–8.0 keV band using the 2–10 keV band AGN luminosity function found by Mushotzky et al. (2000), but in this energy range only ~ 18 of the objects detected with $\sigma > 4.5$ are not background AGNs. The slope of the $\log N$ – $\log S$ relation is quite uncertain (0.69 ± 0.18 , where the quoted uncertainty is no more than a formal uncertainty); to within the uncertainties, the 2.0–8.0 keV band is consistent with the 0.5–2.0 band.

That the slope of our 0.5–2.0 keV luminosity function does not agree with that of Wang et al. (1999) ($1.9^{+1.3}_{-0.3}$) is hardly surprising; the HRI data were over an order of magnitude less sensitive, many of the sources detected are actually composed of several point sources, and some of the brighter sources are embedded in diffuse emission that would have fallen within the HRI PSF. Further, the Wang et al. (1999) luminosity function was formed over a much larger area (a radius of $12'$ compared to our $\leq 6'$) and thus may be composed of a different population of sources.

5. SOURCE IDENTIFICATION

About $\frac{1}{4}$ of the *Chandra* X-ray sources can be cross-identified with previously observed X-ray sources or are positionally coincident with other cataloged objects as listed in Table 2. The previous *ROSAT* HRI and PSPC observations of M101 (Wang et al. 1999) detected 19 sources within our field of view, and all but three of these are detected in the *Chandra* observation. A 20th source (their source H42) is just visible on the edge of the field, but it is too highly distorted and affected by the edge of the detector to be included in our final list of sources. Of the three nondetected sources, two of them were classified as

³ The point-source detection limit is spatially variable, as both the size of the PSF and the background rate vary. Although the variation of the size of the PSF dominates, since some point sources are embedded in diffuse emission, the background rate can be significant in determining the point-source detection limit. Both effects, as well as the effect of the vignetting, were included in this analysis.

TABLE 2
CROSS-IDENTIFICATIONS

Source	<i>ROSAT</i> ^a	MF ^b	Other IDs
5	H18 (AGN)
6	2MASS
11	MF33	...
15	MF34	...
37	H22	...	NGC 5458, 2MASS, GSC
38	H23	...	3" south of nucleus
40	H23	...	Nucleus, 2MASS, GSC
42	MF46	...
43	MF49	...
45	H24
51	H25
52	MF50	...
63	H26
64	H27
67	MF54	...
70	H29
76	H30
81	MF61	...
85	MF65	...
95	2MASS
98	H32 (var)
99	H33
103	H35
104	H36	MF83	...
107	H37	...	NGC 5461, 2MASS
108	2MASS
110	H40	...	NGC 5462, 2MASS

^a *ROSAT* source identification from Wang et al. 1999.

^b SNR identification from Matonick & Fesen 1997.

variable by Wang et al. (1999) (their sources H28 and H31), so they may have been in a low flux state during our observation. The other nondetected object (their H41 source) is located near the east corner of the S3 chip, where our detection threshold is relatively high.

The nuclear source seen in the *ROSAT* HRI image (source H23) is resolved into two nearly identical point sources in the *Chandra* image. The northern component (source 40) appears to coincide with the nucleus at 14^h03^m12^s.55, 54°20'56".5 but is otherwise unremarkable compared to the other X-ray sources in the field. The other nuclear component is 3".1 (~110 pc) to the south and coincides with a loose cluster of bright stars as seen in a *Hubble Space Telescope* (*HST*) WFPC1 optical image (after applying an ~1" offset to the nominal *HST* coordinates to co-align the nucleus in both images). The nuclear X-ray source is slightly brighter than the southern component (they are ranked ninth and 12th in relative X-ray count rate in Table 1) and has a slightly softer spectrum, although both sources have fairly neutral X-ray hardness ratios.

Matonick & Fesen (1997) identified 54 optical SNRs within the *Chandra* field of view, and nine of these were detected in the X-ray band and listed in Table 2. Another three SNR sources (their SNR numbers 24, 32, and 43) are faintly visible on the adaptively smoothed *Chandra* image, but the detections were not significant enough to be included in our final list of 110 sources. All nine of the X-ray-detected SNR sources have soft X-ray spectra, as would be expected from a hot plasma. The two brightest SNR sources (our sources 85 and 104) were variable during the *Chandra* observation, so in at least these cases the X-ray

flux must be coming from a compact source and not from the extended SNR itself. In a companion paper (Snowden et al. 2001) we present a more detailed investigation into the properties of several of these SNRs.

To complete the Table 2 list of cross identifications, two of the five giant H II regions in M101 (Williams & Chu 1995) are located near the edge of the S3 chip (source 107 is NGC 5461 and source 110 is NGC 5462). The supernova SN 1951H was located about 12" southeast of NGC 5462 (Israel, Goss, & Allen 1975), but it is not detected in the *Chandra* observation. Finally, seven of the *Chandra* sources are coincident with near-IR objects in the 2MASS catalog, and two of these are also cross-identified with sources in the *HST* Guide Star Catalog (one being the nucleus, the other being NGC 5458, a bright H II region).

About 10% of globular clusters in our Galaxy and other nearby galaxies contain X-ray-emitting neutron stars, so a population of globular clusters might account for some of the X-ray sources in M101. The only published list of globular type clusters in M101 (Bresolin, Kennicutt, & Stetson 1996) contains 41 visually identified clusters within one relatively small (2' square) *HST* WFPC2 observation. It turns out, however, that most of these clusters are very blue and hence presumably young, like those observed in the Magellanic Clouds. Out of the 14 X-ray sources in this same region of M101, only one of them (source 80, one of the SSSs discussed in § 9) is located within 2" of any of the optical clusters, which has an ~40% probability of occurring by chance.

6. X-RAY COLORS

The X-ray events located within a circular region around each of the 110 X-ray sources were used to calculate the count rate within standard energy bands and to generate spectra and light curves for each source. The radius of the region was chosen as a function of the distance of the source from the optical axis so as to enclose greater than 95% of the source flux as predicted from the model PSFs and verified by numerical experiments on the brightest sources. The area of each source region is given by the "Area" column in Table 1, in units of the 0".49 *Chandra* CCD pixels.

Since most of the sources are too faint for detailed spectral analysis, we have characterized the general spectral properties of the sources by using the ratios of the X-ray count rates in two sets of soft, medium, and hard energy X-ray bands. The energy ranges for the first set of bands (S1, M1, and H1) were selected to produce roughly equal count rates in each band for the typical sample of sources seen in M101. For the second set of bands (S2, M2, and H2) a more standard set of astrophysically relevant energy ranges was used to facilitate the comparison of our results with those from previous X-ray missions. The adopted energy range for each of these bands is given in Table 3 along with the observed mean background flux level in each band. (The slight variation in background level and diffuse emission across the field is not significant for these purposes). The net counts, after background subtraction, for each source in each of these six bands are listed in Table 1. A pair of soft and hard X-ray flux ratios (or X-ray colors) were then calculated from the net counts in the three bands, where $SR1 = (M1 - S1)/(S1 + M1)$, $HR1 = (H1 - M1)/(M1 + H1)$, and similarly for the SR2 and HR2 ratios. In principle, the measured colors will depend on the position of the source in the field because the sensitivity of the ACIS

TABLE 3
ENERGY BANDS

Band	Low (keV)	High (keV)	Background (counts pixel ⁻¹)
S1	0.125	0.8	0.034
M1	0.8	1.3	0.007
H1	1.3	8.0	0.032
S2	0.125	0.5	0.023
M2	0.5	2.0	0.025
H2	2.0	8.0	0.025

instrument (i.e., the effective area of the telescope) decreases with distance from the aim point at slightly different rates depending on the energy band. This effect is small over the area covered by the S1 chip, however, and would cause the apparent X-ray colors to shift by only about 0.01 as the source is moved from the aim point to the far corner of the chip.

We compared the *Chandra* 0.125–2.0 keV (S2 + M2 band) count rates with the *ROSAT* HRI count rates as measured by Wang et al. (1999). Figure 5 shows that there is a very good correlation between the HRI and *Chandra* count rates for 15 of the 16 sources in common, which indicates that they have not varied significantly between the epochs of the observations (80% of the HRI data were obtained in 1996 and 20% in 1992). The one very discrepant point is source 98, which is highly variable (see § 7) and is now about 25 times brighter in mean flux level than during the *ROSAT* observations. It has been plotted at 1/10 of its actual mean *Chandra* count rate in order to fit onto the scale of Figure 5. The other bright source in Figure 5 (NGC 5462) has a slightly higher *ROSAT* count rate than would be predicted by the relation defined by the other points, but




FIG. 5.—Comparison of the *Chandra* and *ROSAT* count rates. The bright source 98 (plotted at 1/10 of its mean *Chandra* count rates in order to fit it on the graph) is highly variable and now has a mean flux level ~ 25 times brighter than when observed by *ROSAT*. The arrows at the bottom of the figure show the *Chandra* count rates for sources that were not previously detected by *ROSAT*.

the comparison is complicated because the higher resolution *Chandra* image shows that it is a blend of several sources. Also shown at the bottom of Figure 5 are the *Chandra* count rates for 11 sources that were not detected in the *ROSAT* observations. The brightest two of these sources (sources 93 and 71) were variable during the *Chandra* observation (and hence could have been in a low state during the HRI observations), and the other sources are close to the *ROSAT* detection threshold, so it is not particularly surprising that these sources were not previously detected by the *ROSAT* data.

The SR1 and HR1 X-ray colors provide a sensitive discriminator between sources with different types of X-ray spectra as shown in Figure 6. (Note that the SR2 and HR2 colors are less useful for this purpose because most of the flux lies in the M2 band and hence most of the points end up crowded into the lower right quadrant of the HR2 vs. SR2 diagram.) The 45 brightest sources, which have more than 40 net X-ray counts, have been plotted with their calculated 1σ error bars. The fainter sources have been omitted for clarity. Any points that have an SR1 or HR1 value lying outside the physically allowed range of -1 to 1 (as a result of uncertain background subtraction) have been plotted at the limiting value. It can be seen that most of the points in Figure 6 lie within a broad diagonal band ranging from the softest sources in the lower left to the hardest sources in the upper right.

The calculated SR1 and HR1 flux ratios for several model X-ray source distributions are also plotted in Figure 6. The upper three curves show models with power-law flux distributions with photon index slopes of 1.5, 2.0, and 4.0,




FIG. 6.—Observed X-ray color-color plot for the 45 brightest M101 sources that have more than 40 net X-ray counts, shown with the 1σ error bars. The six likely AGN sources located in the interarm regions of M101 are plotted with open triangles. The lines connecting the open squares show the predicted colors for five different sets of models. The upper three models have power-law flux distributions with the indicated photon index slopes, and the lower two models are blackbodies with the indicated temperatures (in units of keV). For each model the open squares are plotted at N_H values of, from left to right, 10^{20} , 5×10^{20} , 10^{21} , and 10^{22} atoms cm^{-2} .

respectively, and the two lower curves show blackbody models with temperatures of $kT = 0.1$ and 0.2 keV. In each case the lines connect the calculated SR1 and HR1 values for four N_H hydrogen column densities of, from left to right, 10^{20} , 5×10^{20} , 10^{21} , and 10^{22} atoms cm^{-2} . These models show that the SR1 color is most sensitive to the N_H parameter such that the more heavily obscured sources lie toward the right-hand side of the figure. The HR1 color, on the other hand, is a better indicator of the intrinsic shape of the spectral flux distribution such that the flatter spectrum sources lie toward the top of the figure.

The representative models that are plotted in Figure 6, while not unique, provide an indication of the physical nature of the objects that lie in different areas of the color-color diagram. The very soft sources in the lower left corner form a distinct subpopulation (see § 9). The expected population of obscured background AGNs seen through the disk of M101 will generally have power-law spectra with photon index slope in the range $1.3 < \Gamma < 4.0$ and total H I columns greater than 5×10^{20} . There are more than enough sources within this range of Figure 6 to account for the expected number of AGNs in the field, and the remainder are likely to be low-mass X-ray binaries in M101 itself. The six AGN candidates located in the interarm regions of M101, previously discussed in § 3, fall within this region of the figure (although one of them, source 39, lies quite close to the 4.0 slope limit) and are plotted with open triangles.

The sources below the $\Gamma = 4.0$ power-law model line are inconsistent with the spectrum of a typical AGN and are therefore more likely to be physically associated with M101.

7. TEMPORAL VARIABILITY

Out of the 37 brightest sources that have more than 60 net counts, four of them (sources 85, 93, 98, and 104) showed definite intensity variability during the *Chandra* observation (less than 10^{-5} probability of being a constant source based on the Kolmogorov-Smirnov test), and four others (sources 51, 66, 71, and 108) are probably variable (with probabilities ranging from 0.01 to 0.06 of being a constant source). The flux curves for these sources are shown in Figure 7. Two of these variable sources (sources 93 and 71) are now brighter than the *ROSAT* detection threshold, but they were not detected in the previous *ROSAT* observations taken in 1992 and 1996. In addition to these eight variable sources, there were also two variable sources previously observed with *ROSAT* (sources H28 and H31) that apparently have now faded below the *Chandra* detection limit.

Source 98 is currently the brightest and also the most dramatically variable source in M101. It has increased by about a factor of 25 in mean flux level from when it was previously observed by *ROSAT*, and the count rate varied by more than a factor of 6 during the *Chandra* observation alone.




FIG. 7.—X-ray light curves for the variable sources. The upper four sources are definitely variable, while the lower four sources are only marginally variable according to the Kolmogorov-Smirnov probability test for a constant source.

8. X-RAY MODELS AND FLUXES

The 29 brightest sources have enough flux (greater than 100 counts) to perform at least crude spectral modeling using the HEASARC XSPEC (Arnaud 1996) model fitting program. Most of the sources could be satisfactorily fitted with a simple absorbed power-law spectrum model; however, eight of the softest sources were better fitted by an absorbed blackbody model. Table 4 lists the best-fitting model parameters for these 29 sources. Also tabulated are the apparent X-ray fluxes calculated by integrating the model spectrum over the energy range of the S2 (0.125–0.5 keV), M2 (0.5–2.0 keV), and H2 (2.0–8.0 keV) energy bands as well as the intrinsic X-ray luminosity of the source in each of the three bands, correcting for the model N_H absorption and assuming a distance of 7.2 Mpc to M101. In general, the models provided a satisfactory fit to the observed spectra, as shown by the reduced χ^2 values given in the table. The main exception to this is the brightest variable source 98, whose integrated spectrum could not be well fitted by any simple model. The X-ray flux from this source is generally very soft, but the spectrum changes significantly with the flux level. As a consequence, the flux and luminosity values given for this object are only representative of its true nature. This unusual object will be studied in more detail in a separate paper, but it is already apparent that this is a super-Eddington source with a maximum instantaneous luminosity in the range of 10^{39} – 10^{40} ergs s^{-1} .

The ratio between the total net X-ray counts and the model X-ray flux in the S2, M2, and H2 bands for the 29 brightest sources was used to derive a mean counts-to-flux conversion factor that could be applied to the fainter sources to estimate their fluxes. Figure 8 shows the flux-per-count ratio plotted as a function of the hardness ratio, HR2, for each source. The adopted multiplication factors, which convert from total net counts in our 98.2 ks exposure into units of ergs $cm^{-2} s^{-1}$, are

$$C_{S2} = (2.37 \pm 0.39) \times 10^{-17},$$

$$C_{M2} = [(3.5 + 0.9HR2) \pm 0.24] \times 10^{-17},$$

and

$$C_{H2} = (1.89 \pm 0.34) \times 10^{-16},$$

where the M2 band conversion factor has a slight dependence on the hardness ratio of the source. These conversion factors were used to compute the X-ray flux values given in Table 1 for all the sources. Note that all the X-ray flux and luminosity values (but not the net counts values) quoted in this paper have been corrected for several instrumental effects. Every flux and luminosity value has been multiplied by a nominal factor of 1.02 (or 1.05 in the case of the slightly smaller apertures that were used for the two nuclear sources) to correct for the flux that was excluded by the circular integration apertures. In addition, six of the sources needed an $\sim 5\%$ correction factor because they were projected near one of the “warm” columns next to the CCD

TABLE 4
XSPEC MODEL FITS TO THE 29 BRIGHTEST SOURCES

SOURCE	Pow (α)	BB (kT)	N_H (10^{21})	χ^2_v	$\log F_x$ (ergs $cm^{-2} s^{-1}$)			$\log L_x$ (ergs s^{-1})		
					S2	M2	H2	S2	M2	H2
5	1.29	...	0.65	1.46	−14.92	−13.88	−13.38	37.56	37.99	38.41
13	0.10	0.10	2.06	−14.50	−14.87	−20.06	37.48	36.95	31.74
17	1.06	...	3.66	1.00	−17.06	−14.67	−13.84	36.83	37.40	37.96
19	1.73	...	0.62	1.40	−15.35	−14.52	−14.28	37.19	37.35	37.51
25	1.51	...	5.97	0.75	−17.19	−14.25	−13.53	37.70	37.99	38.29
29	2.07	...	4.56	1.13	−16.65	−14.44	−14.06	37.84	37.79	37.75
37	0.13	1.92	1.74	−15.40	−14.63	−17.75	37.68	37.58	34.07
38	1.56	...	0.35	1.30	−14.97	−14.25	−13.95	37.32	37.58	37.85
40	2.02	...	0.34	1.30	−14.72	−14.22	−14.19	37.63	37.62	37.61
45	0.06	0.78	1.34	−14.34	−14.89	−23.68	38.50	37.14	28.12
47	0.15	4.96	1.39	−16.35	−14.55	−16.67	37.95	38.04	35.20
51	2.08	...	0.84	1.13	−15.28	−14.48	−14.42	37.47	37.42	37.38
57	2.08	...	10.14	0.76	−18.46	−14.47	−13.83	38.10	38.06	38.01
63	1.44	...	0.39	0.58	−14.95	−14.14	−13.76	37.36	37.70	38.03
64	2.16	...	3.11	1.57	−16.26	−14.58	−14.34	37.66	37.56	37.46
66	2.71	...	19.56	1.24	−21.25	−14.69	−14.00	38.76	38.33	37.90
70	1.88	...	2.02	1.54	−15.11	−13.71	−13.42	38.24	38.31	38.38
71	1.38	...	0.47	1.59	−15.21	−14.32	−13.90	37.15	37.53	37.90
76	1.64	...	4.60	1.11	−16.13	−13.70	−13.12	38.26	38.48	38.69
85	2.31	...	0.54	1.14	−15.21	−14.69	−14.80	37.37	37.18	36.99
93	4.00	...	7.57	1.38	−16.30	−13.99	−14.26	40.01	38.80	37.60
98	0.14	0.53	4.11	−13.17	−12.78	−15.77	39.11	39.12	36.03
99	0.08	0.63	1.81	−14.44	−14.73	−21.32	38.11	37.24	30.48
103	1.61	...	1.04	0.75	−15.36	−14.26	−13.91	37.42	37.65	37.89
104	2.70	...	1.23	1.19	−14.54	−13.80	−14.04	38.60	38.18	37.76
105	2.21	...	9.74	0.73	−18.31	−14.51	−13.94	38.15	38.02	37.90
107	0.13	3.75	1.12	−15.41	−14.13	−16.89	38.46	38.38	34.96
109	0.20	0.20	0.98	−15.25	−14.68	−16.56	36.75	37.15	35.24
110	2.90	...	2.27	1.39	−14.79	−13.71	−13.93	38.96	38.42	37.87




FIG. 8.—Derived counts-to-flux conversion factors for the S2, M2, and H2 energy bands, based on the model spectra fit to the 29 brightest sources.

readout amplifier that were excluded during the initial data cleaning procedure. Finally, two of the sources were located near the edge of the chip and required an additional vignetting correction factor because the deliberate wobbling of the telescope pointing axis during the exposure caused the source to move periodically on and off the chip.

Figure 9 shows the distribution of $\log F_X$ for all 110 sources (using the values from Table 1), and Figure 10 shows the distribution of $\log L_X$ for the 29 brightest sources from Table 4. Note that it is not possible to compute reliably the luminosity for the fainter sources because the individual N_H values are unknown. Several of the sources, especially in the softest band, appear to have luminosities greater than the Eddington limit of $\sim 1.3 \times 10^{38} \text{ ergs s}^{-1}$ for a $1 M_\odot$ object, but the uncertainties in the calculated luminosities are sometimes quite large. The brightest S2




FIG. 10.—X-ray luminosity histograms for the 29 brightest sources in the three energy bands and the total luminosity. The positions in the histograms of the four brightest interarm AGN candidates, discussed in § 3, are indicated by the crosses.

band source, source 93, for example, has a high luminosity mainly because the best-fitting model has a very steep spectrum and a large hydrogen column density of $7.5 \times 10^{21} \text{ cm}^{-2}$, which leads to a large correction factor to the apparent flux. The next most luminous S2 band source is source 98, which, as mentioned previously, is highly variable and has a spectrum that is not well fitted by simple flux models. It has been plotted at its mean *Chandra* flux level, but its instantaneous luminosity has been observed to vary by at least 2 orders of magnitude.

In principle, the observed color dependence of the counts-to-flux conversion factor for the M2 band (Fig. 8)




FIG. 9.—X-ray flux histograms for all 110 sources in the three energy bands and the total flux. The positions in the histograms of the six interarm AGN candidates, discussed in § 3, are indicated by the crosses.




FIG. 11.—Mean HR2 color as a function of source count rate. The sources were sorted in order of M2 band count rate, then the mean HR2 color was computed for bins containing 10 sources each. The error bars show the mean statistical error on the HR2 colors for the individual sources in each bin. The nearly horizontal line is the least-squares fit to the points, which shows that there is no significant trend in mean HR2 color as a function of count rate.

could affect the analysis of the $\log N$ – $\log S$ relation in § 4. In that analysis we assumed a constant conversion factor between counts and flux, but if the mean X-ray color of the sources changes with count rate (i.e., if the fainter sources are significantly harder or softer in the mean than the brighter sources), then this would cause a systematic difference between the slope of the $\log N$ – \log (counts) and the $\log N$ – \log (flux) relations. This difference could be important because most observational source population studies have used $\log N$ – \log (counts), whereas most theoretical interpretation is done in terms of $\log N$ – \log (flux).

In order to judge the importance of this effect, we computed the mean HR2 color of sources as a function of count rate, as shown in Figure 11. The fact that there is no significant trend in the figure (the best-fitting line has a slope of -0.013 ± 0.16) means that our assumption of a constant counts-to-flux conversion factor has not biased the derived slope of the $\log N$ – $\log S$ relation.

9. SUPERSOFT SOURCES

Ten of the *Chandra* X-ray sources have very soft spectra similar to the SSSs that have been seen in other nearby galaxies. These M101 sources can be divided into two distinct subclasses: the brighter class consists of three sources (sources 13,⁴ 45, and 99), and the fainter and softer class consists of seven sources (sources 8, 16, 30, 72, 80, 96, and 101). The brighter class of SSSs has a mean count rate of 2.0 counts ks^{-1} and has an S2 band (0.125–0.5 keV) to M2 band count ratio of 3:1. The fainter class of SSSs, on the other hand, is an order of magnitude fainter with a mean flux of 0.26 counts ks^{-1} and has such a soft spectrum that essentially all the flux is emitted in the S2 band.

The three brighter SSSs have very similar spectra as demonstrated by the good agreement between the blackbody model fits given in Table 4 for the sources: the best-fitting model temperatures range from 60 to 100 eV (equivalent to 0.7 – 1.1×10^6 K), and the hydrogen column densities lie in the range of 1.0 – $8.0 \times 10^{20} \text{ cm}^{-2}$, which is consistent with the Galactic column density of $\sim 1.2 \times 10^{20} \text{ cm}^{-2}$ in the direction of M101 (Stark et al. 1992; Hartmann & Burton 1997) plus an additional absorption component local to the sources in M101. Because the spectra are so similar, we combined the counts from the three sources to obtain the better signal-to-noise spectrum shown in Figure 12. The best-fitting blackbody model to this combined spectrum has a temperature of 72 ± 2 eV [equivalent to $(8.4 \pm 0.2) \times 10^5$ K] and $N_{\text{H}} = (6.0 \pm 0.7) \times 10^{20} \text{ atoms cm}^{-2}$. Note that the formal errors on the model parameters are unrealistically small and do not take into account the unknown systematic errors in the calibration of the instrumental response below 0.3 keV. The mean luminosity of these three sources over the 0.125–2.0 keV band is $1.4 \times 10^{38} \text{ ergs s}^{-1}$ with most of this flux emitted below 0.5 keV. None of these three sources showed any obvious signs of variability, either during the *Chandra* exposure or between the epochs of the *ROSAT* and *Chandra* observations.

⁴ The only source detected in M101 in the *Einstein* IPC ultrasoft survey, at $14^{\text{h}}03^{\text{m}}00^{\text{s}}$, $54^{\circ}22'25''$ (Thompson, Shelton, & Arning 1998), may coincide with our source 13. The positions differ by 1.3, which is about twice the typical error in the *Einstein* source positions but is still well within the $3'$ integration radius that was used to extract the *Einstein* source counts.




FIG. 12.—Combined X-ray spectrum of the three SSSs cited in the text. The line shows the best-fitting model spectrum with a blackbody temperature of 72 eV and $N_{\text{H}} = 5.6 \times 10^{20} \text{ cm}^{-2}$.

The combined spectrum of the seven fainter SSSs in M101 is shown in Figure 13. In spite of the statistically large uncertainties in the count rates and the large systematic uncertainties in the energy calibration of the *Chandra* detector at these low energies, it is clear that this fainter class of SSSs has a much softer spectrum than the brighter class shown in Figure 12. Since the N_{H} value is not well constrained by the model fits, we assumed a conservative value equal to the Galactic column of $1.2 \times 10^{20} \text{ cm}^{-2}$. Under this assumption the best-fitting blackbody model to the mean spectrum has a temperature of 47 ± 2 eV and a 0.125–2.0 keV luminosity of $1.1 \times 10^{37} \text{ ergs s}^{-1}$ (85% of this flux is emitted below 0.3 keV). Assuming a larger N_{H} value decreases the best-fitting model temperature only slightly, but it greatly increases the calculated luminosity (by a factor




FIG. 13.—Combined X-ray spectrum of the seven extremely soft sources cited in the text. The line shows the best-fitting model spectrum with a blackbody temperature of 47 eV and an assumed value of $N_{\text{H}} = 1.2 \times 10^{20} \text{ cm}^{-2}$.

of 7 for $N_{\text{H}} = 6 \times 10^{20} \text{ cm}^{-2}$) because even a small increase in column density produces a large increase in extinction for these soft sources.

The SSSs were first detected in other nearby galaxies by *Einstein* (Long, Helfand, & Grabelsky 1981) and later studied more extensively with *ROSAT* (Trumpler et al. 1991; Greiner, Hasinger, & Kahabka 1991). Supper et al. (1997) classified 3.8% (15 out of 396) of the *ROSAT* PSPC sources in M31 as being SSSs. This is not necessarily inconsistent with the higher rate of SSSs we found in M101 (9%), however, because the higher Galactic N_{H} absorption toward M31 ($6 \times 10^{20} \text{ cm}^{-2}$) compared to M101 will absorb much of the soft flux from these sources. For this same reason, the SSSs are difficult to detect in our Galaxy if they are more than $\sim 1 \text{ kpc}$ away in the plane of the disk. Clearly, it will be of great interest to see how the properties of the SSS population vary between different types of galaxies as more data from the *Chandra* and *XMM* observations become available in the near future.

While the diversity of observed properties argues against lumping all SSSs into a single class of objects, the generally accepted model for the classical SSSs is that they are white dwarf stars with steady nuclear burning in their envelopes (van den Heuvel et al. 1992). It is thought that high accretion rates (greater than $10^{-7} M_{\odot} \text{ yr}^{-1}$) from a low-mass secondary onto the white dwarf primary fuel the burning of the hydrogen near the surface of the white dwarf. The predicted luminosities range from 6×10^{36} to $1 \times 10^{38} \text{ ergs s}^{-1}$, with peak energy fluxes in the 30–50 eV range, in good agreement with the sources seen in M101. The steep drop in flux above about 0.8 keV seen in Figure 12 is typical of most SSSs; recent high signal-to-noise *ASCA* observations (Asai et al. 1998) have shown that this break can be explained by an absorption edge feature probably due to O VIII at about 0.87 keV. This and other edges originating from hydrogen-like or helium-like ions of carbon, nitrogen, and oxygen would be expected for the emission from a hot white dwarf with a stable hydrogen-burning envelope.

The SSSs are also related to classical novae (CNs) in that SSSs are quasi-steady nuclear burning, whereas CNs are the result of thermonuclear runaways. Indeed, many CNs go through an SSS phase several months after the peak of optical outburst. In one case, the SSS phase lasted for nearly a decade (Shanley et al. 1995). We looked for a similar correlation in M101 by comparing the positions of the seven known optical novae that occurred within our M101 field between 1994 and 1997 (Shafter, Ciardullo, & Pritchett 2000) with our X-ray source catalog and with the position of even fainter possible sources seen in the adaptively smoothed X-ray image, but we found no matches. Thus, at least none of these novae had a long-lasting, bright, SSS phase after their optical outburst.

10. CONCLUSIONS

This paper provides a preview of what the studies of the X-ray source populations in external galaxies will reveal and what they will teach us about the X-ray sources in our own Galaxy. The main result of this paper is the catalog of 110 X-ray sources seen in the *Chandra* observation of the face-on spiral galaxy M101. The detection threshold is about $10^{36} \text{ ergs s}^{-1}$ but depends on position because of the variable angular resolution of the X-ray telescope across the field of view. Since this flux threshold is several orders of magnitude higher than the level of coronal X-ray emission

expected from normal stars, most of these sources must be more exotic objects, including X-ray binaries, supersoft sources (SSSs), and supernova remnants (SNRs). The field of view of the current observation only covers about 50% of the prominent spiral arms in M101, so a complete census of all the X-ray sources visible with *Chandra* in M101 at this flux limit could contain twice as many objects.

Only about $\frac{1}{4}$ of the sources are estimated to be background AGNs, so most of our sources are physically associated with M101. The bulk of the point sources are related to spiral arms, suggesting that the population of X-ray sources is derived from young objects in regions of active star formation and thus may be dominated by high-mass X-ray binaries. Conversely, spiral arms have higher stellar densities for all stellar types, so there may not be an overabundance of high-mass X-ray binaries. Most of the sources seen in the interarm regions are probably AGNs because their surface density is close to the expected number of background sources and because the X-ray colors of most of the interarm sources are consistent with that from a power-law spectrum usually seen in AGNs.

At a distance of 7.2 Mpc *Chandra* can provide a luminosity function down to $\sim 10^{36} \text{ ergs s}^{-1}$, so measuring the luminosity functions for the nearby galaxies will do much to explore the link between the X-ray source populations and the star formation history and will help to interpret the X-ray source luminosity functions being developed for the Milky Way. The derived $\log N$ – $\log S$ relation for the sources in M101 is nearly linear with a slope of -0.80 ± 0.05 over the flux range $\sim 10^{36}$ – $10^{38} \text{ ergs s}^{-1}$. We observed no difference in the slope of the relations between bulge ($R < 1'$) and disk sources ($1' < R < 4.5'$); however, there are few sources in the bulge so the uncertainties in the bulge slope are relatively large. Another recently published $\log N$ – $\log S$ study similar in scope to the present work is that of Tennant et al. (2001) for the galaxy M81 (NGC 3031) at a distance of 3.63 Mpc (Freedman et al. 1994). M81 is an Sab galaxy with a substantially larger bulge than M101. For the 0.2–8.0 keV band Tennant et al. (2001) find the $\log N$ – $\log S$ of the disk sources to have a slope of -0.50 over the flux range from 4×10^{36} to $2 \times 10^{39} \text{ ergs s}^{-1}$, somewhat flatter than our relation. Given the difference in energy range, as well as possible differences in the slope of the AGN luminosity functions used, the two values are consistent with each other. At a limiting flux of $10^{37} \text{ ergs s}^{-1}$, however, the surface density of X-ray sources is nearly 4 times greater for M81 (0.2 kpc^{-2}) than for M101 (0.05 kpc^{-2}), presumably reflecting the differences in the star formation rate between the two galaxies.

Eight of the brighter sources show evidence for short-term temporal variation over the 98.2 ks period of our observation, and two other sources previously observed with *ROSAT* are no longer visible with *Chandra*. The brightest *Chandra* source is also the most variable source, varying by more than a factor of 6 during the observation. The current mean flux level is about 25 times higher than when previously observed with *ROSAT*. This source has a soft spectrum that varies significantly with the flux level. Simple spectral models do not provide a very good fit to the spectrum, but it is clear that it has a super-Eddington luminosity that sometimes exceeds $10^{39} \text{ ergs s}^{-1}$. This unusual source will be studied in more detail in a subsequent paper.

The nuclear X-ray source is nonexceptional and looks very much like the other sources in the field. It is not spa-

tially resolved, although there is evidence for enhanced diffuse X-ray emission near the nucleus (to be studied in a separate paper). While this source may well be a weak AGN similar to that detected in most other nearby galaxies (Ho et al. 2001), it could also simply be a typical X-ray binary located in the dense stellar environment close to the nucleus. There is another nearly identical X-ray source just $3''.1$ to the south of the nucleus that coincides with a loose cluster of bright stars. A power-law spectral model gives a reasonable fit to both the nuclear source and the southern companion, resulting in photon indices of 2.02 and 1.56 and luminosities of 4.2×10^{37} and 3.8×10^{37} ergs s^{-1} in the 0.5–2.0 keV band, respectively.

About 9% of the M101 sources have extremely soft spectra similar to the SSSs seen in other nearby galaxies. The 10 SSSs in M101 can be naturally divided into two classes: the three brightest SSSs have a mean luminosity of 1.4×10^{38} ergs s^{-1} and a model blackbody temperature of 72 ± 2 eV, while the other seven SSSs have an order of

magnitude smaller X-ray count rate, most of which is emitted below 300 eV. The model temperature and luminosity of these fainter sources are dependent on the assumed hydrogen column density but are in the neighborhood of 50 eV and 1.1×10^{37} ergs s^{-1} , respectively. The observed luminosities and temperatures of both these classes of SSSs are within the range predicted by the standard models of a white dwarf star with steady nuclear burning in its envelope fueled by accretion from a low-mass secondary.

We thank the referee, Philip Kaaret, for his careful review and useful suggestions that improved this manuscript. This research was supported in part by NASA grant 01900441. We have made use of data and software obtained from the High Energy Astrophysics Science Archive Research Center (HEASARC), provided by NASA's Goddard Space Flight Center.

REFERENCES

- Arnaud, K. A. 1996, in ASP Conf. Ser. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. Jacoby & J. Barnes (San Francisco: ASP), 17
- Asai, K., Dotani, T., Nagase, F., Ebisawa, K., Mukai, K., Smale, A. P., & Kotani, T. 1998, *ApJ*, 503, L143
- Bresolin, F., Kennicutt, R. C., Jr., & Stetson, P. B. 1996, *AJ*, 112, 1009
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H., Jr., Buta, R., Paturel, G., & Fouque, P. 1991, *Third Reference Catalogue of Bright Galaxies* (Berlin: Springer)
- de Vaucouleurs, G., & Pence, W. D. 1978, *AJ*, 83, 1163
- Freedman, W. L., et al. 1994, *ApJ*, 427, 628
- Greiner, J., Hasinger, G., & Kahabka, P. 1991, *A&A*, 246, L17
- Hartmann, D., & Burton, W. B. 1997, *Atlas of Galactic Neutral Hydrogen* (Cambridge: Cambridge Univ. Press)
- Ho, L. C., et al. 2001, *ApJ*, 549, L51
- Israel, F. P., Goss, W. M., & Allen, R. J. 1975, *A&A*, 40, 421
- Kamphuis, J. J. 1993, Ph.D. thesis, Rijksuniversiteit, Groningen
- Lampton, M., Margon, B., & Bowyer, S. 1976, *ApJ*, 208, 177
- Long, K. S., Helfand, D. J., & Grabelsky, D. A. 1981, *ApJ*, 248, 925
- Matonick, D. M., & Fesen, R. A. 1997, *ApJS*, 112, 49
- McCammon, D., & Sanders, W. T. 1984, *ApJ*, 287, 167
- Mushotzky, R. F., Cowie, L. L., Barger, A. J., & Arnaud, K. A. 2000, *Nature*, 404, 459
- Okamura, S., Kanazawa, T., & Kodaira, K. 1976, *PASJ*, 28, 329
- Shafter, A. W., Ciardullo, R., & Pritchet, C. 2000, *ApJ*, 530, 193
- Shanley, L., Ögelman, H., Gallagher, J. S., Orio, M., & Krautter, J. 1995, *ApJ*, 438, L95
- Snowden, S. L., Mukai, K., Pence, W., & Kuntz, K. D. 2001, *AJ*, 121, 3001
- Snowden, S. L., & Pietsch, W. 1995, *ApJ*, 452, 627
- Stark, A. A., et al. 1992, *ApJS*, 79, 77
- Stetson, P. B., et al. 1998, *ApJ*, 508, 491
- Supper, R., Hasinger, G., Pietsch, W., Trumper, J., Jain, A., Magnler, E. A., Lewin, W. H. G., & van Paradijs, J. 1997, *A&A*, 317, 328
- Tennant, A. F., Wu, K., Ghosh, K. K., Kolodziejczak, J. J., & Swartz, D. A. 2001, *ApJ*, 549, L43
- Thompson, R. J., Jr., Shelton, R. G., & Arning, C. A. 1998, *AJ*, 115, 2587
- Trinchieri, G., Fabbiano, G., & Romaine, S. 1990, *ApJ*, 356, 110
- Trumpler, J., et al. 1991, *Nature*, 349, 579
- van den Heuvel, E. P. J., Bhattacharya, D., Nomoto, K., & Rappaport, S. A. 1992, *A&A*, 262, 97
- Wang, Q. D., Immler, S., & Pietsch, W. 1999, *ApJ*, 523, 121
- Williams, R., & Chu, Y.-H. 1995, *ApJ*, 439, 132