

## FIRST RESULTS FROM THE CATALINA REAL-TIME TRANSIENT SURVEY

A. J. DRAKE<sup>1</sup>, S. G. DJORGOVSKI<sup>1</sup>, A. MAHABAL<sup>1</sup>, E. BESHORE<sup>2</sup>, S. LARSON<sup>2</sup>, M. J. GRAHAM<sup>1</sup>, R. WILLIAMS<sup>1</sup>, E. CHRISTENSEN<sup>3</sup>,  
 M. CATELAN<sup>4,5,6</sup>, A. BOATTINI<sup>2</sup>, A. GIBBS<sup>2</sup>, R. HILL<sup>2</sup>, AND R. KOWALSKI<sup>2</sup>

<sup>1</sup> California Institute of Technology, 1200 E. California Blvd., CA 91225, USA

<sup>2</sup> Department of Planetary Sciences, The University of Arizona, Lunar and Planetary Laboratory, 1629 E. University Blvd., Tucson, AZ 85721, USA

<sup>3</sup> Gemini Observatory, Casilla 603, La Serena, CL, Chile

<sup>4</sup> Pontificia Universidad Católica de Chile, Departamento de Astronomía y Astrofísica, Av. Vicuña Mackena 4860, 782-0436 Macul, Santiago, Chile

<sup>5</sup> Catholic University of America, Department of Physics, 200 Hannan Hall, Washington, DC 20064, USA

Received 2008 September 3; accepted 2009 February 12; published 2009 April 20

### ABSTRACT

We report on the results from the first six months of the Catalina Real-Time Transient Survey (CRTS). In order to search for optical transients (OTs) with timescales of minutes to years, the CRTS analyses data from the Catalina Sky Survey which repeatedly covers 26,000 of square degrees on the sky. The CRTS provides a public stream of transients that are bright enough to be followed up using small telescopes. Since the beginning of the survey, all CRTS transients have been made available to astronomers around the world in real time using HTML tables, RSS feeds, and VOWEvents. As part of our public outreach program, the detections are now also available in Keyhole Markup Language through Google Sky. The initial discoveries include over 350 unique OTs rising more than 2 mag from past measurements. Sixty two of these are classified as supernovae (SNe), based on light curves, prior deep imaging and spectroscopic data. Seventy seven are due to cataclysmic variables (CVs; only 13 previously known), while an additional 100 transients were too infrequently sampled to distinguish between faint CVs and SNe. The remaining OTs include active galactic nucleus, blazars, high-proper-motions stars, highly variable stars (such as UV Ceti stars), and transients of an unknown nature. Our results suggest that there is a large population of SNe missed by many current SN surveys because of selection biases. These objects appear to be associated with faint host galaxies. We also discuss the unexpected discovery of white dwarf binary systems through dramatic eclipses.

**Key words:** BL Lacertae objects: general – galaxies: general – novae, cataclysmic variables – stars: flare – supernovae: general

*Online-only material:* color figures

### 1. INTRODUCTION

Time-domain astronomy is one of the most rapidly emerging areas of astronomy (Paczynski 2000). Several large experiments such as the Large Synoptic Survey Telescope (LSST; Ivezić et al. 2008), the Panoramic Survey Telescope and Rapid Response System (PanSTARRS; Hodapp et al. 2004), and SkyMapper (Keller et al. 2007), are set to make a major impact in the near future by covering thousands of square degrees with targets ranging from distant (supernovae) SNe to near-Earth asteroids. Past time-domain surveys have mainly concentrated on areas of tens to hundreds of square degrees and have typically searched for specific types of astronomical transients such as microlensing (MACHO; Alcock et al. 2000; OGLE, Udalski et al. 1994; EROS, Aubourg et al. 1995), gamma-ray burst (GRB) afterglows (ROTSE; Akerlof et al. 2000), and SNe (Lick Observatory Supernova Search (LOSS); Filippenko et al. 2001).

Deep surveys for variability have been carried out over small areas ( $< 25$  deg<sup>2</sup>) by the Subaru/XMM-Newton Deep Survey (SXDS; Morokuma et al. 2008), the Deep Lensing Survey (DLS; Becker et al. 2004), and the Faint Sky Variability Survey (FSVS; Huber et al. 2006) whereas large-area surveys, such as the Sloan Digital Sky Survey (SDSS; York et al. 2002), the Two Micron All Sky Survey (2MASS; Skrutskie et al. 2006), and the *Galaxy Evolution Explorer* (GALEX; Martin et al. 2005) have generally not been synoptic. One survey intermediate between past deep surveys and wide future surveys was carried out by the SDSS

consortium to discover Type Ia SNe. This survey covered 300 deg<sup>2</sup> in five optical bands to  $r \sim 23$  during nine months spaced over 2005–2007 (Sesar et al. 2007).

#### 1.1. Current Transient Surveys

Current wide-field transient surveys include the Robotic Optical Transient Search Experiment (ROTSE-III; Akerlof et al. 2003) and the All Sky Automated Survey (ASAS-3; Pojmanski 2001). However, like many other current surveys, ROTSE-III is targeted toward the discovery and characterization of GRB afterglows. The ASAS-3 survey can cover the visible southern sky every two nights and covers  $\sim 30,000$  deg<sup>2</sup> to  $V \sim 13.5$  during the year (Pojmanski 2001), while the four telescopes of the ROTSE-III survey routinely cover 1260 deg<sup>2</sup> to  $R \sim 17.5$  (Rykoff et al. 2005; Yost et al. 2007).

In the time between targeted surveys, and future wide, deep, transient surveys, two experiments have been searching tens of thousands of square degrees of the sky for transients; the Palomar-Quest (PQ)<sup>7</sup> digital synoptic sky survey (Djorgovski et al. 2008a), and the Catalina Real-Time Transient Survey (CRTS). The PQ survey started analyzing drift scan data for optical transients (OTs) in real time in August 2006, while the CRTS began in 2007 November. The transients discovered in PQ data will be presented in A. Mahabal et al. (2009a, in preparation).

Both CRTS and PQ experiments use a purpose-built pipeline for real-time transient detection, analysis and distribution (A. J.

<sup>6</sup> John Simon Guggenheim Memorial Foundation Fellow.

<sup>7</sup> <http://palquest.org>

Drake et al. 2009, in preparation). These surveys are now set to provide estimates of the rates and types of OTs that can be expected from future synoptic surveys.

In these experiments, unlike most current and past OT surveys, the results are made public within minutes of discovery using the International Virtual Observatory Alliance (IVOA) standard VOEvent protocol and the VOEventNet astronomical event distribution network (Drake et al. 2007a). In addition, VOEventNet, in collaboration with Google, makes transient astronomy available to the world in real time via a dedicated layer in Google Sky.<sup>8</sup> Current events available through VOEventNet also include MOA and OGLE microlensing, GCN GRB alerts, and PQ transients.

The CRTS routinely searches for OTs in data from the Catalina Sky Survey (CSS) Schmidt Telescope. In this paper, we present results from the first six months. First, we will discuss the observations and data analysis (Section 2). We will then list the current results (Section 3) and discuss some of the types of transients discovered. Finally, we will summarize the current findings (Section 4).

## 2. OBSERVATIONS AND DATA REDUCTION

The CSS<sup>9</sup> (Larson et al. 2003) uses the 0.7 m *f*/1.9 Catalina Schmidt Telescope north of Tucson, Arizona to discover near-Earth objects (NEOs) and potentially hazardous asteroids (PHAs). This survey uses a single unfiltered 4k × 4k CCD with 2".5 pixels, giving an 8 deg<sup>2</sup> field of view. To date, CSS has found hundreds of PHAs, comets and other solar system objects, and currently leads the rate of NEA discoveries.<sup>10</sup> On a clear night, the Catalina Schmidt typically covers ∼ 1200 deg<sup>2</sup> of sky in a sequence of four 30 s exposures. For each observation, there is an additional 18 s overhead for readout, step and settle time.

Currently observed fields cover ∼ 26,000 deg<sup>2</sup> in the declination range  $-30^\circ < \delta < 70^\circ$ . Observations are generally limited to Galactic latitudes  $|b| > 10^\circ$  to prevent confusion caused by crowding. Weather permitting, observations are made on 21–24 nights per lunation and typically reach *V* magnitudes from 19 to 20. Observations are made in a grid of adjacent fields so that sets of four observations of a field are taken at equal intervals over ∼ 30 minutes. This allows us to detect transients varying on timescales from minutes to years. In addition, the four image sequence provides a significant veto for asteroids and artifacts (due to cosmic rays and hot pixels) that often cannot be distinguished from genuine rapid transients in pairs of exposures.

In the near future, we hope to include two additional dedicated CSS telescopes in OT searches. The 1.5 m Mt. Lemmon reflector near Tucson, with a 1 deg<sup>2</sup> field of view, and the 0.5 m Uppsala Schmidt at Siding Spring, Australia, with a 4.2 deg<sup>2</sup> field of view. All three telescopes operate in the same mode with similar cameras and reduction software.

### 2.1. Transient Detection

The main goal of the CRTS thus far has been to discover and characterize variable source populations that exhibit high-amplitude variability on the timescales probed by CSS data. In this paper, we define OTs as objects that vary in brightness by greater than 2 mag between past catalogs derived from high signal-to-noise ratio (S/N) co-added images. The OT characterization process undertaken here includes understanding known

types of variable objects, as well as searching for new kinds of OTs. At the same time, the results of this survey have been designed to deliver a transient discovery stream that is available to the entire astronomical community as a testbed for VOEventNet technology.

The CSS uses standard SExtractor photometry software to produce object catalogs for each image as it is read out. To find transients, we compare new detections against deep, clean source catalogs, rather than simply comparing these to catalogs from earlier observations. This is necessary as image artifacts and differences in image depth lead to an overwhelming number of unmatched detections. The CSS detections are primarily matched with objects detected in past CSS co-added data. Each co-add consists of the median combination of at least 20 images. These images reach sources approximately 2 mag fainter than the deepest individual CSS images (mag ∼ 22). Thus, very few objects are missed in the reference catalog because of image artifacts or changes in depth and transparency. However, some objects are missing in the catalogs because they are blended in the co-added images. For this reason, the OT candidates are also matched against objects in the higher resolution USNO-B, SDSS, and PQ surveys.

An alternate method to catalog searches for finding transients is image subtraction. The image subtraction technique involves matching new observations to a high signal-to-noise template image before subtracting one from the other (Tomaney & Crotts 1996; Drake et al. 1999). The resulting difference image is then searched for transients. Although this increases the noise associated with detections, this is a very effective way of revealing transient objects in the presence of significant flux from blended constant sources. For this reason, subtraction has been particularly successful for dense stellar fields observed in SN searches such as used by the Nearby Supernova Factory (SNfactory; Aldering et al. 2002). However, this technique does not work well at blue wavelengths in broad filters because of atmospheric differential refraction (Drake et al. 1999). The resulting difference images have bipolar residuals at the locations of the stars and often lead to false detections and missed transients. The image subtraction process is also prone to fail, or produce artifacts, when images are taken in bad observing conditions. As catalog-based searches simply match lists of sources, they do not introduce new artifacts into the images, making them more suitable for unsupervised real-time transient searches.

In the PQ survey, we found that the detection of OTs in a single scan was hampered by the presence of large numbers of previously undiscovered asteroids (A. Mahabal et al. 2009a, in preparation). By design, the longer temporal separation of CSS images means that almost all asteroids can be detected from their motion between images. We currently require that a transient is detected in at least two observations in addition to requiring positional coincidence between images. This aids the removal of most false detections due to image artifacts such as hot pixels and cosmic rays. The four image sequences used by CSS to discover NEOs also allow us to search for objects that vary on timescales of minutes.

The structure of the CRTS data processing pipeline largely follows the PQ pipeline (A. J. Drake et al. 2009, in preparation). In short, all the CSS catalogs are processed as the images are taken, on site. Candidate OTs are filtered to remove detections associated with artifacts (such as saturation spikes) and moving objects (such as asteroids and satellite trails). After filtering, the transient cutout images, light curves, and associated metadata

<sup>8</sup> <http://earth.google.com/sky>

<sup>9</sup> <http://www.lpl.arizona.edu/css/>

<sup>10</sup> <http://neo.jpl.nasa.gov/stats/>


are posted to VOEventNet where the OTs are posted on webpages and an RSS feeds.<sup>11</sup> Additional image cutouts (from SDSS, PQ, and the DSS) are added to the information about each OT candidate by VOEventNet. As one of the goals of CRTS is to enable rapid automated follow-up of short-timescale transients by robotic telescopes, VOEvent alerts are sent from the observatory as soon as the data is processed (approximately 5 minutes after the fourth observation in a sequence is obtained).

After the filtering process  $\sim 1$  source in 200,000 is detected as a significant OT. All of these candidates are sent as VOEvents. Currently  $\sim 50\%$  of these are associated with clearly genuine OTs while others are mainly detections of coincident image artifacts, such as saturation spikes. The handful of transient candidates are screened by eye within minutes to hours of discovery and an updated list of transients is posted.<sup>12</sup> We expect to remove these image artifacts employing artificial neural networks as we used in the PQ survey.

### 3. RESULTS

In the first six months of the CRTS survey, images were searched covering a total  $\sim 450,000 \text{ deg}^2$  from  $\sim 2000$ ,  $8 \text{ deg}^2$  fields. Over 350 OTs passed selection by eye. These included dozens of newly discovered cataclysmic variables (CVs; Drake et al., 2007b, 2008a, 2008c, 2008e, 2008g; Djorgovski et al. 2008b, 2008d; Glikman et al. 2008), flaring events (Drake et al., 2008d; Djorgovski et al. 2007), nearby high-proper-motion stars, blazars (Mahabal et al. 2008a), asteroids, and comets. In addition, although CRTS has not focused on SNe, many OTs have been confirmed as such, including SN 2007sr (Drake et al. 2007b), a bright SN in the Antennae galaxies. Fainter SNe have also been discovered (SN 2008au, 2008av, 2008al, 2008ba, 2008bb, 2008bm, 2008cg, 2008ck, 2008dc, 2008dd, 2008de, 2008df, 2008dk (Drake et al. 2007a, 2007b, 2008b, 2008d, 2008e, 2008f, 2008g, 2008h; and Mahabal et al. 2008a). Nine known SNe were also independently rediscovered by the CRTS pipeline (SNe 2006tf, 2007nm, 2007no, 2007pu, 2007qv, 2008aq, 2008aw, 2008ax, 2008dk).

Many types of OTs are readily distinguished in a small number of images. For instance, the presence of motion between images is the certain sign a transient is a nearby object. However, there are a number of cases where classification based on a single set of images is difficult. For instance, dwarf novae (DNe) type CVs can produce outbursts as large as 8 mag, rising to an absolute  $V$ -band magnitude of 2 (Harrison et al. 2004). The presence of such large outbursts means that faint CV systems may be undetectable at quiescence. In a similar way, SN can rise more than 20 mag and range between  $-20 < M_V < -16$  at peak brightness. SNe are a common endpoint of stellar evolution and can occur in undetected dwarf galaxies many magnitudes fainter than the SN itself. Although the absolute magnitudes of DNe and SNe are markedly different, in sparsely sampled data, both DNe and SNe associated with faint sources can be difficult to distinguish at discovery. For this reason, transients associated with bright galaxies are more likely to be followed as SNe candidates. Clearly, this kind of selection bias means that many SNe are never followed, and this may also have led to errors in determinations of SN rates. For instance, core collapse SNe (II, Ib/c) and Type Ia SN are well known to occur at different rates depending on galaxy morphology (Mannucci et al. 2008). The



**Figure 1.** Color-color diagram of CRTS transients matching SDSS-DR6 sources. Triangles: likely and confirmed SN. Squares: confirmed and likely dwarf novae. Crosses: all other transients including possible SNe and DNe. The dots show the de-reddened stellar locus.

(A color version of this figure is available in the online journal.)

fact that few SNe are followed in faint galaxies may mean that the types of SNe occurring in these settings are underestimated.


DNe can readily be distinguished from SNe with spectroscopic observations. However, the number of telescopes capable of photometrically observing faint sources far outnumbers those which can spectroscopically monitor such sources. When spectroscopic observations are unavailable, other differences can be investigated. For instance, DNe outbursts typically last from few days to a couple of weeks, while SNe are bright for months. DNe outbursts take 1–3 days to rise to peak brightness while most SNe take 10–20 days. DNe recur on scales of days to years while SNe are one-time events. Both the DNe system and their outbursts are usually very blue while SNe pass through a range of colors as they age (depending on their type and host galaxy extinction). DNe can also be distinguished from SNe using accurate measurement of source color over time (Poznanski et al. 2002). Most CV systems can also be distinguished from variable stars on the giant branch and main-sequence stars in multiband photometry. For instance, from blue ( $g - r$ ) and ( $r - i$ ) colors in SDSS data (Kricinunas et al. 1998).

In Figure 1, we present the de-reddened color-color diagram of all the CRTS transients that match SDSS sources within  $2''.5$ . In this figure, we have separated SNe and DNe candidates with different symbols so that differences may be clearly seen. This figure shows that CVs are generally bluer than the SNe hosts in  $(u-g)$  colours. The  $(u-g)$  colours are consistent with those expected for white dwarfs (Eisenstein et al. 2006) and CVs discovered in SDSS data (Szkody et al. 2007). Since this figure only includes matches within  $2''.5$ , it does not include the SNe discovered which were associated with large nearby galaxies. For these objects, the offsets from the galactic centers are much larger. The color separation of the two populations seen in Figure 1 tells us that if we see a source with  $(u-g) < 0.7$  it is much more likely to be due to a CV than an SN. Indeed of the 213 new and known CVs measured by Szkody et al.

<sup>11</sup> <http://voeventnet.caltech.edu/feeds/Catalina.shtml>

<sup>12</sup> <http://nesssi.cacr.caltech.edu/catalina/AllIns.html>





**Figure 2.** Palomar 200 inch DBMS spectrum of supernova SN 2008ba.  
(A color version of this figure is available in the online journal.)

(2002, 2003, 2004, 2005, 2006, 2007) in SDSS data, only 10 had  $(u - g) > 0.7$ .

### 3.1. Supernovae

SNe are explosive endpoints of stellar evolution. In some cases, these objects can be used as cosmological standard candles that probe the depths of the expanding universe and aid our understanding dark energy. Many groups are currently searching for nearby SNe (LOSS; Filippenko et al. 2001, theSNfactory; Aldering et al. 2002, Chilean Automatic Supernova Search (CHASE); Pignata et al. 2007; Monard 2003; Puckett et al. 2003). Most of these groups only search for low- $z$  SNe clearly associated with large, bright, galaxies (Prieto et al. 2008)<sup>13</sup>. However, the SNfactory uses the wide-field images from the PQ survey to cover thousands of squares degrees (Wood-Vasey et al. 2004), like the CRTS.

In addition to the well known types of transients listed above, the CRTS discovered a large fraction of OTs that were either spatially associated with very faint blue galaxies, or had no visible source associated with them to the limits of PQ and SDSS surveys (greater than 3.5 mag fainter than the discovery images). One well known source of transients associated with galaxies is active galactic nucleus (AGN) flares (Totani et al. 2005). AGNs are known to contaminate SNe surveys (Gal-Yam et al. 2008; Sand et al. 2008) and vary on timescales from hours to years, with outbursts generally being up to  $\sim 1$  mag (Webb & Malkan 2000). We expected to find a number of events from such objects. In contrast, the transients we discovered became many magnitudes brighter than their hosts over a period of 2 weeks and were visible for more than a month, consistent with SNe. We carried out photometric follow-up of a number of these candidates and found that their colours and decline rates were indeed consistent with SNe, rather than AGN, CVs, or other variables stars (Poznanski et al. 2002). The most likely sources of such transients are SNe in very faint host galaxies or associated with intracluster stars.

To select SNe from among CRTS transients, we removed asteroids near stationary points, flare stars, and DNe with faint sources. This was achieved by only selecting transients present at the same location on multiple nights spanning more than 2 weeks. In addition, we required that the objects were at least 2 mag brighter than any possible host, and that the transients faded slowly (as expected for SNe). To investigate the accuracy of the classification based on six months of data, we reviewed archival CSS photometry going back more than three years and found no evidence of contamination by repeated outbursts of variable objects. The SNe candidates are given in Table 1.

Follow-up spectroscopy of two of the CRTS candidates associated with faint galaxies by Djorgovski et al. (2008c) revealed a Type Ia SN (SN 2008ba,  $z \sim 0.03$ ) and a Type II SN (SN 2008bb,  $z \sim 0.03$ ). In Figure 2, we present the spectrum of SN 2008ba. In Figure 3, we show the locations of the SNe in much deeper SDSS  $r$ -band images. SN 2008ba is  $\sim 4''$  from an SDSS-identified galaxy with magnitude  $m_r = 21.5$ . There are no other bright galaxies within  $2'$  of this object. As the redshift of this SN is 0.03, the host galaxy must be fainter than  $M_r \sim -13.9$ . For SN 2008bb, there are also no bright ( $m_r < 19$ ) galaxies within  $2'$  and no galaxies with  $m_R < 23$  within  $0.5'$  (Figure 2, right). Assuming the brightest galaxy missed by SDSS to have  $m_r \sim 23$ , the host would have to be  $M_r > -12.9$  at the SN's redshift (0.03).

In addition to the confirmed SNe, we have discovered dozens of similar, long-timescale OTs. Seventeen of these have no sources present in CSS, PQ, or SDSS images overlapping the discovery locations (to magnitudes  $\sim 21$ – $23$ ). The remaining 24 match with faint galaxies that are  $> 2$  mag fainter than the transients associated with them. In Figure 4, we present an example of an SN where a faint host galaxy is detected, and in Figure 5, an SN where no host is seen to  $r \sim 23$ . The light curves corresponding to these OTs are given in Figure 6.

SNe with faint host galaxies are not unknown. For example, Prieto et al. (2008) studied recent nearby SN and found the SN Ic, 2007bg, to be associated with a faint ( $M_B \sim -12$ ), very metal-poor ( $\sim \frac{1}{20}$  solar) host. Similarly, the PQ experiment discovered a Type Ic SN (2007nm) with an  $m_R \sim 22$  host in DPOSS

<sup>13</sup> <http://www.supernovae.net>

**Table 1**  
Confirmed and Candidate Supernovae

ID	R.A. (h:m:s)	Dec (° ′ ″)	Date MJD	Mag	Mag <sub>H</sub>	Designation	Type
CSS080514:145613+185115	14:56:13.27	+18:51:15.4	54600.18572	18.4	23	2008dk	Ia <sup>a</sup>
CSS080508:112311+341251	11:23:10.90	+34:12:51.0	54594.27706	18.4	16.4	2008cz	Ia
CSS080505:160625+100521	16:06:25.06	+10:05:21.4	54591.37061	18.8	21.8	2008df	...
CSS080505:155415+105825	15:54:15.15	+10:58:25.0	54591.37006	16.7	16.8	2008cg	Ia
CSS080429:132846+251644	13:28:46.47	+25:16:44.2	54585.25061	18.6	21.2	...	...
CSS080427:152242+302208	15:22:42.30	+30:22:08.0	54583.39125	18.5	21.7	2008ck	Ia
CSS080426:084328+302346	08:43:27.86	+30:23:45.6	54582.15121	18.2	>23	...	...
CSS080417:155525+094147	15:55:24.97	+09:41:47.1	54573.37332	16.7	21 <sup>c</sup>	2008de	II
CSS080415:122958+283538	12:29:57.71	+28:35:38.1	54571.16946	19.5	>23	...	...
CSS080414:134142+162022	13:41:42.18	+16:20:21.8	54570.31002	18.4	21 <sup>c</sup>	...	...
CSS080409:170404+213542	17:04:03.60	+21:35:42.0	54565.34357	17.7	20.1	2008dd	...
CSS080406:164729+091826	16:47:29.45	+09:18:25.8	54562.46315	18.1	21 <sup>c</sup>	2008dc	Ib/c
CSS080406:122515+064526	12:25:15.03	+06:45:26.2	54562.35553	18.9	>23	...	...
CSS080405:155205+000627	15:52:04.82	+00:06:27.0	54561.46586	18.7	21.2	...	...
CSS080404:130414+101913	13:04:14.11	+10:19:12.9	54560.34851	15.1	11.8	2008aw	II
CSS080404:125030+105201	12:50:30.36	+10:52:01.3	54560.34718	15.9	13.1	2008aq	IIf
CSS080403:123041+413816	12:30:40.80	+41:38:16.1	54559.20966	13.3	9.4	2008ax	IIn
CSS080401:092227+282421	09:22:27.05	+28:24:20.8	54557.22644	18.7	23*	...	...
CSS080330:121539+033613	12:15:39.10	+03:36:12.9	54555.30168	18.8	20.6	...	...
CSS080330:124441+051438	12:44:41.47	+05:14:37.6	54555.30276	19.2	21.4	...	...
CSS080329:130259+103027	13:02:58.75	+10:30:27.0	54554.30700	17.2	15.8	2008bm	IIn
CSS080313:132560+020027	13:25:59.69	+02:00:27.1	54538.36493	18.6	22.1	...	...
CSS080312:140609+242013	14:06:08.61	+24:20:12.5	54537.39966	19.4	>23	...	...
CSS080312:102245+021753	10:22:44.65	+02:17:52.5	54537.27925	18.9	21.7	...	...
CSS080310:112544+182317	11:25:44.03	+18:23:17.1	54535.36676	20.0	>23	...	...
CSS080308:090908+111115	09:09:07.99	+11:11:15.3	54533.20757	18.4	21.2	...	...
CSS080303:081413+271939	08:14:13.15	+27:19:39.0	54528.11129	18.1	23*	...	...
CSS080303:075520+203908	07:55:19.52	+20:39:08.4	54528.10484	19.3	>23	2008bb	II
CSS080302:124132+332203	12:41:31.98	+33:22:03.2	54527.29264	19.2	22.5*	...	...
CSS080302:145726+232348	14:57:26.21	+23:23:48.2	54527.47633	18.0	>22	2008av	...
CSS080228:142427+063345	14:24:26.63	+06:33:44.8	54524.48530	18.3	21*	...	...
CSS080228:105301+075656	10:53:00.81	+07:56:55.9	54524.30812	19.2	>22	...	...
CSS080227:160549+192717	16:05:49.17	+19:27:17.2	54523.47442	17.8	22	2008ba	Ia
CSS080227:113034+130905	11:30:33.96	+13:09:05.3	54523.31151	18.7	>23	2008au	...
CSS080219:151457+234110	15:14:56.75	+23:41:10.1	54515.45977	16.6	>23	...	...
CSS080218:134903+315237	13:49:03.35	+31:52:37.1	54514.41836	19.0	21.8	...	...
CSS080213:024607+073834	02:46:07.23	+07:38:33.8	54509.09948	17.3	14.4	2008al	II
CSS080210:080306+115121	08:03:06.09	+11:51:20.8	54506.19556	18.8	>23	...	...
CSS080206:130745+060805	13:07:44.64	+06:08:05.0	54502.43974	19.8	>23	...	...
CSS080111:070447+412518	07:04:47.34	+41:25:18.0	54476.25764	19.0	>21	...	...
CSS080111:145436+051254	14:54:35.97	+05:12:53.8	54476.49167	18.1	21*	...	...
CSS080101:035151+062700	03:51:50.64	+06:27:00.5	54466.25782	18.3	>23	...	...
CSS080101:085249+052247	08:52:48.90	+05:22:46.8	54466.37420	19.1	20.5	...	...
CSS071230:082550+220041	08:25:50.48	+22:00:41.3	54464.35765	17.6	21.3	...	...
CSS071230:081807+200719	08:18:07.05	+20:07:19.2	54464.32780	18.1	22	...	...
CSS071219:223508+011160	22:35:08.45	+01:11:59.6	54453.08583	18.3	>23	2007qv	II
CSS071218:120153+185822	12:01:52.80	+18:58:21.7	54452.53386	12.9	9.8	2007sr	Ia
CSS071216:121630+102303	12:16:29.63	+10:23:02.9	54450.44569	18.9	21.5	...	...
CSS071206:102130+200007	10:21:30.15	+20:00:07.4	54440.42300	18.6	21 <sup>c</sup>	...	...
CSS071204:035956+233018	03:59:56.34	+23:30:18.2	54438.21532	18.5	>22	...	...
CSS071204:002043+102044	00:20:42.71	+10:20:44.1	54438.13146	18.2	>22	...	...
CSS071117:224558+003854	22:45:58.17	+00:38:54.3	54421.10263	18.6	20.7	2007pu	Ia
CSS071112:012312+120818	01:23:12.27	+12:08:18.0	54416.17262	19.0	22 <sup>c</sup>	...	...
CSS071111:095436+045612	09:54:35.98	+04:56:11.8	54415.46121	19.0	22.6	...	...
CSS071109:003334+214127	00:33:34.36	+21:41:26.7	54413.27788	18.8	16.1	...	...
CSS071107:093912+301341	09:39:11.62	+30:13:41.5	54411.43569	18.2	20.0	...	...
CSS071103:224527+103933	22:45:27.41	+10:39:32.7	54407.18815	17.8	>21	2007nm	Ia
CSS071102:024106+033814	02:41:05.71	+03:38:13.6	54406.26445	19.6	>22	2007no	Ia
CSS071102:024457+044918	02:44:57.22	+04:49:17.7	54406.26445	18.8	>22	...	...
CSS071101:015607+203404	01:56:06.60	+20:34:03.5	54405.26126	18.8	>22	...	...
CSS070320:124616+112555	12:46:15.81	+11:25:55.4	54179.37725	17.6	20.8	2006tf	IIn <sup>b</sup>

**Notes.** Column 1, CSS ID; Columns 2 and 3, right ascension and declination J2000; Column 4, modified Julian date of detection; Column 5, detection magnitude; Column 6, approximate source galaxy Gunn *r* magnitude; Column 7, IAU designation.


<sup>a</sup> Sand et al. (2008, <http://www.cfa.harvard.edu/iau/cbet/001400/CBET001410.txt>). However, first announced and spectroscopically confirmed by the Supernova Factory as SNF20080510–001.

<sup>b</sup> Found in archival data.


<sup>c</sup> Estimated *r*-band magnitude.

images and redshift  $z = 0.04$ . At this redshift, the host galaxy would have to be  $M_R > -13.8$ . Another possibility is that SNe without visible hosts could be associated with intracluster

stars. Past searches for such SNe have been carried out by Sand et al. (2008) and Gal-Yam et al. (2008). As part of the WOOTs survey, Gal-Yam et al. (2003, 2008) discovered 12 SNe in Abell



**Figure 3.** Left: SDSS predisccovery Gunn  $r$ -band image of the location of SN2008ba. The circle shows the location of the probable host, while the cross marks the location of the supernova. Right: SDSS predisccovery Gunn  $r$ -band image of the location of SN2008bb. The box shows the location of the SN, while the circle shows the location of the nearest SDSS galaxy.



**Figure 4.** Left: 2007 December 30 CSS discovery image, showing the location of OT CSS071230:082550+220041 (circle). Right: SDSS Gunn  $r$  image, showing the location of CSS071230:082550+220041 and the corresponding faint galaxy host.

clusters selected according to  $0.06 < z < 0.2$ . Two of these were identified as being intracluster SN Ia (SN 1998fc and SN 2001al). They found upper limits to possible dwarf hosts of  $M_r > -14$  and  $-11.8$ , respectively. Sand et al. (2008) discovered four photometrically selected intracluster SN with hosts  $M_g > -14.3$ . Based on this, they found 20% of stellar mass may reside in intracluster stars and thus 20% of the SN Ia parent stellar population could be intergalactic.

Interestingly, Stanek et al. (2006) and Savaglio et al. (2008) found that the long-duration GRBs associated with SNe are in faint, metal-poor, star-forming dwarf galaxies. This result was supported by Modjaz et al. (2008) who found that the galaxies containing Type Ic SNe that were not associated with GRBs, were significantly more metal rich than those with associated GRBs. Savaglio et al. (2008) also found that the host galaxies associated with GRBs are low-mass star-forming objects. The rate of SNe in such low-mass galaxies is expected to be low, as Kauffmann et al. (2003) found that only  $\sim 13\%$  of the stellar mass in galaxies is found in objects less massive than  $10^9 M_\odot$ . Such galaxies have only tens of millions of stars in contrast to large galaxies like our own with hundreds of billions of stars. For the hosts of SNe candidates not to be visible, most of these must be associated with low stellar-mass galaxies.


### 3.1.1. Supernovae with Faint Galaxy Hosts?

In order to test whether some of our SNe candidates are associated with galaxy clusters, we compared their locations to Abell clusters (Abell et al. 1989) and DPOSS clusters (Gal-Yam et al. 2003). Two of the 48 SNe in the DPOSS catalog region (with Galactic latitude  $b > 30^\circ$ ) matched the locations of DPOSS clusters and two matched Abell clusters. With only four matches to 12,000 clusters it appears very unlikely that the SNe are associated with such clusters. The SNfactory, like CRTS, uses wide-field images to find SNe rather than concentrating on bright galaxies like most other nearby SNe surveys. Therefore, both surveys have similar spatial sensitivity. We compared our results with 95 confirmed SNe from the Supernova Factory.<sup>14</sup> Once again only two SNfactory SNe of the 63 SNe with  $b > 30^\circ$  matched DPOSS clusters and two matched Abell clusters.


To determine whether the galaxies associated with our SNe candidates were indeed low-luminosity hosts, we calculated the difference between the peak measured magnitude in CRTS data and the host  $R$ -band magnitude. To provide upper limits to the host galaxies brightness where no host was visible,

<sup>14</sup> [http://snfactory.lbl.gov/snf/open\\_access/login.php](http://snfactory.lbl.gov/snf/open_access/login.php)





**Figure 5.** Left: 2008 January 1 CSS image, showing the location of OT CSS080219:151457+234110 (circle). Right: SDSS Gunn  $r$  image of the same field. At the location of CSS080219:151457+234110, no host galaxy is visible.




**Figure 6.** Left: the CRTS light curve of object CSS071230:082550+220041. Right: the CRTS light curve of object CSS080219:151457+234110. The black points are measurements and blue triangles nondetection upper limits.

(A color version of this figure is available in the online journal.)

the hosts were assumed to reside at the limits of the deepest observations covering them. Next, for the SNe with known spectral types (hence known peak absolute magnitude), we added the CRTS magnitude to the peak expected for the type of SNe. This provides a rough estimate of the host's absolute  $R$ -band magnitude. To test the SNe hosts where we do not have known spectral types we carried out the same process and assumed the SNe were all the brightest, common SN type, Ia. As the actual SNe types should be the same or fainter than our assumption, we obtain an approximate upper limit to the brightness of the associated host galaxies. For comparison, we took the SNfactory SNe and matched them with SDSS DR6 to find their host magnitudes. Fifty four lie within the region covered by SDSS DR6 and 52 match SDSS objects. Twenty nine of these hosts have spectra from which the redshift has been determined. For the other 23 SNe, we used the redshift of the SNe given by the SNfactory webpage. We then used the SDSS-DR6  $r$ -band magnitudes and redshifts to derive the absolute magnitudes of the host galaxies assuming  $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$ .

In Figure 7, we plot histograms of the host galaxy magnitudes, for those with confirmed types and these combined with candidates (assumed Type Ia) as well as those with host from the SNfactory. This figure strongly suggests that the host galaxies must be intrinsically faint. The spectroscopically confirmed SNe, that provide the most accurate host galaxy magnitudes, clearly reside in very faint hosts. The fact that the unconfirmed SN–host distribution peaks at a brighter magnitude is expected because of our assumption that these SNe were all bright Type Ia, rather than a mixture of types. Also, it is apparent that the SNfactory finds SNe in intrinsically brighter galaxies.

It is clear that by only selecting transients that are 2 mag brighter than the sources associated with them, we will introduce a bias toward discovering SNe in faint galaxies, opposite to SNe surveys that have chosen only to follow bright galaxies. However, the CRTS SNe can also be discovered in large galaxies when they rise more than 2 mag above the local background. For intermediate magnitude hosts, the incremental increase in brightness is insufficient for SNe to be detected as separate sources. Detection of these would require the use of image



**Figure 7.** Histograms of inferred host galaxy magnitudes. Solid line: all CRTS host galaxy magnitudes. Dashed line: host galaxy magnitudes for CRTS SNe with known types. Dotted line: host galaxy magnitudes for SNe discovered and confirmed by the SNfactory. See the text for details. The numbers in SNE histogram are offset by +0.1 for clarity.

(A color version of this figure is available in the online journal.)


subtraction. As the host galaxies discovered by CRTS are generally much fainter than the SNe they do not add significant flux to follow-up spectra, making confirmation easier.

Two additional possibilities might explain some of the observed SNe candidates. One possibility is that some of the SNe discovered may be brighter than SN Ia. There is increasing evidence for such rare types of bright SNe (Miller et al. 2009). An example of such an event is 2006gy, that had a peak luminosity of  $M_R \sim -22$  (Smith et al. 2007). Based on the small number of discoveries these SNe must be extremely scarce. A second possibility is that some of the candidates are not SNe, but rather part of a rare group of variable stars. Based on the discovery rate such hypothetical variable stars would have to be less common than DNe and fainter on average. Spectroscopic evidence suggests that many SNe occur in faint hosts, but cannot rule out other possibilities for all candidates.

One of the SNe discovered by CRTS (SN 2007sr, Drake et al. 2007b) resides in NGC 4089/39, the Antennae. This galaxy is sufficiently nearby ( $22 \pm 3$  Mpc; Schweizer et al. 2008) that Cepheid variables can be discovered using the *Hubble Space Telescope* (HST). The HST Key Project (Freedman et al. 2001) used Cepheid distances to measure the Hubble constant. However, only eight Type Ia SN, occurring between 1937 and 1998, are known in galaxies where Cepheid distances were measured by the HST Key Project. The combination of a Cepheid distance determination to the Antennae along with well-sampled light curves of SN 2007sr can thus play an important role in independently calibrating the cosmological distance scale.

### 3.2. Cataclysmic Variables

In this survey, we discovered a number of OTs that have abrupt outbursts that lasted for less than a couple of weeks. These are mainly due to DNe type CVs. Longer outbursts can



**Figure 8.** Palomar 200 inch DBMS spectrum of CV eclipsing CV CSS080227:112634-100210. The Balmer emission lines are clearly seen as well as strong helium lines.

(A color version of this figure is available in the online journal.)

occur for DNe that undergo superbursts and standstills. These outbursts can easily be distinguished from classical novae within the Galaxy because such events have an extreme peak brightness  $M_V = -7.5$ . Novae are much more common than SNe, but 10 or more magnitudes fainter. Novae are much less common than CV outbursts and less luminous than the host galaxies mentioned in the previous section, so few are expected in CRTS data.

The brightest outbursts of CVs are closely followed by a large group of astronomers with modest sized telescopes. For this reason, most CV follow-up concentrates on objects brighter than  $V = 17$  during outburst. As LSST images will saturate at 16th magnitude, the CV outbursts discovered by CRTS are much better suited for follow-up by large numbers of small telescopes.

The CRTS transient pipeline has initially been tuned to discover objects brightening by 2 or more magnitudes between a catalog magnitude and subsequent observations. This threshold was chosen to discover highly variable sources while suppressing noise and well known types of low-amplitude variable stars. However, this threshold makes CRTS ideal for the detection of DN type CVs (which typically brighten by 2–8 mag). In the first six months, the CRTS discovered 64 new DNe. All the CV candidates discovered in the CRTS data are presented in Table 2. The spectrum of newly discovered eclipsing CV CSS080227:112634-100210 is given in Figure 8.

In recent years, many bright CVs have been identified in SDSS data from the spectra (Szkody et al. 2002, 2003, 2004, 2005, 2006, 2007). To date, the SDSS surveys have taken more than 1 million spectra.<sup>15</sup> However, this is less than 0.5% of the objects observed photometrically and limited to objects brighter than  $i = 20.1$ . Thus, most CVs residing in areas covered by SDSS will not have been identified. Indeed, a large fraction of the CVs presented in Table 2 reside within regions observed by SDSS. As expected, most of these sources clearly match blue point sources in the SDSS data. Transient CSS080427:124418+300401 matches SDSS star J124417.89+300401.0 and has a spectrum that clearly exhibits the strong emission lines associated with CVs but has not previously been identified as a CV by the SDSS team. However, many of the DNe we discovered (without spectra) are noted

<sup>15</sup> <http://www.sdss.org>



**Table 2**  
Cataclysmic Variable Candidates

ID	R.A. (h:m:s)	Dec ( $^{\circ}$ ' ")	Date <i>MJD</i>	Mag	Mag <sub>Q</sub>	FUV	NUV
CSS080514:162606+225044	16:26:05.71	+22:50:44.4	54600.36523	18.5	22.5	...	...
CSS080514:151021+182303	15:10:20.74	+18:23:02.5	54600.22210	17.8	20.7	...	...
CSS080513:164002+073208	16:40:02.11	+07:32:07.8	54599.31544	18.0	21 <sup>a</sup>	21.4	...
CSS080512:064608+403305	06:46:08.23	+40:33:05.1	54598.14377	16.2	21 <sup>a</sup>	...	22.7
CSS080512:173860+344023	17:38:59.82	+34:40:23.1	54598.35924	18.7	>21	...	...
CSS080511:212555-032406	21:25:55.07	-03:24:05.9	54597.43009	17.8	>21	...	23.3
CSS080506:085409+201339	08:54:09.41	+20:13:39.2	54592.15678	17.7	20.5	21.5	20.3
CSS080505:105836+120049	10:58:35.93	+12:00:48.5	54591.20280	18.4	>23	...	... <sup>b</sup>
CSS080505:163121+103134	16:31:20.89	+10:31:33.9	54591.37168	14.2	19.0	...	...
CSS080505:214804+080951	21:48:04.40	+08:09:50.5	54591.43395	18.3	20.5	22.1	21.6
CSS080502:141002-124809	14:10:02.21	-12:48:08.7	54588.32033	16.2	21 <sup>a</sup>	...	...
CSS080501:223058+210147	22:30:58.32	+21:01:47.0	54587.46701	16.7	21 <sup>a</sup>	20.9	21.4
CSS080428:162502+390926	16:25:01.72	+39:09:26.3	54584.28928	13.5	17.1	18.2	17.9 <sup>c</sup>
CSS080428:163805+083758	16:38:05.39	+08:37:58.5	54584.43007	15.4	19.0	21.2	20.8 <sup>d</sup>
CSS080428:160524+060816	16:05:24.14	+06:08:15.8	54584.39153	19.4	22.5	...	...
CSS080427:153634+332852	15:36:34.41	+33:28:52.1	54583.41750	15.2	19.0	...	... <sup>e</sup>
CSS080427:124418+300401	12:44:17.88	+30:04:01.2	54583.32956	15.0	18.5	18.7	18.2
CSS080427:131626-151313	13:16:25.68	-15:13:13.5	54583.23364	16.9	20 <sup>a</sup>	20.6	20.6
CSS080426:162657-002549	16:26:56.80	-00:25:48.6	54582.41287	18.0	22.2	...	...
CSS080425:141712-180328	14:17:11.98	-18:03:27.7	54581.33298	15.0	20.5	...	...
CSS080425:143143-032520	14:31:43.05	-03:25:20.5	54581.41153	18.5	>21	...	...
CSS080424:160232+161732	16:02:32.15	+16:17:32.5	54580.44400	18.4	21.8	...	...
CSS080424:155326+114437	15:53:25.67	+11:44:36.8	54580.44238	18.2	23.4	...	...
CSS080417:081543-004209	08:15:43.16	-00:42:08.6	54573.11929	17.8	21.8	...	...
CSS080416:080854+355053	08:08:53.73	+35:50:53.3	54572.15297	16.4	19.6	19.8	20.0
CSS080415:162012+115257	16:20:12.00	+11:52:57.1	54571.41746	19.3	22.3	...	...
CSS080411:154258-020705	15:42:58.36	-02:07:04.8	54567.39688	16.8	23	...	...
CSS080409:174714+150048	17:47:14.34	+15:00:47.7	54565.37312	17.0	21 <sup>a</sup>	...	...
CSS080409:081419-005022	08:14:18.90	-00:50:22.1	54565.14409	14.8	19.0	19.0	19.0
CSS080406:075648+305805	07:56:48.02	+30:58:05.4	54562.16871	17.0	21 <sup>a</sup>	21.3	21.6
CSS080405:161851-052509	16:18:50.76	-05:25:08.6	54561.43750	16.1	21 <sup>a</sup>	...	...
CSS080404:213309+155004	21:33:09.43	+15:50:04.3	54560.50262	15.7	>21	18.8	19.0
CSS080403:160004+331114	16:00:03.71	+33:11:13.9	54559.36671	15.2	19.5	15.8	16.0 <sup>f</sup>
CSS080401:085113+344449	08:51:13.43	+34:44:48.7	54557.18634	16.4	20	...	...
CSS080401:153151+152447	15:31:50.80	+15:24:46.8	54557.43700	18.9	22.6	...	...
CSS080331:160205+031632	16:02:04.80	+03:16:31.8	54556.32329	17.1	22.8	...	...
CSS080329:143500-004606	14:35:00.23	-00:46:06.3	54554.41592	15.0	18.4	18.7	18.6 <sup>g</sup>
CSS080324:122060-102735	12:20:59.77	-10:27:35.1	54549.34105	16.8	19.7	18.3	18.6
CSS080309:084358+425037	08:43:58.06	+42:50:37.2	54534.25796	18.1	19.9	21.0	20.0
CSS080309:070501+372505	07:05:01.06	+37:25:05.2	54534.18927	18.1	>22	...	22.7
CSS080307:090624-085141	09:06:23.52	-08:51:40.5	54532.17312	16.5	>21	...	...
CSS080306:082655-000733	08:26:54.68	-00:07:32.9	54531.19113	16.4	19.5	...	... <sup>h</sup>
CSS080305:102938+414046	10:29:37.71	+41:40:46.4	54530.30353	17.3	22.3	...	...
CSS080305:080846+313106	08:08:46.19	+31:31:06.1	54530.12195	14.8	18.7	19.4	19.9 <sup>i</sup>
CSS080304:164002+073208	16:40:02.14	+07:32:07.6	54529.44911	17.9	>21	21.4	...
CSS080304:090240+052501	09:02:39.70	+05:25:00.6	54529.20407	16.2	23.1	...	23.7 <sup>j</sup>
CSS080303:073921+222454	07:39:21.16	+22:24:54.0	54528.10431	18.5	22.4	...	...
CSS080302:160845+220610	16:08:44.79	+22:06:10.0	54527.50294	17.9	21.0	...	...
CSS080227:132103+015329	13:21:03.18	+01:53:29.2	54523.43337	14.1	19.2	...	... <sup>k</sup>
CSS080227:112634-100210	11:26:33.98	-10:02:10.1	54523.35480	16.2	18.9	...	20.2 <sup>l</sup>
CSS080211:024602+345508	02:46:02.38	+34:55:08.2	54507.11196	15.0	19	...	... <sup>m</sup>
CSS080209:090628+052657	09:06:28.25	+05:26:56.9	54505.30194	15.6	18.5	16.4	16.2 <sup>n</sup>
CSS080209:084555+033929	08:45:55.05	+03:39:29.3	54505.26151	18.5	20.9	...	... <sup>o</sup>
CSS080208:103317+072119	10:33:17.25	+07:21:18.7	54504.37302	15.5	19.8	20.2	20.0
CSS080207:060038-080945	06:00:37.59	-08:09:44.7	54503.20131	16.3	19.8	...	...
CSS080207:052034-000530	05:20:33.85	-00:05:30.1	54503.20510	16.8	19.7	...	...
CSS080202:081415+080450	08:14:14.92	+08:04:50.2	54498.23365	18.5	22	...	...
CSS080202:084608+011743	08:46:07.73	+01:17:42.9	54498.23741	18.2	22.5	23.4	...
CSS080201:115330+315836	11:53:30.23	+31:58:36.0	54497.33687	17.5	19.9	...	...
CSS080131:163943+122414	16:39:42.70	+12:24:14.4	54496.51670	17.4	19.2	...	...
CSS080130:033056+251255	03:30:55.66	+25:12:55.2	54495.15831	16.5	21 <sup>a</sup>	...	...
CSS080130:090951+184947	09:09:50.52	+18:49:47.1	54495.31194	12.6	15.7	17.0	17.0 <sup>p</sup>
CSS080130:021110+171624	02:11:10.22	+17:16:24.3	54495.13080	14.3	19.0	19.8	19.8
CSS080130:105550+095621	10:55:50.08	+09:56:20.5	54495.44905	15.4	18.5	...	...
CSS080110:032627+070744	03:26:27.26	+07:07:44.4	54475.19676	17.7	21	...	...

**Table 2**  
(Continued)

ID	R.A. (h:m:s)	Dec (° ' ")	Date <i>MJD</i>	Mag	Mag <sub>Q</sub>	FUV	NUV
CSS080101:232519–081819	23:25:19.20	−08:18:18.8	54466.08479	14.9	18.6	19.9	19.3 <sup>q</sup>
CSS071231:082822+105344	08:28:21.75	+10:53:44.5	54465.38067	17.4	22.2	22.0	...
CSS071215:041456+215643	04:14:55.71	+21:56:43.1	54449.23934	17.1	21 <sup>a</sup>	...	...
CSS071214:090904+091714	09:09:04.38	+09:17:13.5	54448.44999	16.9	22	...	...
CSS071116:214843–000723	21:48:42.53	−00:07:23.5	54420.09361	15.8	22.4	23.0	23.3
CSS071115:172406+411410	17:24:06.32	+41:14:10.1	54419.06981	17.1	18.5	19.7	19.1 <sup>r</sup>
CSS071115:044216–002334	04:42:16.03	−00:23:33.9	54419.28884	17.0	22 <sup>a</sup>	...	21.9
CSS071112:024457+352249	02:44:57.42	+35:22:49.3	54416.20408	18.4	21 <sup>a</sup>	...	...
CSS071112:085823–003729	08:58:22.86	−00:37:29.0	54416.42038	17.5	22.0	...	...
CSS071106:075747+305307	07:57:46.97	+30:53:07.2	54410.40003	18.5	>23	...	...

**Notes.** Column 1, CSS ID; Columns 2 and 3, right ascension and declination J2000; Column 4 modified Julian date of detection; Column 5 Catalina detection magnitude; Column 6 approximate Gunn *r*-band quiescent magnitude; Columns 7 and 8, *GALEX*, FUV, and NUV magnitudes.

<sup>a</sup> Estimated *r*-band magnitude.

<sup>b</sup> Mahabal et al. (2008a), Atel 1520.

<sup>c</sup> V844 Her.

<sup>d</sup> V544 Her.

<sup>e</sup> SDSS J153634+332851.

<sup>f</sup> VW CrB.

<sup>g</sup> OU Vir.

<sup>h</sup> SDSS QSO candidate J082654.69–000733.1 (Richards et al. 2004).

<sup>i</sup> Cnc SDSS J080846+313106.

<sup>j</sup> Djorgovski et al. (2008d, 2008b), Atel 1418 and 1411.

<sup>k</sup> HV-Vir.

<sup>l</sup> Eclipsing system.

<sup>m</sup> Antipin V80.

<sup>n</sup> Hya SDSS J090628+052656.

<sup>o</sup> First appeared in CBAT Jan 20th.

<sup>p</sup> GY Cnc.

<sup>q</sup> EG Aqr.

<sup>r</sup> V1007 Her.

as SDSS QSO targets. These were excluded in the main SDSS QSO survey (Richards et al. 2002), but were later included in the faint SDSS QSO extension survey. Clearly, the reason that these are QSO candidates is simply that CVs and QSOs overlap in the color–color space of SDSS images. Since CVs are strong UV sources, we matched DNe candidates within the far ultraviolet and near ultraviolet observations of *GALEX*. White dwarfs and QSOs also overlap when the *GALEX* UV observations are combined with SDSS observations (Bianchi et al. 2007). The detection of these objects in *GALEX* FUV and NUV data provides additional evidence that these are CVs. At the current CRTS CV discovery rate, using archival CSS data, we expect to find ~300 new CVs. In comparison, Szkody et al. (2002, 2003, 2004, 2005, 2006, 2007) discovered 177 new CVs in six years of SDSS data, while Rykoff et al. (2005) discovered four CVs in ROTSE-III data taken between 2003 September and 2005 March.

In this survey, we selected CVs by matching the object locations with SDSS-DR6 data (Adelman-McCarthy et al. 2008). Most CVs overlapping SDSS clearly match blue objects marked as stars in SDSS observations. For those CV candidates not covered by SDSS-DR6 data, we checked DSS1, DSS2, and PQ Johnson and Gunn images looking for prior outbursts. For CV candidates without SDSS spectra, matches with blue sources, or evidence of prior outbursts in archival images, we required that the CRTS detection was made at the same location on more than one night. We also required that there should be additional photometry showing that the object had returned to

quiescence within a few weeks. In a small number of cases, the CV locations were covered by DSS as well as SDSS or PQ data, yet no sources were present in any of the past images. These objects were classified based on their light curves.


Many different types of CV systems exist. SU UMa type CVs exhibit superbursts of up to ~8 mag. Among SU UMa CVs, the WZ Sge subtypes have large outbursts with intervals between bursts of up to decades. In order to alert astronomers of outbursts, we initially posted alerts for new discoveries in Astronomer’s Telegrams (ATels).<sup>16</sup> As the classification process progressed, we posted candidates to the CVNet group.<sup>17</sup> However, many of the CVs were also independently followed by members of VSNET<sup>18</sup> within hours based on the real-time alerts posted on the VOEventNet webpages. Subsequent observations of some of the CVs by members of these groups showed the presence of superhumps (seen in SU UMa type systems). The timing of these humps gives the orbital period of the systems. In addition, some of the CVs we discovered were later found to exhibit outbursts in archived ASAS-3, DSS, and PQ images.

Recently, Rau et al. (2007) undertook follow-up observations of four OTs discovered in two wide-field surveys. Of these one was discovered to be an asteroid and the other three were CVs. In addition, they determined limits on the number of faint CVs expected at high Galactic latitude. Based on their model,

<sup>16</sup> <http://www.astronomersteletgram.org>

<sup>17</sup> <http://tech.groups.yahoo.com/group/cvnet-discussion/>

<sup>18</sup> <http://ooruri.kusastro.kyoto-u.ac.jp/pipermail/vsnet-alert/>



**Figure 9.** CSS images of object CSS080502:090812+060421 in and out of eclipse. The baseline CSS magnitude is  $17.0 \pm 0.1$ , while the individual CSS image magnitudes are  $18.72 \pm 0.09$ ,  $18.91 \pm 0.10$ ,  $17.23 \pm 0.04$ , and  $17.15 \pm 0.04$ . This system matches SDSS object J090812.04+060421.2 with magnitudes  $u = 17.34$ ,  $g = 17.08$ ,  $r = 17.28$ ,  $i = 17.08$ , and  $z = 16.69$ . This object was found to be a DA white dwarf of subtype 2.8, plus an M3 dwarf by Silvestri, et al. (2006) also matches 2MASS09081205+0604211 (mags:  $J = 15.5$ ,  $H = 14.9$ ,  $K = 14.7$ ) and GALEX source J090812.0+060420 (FUV = 17.1 and NUV = 17.2).

they predicted that CVs with quiescent magnitudes fainter than  $R = 22$  should not be discovered at Galactic latitude  $|b| > 45^\circ$ . In our survey, we have so far found 2 of the 16 CVs fainter than  $R = 22$  at Galactic latitudes above this.

### 3.3. Eclipses

Apart from searches for objects that newly appeared in a sequence of CSS images, or varied on average from catalog magnitudes, we searched for intrasequence variability. That is, objects with real brightness variations within the 10 minutes span between each image in a sequence of four. This search revealed a number of candidates that dropped by more than a magnitude from the baseline brightness between observations.

These transients are of some interest as the eclipses observed for stellar binary systems are typically much less than a magnitude. Larger eclipses require that the eclipsed star be brighter and smaller than the eclipsing star. As brighter stars are generally larger, they are not fully eclipsed by their companions. One of the exceptions to this situation occurs for white dwarf stars. These objects are much smaller than main-sequence stars of a similar bolometric luminosity. Thus, large eclipses can occur when they are eclipsed by late type dwarf stars. We found SDSS spectra corresponding to two of these objects which clearly show the presence of white dwarf–M dwarf binary systems. In addition, a couple of objects were found to exhibit outbursts and flickering. These are eclipsing CV systems where unlike white dwarf binary systems the drop can be due to the eclipse of an accretion hot spot (Brady et al. 2008).

Eclipses of stars are an effect that can be used to constrain the radii of the eclipsing objects. Steinfadt et al. (2008) recently discovered a partially eclipsing white dwarf (SDSS J143547.87+373338.5) and used it to determine the characteristics of the M dwarf. When secondary eclipses are observed in partially eclipsing binary systems, it is possible to constrain their orbital inclination, eccentricity, and orientation. When no secondary eclipse is seen, the cause of this may be a combination of inclination, orientation, and orbital eccentricity, or the low relative luminosity of the eclipsed star. The existence of full


eclipses provides a strong constraint on the inclination of such systems, whereas the depth and length of the eclipse constrain the relative effective temperatures and the radii. Variations in luminosity during a full eclipse can also be used to map surface features of an eclipsed star, such as spots. For white dwarf binaries, more fully eclipsing systems are expected than partially eclipsing systems, simply because radii of other types of stars are  $>10$  larger than white dwarfs (López-Morales et al. 2007).

Two clear WD–MD binary systems with eclipses  $>1.7$  mag in depth were discovered. In Figure 9, the WD–MD binary system CSS080502:090812+060421 is presented. This is a known WD–MD binary (Silvestri et al. 2006), while CSS080408:142355+240925 is a newly discovered system. In both cases, the M dwarf can be seen in the SDSS spectra and its flux is detected in 2MASS data. The 2MASS magnitudes and colours consistent with M dwarf companions (Wachter et al. 2004). Using three years of archival CSS data, we determined the period of CSS080502:090812+060421 to be 3.58652 hr. In Figure 10, we present its spectrum and phased light curve. The shape of the light curve outside eclipse suggests that the M dwarf is tidally distorted (Drake et al. 2003). The presence of weak  $H_\alpha$  emission in the spectrum suggests that this is probably a pre-CV system.

### 3.4. UV Ceti Variables

One goal of the CRTS is the discovery of transients that may brighten and fade on timescales of minutes. In the course of the analysis, we discovered a number of OTs by searching for significant variations (greater than 2 mag) in two images within the sequence of four images (taken over a span of 30 minutes). In some cases, the source of the events could be seen in the co-added CSS catalog images, although in other cases there was no apparent source. As expected, inspecting SDSS-DR6 images at the locations of these events revealed clear matches with faint M dwarf stars. Thirteen of these events were clearly due to flare stars. An example of this is the “Lynx OT,” the first announced CSS transient (Christensen 2004; Djorgovski et al. 2004). These events are simply flares exhibited by UV Ceti variables. These flares were discovered at Galactic latitudes





**Figure 10.** Left: the five-pixel smoothed SDSS spectrum of CSS080502:090812+060421. The inset shows an expanded view of the M dwarf TiO features. Right: the phased light curve of CSS080502:090812+060421 with period  $3.58652 \pm 0.00002$  hr.

ranging from  $-60^\circ < b < 70^\circ$ . The largest observed flare was 5.5 mag brighter than an associated SDSS source. Aside from stellar flares associated with M dwarfs, there were transients showing repeated large variations. These variations are thus possibly related to active M dwarf stars. The CRTS is in an excellent position to resolve the frequency and general nature of UV Ceti flares, as the timescales of flaring events are tens of minutes, and the CSS survey takes four images separated by 30 minutes.

Past OT surveys have also discovered flare stars. For instance, the DLS discovered three faint short-timescale ( $\sim 1000$  s) transients during the course of their three year survey (Becker et al. 2004). One of these transients was spectroscopically identified as a flaring dM4 star. The other two were later identified as flare stars by Kulkarni & Rau (2006). These authors noted the prevalence of flare star detections in past transient surveys. Additionally, they noted that the presence of a large “fog” of such stars would hamper future efforts to discover fast transients. From the transients discovered in this survey, it is clear that separation of flare events from other fast transients can easily be achieved with observations taken on timescales of flaring events. That is to say, there are distinctive brightness variations between images. In this study, classification is further aided by the presence of much deeper photometry (from PQ, SDSS DR6) where the M dwarf stars can be clearly identified. Future large surveys such as LSST will go much deeper than SDSS (Ivezic et al. 2008). These deep experiments will have difficulty detecting the faint flaring sources without a significant amount of deeper follow-up. However, it is also clear from these results that large flaring events lasting more than 10 minutes are much less common than large outbursts of DNe.

Recently, Welsh et al. (2007) found 52 flares associated with 49 flare stars in *GALEX* data. They discovered two kinds of flare events, namely events lasting  $< 500$  s with quasi-exponential decays and events lasting longer than 500 s with secondary emission peaks. Thirty nine of the *GALEX* flares were greater than 2 mag. This is three times the number of CRTS discoveries, yet the *GALEX* coverage surveyed (in square degree seconds) is only 20% of the CRTS data analyzed here. However, 27 of the *GALEX* flares lasted  $< 500$  s. Such events would appear as a single point and thus not be discovered in CRTS data. In

addition, *GALEX*’s observations reach much fainter magnitudes ( $m_{\text{NUV}} \sim 21$ ) than CRTS.

There is some evidence that the flaring activity of some late M dwarf stars has been associated with the presence of stellar companions (Scholz et al. 2004). One of the transients we discovered (CSS080118:112149–131310) was a  $> 3.5$  mag flare on a nearby high-proper-motion star, LHS-2397a. This star has a proper motion of  $0''.509 \text{ year}^{-1}$  and was serendipitously observed spectroscopically during a flare by Bessell (1991). LHS-2397a is, in fact, a binary system where the primary star is an M8 type star (LHS-2397aA) and the companion (LHS-2397aB), a L7.5 Brown Dwarf orbiting within 4 AU (Freed et al. 2003).

### 3.5. Asteroids and Short-Timescale Optical Transients

The only interest of the CRTS is the discovery of transients beyond our solar system. Moving transients are discovered by the CSS in their search for NEOs. However, to detect faint transients we have to allow for moderate astrometric uncertainties between detections. Thus, our transients selection is limited to objects that move an average of  $< 0''.1 \text{ minute}^{-1}$  between observations. In order to determine how asteroids near stationary points might contaminate extra-solar transient detections, we investigated the distribution of known asteroid apparent motions. We chose the night of 2006 October 21st which contains 21,492 known asteroids in the CSS fields. Of these objects 2.2% were predicted to have apparent motion  $< 0''.1 \text{ minute}^{-1}$  when observed. In our processing, we remove asteroids by comparing the locations of transients with all known asteroids determined from MPC ephemerides. Furthermore, our detections are limited to a magnitude of  $\sim 20$  where a large fraction of asteroids are known. Nevertheless, a small number of OTs were seen in only one sequence of images and many of these are probably unknown asteroids. As objects such as trans-Neptunian objects (TNOs) have motions of a few arcseconds per hour, very bright TNOs could be found in CSS data. However, as there are only  $\sim 10$  known TNOs brighter than  $V \sim 20$ , and extensive surveys have been carried out to find these, it is unlikely that new TNO will be discovered in CSS data.

**Table 3**  
Blazar Candidates

ID	R.A. (h:m:s)	Dec ( $^{\circ}$ $'$ $''$ )	Date <i>MJD</i>	<i>u</i>	<i>g</i>	<i>r</i>	<i>i</i>	<i>z</i>	CSS
CSS080506:120952+181007	12:09:51.75	+18:10:07.0	54592.21514	19.3	18.9	18.6	18.3	18.1	15.8 <sup>a</sup>
CSS080409:154159+235603	15:41:59.97	+23:56:03.3	54565.28634	23.9	23.7	23.6	22.6	22.7	18.8 <sup>a</sup>
CSS080426:165347+164950	16:53:46.61	+16:49:49.5	54582.44017	22.4	22.2	22.4	22.2	21.5	19.0
CSS080306:141549+090354	14:15:48.80	+09:03:54.4	54531.32834	21.2	20.5	20.4	19.9	19.6	18.6
CSS080208:120722+250650	12:07:21.938	+25:06:50.26	54504.32945	...	...	...	...	...	19.3

**Notes.** Column 1, CSS ID; Columns 2 and 3, right ascension and declination J2000; Column 4 modified Julian date of detection; Columns 5–9, SDSS-DR6 magnitudes; Column 10 Catalina detection magnitude.

<sup>a</sup> Spectroscopic candidates (Mahabal et al. 2008a).

The main LSST survey is planning to take pairs of observations in a single filter separated by 15 s. A 1 mag change of a typical flare star over 500 s corresponds to a 0.03 mag change during this time. This will be undetectable for most stars observed by LSST, particularly at the near faint limit where most late M dwarfs can be expected. In addition to stellar flares, over a 15 s time span, faint, unknown asteroids will appear stationary. For example, an asteroid with an apparent motion of  $0''.1 \text{ minute}^{-1}$  will move only  $0''.025$  between observations. The LSST intends to make two pairs of observations for each field on the same night. This cadence enables asteroid detections to be linked together to form arcs that can in turn be linked to form orbits. If the same field was observed again after 500 s (in order to veto UV Ceti flares), the change in brightness of a flare star would be significant. However, the asteroid in our example would only have moved  $0''.8$ . This seems very unlikely to provide an arc that can be uniquely linked with a detection 24 hr later. Deep photometric or spectroscopic follow-up observations could be used to determine the nature of individual rapid transients discovered by LSST. However, with a conservative estimate of hundreds of flare events every night, a significant amount of time will be required to follow even a fraction of these objects. If the LSST was to take a third observation on the same night, flare stars and moving objects could be clearly separated.

Recently, Horesh et al. (2008) used overlaps between two SDSS scans to select SN candidates. They removed asteroids by imposing an  $0''.16 \text{ minute}^{-1}$  limit on motion. Using the CSS data, we found  $\sim 6\%$  of the known asteroids had motions less than this limit. Although these authors removed known asteroids, two-thirds of asteroids at the magnitudes of their SNe candidates ( $r \sim 20\text{--}21.5$ ) are unknown (Ivezic et al. 2002). Clearly, care will be required when selecting short-timescale OTs.

### 3.6. Blazars and Other Variable Sources

Although the main types of transients we have discovered so far are SNe, CVs, and UV Ceti stars, there are other clear populations of transients. These include nearby high-proper-motion stars, blazars, and highly variable stars (such as Miras).

Two possible blazars were discovered in CSS data and spectroscopically followed, CSS080506:120952+181007 and CSS080409:154159+235603 (Mahabal et al. 2008a). CSS080409:154159+235603 was fainter than magnitude 20.5 on 2008 January 1st and when observed on February 9th was mag 18.8. It has been approximately the same brightness since discovery, suggesting the object is being fueled by a significant source of energy. CSS080506:120952+181007 was discovered at magnitude 15.8 on 2008 May 6th and matches the flat spectrum radio source MG1 J120953+1809 (Helmbolt et al. 2007).

Based on the light curves of these objects, we have selected additional transients with similar abrupt and irregular light curves. All these objects are presented in Table 3.

CSS080426:165347+164950 matches a radio source seen in NVSS data and appears to vary by 1 mag in as little as 20 minutes. CSS080306:141549+090354 corresponds to the flat-spectrum radio source RGB J1415+090 (Jackson et al. 2007) and appears to rapidly vary between magnitudes 18.4 and 20.4. CSS080208:120722+250650 matches an uncatalogued source seen in SDSS images. Based on nearby sources, we estimate that before detection it was magnitude  $m_r \sim 23$ . This object faded from magnitude  $\sim 19.2$  to  $\sim 20.2$  over a period of 140 days. Detailed follow-up of these objects is required to understand their true nature.

## 4. SUMMARY AND DISCUSSION

We have presented a few of the types of OTs found in the first six months of CRTS analysis of CSS observations. With the observational parameters of this survey, the transient discoveries are dominated by two types of objects, DNe and SNe. Apart from Miras, few of the many kinds of highly variable stars were discovered. We believe that the main reasons for this are our high-detection threshold (2 mag), use of catalogs based on co-added images (where cyclic variability on short timescales are averaged out), and the natural limit to the number of stars observed by only observing fields with  $|b| > 10^{\circ}$ .

Photometric follow-up was taken a day or two after discovery for 80 of the OTs in the first six months using a combination of Gunn *g*, *r*, *i*, and *z* filters (A. Mahabal et al. 2009b, in preparation). Spectroscopic follow-up was also carried out to classify a small number of OTs. Clearly, one of the most important next steps is that of classifying these and other such transients, such that the expensive spectroscopic resources are used to a minimum. Given that surveys tend to provide a small number of data points, this requires optimizing photometric follow-up in combination with probabilistic classification techniques. Our initial photometric results suggest that young Type Ia SN have much lower *r*–*i* colours than DN outbursts. Thus, even limited photometric follow-up of transients from surveys such as LSST can provide very large numbers of these important cosmological distance indicators. However, we are also making steady progress to tackle the classification problem using advanced mathematical and statistical methodologies (see, e.g., Mahabal et al. 2008b). As the survey progresses, we will also optimize transient discovery by employing machine learning techniques (Borne 2008). This will enable unsupervised rapid follow-up to be carried out on the most interesting objects.

The CRTS has a firm commitment to making discoveries and the associated metadata public as quickly as possible. In this

way, we hope that all aspects and types of OTs can be explored. Many of the CRTS OTs were posted in ATels and a number of these were independently observed after these announcements. We chose to announce most of the obvious SNe with bright hosts in CBETs and these were usually spectroscopically followed and thus confirmed. Although significant follow-up would be required to characterize all the OTs discovered by CRTS, this is possible as the SDSS-II Supernova Survey acquired  $\sim 180$  nights of spectroscopic time on 2.4–11 m telescopes, to follow SNe found during  $\sim 70$  nights of imaging (Frieman et al. 2008). All OTs will continue to be made publicly available from VOEventNet and those events that appear the most interesting will be announced in ATels and CBETs. In addition, we will be using the two additional CSS survey telescopes to search for transients. These will allow us to find similar numbers of transients in the southern sky as well as fainter transients in the north.

In conclusion, while there is little doubt that many of the “major advances in our understanding of the universe have historically arisen by improvements in our ability to see” (Ivezic et al. 2008), we believe that significant advances have, and will continue to be made by also changing how and what is observed. The upcoming next generation of transient surveys (PanSTARRs and LSST) promise a unique ability to discover rare types of OTs in the faint time-domain sky. However, it is clear that the full characterization of many kinds of transients will require follow-up observations and these are most easily achieved when the transients are both accessible to the entire astronomical community, and bright enough to be followed by the large numbers of existing small telescopes.

We thank J. Greaves and members of the CVNet for the contributions and discussions about CVs.

This work is supported by the National Science Foundation under grant CNS-0540369. The CSS survey is funded by the National Aeronautics and Space Administration under grant NNG05GF22G issued through the Science Mission Directorate Near-Earth Objects Observations Program. The PQ digital sky survey is a collaborative venture between California Institute of Technology (Caltech) and Yale University. The data are obtained at the Samuel Oschin telescope at Palomar Observatory, and processed at the Center for Advanced Computing Research (CACR) at Caltech, using techniques developed in part for the U.S. National Virtual Observatory (NVO). The PQ survey is supported by the U.S. National Science Foundation under grants AST-0407448 and AST-0407297. Support for M.C. is provided by Proyecto Basal PFB-06/2007, by FONDAP Centro de Astrofísica 15010003, and by a John Simon Guggenheim Memorial Foundation Fellowship.

This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This research has made use of the NASA/IPAC Infrared Science Archive and NASA/IPAC Extragalactic Database (NED), which are operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. *GALEX* is a NASA Small Explorer, operated for NASA by California Institute of technology under NASA contract NAS-98034. Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS web site is <http://www.sdss.org/>.

## REFERENCES

- Abell, G. O., Corwin, H. G., Jr., & Olowin, R. P. 1989, *ApJS*, **70**, 1
- Adelman-McCarthy, J. K., et al. 2008, *ApJS*, **175**, 297
- Akerlof, C., et al. 2000, *ApJ*, **119**, 1901
- Akerlof, C. W., et al. 2003, *PASP*, **115**, 120
- Alcock, C., et al. 2000, *ApJ*, **542**, 281
- Aldering, G., et al. 2002, in Proc. SPIE 4836, 61 Survey and Other Telescope Technologies and Discoveries, ed. T. J. Anthony & W. Sidney (Bellingham, WA: SPIE)
- Aubourg, E., et al. 1995, **301**, 1
- Becker, A. C., et al. 2004, *ApJ*, **611**, 418
- Bessell, M. S. 1991, *AJ*, **101**, 662
- Bianchi, L., et al. 2007, *ApJS*, **173**, 659
- Borne, K. 2008, *Astron. Nachr.*, **329**, 255
- Brady, S., et al. 2008, *PASP*, **120**, 301
- Christensen, E. 2004, GCN Obs. Rep. 2849 (Greenbelt, MD: GSFC)
- Djorgovski, S. G., Gal-Yam, A., & Price, P. 2004, GCN Obs. Rep. 2851 (Greenbelt, MD: GSFC)
- Djorgovski, S. G., et al. 2007, *Atel* 1262
- Djorgovski, S. G., et al. 2008a, *Astron. Nachr.*, **329**, 263
- Djorgovski, S. G., et al. 2008b, *Atel*, 1411
- Djorgovski, S. G., et al. 2008c, *Atel*, 1416
- Djorgovski, S. G., et al. 2008d, *Atel*, 1418
- Drake, A. J., et al. 1999, *ApJ*, **521**, 602
- Drake, A. J., et al. 2003, *ApJ*, **589**, 1020
- Drake, A. J., et al. 2007a, *BAAS*, **39**, 805
- Drake, A. J., et al. 2007b, *Atel*, 1337
- Drake, A. J., et al. 2008a, *Atel*, 1374
- Drake, A. J., et al. 2008b, *Atel*, 1388
- Drake, A. J., et al. 2008c, *Atel*, 1399
- Drake, A. J., et al. 2008d, *Atel*, 1404
- Drake, A. J., et al. 2008e, *Atel*, 1416
- Drake, A. J., et al. 2008f, *Atel*, 1447
- Drake, A. J., et al. 2008g, *Atel*, 1479
- Drake, A. J., et al. 2008h, *Atel*, 1510
- Eisenstein, D. J., et al. 2006, *ApJS*, **167**, 40
- Filippenko, A., Li, W. D., Treffers, R. R., & Modjaz, M. 2001, in ASP Conf Ser. 246, Small-Telescope Astronomy on Global Scales ed. W. P. Chen, C. Lemme, & B. Paczynski (San Francisco, CA: ASP), **121**
- Freed, M., Close, L. M., & Siegler, N. 2003, *ApJ*, **584**, 453
- Freedman, W. L., et al. 2001, *ApJ*, **553**, 47
- Frieman, J. A., et al. 2008, *AJ*, **135**, 338
- Gal-Yam, A., et al. 2003, *AJ*, **125**, 1087
- Gal-Yam, A., et al. 2008, *ApJ*, **680**, 550
- Glikman, E., et al. 2008, *Atel*, 1413
- Harrison, T., et al. 2004, *ApJ*, **127**, 460
- Helmbolt, J. F., et al. 2007, *ApJ*, **658**, 2003
- Hodapp, K. W., et al. 2004, *Astron. Nachr.*, **325**, 636
- Hosh, A., Poznanski, D., Ofek, E., & Moaz, D. 2008, *arXiv:0805.1922*
- Huber, M., et al. 2006, *AJ*, **132**, 633
- Ivezic, Z., et al. 2002, *AJ*, **124**, 2364
- Ivezic, Z., et al. 2008, *arXiv:0805.2366*
- Jackson, N., et al. 2007, *MNRAS*, **76**, 371
- Kauffmann, G., et al. 2003, *MNRAS*, **341**, 33
- Keller, S. C., et al. 2007, *PASA*, **24**, 1
- Kricinunas, K., Margon, B., & Szkody, P. 1998, *PASP*, **110**, 1342
- Kulkarni, S. R., & Rau, A. 2006, *ApJ*, **644**, L63
- Larson, S., et al. 2003, *DPS*, **35**, 3604
- López-Morales, M. 2007, *ApJ*, **660**, 732
- Mahabal, A., et al. 2008a, *Atel*, 1520
- Mahabal, A., et al. 2008b, *Astron. Nachr.*, **328**, 288
- Mannucci, F., et al. 2008, *MNRAS*, **383**, 1121
- Martin, D. C., et al. 2005, *ApJ*, **619**, 1
- Miller, A. A., et al. 2009, *ApJ*, **690**, 1303
- Modjaz, M., et al. 2008, *AJ*, **135**, 1136
- Monard, B. 2003, *Mon. Notes Astron. Soc. South Africa*, **62**, 131
- Morokuma, T., et al. 2008, *ApJ*, **676**, 163
- Paczynski, B. 2000, *PASP*, **112**, 1281
- Pignata, G., et al. 2007, *CBET*, **1114**, 1
- Pojmanski, G. 2001, in ASP Conf Ser. 246, Small-Telescope Astronomy on Global Scales, ed. W. P. Chen, C. Lemme, & B. Paczynski (San Francisco, CA: ASP), **53**
- Poznanski, D., et al. 2002, *ApJ*, **114**, 833
- Prieto, J. L., Stanek, K. S., & Beacom, J. F. 2008, *ApJ*, **673**, 999
- Puckett, T., Langoussis, A., & Marcus, M. 2003, *IAU Circ.*, **8154**, 1
- Rau, A. R., et al. 2007, *ApJ*, **664**, 474



- Richards, G., et al. 2002, [AJ](#), **123**, 2945  
Rykoff, E. S., et al. 2005, [ApJ](#), **631**, 1032  
Sand, D. J., et al. 2008, [AJ](#), **135**, 1917  
Savaglio, S., Glazebrook, K., & LeBorgne, D. 2008, arXiv:0803.2718  
Sesar, B., et al. 2007, [AJ](#), **134**, 2236  
Scholz, R.-D., et al. 2004, [MNRAS](#), **347**, 685  
Schweizer, F., et al. 2008, [AJ](#), **136**, 1482  
Silvestri, N., et al. 2006, [AJ](#), **131**, 1674  
Skrutskie, M. F., et al. 2006, [AJ](#), **131**, 1163  
Smith, N., et al. 2007, [ApJ](#), **666**, 1116  
Stanek, K. Z., et al. 2006, *Acta Astron.*, **56**, 333  
Steinfadt, J. D. R., Bildsten, L., & Howell, S. B. 2008, [ApJ](#), **677**, L113  
Szkody, P., et al. 2002, [AJ](#), **123**, 430  
Szkody, P., et al. 2003, [AJ](#), **126**, 1499  
Szkody, P., et al. 2004, [AJ](#), **128**, 1882  
Szkody, P., et al. 2005, [AJ](#), **129**, 2386  
Szkody, P., et al. 2006, [AJ](#), **131**, 973  
Szkody, P., et al. 2007, [AJ](#), **134**, 185  
Tomaney, A., & Crotts, 1996, [AJ](#), **112**, 2872  
Totani, T., et al. 2005, [ApJ](#), **621**, 9  
Udalski, A., et al. 1994, [ApJ](#), **426**, 69  
Wachter, L. M., Hawley, S. L., & West, A. A. 2004, [PASP](#), **116**, 1105  
Webb, W., & Malkan, M. 2000, [ApJ](#), **540**, 652  
Welsh, B. Y., et al. 2007, [ApJS](#), **173**, 673  
Wood-Vasey, W. M., et al. 2004, [New Rev. Astron.](#), **48**, 637  
York, D. G., et al. 2002, [AJ](#), **120**, 1579  
Yost, S. A., et al. 2007, [ApJ](#), **669**, 1107