

Operator Mixing and the BMN Correspondence

To cite this article: Neil R. Constable *et al* JHEP10(2002)068

View the [article online](#) for updates and enhancements.

You may also like

- [Ultrasonic mist chemical vapor deposition and dielectric properties of cubic pyrochlore bismuth magnesium niobate thin films](#)
Pingfan Ning, Didi Wang, Yuqiang Li et al.
- [Interface stability of electrode/Bi-containing relaxor ferroelectric oxide for high-temperature operational capacitor](#)
Takahiro Nagata, Somu Kumaragurubaran, Yoshifumi Tsunekawa et al.
- [Transparent Capacitor for the Storage of Electric Power Produced by Transparent Solar Cells](#)
Cheng-Ji Xian and Soon-Gil Yoon

Operator mixing and the BMN correspondence

Neil R. Constable

*Center for Theoretical Physics and Laboratory for Nuclear Science
Massachusetts Institute of Technology
Cambridge, Ma. 02139, USA
E-mail: constabl@lns.mit.edu*

Daniel Z. Freedman

*Center for Theoretical Physics and Laboratory for Nuclear Science
Massachusetts Institute of Technology, Department of Mathematics
Cambridge, Ma. 02139, USA
E-mail: dzf@math.mit.edu*

Matthew Headrick and Shiraz Minwalla

*Jefferson Physical Laboratory, Department of Physics
Harvard University, Cambridge MA 02138, USA
E-mail: headrick@pascal.harvard.edu, minwalla@fas.harvard.edu*

ABSTRACT: In this note we update the discussion of the BMN correspondence and string interactions in [4] to incorporate the effects of operator mixing. We diagonalize the matrix of two point functions of single and double trace operators, and compute the eigen-operators and their anomalous dimensions to order $g_2^2 \lambda'$. Operators in different R symmetry multiplets remain degenerate at this order; we propose this is a consequence of supersymmetry. We also calculate the corresponding energy shifts in string theory, and find a discrepancy with field theory results, indicating possible new effects in light-cone string field theory.

KEYWORDS: Penrose limit and pp-wave background, AdS/CFT and dS/CFT Correspondence.

Contents

1. Introduction	1
2. Anomalous dimensions of BMN eigen-operators	3
2.1 Two-point functions, three-point functions, and mixing	3
2.2 Construction of eigen-operators	5
2.3 Consistency conditions from three-point functions	6
2.4 Anomalous dimensions of BMN eigen-operators	7
3. String field theory revisited	8
A. The space-time structure of two- and three-point functions	10
B. Absence of mixing with BPS double-trace operators	10
C. Diagonalization procedure	11
D. Deconstruction identities	12
E. Degenerate operators and mixing	14

1. Introduction

Several aspects of the correspondence [1] between operators of $\mathcal{N} = 4$ $SU(N)$ super-Yang-Mills theory at large R-charge J and type-IIB string theory in a pp-wave background geometry [2, 3] have been investigated in the recent literature. The map between field theory operators and free string theory in the pp-wave background was established by [1]. In our paper [4] we attempted to extend this map to a correspondence between Yang-Mills correlators and interactions in the pp-wave background. In this note we will update and correct the discussion presented in [4].

Field theory in the BMN limit, $N \rightarrow \infty$, $J \rightarrow \infty$ with J^2/N fixed, appears to be governed by two parameters [1, 4, 5, 6]: an effective gauge coupling $\lambda' = g_{\text{YM}}^2 N/J^2$, and an effective genus counting parameter $g_2 = J^2/N$. These quantities can be expressed in terms of the string scale α' and coupling g_s , light-cone momentum p^+ , and transverse string mass μ as

$$\lambda' = \frac{1}{(\mu p^+ \alpha')^2}, \quad g_2 = 4\pi g_s (\mu p^+ \alpha')^2. \quad (1.1)$$

In [4] we proposed that free three-point functions of the BMN operators

$$O_n^J \equiv \frac{1}{\sqrt{JN^{J+2}}} \sum_{l=0}^J e^{2\pi i n l / J} \text{Tr} \left(\phi Z^l \psi Z^{J-l} \right) \quad (1.2)$$

are related to matrix elements of the string field theory light-cone hamiltonian according to

$$\langle i | H_{\text{int}} (|j\rangle \otimes |k\rangle) = \mu g_2 (\Delta_i - \Delta_j - \Delta_k) C_{ijk}, \quad (1.3)$$

where $|i\rangle, |j\rangle, |k\rangle$ are the dual string states and $\langle O_i \bar{O}_j \bar{O}_k \rangle = g_2 C_{ijk}$ (we have factored out the trivial space time dependence). This proposal was intended to describe string transitions in which the initial and final states have the same number of excitations, so that $\Delta_i - \Delta_j - \Delta_k$ is of order λ' . The proposal applies in the limit of large μ , corresponding to weak coupling in field theory. The r.h.s. of (1.3) was computed in [4] by a direct evaluation of the free three-point coupling C_{ijk} . Subsequently, the l.h.s. of (1.3) was obtained in [7] (see also [10, 11, 12]) as the large- μ limit of the string field theory interaction H_{int} in the pp-wave background [13]. Results of [7] confirm the proposal (1.3), which thus appears to be on firm ground.

We claimed in [4] that the proposal (1.3) obeys a non-trivial consistency check, which we now review. Utilizing hamiltonian matrix elements from (1.3), and second-order non-degenerate perturbation theory, we computed the one loop mass renormalization of excited pp-wave string states. We then compared the result to an explicit computation of the order $\lambda' g_2^2$ correction to the anomalous dimension of the corresponding BMN operator. In [4] we reported agreement between these two computations.

In this note we will point out that this agreement is in fact spurious.¹ Firstly our result for the anomalous dimension computed in our paper had the wrong sign (this was pointed out to us by the authors of [5]). Secondly our computation was incomplete, in that it did not take into account mixing between single- and double-trace BMN operators. As we will describe in this paper, this mixing alters the result for the anomalous dimension at the order under consideration. In this note we will present a corrected result for the anomalous dimension of BMN operators at order $\lambda' g_2^2$.²

Surprisingly, this corrected anomalous scaling dimension does not match the result of the analogous perturbative calculation on the string theory side, as we show in section 3. The discrepancy could be due to presence of a quadratic contact term of order g_2^2 in the string field theory hamiltonian. This is an important issue.

Note added: after submission of this paper for publication we were informed that the results of [7] (version 2), which claimed to support our proposal (1.3), were erroneous. Subsequently the correct expression for the three-string vertex (in the large μ limit) has

¹In appendix D we show that the relation referred to in [4] as the “unitarity check” (with the sign corrected) in fact follows purely from field theory considerations, without making use of the BMN duality.

²Operator mixing was first suggested in the BMN context in [14]. A preliminary version of the present work was presented by one of us (S.M.) at the conference Strings 2002. As we were drafting this note, the paper [15] was submitted to the archive with similar methodology and results, for a larger class of operators. One of us (D.Z.F.) acknowledges useful discussions with G. Semenoff at the Aspen Center for Physics. Mixing is also discussed in the very recent paper [16].

appeared in [8, 9] as well as version 3 of [7]. This does not effect any of the key results presented in this paper. In particular our computation of the torus contributions to the anomalous dimensions of the BMN operators is, of course, independent of the string theory vertex. We have, however, updated section 3 below in which we compare the results obtained in section 2 of this paper with the string theory prediction for the one loop mass renormalization of the string states which are dual to the BMN operators considered here. We again find that there is a mismatch between the two approaches and we maintain our position that this provides evidence for a non-trivial contact term in the string field theory Hamiltonian.

2. Anomalous dimensions of BMN eigen-operators

The BMN limit preserves an $SO(4)$ R-symmetry group. The complex impurity fields ϕ and ψ transform in the fundamental representation of an $SU(2)$ subgroup, and operators containing two such impurities transform in the $2 \otimes 2 = 1 \oplus 3$ representation. Our calculations thus split into independent sectors for the two representations.³ For example, if we form the linear combinations

$$O_n^{\pm J} = \frac{1}{\sqrt{2}} (O_n^J \pm O_{-n}^J) \tag{2.1}$$

of the operators (1.2), then O_n^{+J} is a member of a triplet, while O_n^{-J} is a singlet. In this section we will determine, to order $\lambda' g_2^2$, the anomalous dimension of the eigen-operators $\tilde{O}_n^{\pm J}$ that reduce to $O_n^{\pm J}$ at $g_2 = 0$. A priori the results could have been different for the two representations. We find instead that the degeneracy persists to this order in g_2 ; at the end of this subsection we will argue that this degeneracy is a consequence of supersymmetry.

The mixing problem requires the diagonalization of the matrix of two-point functions $\langle O_{(1)} \bar{O}_{(1)} \rangle, \langle O_{(1)} \bar{O}_{(2)} \rangle, \langle O_{(2)} \bar{O}_{(2)} \rangle$ of single- and double-trace operators including free and order λ' terms. The mixed two-point functions are of order g_2 . To obtain the eigen-operators to this order we need only the order g_2^0 parts of $\langle O_{(1)} O_{(1)} \rangle$ and $\langle O_{(2)} O_{(2)} \rangle$. The eigen-operators are obtained in subsection 2.1, in a treatment which emphasizes the relation of two- and three-point functions and the issue of correct conformal behavior. In subsection 2.2 we go on to find the order $\lambda' g_2^2$ correction to the anomalous dimension of the eigen-operators $\tilde{O}_n^{\pm J}$ which requires the known [4, 5] order g_2^2 parts of $\langle O_{(1)} O_{(1)} \rangle$. As discussed in appendix E, in these calculations we make the assumption that the single-trace operators do not mix with degenerate triple-trace operators, which is not yet justified by calculation of the relevant two-point functions. However, the mode $n = 1$ is not degenerate with any multi-trace operators, so results for $n = 1$ are safe from this danger.

2.1 Two-point functions, three-point functions, and mixing

The two-point functions between single-trace BMN operators O_n^J was computed to lowest order in g_2 in [1]:

$$\langle O_n^{\pm J} \bar{O}_m^{\pm J} \rangle = \delta_{nm} (1 - \lambda' m^2 \ln(x^2 \Lambda^2)), \quad \langle O_n^{+J} \bar{O}_m^{-J} \rangle = 0. \tag{2.2}$$

³We thank H. Verlinde and L. Motl for pointing this out to us.

In order to establish some notation that will be useful below we may express two-point functions between BMN operators of charge $J_1 < J$ as

$$\langle O_n^{\pm J_1} \bar{O}_m^{\pm J_1} \rangle = \delta_{nm} (1 - \lambda' k^2 \ln(x^2 \Lambda^2)), \quad (2.3)$$

where $k = Jm/J_1 = m/s$, and we have defined $s = J_1/J$ (this was denoted y in [4]).

We now turn to the construction of the matrix of two-point functions between single-trace and double-trace BMN operators. We will find it convenient to first compute, to order $\lambda' g_2$, the three-point functions of three single-trace BMN operators located at distinct spatial points; the requisite two-point functions may then be obtained allowing two of the insertions in the three-point function to approach each other.

The Feynman graphs that contribute to the three-point function between O_n^J , $O_m^{J_1}$, and $O^{J-J_1} \equiv \text{Tr}(Z^{J-J_1})/\sqrt{(J-J_1)N^{J-J_1}}$ at order λ' are of two kinds, with different space time dependence. This gives the following general structure for the three-point function:

$$\begin{aligned} \langle O_n^J(x_1) \bar{O}_m^{J_1}(x_2) \bar{O}^{J-J_1}(x_3) \rangle &= \\ &= g_2 C_{nms} \left[1 - \lambda' \left(a_{nk} \ln(x_{12}\Lambda)^2 + b_{nk} \ln\left(\frac{x_{13}x_{12}\Lambda}{x_{23}}\right) \right) \right]. \end{aligned} \quad (2.4)$$

Here C_{nms} is the free three-point function between the relevant operators which was computed in [4],

$$C_{nms} = \sqrt{\frac{1-s}{J_s}} \frac{\sin^2(\pi ns)}{\pi^2(n-k)^2}. \quad (2.5)$$

Evaluation of the Feynman diagrams gives the values

$$a_{nk} = k^2, \quad b_{nk} = n(n-k). \quad (2.6)$$

but we find it useful to regard a_{nk} and b_{nk} as unspecified parameters in much of this section.

Note that (2.4), with the values of a_{nk} and b_{nk} given in (2.6) *does not* take the form dictated by conformal invariance for the three-point functions of three operators of anomalous dimension $n^2\lambda'$, $k^2\lambda'$, and 0, respectively. This already indicates that O_n^J does not have well defined scaling dimension; we will see this in more detail below.

Transforming (2.4) to the $O_n^{\pm J}$ basis, and using the properties $a_{nk} = a_{-n,-k}$, $b_{nk} = b_{-n,-k}$, we can rewrite the above three-point function as

$$\begin{aligned} \langle O_n^{\pm J}(x_1) \bar{O}_m^{\pm J_1}(x_2) \bar{O}^{J-J_1}(x_3) \rangle &= \\ &= g_2 C_{nms}^{\pm} \left[1 - \lambda' \left(a_{nk}^{\pm} \ln(x_{12}\Lambda)^2 + b_{nk}^{\pm} \ln\left(\frac{x_{13}x_{12}\Lambda}{x_{23}}\right) \right) \right] \end{aligned} \quad (2.7)$$

and

$$\langle O_n^{\pm J}(x_1) \bar{O}_m^{\mp J_1}(x_2) \bar{O}^{J-J_1}(x_3) \rangle = 0, \quad (2.8)$$

where

$$\begin{aligned} C_{nms}^+ &= C_{nms} + C_{-n,m,s} = H_{nks} \frac{2(n^2 + k^2)}{(n^2 - k^2)^2}, \\ C_{nms}^- &= C_{nms} - C_{-n,m,s} = H_{nks} \frac{4nk}{(n^2 - k^2)^2}, \\ H_{nks} &= \sqrt{\frac{1-s}{J_s}} \frac{\sin^2(n\pi s)}{\pi^2}. \end{aligned} \quad (2.9)$$

When the values in (2.6) are used

$$\begin{aligned} a_{nk}^\pm &= k^2, \\ b_{nk}^+ &= \frac{n^2(n^2 - k^2)}{n^2 + k^2}, \quad b_{nk}^- = \frac{n^2 - k^2}{2}. \end{aligned} \quad (2.10)$$

We now turn to the construction of two-point functions between double-trace and single-trace operators. In appendix A we demonstrate that the two-point function

$$\langle O_n^J(0) : \bar{O}_m^{J_1} \bar{O}^{J-J_1} : (x) \rangle \equiv \langle O_n^J(0) \bar{O}_{ms}^J(x) \rangle \quad (2.11)$$

may be obtained from (2.4) by the replacement $x_{12} = x_{13} \rightarrow x$, $x_{23} \rightarrow 1/\Lambda$ so that

$$\langle O_n^J(0) \bar{O}_{ms}^J(x) \rangle = g_2 C_{nms} (1 - \lambda' \ln(x\Lambda)^2 (a_{nk} + b_{nk})). \quad (2.12)$$

The final two-point function that we need involves two double-trace operators. It is straightforward to show that the result up to order g_2 is

$$\langle : O_m^{J_1} O^{J-J_1} : (0) : \bar{O}_n^{J_1} \bar{O}^{J-J_1} : (x) \rangle = \delta_{mn} (1 - \lambda' k^2 \ln(x^2 \Lambda^2)). \quad (2.13)$$

The complete matrix of two-point functions can now be assembled from (2.2) and (2.12) and (2.13). If we introduce the index $A = i, j, m, n$ where m, n index the single-trace operators O_n^J and i, j index the double-trace operators O_{ms}^J so that $i \equiv (m, s)$ then the matrix of two-point functions found in this section can be summarized as

$$\langle O_A \bar{O}_B \rangle = g_{AB} - \lambda' h_{AB} \ln x^2 \Lambda^2. \quad (2.14)$$

Here g_{AB} gives an inner product on the space of operators (in fact it is just the Hilbert space inner product, according to the state-operator mapping), while h_{AB} is the matrix of anomalous dimensions (obtained by applying the dilatation operator $D = x^\nu \partial_\nu$). See appendix C for a schematic discussion of the diagonalization of $\langle O_A \bar{O}_B \rangle$.

2.2 Construction of eigen-operators

In order to find eigen-operators we must now diagonalize the matrix of two-point functions. The eigen-operators $\tilde{O}_n^{\pm J}$ and \tilde{O}_{ms}^J must, by definition, reduce to $O_n^{\pm J}$ and $: O_m^{J_1} O^{J-J_1} :$, respectively at $g_2 = 0$. These operators take the form

$$\begin{aligned} \tilde{O}_n^{\pm J} &= O_n^{\pm J} + g_2 \sum_{m=0}^{\infty} \sum_{J_1=0}^J C_{nms}^\pm M_{nm} : O_m^{\pm J_1} O^{J-J_1} : + \dots, \\ \tilde{O}_{ms}^{\pm J} &= : O_m^{\pm J_1} O^{J-J_1} : - g_2 \sum_{m=0}^{\infty} C_{nms}^\pm N_{nm} O_n^{\pm J} + \dots, \end{aligned} \quad (2.15)$$

where M_{nk} and N_{nk} are coefficients to be determined.⁴ The two-point function between $\tilde{O}_n^{\pm J}$ and $\tilde{O}_{ms}^{\pm J}$ is easily computed using the results of the previous subsection

$$\begin{aligned} (4\pi^2 x^2)^{J+2} \langle \tilde{O}_n^{\pm J}(0) \tilde{O}_{ms}^{\pm J}(x) \rangle &= \\ &= g_2 C_{nms}^\pm [(1 + M_{nm} - N_{nm}) - \lambda' \ln(\Lambda x)^2 (a_{nk}^\pm + b_{nk}^\pm + M_{nm} k^2 - N_{nm} n^2)]. \end{aligned} \quad (2.16)$$

⁴The transformation to the \pm basis implies that the summation symbol in (2.15) and other formulas in this basis should be defined as $\sum_{m=0}^{\infty} \equiv \frac{1}{2} \sum_{m=-\infty}^{\infty}$.

As $\tilde{O}_n^{\pm J}$ and \tilde{O}_{ms}^J are eigen-operators in a conformal field theory, (2.16) must vanish identically; this yields the set of simultaneous equations

$$\begin{aligned} 1 + M_{nm} - N_{nm} &= 0, \\ a_{nk}^{\pm} + b_{nk}^{\pm} + M_{nm}k^2 - N_{nm}n^2 &= 0 \end{aligned} \tag{2.17}$$

which may easily be solved for M_{nm} and N_{nm} . Carrying out the algebra we find

$$\begin{aligned} \tilde{O}_n^{\pm J} &= O_n^{\pm J} + g_2 \sum_{m=0}^{\infty} \sum_{J_1=0}^J C_{nms}^{\pm} \frac{a_{nk}^{\pm} + b_{nk}^{\pm} - n^2}{n^2 - k^2} O_m^{\pm J_1} O^{J-J_1} + \dots, \\ \tilde{O}_{ms}^{\pm J} &= : O_m^{\pm J_1} O^{J-J_1} : - g_2 \sum_{m=0}^{\infty} C_{nms}^{\pm} \frac{a_{nk}^{\pm} + b_{nk}^{\pm} - k^2}{n^2 - k^2} O_n^{\pm J} + \dots \end{aligned} \tag{2.18}$$

2.3 Consistency conditions from three-point functions

Having determined the BMN eigen-operators to first order in $g_2\lambda'$, we now turn to the determination of their three-point functions. We will demonstrate that the three-point functions of these operators do indeed take the form required by conformal invariance. We find that this is true for any value of b_{nk} provided the coefficient $a_{nk} = k^2$. This is consistent with (2.6) and provides a check on our algebra.

Combining (2.2), (2.3) and (2.4) and (2.18) (and using the fact that double trace correlators factorize to lowest order in g_2) we find

$$\begin{aligned} \left\langle \tilde{O}_n^{\pm J}(x_1) \tilde{O}_m^{\pm J_1}(x_2) \tilde{O}^{J-J_1}(x_3) \right\rangle &= \\ &= g_2 C_{nms}^{\pm J} \frac{a_{nk}^{\pm} + b_{nk}^{\pm} - k^2}{n^2 - k^2} \left(1 - \lambda' \left[\ln |x_{12}\Lambda| \frac{2n^2k^2 - b_{nk}^{\pm}(k^2 + n^2) - 2a_{nk}^{\pm}n^2}{k^2 - a_{nk}^{\pm} - b_{nk}^{\pm}} + \right. \right. \\ &\quad \left. \left. + x \ln \left| \frac{x_{13}x_{12}\Lambda}{x_{23}} \right| \left[b_{nk}^{\pm} \frac{k^2 - n^2}{k^2 - a_{nk}^{\pm} - b_{nk}^{\pm}} \right] \right) \right). \end{aligned} \tag{2.19}$$

(2.19) takes the standard CFT form for a three-point function between operators of anomalous dimension n^2 , k^2 , and 0, respectively, provided that

$$\begin{aligned} \frac{2n^2k^2 - b_{nk}^{\pm}(k^2 + n^2) - 2a_{nk}^{\pm}n^2}{k^2 - a_{nk}^{\pm} - b_{nk}^{\pm}} &= k^2 + n^2, \\ b_{nk}^{\pm} \frac{k^2 - n^2}{k^2 - a_{nk}^{\pm} - b_{nk}^{\pm}} &= k^2. \end{aligned} \tag{2.20}$$

It is easily verified that the two eqs. (2.20) are not independent; they are both satisfied for any value of b_{nk} if and only if

$$a_{nk}^{\pm} = k^2, \tag{2.21}$$

as promised at the beginning of this subsection.

Notice that the three-point coupling between three normalized eigen-operators involves the coefficient

$$\tilde{C}_{nms}^{\pm} = C_{nms}^{\pm} \frac{b_{nk}^{\pm}}{n^2 - k^2}$$

which is distinct from the free three-point coupling C_{nms}^\pm even at lowest order in λ' ! Even though this modification is independent of λ' , it is clearly a quantum effect as the modified three-point function depends on b_{nk} , the coefficient of the order λ' term in (2.7).

2.4 Anomalous dimensions of BMN eigen-operators

It is now a simple matter to compute the two-point function of the operator (2.18) and thereby determine its anomalous dimension to order $g_2^2\lambda'$.⁵ The two-point functions of single-trace BMN operators with double-trace operators were presented above while the complete order $\lambda'g_2^2$ two-point functions of single-trace BMN operators were presented in [4, 5] and we reproduce here (with different relative signs than [4]) the relevant equations,

$$\langle O_n^J(0)\bar{O}_m^J \rangle = (\delta_{nm} + g_2^2 A_{nm}) (1 - (n^2 - nm + m^2)\lambda' \ln x^2 \Lambda^2) - \frac{g_2^2 \lambda'}{4\pi^2} B_{mn} \ln x^2 \Lambda^2, \quad (2.22)$$

where the matrices A_{mn} and B_{mn} are given by

$$A_{nm} = \begin{cases} \frac{1}{24}, & \text{if } m = n = 0, \\ 0, & \text{if } m = 0, n \neq 0 \text{ or } n = 0, m \neq 0, \\ \frac{1}{60} - \frac{1}{6u^2} + \frac{7}{u^4}, & \text{if } m = n \neq 0, \\ \frac{1}{4u^2} \left(\frac{1}{3} + \frac{35}{2u^2} \right), & \text{if } m = -n \neq 0, \\ \frac{1}{(u-v)^2} \left(\frac{1}{3} + \frac{4}{v^2} + \frac{4}{u^2} - \frac{6}{uv} - \frac{2}{(u-v)^2} \right), & \text{all other cases,} \end{cases} \quad (2.23)$$

$$B_{nm} = \begin{cases} 0, & n=0 \text{ if } m = 0, \\ \frac{1}{3} + \frac{10}{u^2}, & \text{if } n = m \neq 0, \\ -\frac{15}{2u^2}, & \text{if } n = -m \neq 0, \\ \frac{6}{uv} + \frac{2}{(u-v)^2}, & \text{all other cases,} \end{cases} \quad (2.24)$$

and

$$u = 2\pi m, \quad v = 2\pi n.$$

As explained in appendix C, the anomalous dimension of the $\tilde{O}_n^{\pm J}$ can be readily computed from the diagonal two-point functions $\langle \tilde{O}_n^\pm \bar{\tilde{O}}_n^\pm \rangle$. The required algebra is straightforward using the ingredients (2.12), (2.18), (2.22), (2.23) and (2.24). The result is

$$\frac{\Gamma_n^\pm}{\lambda' g_2^2} = \mp 2n^2 A_{n,-n} + \frac{1}{4\pi^2} (B_{nn} \pm B_{n,-n}) - \sum_{m=0}^{\infty} \sum_{J_1=0}^J (C_{nms}^\pm)^2 \frac{(b_{nk}^\pm - n^2 + k^2)^2}{n^2 - k^2}. \quad (2.25)$$

where we use the value for b_{nk}^\pm given in (2.10). Using the identities in appendix D this may be re-written as

$$\Gamma_n^\pm = g_2^2 \lambda' n^2 A_{n-n} = \frac{g_2^2 \lambda'}{4\pi^2} \left(\frac{1}{12} + \frac{35}{32\pi^2 n^2} \right). \quad (2.26)$$

⁵We do not need to know the form of the operator to order g_2^2 in order to perform this computation. The only order g_2^2 correction to this operator that can contribute to its two-point function at order g_2^2 is a piece proportional to O_n^J ; such an addition can be absorbed into a normalization of the operator at this order, and so does not contribute to its scaling dimension.

The full anomalous dimension of this operator (including the planar contribution) is

$$\Delta_n^\pm - J - 2 = \lambda' n^2 (1 + g_2^2 A_{n,-n}) = \lambda' \left[n^2 + \frac{g_2^2}{4\pi^2} \left(\frac{1}{12} + \frac{35}{32\pi^2 n^2} \right) \right]. \quad (2.27)$$

This is the principal result of this note.

It is striking that the degeneracy between the even and odd operators (obvious at $g_2 = 0$) persists to order g_2^2 . It appears that this degeneracy is a consequence of supersymmetry;⁶ all two-impurity BMN operators lie in a single representation of the pp-wave supersymmetry algebra and so are guaranteed to have equal anomalous dimensions. Note that 16 of the 32 supercharges of the $\mathcal{N} = 4$ $d = 4$ superconformal algebra commute with $p^- = \frac{\mu}{2}(\Delta - J)$; further all supercharges commute with p^+ which is a central element of this algebra in the pp-wave limit. Consequently, states in long representations of the pp-wave superalgebra appear in multiplets whose minimum degeneracy is $2^{16/2} = 16 \times 16$. But this is precisely the number of two-impurity BMN operators; it is natural to conjecture that the two-impurity BMN operators may be related to one another by the action of the 16 supercharges that commute with p^- ; their degeneracy is thus a consequence of supersymmetry. Special examples of such relations have been worked out directly in the gauge theory in [17, 18]. We hope to return to this issue in more detail in future work.

3. String field theory revisited

In this section we will use quantum mechanical perturbation theory to compute the energy shifts of a particular set of excited string states in the pp-wave background. This section is almost a direct transcription of section 5 of [4], now adapted to the $O_n^{\pm J}$ basis.⁷

Utilizing the large μ expression for the three string vertex [7, 8, 9]

$$\frac{1}{\mu} \langle n | H_{\text{int}} | i \rangle = \frac{1}{2} (n - k)^2 C_{ni} \quad (3.1)$$

and second order non-degenerate perturbation theory (see section 5 of [4]) we find,

$$\frac{1}{\mu} \sum_i \frac{|\langle n | H_{\text{int}} | i \rangle|^2}{E_n - E_i} = \frac{g_2^2 \lambda'}{4} \sum_i \frac{(n - k)^4}{n^2 - k^2} C_{ni}^2 = -\frac{1}{4} \frac{g_2^2 \lambda'}{4\pi^2} B_{n,-n} = \frac{g_2^2 \lambda'}{4\pi^2} \frac{15}{32\pi^2 n^2} \quad (3.2)$$

and

$$\begin{aligned} \frac{1}{\mu} \sum_i \frac{\langle n | H_{\text{int}} | i \rangle \langle i | H_{\text{int}} | -n \rangle}{E_n - E_i} &= \frac{g_2^2 \lambda'}{4} \sum_i \frac{(n^2 - k^2)^2}{n^2 - k^2} C_{ni} C_{-n,i} = -\frac{1}{4} \frac{g_2^2 \lambda'}{4\pi^2} B_{n,-n} \\ &= \frac{g_2^2 \lambda'}{4\pi^2} \frac{15}{32\pi^2 n^2} \end{aligned} \quad (3.3)$$

where we have used i as a collective index for (m, s) as well as for the states in which the ϕ and ψ excitations are on different strings, corresponding to the BPS operator $: O_\phi^{J_1} O_\psi^{J-J_1} :$.

⁶This explanation was suggested to us by J. Maldacena and M. Van Raamsdonk in response to an earlier version of this paper.

⁷This section has been updated to account for the corrected three string vertex which now appears in [8, 9] and version 3 of [7]. Our results agree with those obtained in [8, 9].

Unlike in the Yang-Mills calculation of the previous section, the degeneracy between the + and - sectors of the string theory is thus broken: the correct zeroth-order eigenvectors are

$$|n, \pm\rangle \equiv \frac{1}{\sqrt{2}} (|n\rangle \pm |-n\rangle), \tag{3.4}$$

and their second order energy shifts are

$$\begin{aligned} \frac{1}{\mu} \sum_i \frac{|\langle n, + | H_{\text{int}} | i, + \rangle|^2}{E_{n,+} - E_{i,+}} &= -\frac{1}{2} \frac{g_2^2 \lambda'}{4\pi^2} B_{n,-n} = \frac{g_2^2 \lambda'}{4\pi^2} \frac{15}{16\pi^2 n^2} \\ \frac{1}{\mu} \sum_i \frac{|\langle n, - | H_{\text{int}} | i, - \rangle|^2}{E_{n,-} - E_{i,-}} &= \frac{1}{2} \frac{g_2^2 \lambda'}{4\pi^2} (B_{n,-n} - B_{n,-n}) = 0. \end{aligned} \tag{3.5}$$

Sadly, neither of these energy shifts agrees with the anomalous dimension computed in the previous section.

It is possible that this disagreement is resolved by the presence of an explicit quadratic contact term in the string field theory Hamiltonian. In order to restore the degeneracy of string states and resolve the discrepancy, this contact term would have to have matrix elements

$$\begin{aligned} \frac{1}{\mu} \langle n, + | V_2 | n, + \rangle &= \frac{g_2^2 \lambda'}{4\pi^2} \left(\frac{1}{12} + \frac{5}{32\pi^2 n^2} \right) \\ \frac{1}{\mu} \langle n, - | V_2 | n, - \rangle &= \frac{g_2^2 \lambda'}{4\pi^2} \left(\frac{1}{12} + \frac{35}{32\pi^2 n^2} \right). \end{aligned} \tag{3.6}$$

Since these expressions are positive it is at least possible that they come from a contact term in the string field theory Hamiltonian, which would be determined by the anti-commutator of two supersymmetry generators [19, 20, 21]. It would be very interesting to compare 3.6 with a direct computation of contact terms in string field theory.

It should also be mentioned that the energy shifts could possibly receive contributions at this order from states with 4 or more impurities, due to enhanced matrix elements between states with different numbers of impurities, as reported in [7]. Note, however, that this effect could not alone cure the apparent discrepancy between string theory and gauge theory, since the contributions, if any, due to such massive intermediate states would be negative.

Acknowledgments

We would like to thank V. Khöze, C. Kristjansen, L. Motl, J. Plefka, A. Postnikov, G. Semenoff, W. Skiba, M. Spradlin, M. Staudacher, A. Volovich and especially J. Maldacena, M. Van Raamsdonk and H. Verlinde for useful discussions and correspondence. S. M. would like to thank the Tata Institute for Fundamental Research, Mumbai for hospitality while this work was being completed. S.M. and D.F.Z would like to thank the organizers of the Amsterdam Summer workshop for hospitality while this work was in progress. N.R.C. and D.Z.F. are supported by the NSF under PHY 00-96515 and the DOE under DF-FC02-94ER40818. N.R.C. is also supported by NSERC of Canada. M.H. is supported in part by the DOE under grant DE-FG01-91ER40654 and in part by the Packard Foundation. S.M. is supported by a Harvard Junior Fellowship and by the DOE grant DE-FG01-91ER40654.

A. The space-time structure of two- and three-point functions

In this appendix we will justify the space time dependence of the order λ' terms in the eq. (2.4) for the three-point function $\langle O_n^J(x_1) \bar{O}_{m_1}^{J_1}(x_2) \bar{O}^{J_2}(x_3) \rangle$, and discuss its relation to mixed two-point functions as in (2.12). There are two distinct structures coming from the quartic interactions in contributing Feynman diagrams. The first structure occurs in diagrams in which one pair of lines from the operator $O_n^J(x_1)$ and one pair from $\bar{O}_{m_1}^{J_1}(x_2)$ terminate at the interaction vertex at z . This leads to the the same space time integral which occurs in two-point functions, namely

$$\int \frac{d^4 z}{z^4(z-x_{12})^4} = 2\pi^2 \frac{\ln(x_{12}^2 \Lambda^2)}{x_{12}^4}. \quad (\text{A.1})$$

The second structure occurs in diagrams where lines from all three operators are connected to the interaction vertex. This leads to the integral

$$\int \frac{d^4 z}{z^4(x_{12}-z)^2(x_{13}-z)^2} = \pi^2 \ln \left(\frac{x_{12}^2 x_{13}^2 \Lambda^2}{x_{23}^2} \right). \quad (\text{A.2})$$

This integral was evaluated using dimensional regularization in [22] and with differential regularization [23] in our work. The integrals above determine the general structure in (2.4), and the specific values of the parameters a_{nk}, b_{nk} which come from our summation of the contributing diagrams are given in (2.6).

The two-point function $\langle O_n^J(x_1) : \bar{O}_{m_1}^{J_1} \bar{O}^{J_2} : (x_2) \rangle$ can be obtained from the three-point function in the OPE limit $x_3 \rightarrow x_2$. See [24]. It can also be found simply by setting $x_{13} = x_{12}$ and $x_{23} = 1/\Lambda$, since (A.2) reduces to (A.1) if this is done. Two-point functions were also obtained directly from summation of Feynman diagrams in our work.

B. Absence of mixing with BPS double-trace operators

In [4] — see also [22, 25] — non-vanishing three-point functions $\langle O_n^J \bar{O}_\phi^{J_1} \bar{O}_\psi^{J_2} \rangle$ were computed, where $O_\phi^J = \text{Tr}(\phi Z^J)/\sqrt{N^{J+1}}$ and $O_\psi^J = \text{tr}(\psi Z^J)/\sqrt{N^{J+1}}$ are BPS operators. The corresponding dual two-string state contributed to the sum over intermediate states in the consistency check in [4]. One may then wonder why the double-trace $: \bar{O}_\phi^{J_1} \bar{O}_\psi^{J_2} :$ was not included in the initial ansatz (2.18) for our mixing calculation, and we now justify its omission.

Let $O^J = \text{Tr}(Z^J)/\sqrt{JN^J}$ denote the chiral primary operator. It is quite easy to verify that the linear combination

$$O_{\text{susy}}^J \equiv \left(\sqrt{\frac{J_1-1}{J_2+1}} : O_0^{J_1-1} O^{J_2+1} : + : O_\phi^{J_1} O_\psi^{J_2} : \right) + (J_1 \leftrightarrow J_2) \quad (\text{B.1})$$

is an SU(4) descendant of the chiral primary $: \text{Tr}(Z^{J_1+1}) \text{Tr}(Z^{J_2+1}) :$. Thus the operator O_{susy}^J decouples from O_n^J and from $: O_m^{J_1} O^{J_2} :$ for $n, m \neq 0$. The decoupling can easily be observed; the appropriate combination of mixed two-point functions which can be obtained

from the three-point functions given in [4, (3.10) and (3.11)] vanish.⁸ The decoupling means that we can simply omit the operators $:\bar{O}_\phi^{J_1}\bar{O}_\psi^{J_2}:$ from the mixing calculation, if we keep $:\mathcal{O}_0^{J_1}\mathcal{O}^{J_2}:$. However, one may also observe directly from the solution for \tilde{O}_n^J in (2.18) that $:\mathcal{O}_0^{J_1}\mathcal{O}^{J_2}:$ also decouples.

C. Diagonalization procedure

In this appendix we provide an overview of the diagonalization procedure used to obtain the eigen-operators in (2.18) and their scale dimension.

The matrix of two point functions is given in (2.14). The problem is to diagonalize the matrix of anomalous scaling dimensions h_{AB} relative to the inner product g_{AB} . Both h_{AB} and g_{AB} are functions of g_2 and we will perform our analysis up to and including contributions at order g_2^2 . To explain the notion of relative diagonalization, we note that the goal is to find a set of operators,

$$\tilde{O}_A = O_B S^B{}_A \tag{C.1}$$

and numbers γ_A such that

$$(4\pi^2 x^2)^{J+2} \langle \tilde{O}_A(0) \tilde{O}_B(x) \rangle = \delta_{AB} (1 - \lambda \gamma_A \ln x^2 \Lambda^2). \tag{C.2}$$

Hence the matrix S must satisfy

$$S^\dagger g S = I, \quad S^\dagger h S = \gamma, \tag{C.3}$$

where $\gamma_{AB} = \gamma_A \delta_{AB}$ is the diagonalized matrix of anomalous scaling dimensions. Together these imply

$$S^{-1} g^{-1} h S = \gamma. \tag{C.4}$$

Thus we must find the eigenvalues and eigenvectors of the matrix

$$h^A{}_C = g^{AB} h_{BC}, \tag{C.5}$$

where g^{AB} is the inverse matrix of g_{AB} . This diagonalization process is essentially equivalent to non-degenerate quantum mechanical perturbation theory. We use a two-step process in which we first find the eigenvectors to order g_2 , and then use the two-point function of the eigen-operators to compute the scale dimension to order g_2^2 .

To make the method clear consider the toy eigen-value problem

$$M^a{}_b V_{(c)}^b = \lambda_{(c)} V_{(c)}^a. \tag{C.6}$$

Diagonal elements are given by $M^a{}_a = \rho_{(a)}$, and we wish to work to first order in all off-diagonal elements. To zero order, the eigenvalues are $\lambda_{(c)} = \rho_{(c)}$ and eigenvectors are $V_{(c)}^a = \delta^a{}_c$. It is then trivial to see that the first order eigenvectors are

$$V_{(c)}^a = 1, \quad \text{if } a = c, \\ V_{(c)}^a = \frac{M^a{}_c}{\rho_{(c)} - \rho_{(a)}}, \quad \text{if } a \neq c. \tag{C.7}$$

⁸To correct an error the right side of (3.11) in [4] should be multiplied by -1 .

The correction is well-defined if there is no degeneracy. This toy model is pertinent to our block matrix h^A_B because the non-diagonal corrections to h^n_m and h^i_j are of order g_2^2 and can be temporarily ignored. One immediately finds the (dominantly single-trace) eigenvectors to first order in g_2 :

$$\tilde{O}_n = O_n + \sum_i \frac{O_i h^i_n}{n^2 - k^2} = O_n - g_2 \sum_i \frac{k}{n+k} C_{n,i} O_i. \tag{C.8}$$

One may also write an analogous expression for the dominantly double-trace eigenvectors of the structure $\tilde{O}_i = O_i + \mathcal{O}(g_2 O_m)$. From linear combinations of O_n and O_{-n} one can easily obtain the eigen-operators in (2.18) (with (2.6) inserted).

The second step of our procedure is based on the matrix of two-point functions $\langle \tilde{O}_A \tilde{O}_B \rangle$ (in the \pm basis). Because of the approximate diagonalization, off-diagonal entries $\langle \tilde{O}_n \tilde{O}_i \rangle$ are of order g_2^3 and can be dropped. We can confine our attention to the upper block of the matrix which is again of the form

$$\langle \tilde{O}_n^\pm \tilde{O}_m^\pm \rangle = G_{nm} - \lambda H_{nm} \ln x^2. \tag{C.9}$$

(with superscripts omitted on the r.h.s. to avoid clutter). We need to think about diagonalizing the matrix $H_m^n = G^{np} H_{pm}$, in which G_{nm}, H_{nm} have the structure

$$\begin{aligned} G_{nm} &= \delta_{nm} + g_2^2 g'_{mn}, \\ H_{nm} &= n^2 \delta_{nm} + g_2^2 h'_{nm}. \end{aligned} \tag{C.10}$$

The off-diagonal elements of H_m^n are of order g_2^2 . There is no degeneracy so these affect eigenvalues beginning only in order g_2^4 . Much as in the toy model above, the anomalous dimension of the eigen-operator \tilde{O}_n^\pm is then simply the diagonal element

$$H_n^n = n^2 + g_2^2 (h'_{nn} - n^2 g'_{nn}). \tag{C.11}$$

It is this quantity in the \pm channels that is computed in (2.25).

D. Deconstruction identities

In this appendix we note several useful identities which relate the three point and two-point coefficients that appear in this paper. We follow the notation of section 3 in which i is used as a collective index labeling both types of double-trace operators: $:O_m^{J_1} O^{J-J_1}:$ (for which $s = J_1/J$ and $k = m/s$) and $:O_\phi^{J_1} O_\psi^{J-J_1}:$ (for which $s = J_1/J$ and $k = 0$). We then have:

$$\begin{aligned} \sum_i C_{ni} C_{mi} &= 2A_{nm}, \\ \sum_i k C_{ni} C_{mi} &= (n+m)A_{nm}, \\ \sum_i k^2 C_{ni} C_{mi} &= (n^2 + m^2) A_{nm} + \frac{1}{4\pi^2} B_{nm}. \end{aligned} \tag{D.1}$$

The first of these was pointed out in [10]. The first and last can be combined to obtain the relation

$$\sum_i (n^2 - k^2) C_{ni}^2 = -\frac{1}{4\pi^2} B_{nn}, \tag{D.2}$$

which is the content (up to a sign) of the “unitarity check” of [4].

These formulas are not numerical coincidences, nor do they in and of themselves provide evidence for the BMN correspondence, since they can be derived purely within the gauge theory. Nonetheless the derivation, which we will explain in this appendix, naturally calls to mind a picture of interacting strings, and may in the future be useful in gaining a deeper understanding of the correspondence.

We imagine the string of Z s in the operator O_n^J to be a circle on which the ϕ and ψ impurities are quantum mechanical particles moving with momentum n and $-n$, respectively, — their wave function is the weight of each possible configuration in the definition of the operator (1.2). (Morally this point of view is a “de-second quantization” of the worldsheet field theory, which is appropriate in the large- μ limit where the excitations are very massive and we are fixing their number and type.) A free torus diagram contributing to A_{nm} can be represented by a unitary operator $T(s, j_1, j_3)$ which acts on this 2-particle Hilbert space by dividing the circle (of length 1, for simplicity) into 4 blocks of lengths j_1, j_2, j_3, j_4 (where $j_2 = s - j_1, j_4 = 1 - s - j_3$) and transposing blocks 1 and 2 and blocks 3 and 4 (see [4, figure 3], for example). We thus have

$$A_{nm} = \phi \langle n |_{\psi} \langle -n | R | n \rangle_{\phi} | -n \rangle_{\psi}, \tag{D.3}$$

where

$$R = \frac{1}{4} \int_0^1 ds \int_0^s dj_1 \int_0^{1-s} dj_3 T(s, j_1, j_3) \tag{D.4}$$

(the integrals count each diagram 4 times, hence the 1/4). But if we imagine pinching the circle into two circles of lengths s and $1 - s$, $T(s, j_1, j_3)$ acts by rotating the s circle through an angle $2\pi j_1$ and the $1 - s$ circle through an angle $2\pi j_3$. The integration over these rotation angles will ensure a factorization over physical intermediate states. Let us see this more explicitly.

In such a pinching, the 2-particle Hilbert space decomposes into a direct sum of four Hilbert spaces depending on which little circle each impurity is on:

$$\mathcal{H}_{\phi}(1)\mathcal{H}_{\psi}(1) = \mathcal{H}_{\phi}(s)\mathcal{H}_{\psi}(s) \oplus \mathcal{H}_{\phi}(1-s)\mathcal{H}_{\psi}(1-s) \oplus \mathcal{H}_{\phi}(s)\mathcal{H}_{\psi}(1-s) \oplus \mathcal{H}_{\phi}(1-s)\mathcal{H}_{\psi}(s). \tag{D.5}$$

The integral over the rotation angle j_1 projects out any state with non-zero total momentum on the s circle, and similarly the integral over j_3 projects out any state with non-zero total momentum on the $1 - s$ circle. Hence we have

$$\begin{aligned} & \int_0^s dj_1 \int_0^{1-s} dj_3 T(s, j_1, j_3) = \\ & = s(1-s) \left(\sum_m P_m(s) + \sum_m P_m(1-s) + P_{0'}(s) + P_{0'}(1-s) \right), \end{aligned} \tag{D.6}$$

where $P_m(s)$ projects onto the state $|m, s\rangle_\phi | -m, s\rangle_\psi$ corresponding to the operator $: O_m^{J_1} \times O^{J-J_1} :$, and $P_{0'}(s)$ projects onto the state $|0, s\rangle_\phi |0, 1-s\rangle_\psi$ corresponding to the operator $: O_\phi^{J_1} O_\psi^{J-J_1} :$. Using

$$C_{ni} = \sqrt{s(1-s)} \begin{cases} \phi \langle n |_\psi \langle -n | m, s \rangle_\phi | -m, s \rangle_\psi, & \text{for } : O_m^{J_1} O^{J-J_1} : , \\ \phi \langle n |_\psi \langle -n | 0, s \rangle_\phi | 0, 1-s \rangle_\psi, & \text{for } : O_\phi^{J_1} O_\psi^{J-J_1} : \end{cases} \quad (\text{D.7})$$

we recover the first identity of (D.1).

Inserting $k^2/2$ in the sum means including interaction vertices in the diagrams. However, the vertex is included only if it is between two lines on the same small circle. Thus (in the language of [4]) all the nearest neighbor interactions are counted, but only half the semi-nearest neighbor interactions. The non-nearest neighbor interactions are also under counted by a factor of 2, but this is a little more complicated: only 1/3 of them are counted, but overall diagrams with three blocks are over counted by a factor of 3/2. Hence we have

$$\frac{1}{2} \sum_i k^2 C_{ni} C_{mi} = nm A_{nm} + \frac{1}{2} (n-m)^2 A_{nm} + \frac{1}{8\pi^2} B_{nm}, \quad (\text{D.8})$$

which is equivalent to the last identity in (D.1).

The second identity of (D.1) can also be derived by similar but more involved reasoning.

E. Degenerate operators and mixing

The diagonalization process is an application of non-degenerate quantum mechanical perturbation theory. A potential complication occurs because single and double-trace operators have equal zeroth-order eigenvalues when $n = k$, threatening a divergence in the calculation of eigenvectors to order g_2 . This would signal the breakdown of non-degenerate perturbation theory and the necessity of using the degenerate theory. As one can see from the right-hand side of (2.18), such a divergence does not occur, thanks to some rather delicate cancellations. These cancellations have an interesting reflection in the dual string theory [4]. The amplitude for an excited string state corresponding to a 2 impurity BMN operator to decay into two particles vanishes onshell at lowest order in $1/\mu$; thus these excited string states are stable to lowest order in g_2 .

Let us now consider higher rank $r \geq 3$ multi-trace operators which we call O_r for short. The two-point functions $\langle \bar{O}_n O_r \rangle$ are of order g_2^{r-1} . Generically such operators contribute to eigenvalues beginning in higher order than g_2^2 . There could be a complication for triple-trace operators $O_3 = : O_m^{J_1} \text{Tr}(Z^{J_2}) \text{Tr}(Z^{J_3}) :$ which are degenerate with single-traces when $n = m/s$. If the relevant interaction matrix elements do not vanish, eigen-operators change at zeroth order and scale dimensions at order g_2^2 . The vanishing or otherwise of these interaction matrix elements is linked to the vanishing or otherwise of the onshell one-three particle decay amplitudes of string field theory.

Degeneracy cannot occur for momentum $n = 1$ since $m \in Z^+$ and $0 < J_1 < J$. A calculation of two-point functions $\langle \bar{O}_n O_3 \rangle$ to order g_2^2 and $\lambda' g_2^2$ would be needed to see if the interaction matrix elements vanish, as they do for double-traces. The calculation has

not been done yet, so a conservative position is that present mixing results are valid for momentum $n = 1$, where degeneracy cannot occur, but may need modification for $n \geq 2$ if such mixing does occur.

References

- [1] D. Berenstein, J.M. Maldacena and H. Nastase, *Strings in flat space and PP waves from $N = 4$ super Yang-Mills*, *J. High Energy Phys.* **04** (2002) 013 [[hep-th/0202021](#)].
- [2] C.M. Hull, *Killing spinors and exact plane wave solutions of extended supergravity*, *Phys. Rev.* **D 30** (1984) 334;
M. Blau, J. Figueroa-O'Farrill, C. Hull and G. Papadopoulos, *A new maximally supersymmetric background of IIB superstring theory*, *J. High Energy Phys.* **01** (2002) 047 [[hep-th/0110242](#)]; *Penrose limits and maximal supersymmetry*, *Class. and Quant. Grav.* **19** (2002) L87 [[hep-th/0201081](#)].
- [3] R.R. Metsaev, *Type iib Green-Schwarz superstring in plane wave Ramond-Ramond background*, *Nucl. Phys.* **B 625** (2002) 70 [[hep-th/0112044](#)];
R.R. Metsaev and A.A. Tseytlin, *Exactly solvable model of superstring in plane wave Ramond-Ramond background*, *Phys. Rev.* **D 65** (2002) 126004 [[hep-th/0202109](#)].
- [4] N.R. Constable, D. Z. Freedman, M. Headrick, S. Minwalla, L. Motl, A. Postnikov and W. Skiba, *PP-wave string interactions from perturbative Yang-Mills theory*, *J. High Energy Phys.* **07** (2002) 017 [[hep-th/0205089](#)].
- [5] C. Kristjansen, J. Plefka, G.W. Semenoff and M. Staudacher, *A new double-scaling limit of $N = 4$ super Yang-Mills theory and PP-wave strings*, *Nucl. Phys.* **B 643** (2002) 3 [[hep-th/0205033](#)].
- [6] D. Berenstein and H. Nastase, *On lightcone string field theory from super Yang-Mills and holography*, [hep-th/0205048](#).
- [7] M. Spradlin and A. Volovich, *Superstring interactions in a PP-wave background, 2*, [hep-th/0206073](#).
- [8] J. Pearson, M. Spradlin, D. Vaman, H. Verlinde and A. Volovich, *Tracing the string: BMN correspondence at finite J^{*2}/N* , [hep-th/0210102](#).
- [9] J. Gomis, S. Moriyama and J.W. Park, *SYM description of SFT hamiltonian in a pp-wave background*, [hep-th/0210153](#).
- [10] M.-X. Huang, *String interactions in PP-wave from $N = 4$ super Yang-Mills*, [hep-th/0206248](#).
- [11] P. Lee, S. Moriyama and J.-w. Park, *Cubic interactions in PP-wave light cone string field theory*, *Phys. Rev.* **D 66** (2002) 085021 [[hep-th/0206065](#)].
- [12] Y.-J. Kiem, Y.-B. Kim, S.-M. Lee and J.-M. Park, *PP-wave/Yang-Mills correspondence: an explicit check*, *Nucl. Phys.* **B 642** (2002) 389 [[hep-th/0205279](#)].
- [13] M. Spradlin and A. Volovich, *Superstring interactions in a PP-wave background*, *Phys. Rev.* **D 66** (2002) 086004 [[hep-th/0204146](#)].
- [14] M. Bianchi, B. Eden, G. Rossi and Y.S. Stanev, *On operator mixing in $N = 4$ SYM*, [hep-th/0205321](#).

- [15] N. Beisert, C. Kristjansen, J. Plefka, G.W. Semenoff and M. Staudacher, *BMN correlators and operator mixing in $N = 4$ super Yang-Mills theory*, [hep-th/0208178](#).
- [16] D.J. Gross, A. Mikhailov and R. Roiban, *A calculation of the plane wave string hamiltonian from $N = 4$ super-Yang-Mills theory*, [hep-th/0208231](#).
- [17] A. Santambrogio and D. Zanon, *Exact anomalous dimensions of $N = 4$ Yang-Mills operators with large R charge*, *Phys. Lett.* **B 545** (2002) 425 [[hep-th/0206079](#)].
- [18] U. Gürsoy, *Vector operators in the BMN correspondence*, [hep-th/0208041](#).
- [19] J. Greensite and F.R. Klinkhamer, *Superstring amplitudes and contact interactions*, *Nucl. Phys.* **B 304** (1988) 108.
- [20] M.B. Green and J.H. Schwarz, *Superstring interactions*, *Nucl. Phys.* **B 218** (1983) 43.
- [21] H. Verlinde, *Bits, matrices and $1/N$* , [hep-th/0206059](#).
- [22] C.-S. Chu, V.V. Khoze and G. Travaglini, *Three-point functions in $N = 4$ Yang-Mills theory and PP -waves*, *J. High Energy Phys.* **06** (2002) 011 [[hep-th/0206005](#)].
- [23] D.Z. Freedman, K. Johnson and J.I. Latorre, *Differential regularization and renormalization: a new method of calculation in quantum field theory*, *Nucl. Phys.* **B 371** (1992) 353.
- [24] C.-S. Chu, V.V. Khoze and G. Travaglini, *PP -wave string interactions from N -point correlators of BMN operators*, *J. High Energy Phys.* **09** (2002) 054 [[hep-th/0206167](#)].
- [25] M.-X. Huang, *Three point functions of $N = 4$ super Yang-Mills from light cone string field theory in PP -wave*, *Phys. Lett.* **B 542** (2002) 255 [[hep-th/0205311](#)].