

Dark matter and $B_s \rightarrow \mu^+ \mu^-$ with minimal SO_{10} soft SUSY breaking

To cite this article: Radovan Dermisek *et al* JHEP04(2003)037

View the [article online](#) for updates and enhancements.

You may also like

- [Neutrino mass models](#)
S F King
- [\$E_6\$ GUT through effects of dimension-5 operators](#)
Chao-Shang Huang, Wen-Jun Li and Xiao-Hong Wu
- [\$SO\(10\)\$ models with flavour symmetries: classification and examples](#)
I P Ivanov and L Lavoura

Dark matter and $B_s \rightarrow \mu^+ \mu^-$ with minimal SO_{10} soft SUSY breaking

Radovan Dermisek

Davis Inst. for High Energy Physics, University of California, Davis, CA 95616, USA

E-mail: dermisek@physics.ucdavis.edu

Stuart Raby*

Department of Physics, The Ohio State University

174 W. 18th Ave., Columbus, Ohio 43210, USA

E-mail: raby@pacific.mps.ohio-state.edu

Leszek Roszkowski

Department of Physics, Lancaster University, Lancaster LA1 4YB, England

E-mail: L.Roszkowski@lancaster.ac.uk

Roberto Ruiz de Austri

Physics Division, School of Technology, Aristotle University of Thessaloniki

GR - 540 06 Thessaloniki, Greece

E-mail: rruiz@gen.auth.gr

ABSTRACT: CMSSM boundary conditions are usually used when calculating cosmological dark matter densities. In this paper we calculate the cosmological density of dark matter in the MSSM using minimal SO_{10} soft SUSY breaking boundary conditions. These boundary conditions incorporate several attractive features: they are consistent with SO_{10} Yukawa unification, they result in a “natural” inverted scalar mass hierarchy and they reduce the dimension 5 operator contribution to the proton decay rate. With regards to dark matter, on the other hand, this is to a large extent an unexplored territory with large squark and slepton masses m_{16} , large A_0 and small $\{\mu, M_{1/2}\}$. We find that in most regions of parameter space the cosmological density of dark matter is considerably less than required by the data. However there is a well-defined, narrow region of parameter space which provides the observed relic density of dark matter, as well as a good fit to precision electroweak data, including top, bottom and tau masses, and acceptable bounds on the branching fraction of $B_s \rightarrow \mu^+ \mu^-$. We present predictions for Higgs and SUSY spectra, the dark matter detection cross section and the branching ratio $BR(B_s \rightarrow \mu^+ \mu^-)$ in this region of parameter space.

KEYWORDS: Supersymmetric Effective Theories, Cosmology of Theories beyond the SM, Dark Matter.

*On leave of absence, School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540, USA

Contents

1. Introduction	1
2. Minimal SO_{10} SUSY Model — $MSO_{10}SM$	2
2.1 Framework	2
2.2 Analysis	4
3. Cosmological Dark Matter Density	5
4. Predictions and summary	10

1. Introduction

The constrained minimal supersymmetric standard model [CMSSM] [1] is a well defined model for soft SUSY breaking with five independent parameters given by m_0 , $M_{1/2}$, A_0 , $\tan\beta$ and $sign(\mu)$. It has been used extensively for benchmark points for collider searches, as well as for astrophysical and dark matter analyses. The economy of parameters in this scheme makes it a useful tool for exploring SUSY phenomena. However the CMSSM may miss regions of soft SUSY breaking parameter space which give qualitatively different predictions. In this paper we consider an alternate scheme, the minimal SO_{10} supersymmetric model [$MSO_{10}SM$], which is well motivated and opens up a qualitatively new region of parameter space.

In the $MSO_{10}SM$ there are 7 soft SUSY breaking parameters μ , $M_{1/2}$, A_0 , $\tan\beta$, m_{16} (a universal squark and slepton mass), m_{10} (a universal Higgs mass) and Δm_H^2 (Higgs up/down mass splitting). Moreover the parameters A_0 , m_{10} , m_{16} must satisfy the constraints [2, 3, 4] $A_0 \approx -2m_{16}$, $m_{10} \approx \sqrt{2}m_{16}$, $m_{16} > 1.2\text{TeV}$ with μ , $M_{1/2} \ll m_{16}$ and $\tan\beta \approx 50$. Note, with these values of the soft SUSY breaking parameters, we can explore SUSY phenomena with qualitatively different behavior than in the CMSSM. This is mainly due to the Higgs splitting (Δm_H^2) which, as is well known [5], enables one to obtain electroweak symmetry breaking with values $m_{16} \gg \mu$, $M_{1/2}$. Also, radiative EWSB with $\tan\beta \approx 50$ requires significantly less fine tuning with Higgs mass splitting (see Rattazzi and Sarid [5]). Furthermore, with 3 Higgs mass parameters μ , m_{10} , and Δm_H^2 we find that the latter two are strongly constrained by EWSB, once we fix the value of μ , which we treat as a free parameter. This is unlike the CMSSM where μ is fixed by EWSB. Also note that small changes in Δm_H^2 lead to big changes in the CP odd Higgs mass m_A [2].

It is not at all obvious that the $MSO_{10}SM$ region of soft SUSY breaking parameter space is consistent with cosmology [3, 4].¹ The dark matter candidate in this model is the lightest neutralino. However, since the scalar masses of the first two families are of

¹See also other recent articles discussing Yukawa unification and dark matter [6, 7].

order $m_{16} > 1.2 \text{ TeV}$, and the third generation sfermions (except for the stops) also tend to be heavy, the usually dominant annihilation channels, for the neutralino LSP to light fermions via t -channel sfermion exchange, are suppressed. On the other hand, the process $\chi\chi \rightarrow f\bar{f}$ via s -channel A exchange becomes important. This is due to the enhanced CP odd Higgs coupling to down-type fermions, which is proportional to $\tan\beta$, and because, in contrast to heavy scalar exchange, the process is not p -wave suppressed. In an earlier analysis, our χ^2 -analysis favored a light CP odd Higgs mass $m_A \sim 100 \text{ GeV}$ [2], although heavier A were also allowed. Such light A are however disfavored for two reasons. In order to provide efficient annihilation for the LSPs, one would be squeezed into a rather low LSP mass region $m_\chi \approx m_A/2$, which would require extreme fine-tuning at best. In addition, such low m_A are anyway inconsistent with the current limits on $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$. In this analysis, we vary the A mass.

We study the cosmology of the MSO_{10}SM in this paper. Obtaining the observed relic abundance of cold dark matter, which along with other cosmological parameters has recently been determined with an unprecedented accuracy [8], will provide a new important constraint on the model. We also compute the branching ratio for the process $B_s \rightarrow \mu^+ \mu^-$ due to A exchange [9]. It is absolutely essential to include this latter constraint in our analysis. Note, the CDF bound $\text{BR}(B_s \rightarrow \mu^+ \mu^-) < 2.6 \times 10^{-6}$ [10]. The cross section for the direct detection of dark matter is also computed. In section 2 we define the MSO_{10}SM , describe its virtues and outline the analysis. In section 3 we compute the cosmological dark matter density and discuss our results. Then in section 4 we discuss our predictions for underground dark matter searches and for collider Higgs and SUSY searches.

2. Minimal SO_{10} SUSY Model — MSO_{10}SM

2.1 Framework

Let us define the minimal SO_{10} SUSY model. Quarks and leptons of one family reside in the **16** dimensional representation, while the two Higgs doublets of the MSSM reside in one **10** dimensional representation. For the third generation we assume the minimal Yukawa coupling term given by

$$\lambda \mathbf{16} \mathbf{10} \mathbf{16} . \quad (2.1)$$

On the other hand, for the first two generations and for their mixing with the third, we assume a hierarchical mass matrix structure due to effective higher dimensional operators. Hence the third generation Yukawa couplings satisfy $\lambda_t = \lambda_b = \lambda_\tau = \lambda_{\nu_\tau} = \lambda$.

Soft SUSY breaking parameters are also consistent with SO_{10} with

- a universal gaugino mass $M_{1/2}$,
- a universal squark and slepton mass m_{16} ,²
- a universal scalar Higgs mass m_{10} ,

² SO_{10} does not require all sfermions to have the same mass. This however may be enforced by non-abelian family symmetries or possibly by the SUSY breaking mechanism.

- and a universal A parameter A_0 .

In addition we have the soft SUSY breaking Higgs mass parameters μ and $B\mu$. $B\mu$ may, as in the CMSSM, be exchanged for $\tan\beta$. Note, not all of these parameters are independent. Indeed, in order to fit the low energy electroweak data, including the third generation fermion masses, it has been shown that A_0 , m_{10} , m_{16} must satisfy the constraints [2]

$$\begin{aligned} A_0 &\approx -2m_{16} \\ m_{10} &\approx \sqrt{2}m_{16} \\ m_{16} &> 1.2 \text{ TeV} \\ \mu, M_{1/2} &\ll m_{16} \end{aligned} \tag{2.2}$$

with

$$\tan\beta \approx 50. \tag{2.3}$$

This result has been confirmed in two recent analyses [3, 4].³ The first property (eq. (2.2)) is necessary to fit the top, bottom and τ masses, in addition to the precision electroweak data [2, 3, 4]. The second property (eq. (2.3)) is a consequence of third generation Yukawa unification, since $m_t(m_t)/m_b(m_t) \sim \tan\beta$.

One loop threshold corrections at the GUT scale lead to two significant parameters we treat as free parameters, although they are calculable in any GUT. The first is a correction to gauge coupling unification given by

$$\epsilon_3 \equiv \frac{[\alpha_3(M_G) - \tilde{\alpha}_G]}{\tilde{\alpha}_G} \tag{2.4}$$

where the GUT scale M_G is defined as the scale where $\alpha_1(M_G) = \alpha_2(M_G) \equiv \tilde{\alpha}_G$. The second is a Higgs splitting mass parameter defined by

$$\Delta m_H^2 \equiv \frac{(m_{H_d}^2 - m_{H_u}^2)}{2m_{10}^2}. \tag{2.5}$$

In order to fit the low energy data we find $\epsilon_3 \approx -4\%$ and $\Delta m_H^2 \approx 13\%$ [2]. The largest corrections to ϵ_3 come from the Higgs and SO_{10} breaking sectors, while the correction to Δm_H^2 is predominantly due to the right-handed τ neutrino. For $M_{\bar{\nu}_\tau} \approx 10^{13-14}$ GeV (appropriate for a light τ neutrino mass ≈ 0.06 eV) we obtain $\Delta m_H^2 \approx 10 - 7\%$.

Finally, as a bonus, these same values of soft SUSY breaking parameters, with $m_{16} \gg \text{TeV}$, result in two very interesting consequences. Firstly, it “naturally” produces an inverted scalar mass hierarchy [ISMH] [12]. With an ISMH squarks and sleptons of the first two generations obtain mass of order m_{16} at M_Z . The stop, sbottom, and stau, on the other hand, have mass less than a TeV. An ISMH has two virtues.

1. It preserves “naturalness” (for values of m_{16} which are not too large), since only the third generation squarks and sleptons couple strongly to the Higgs.

³Note, different regions of parameter space consistent with Yukawa unification have also been discussed in [3, 4, 11].

2. It ameliorates the SUSY CP and flavor problems, since these constraints on CP violating angles or flavor violating squark and slepton masses are strongest for the first two generations, yet they are suppressed as $1/m_{16}^2$. For $m_{16} > \text{a few TeV}$, these constraints are weakened [13].

Secondly, Super-Kamiokande bounds on $\tau(p \rightarrow K^+ \bar{\nu}) > 1.9 \times 10^{33}$ yrs. [14] constrain the contribution of dimension 5 baryon and lepton number violating operators. These are however minimized with $\mu, M_{1/2} \ll m_{16}$ [15].

2.2 Analysis

We use a top-down approach with a global χ^2 analysis [16]. The input parameters are defined by boundary conditions at the GUT scale. The 11 input parameters at M_G are given by — three gauge parameters $M_G, \alpha_G(M_G), \epsilon_3$; the Yukawa coupling λ , and 7 soft SUSY breaking parameters $\mu, M_{1/2}, A_0, \tan \beta, m_{16}^2, m_{10}^2, \Delta m_H^2$. These are fit in a global χ^2 analysis defined in terms of physical low energy observables. Note we keep three parameters ($m_{16}, \mu, M_{1/2}$) fixed; while minimizing χ^2 with the remaining 8 parameters. Below we will plot χ^2 contours as a function of $\mu, M_{1/2}$ for different values of m_{16} . We use two (one) loop renormalization group [RG] running for dimensionless (dimensionful) parameters from M_G to M_Z .⁴ We require electroweak symmetry breaking using an improved Higgs potential, including m_t^4 and m_b^4 corrections in an effective 2 Higgs doublet model below $M_{SUSY} = \sqrt{\frac{1}{2}(m_{t_1}^2 + m_{t_2}^2)}$ [17].

The χ^2 function includes 9 observables; 6 precision electroweak data $\alpha_{EM}, G_\mu, \alpha_s(M_Z), M_Z, M_W, \rho_{NEW}$ and the 3 fermion masses $M_{\text{top}}, m_b(m_b), M_\tau$. In our analysis we fit the central values [18]: $M_Z = 91.188 \text{ GeV}$, $M_W = 80.419 \text{ GeV}$, $G_\mu \times 10^5 = 1.1664 \text{ GeV}^{-2}$, $\alpha_{EM}^{-1} = 137.04$, $M_\tau = 1.7770 \text{ GeV}$ with 0.1% numerical uncertainties; and the following with the experimental uncertainty in parentheses: $\alpha_s(M_Z) = 0.1180 (0.0020)$, $\rho_{new} \times 10^3 = -0.200 (1.1)$ [20], $M_t = 174.3 (5.1) \text{ GeV}$, $m_b(m_b) = 4.20 (0.20) \text{ GeV}$.⁵ We include the complete one loop threshold corrections at M_Z to all observables. In addition we use one loop QED and three loop QCD RG running below M_Z .

The output of this analysis is a set of weak scale squark, slepton, gaugino and Higgs masses. With regards to the calculated Higgs and sparticle masses, the neutral Higgs masses h, H, A are pole masses calculated with the leading top, bottom, stop, sbottom loop contributions; while all other sparticle masses are running masses. This output is then used to compute the cosmological dark matter density of the lightest neutralino which is the LSP. The dark matter analysis is discussed in more detail in section 3.

Using χ^2 penalties⁶ we apply two additional constraints:

⁴Note, we have checked that switching to 2 loop RGEs for dimensionful parameters can be compensated for by small changes in the GUT scale parameters, without significant changes in the low energy results.

⁵Note we take a conservative error for $m_b(m_b)$ [18] in view of recent claims to much smaller error bars [19].

⁶In order to constrain the values of some physical observables in our χ^2 analysis, such as m_{t_1} or m_A , we add a significant contribution to the χ^2 function for values of these observables outside the desired range. We refer to this additional contribution as a χ^2 penalty. Minimization of χ^2 with Minuit, then pushes the fits to the desired range. Of course the χ^2 penalties then vanish.

- $m_{\tilde{t}_1} \geq 300 \text{ GeV}$
- m_A **fixed**.

The first is chosen to be consistent with $\text{BR}(B \rightarrow X_s \gamma)$ [2]. Note, although we do calculate $\text{BR}(B \rightarrow X_s \gamma)$, we do not use it as a constraint in the analysis. This is for two reasons — 1) this decay mode depends on 3-2 generation mixing which is model dependent and 2) it is not difficult to fit $\text{BR}(B \rightarrow X_s \gamma)$ for values of $m_{\tilde{t}_1} \geq 300 \text{ GeV}$. Hence, in order to be generally consistent with the measured value of $\text{BR}(B \rightarrow X_s \gamma)$, we impose $m_{\tilde{t}_1} \geq 300 \text{ GeV}$. With regards to the second constraint, since $\Omega_\chi h^2$ and $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ are both sensitive to the value of m_A , we fix it's value and present our results for different values of m_A .⁷

Finally, we apply the experimental limits:

- lower bound on the lightest chargino mass $m_{\chi^+} > 104 \text{ GeV}$,
- lower bound on the light Higgs mass $m_h > 111 \text{ GeV}$.

Note, because of the theoretical uncertainty in the calculation of m_h ($\sim 3 \text{ GeV}$), we conservatively impose $m_h > 111 \text{ GeV}$, instead of the LEP bound for SM Higgs $m_h > 114.4 \text{ GeV}$.

3. Cosmological Dark Matter Density

We compute the relic abundance $\Omega_\chi h^2$ of the lightest neutralino using exact expressions for neutralino pair annihilation into all allowed final-state channels, which are valid both near and further away from resonances and thresholds [21]. We further treat the neutralino coannihilation with the lightest chargino and next-to-lightest neutralino [22] and with the lighter stau [23] with similar precision. We only neglect the neutralino coannihilation with the stop which would only affect $\Omega_\chi h^2$ in the regions of parameter space which are uninteresting for other reasons, as we comment below. We solve the Boltzmann equation numerically as in [24] and compute $\Omega_\chi h^2$ with an error of a few per cent, which is comparable with today's accuracy on the observational side. The latest determinations of cosmological parameters [8] give $\Omega_M h^2 = 0.135^{+0.008}_{-0.009}$ for the total matter content and $\Omega_b h^2 = 0.0224 \pm 0.0009$ for the baryonic component. The difference, attributed to cold dark matter (CDM), is then

$$\Omega_{\text{CDM}} h^2 = 0.113 \pm 0.009, \quad (3.1)$$

which is significantly narrower than previous ranges. We then apply two constraints on the dark matter abundance:

- the upper bound $\Omega_\chi h^2 < 0.13$,
- 2σ preferred range $0.095 < \Omega_\chi h^2 < 0.13$.

⁷The calculation of $\text{BR}(B \rightarrow X_s \gamma)$ and $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ requires a model for fermion mass matrices. In the absence of such a model we use the observed CKM matrix elements to calculate these flavor violating branching ratios.




Figure 1: Contours of constant χ^2 for $m_{16} = 3$ TeV and $m_A = 500$ GeV. The red regions are excluded by $m_{\chi^+} < 104$ GeV (below and to the left of a black solid curve), $m_h < 111$ GeV (on the right) and by $\Omega_\chi h^2 > 0.13$. To the right of the black broken line one has $m_h < 114.4$ GeV. The green band corresponds to the preferred 2σ range $0.095 < \Omega_\chi h^2 < 0.13$, while the white regions below it correspond to $\Omega_\chi h^2 < 0.095$.

In figures 1–3 we present our results for different values of m_{16} and m_A in the μ , $M_{1/2}$ plane. In particular in figure 1 we present, for $m_{16} = 3$ TeV and $m_A = 500$ GeV, the (magenta) lines of constant χ^2 with the cosmologically preferred dark matter region (shaded green) satisfying $0.095 < \Omega_\chi h^2 < 0.13$. We find significant regions of parameter space which gives $\chi^2 \leq 2$, $\Omega_\chi h^2$ as above, and satisfies all other phenomenological constraints. In addition we have shaded (light red) the regions excluded by collider limits and by $\Omega_\chi h^2 > 0.13$.⁸

In figure 2 we present a more detailed analysis of the same $m_{16} = 3$ TeV, $m_A = 500$ GeV case given in figure 1. We now include lines of constant $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ (upper left), m_h (upper right), $\Omega_\chi h^2$ (lower left), and σ_p^{SI} (lower right). σ_p^{SI} is the spin independent neutralino dark matter cross-section relevant for direct dark matter searches. We now consider each one of these features further.

Recall, the branching ratio $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ is sensitive to the value of the CP odd Higgs mass m_A [9]. For $m_A = 500$ GeV, the branching ratio satisfies $2 \times 10^{-7} < \text{BR}(B_s \rightarrow \mu^+ \mu^-) < 8 \times 10^{-7}$ for acceptable values of $\Omega_\chi h^2$ and $\chi^2 < 2$. In a recent analysis it has been shown that, with an integrated luminosity of 15 fb^{-1} , CDF can discover this process

⁸A similar analysis was performed in the recent paper [4]. However, they were not able to find acceptable cosmological solutions. It appears that they did not find any acceptable solutions because they find Yukawa unification only for $M_{1/2} \sim 100$ GeV and large $\mu \sim 300$ GeV. In this region we probably would not find acceptable solutions for $\Omega_\chi h^2$, no matter what value we take for m_A .




Figure 2: Same as figure 1 with contours of constant $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ (upper left), m_h (upper right), $\Omega_\chi h^2$ (lower left) and σ_p^{SI} (lower right) for $m_{16} = 3$ TeV and $m_A = 500$ GeV.

if $\text{BR}(B_s \rightarrow \mu^+ \mu^-) > 1.2 \times 10^{-8}$ [25]. Hence most acceptable regions of parameter space lead to observable rates for $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$.

In figure 2 (upper right) we see that the light Higgs mass increases as $M_{1/2}$ decreases. In the acceptable regions of parameter space we find $116 < m_h < 121$ GeV. The value of the light Higgs mass is however fairly insensitive to m_{16} .

The cosmological relic abundance of the neutralino $\Omega_\chi h^2$ (figure 2 (lower left)) is primarily determined by the direct s-channel pair-annihilation into SM fermion pairs through the CP odd Higgs. Since all the sfermions are very heavy, their contribution to reducing the neutralino number density is strongly suppressed. In contrast, because of the coupling $A b \bar{b} \propto \tan \beta$ (and similarly for the τ 's), the A -resonance is effective and broad. Near $m_\chi \approx m_A/2$ it reduces $\Omega_\chi h^2$ down to allowed but uninterestingly small values $\ll 0.1$. As one moves away from the resonance, $\Omega_\chi h^2$ grows, reaches the preferred range




Figure 3: The same as in figure 1 but for $m_{16} = 2.5$ TeV and $m_A = 500$ GeV (upper left), $m_{16} = 5$ TeV and $m_A = 500$ GeV (upper right), $m_{16} = 2.5$ TeV and $m_A = 300$ GeV (lower left) and $m_{16} = 3$ TeV and $m_A = 300$ GeV (lower right). Also marked are contours of constant $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$. The blue regions in the lower two panels are excluded by $\text{BR}(B_s \rightarrow \mu^+ \mu^-) > 2.6 \times 10^{-6}$. Note the different mass ranges for $M_{1/2}$ and μ in the different panels.

$0.095 < \Omega_\chi h^2 < 0.13$, before becoming too large $\Omega_\chi h^2 > 0.13$.⁹ (A similar, but much more narrow resonance due to h^0 is also present at $M_{1/2} \approx 150$ GeV and small μ .) When $m_\chi \gtrsim m_t$ ($M_{1/2} \gtrsim 420$ GeV) and the stops are not too heavy, the LSP pairs annihilate to $t\bar{t}$ -pairs. In the region of large $M_{1/2}$, often where m_h is already too low, two additional channels become effective. First, in this region the neutralino becomes almost mass degenerate with the lighter stau which leads to reducing $\Omega_\chi h^2$ through coannihilation. Second, if

⁹Note that at one loop we have $M_1(M_Z) = M_{1/2} * \alpha_1(M_Z)/\alpha_G$ so $M_1(M_Z) \approx 0.4M_{1/2}$. For bino-like neutralino (which is true for larger μ), we thus have $m_\chi \approx 0.4M_{1/2}$. Hence for s-channel annihilation we have $m_A \approx 2m_\chi \approx 0.8M_{1/2}$ or $M_{1/2} \approx (5/4)m_A$ for the position of the “peak suppression.”

m_A is not too large, neutralino pair-annihilation into Higgs boson pairs AA and HH opens up. Finally, at $\mu \ll M_{1/2}$, the relic abundance is strongly reduced due to the increasing higgsino component of the LSP.

Finally, the spin independent neutralino cross-section σ_p^{SI} in the lower right window of figure 2 is predominantly determined by the contribution of the heavy CP even scalar t -channel exchange to both tree-level and one-loop diagrams. Note that in the preferred region of $\chi^2 < 2$ and $0.095 < \Omega_\chi h^2 < 0.13$ we find $10^{-9} \text{ pb} \lesssim \sigma_p^{SI} \lesssim 10^{-7} \text{ pb}$. We will comment further on our predictions for σ_p^{SI} below.

In figure 3 we display the dependence of our constraints, $\Omega_\chi h^2$ and $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ on m_{16} and m_A . By comparing the upper two windows with figure 1 we can see that, as m_{16} increases, the region with $\chi^2 < 2$ rapidly grows. Note, the dominant pull in χ^2 is due to the bottom quark mass. In order to fit the data, the total SUSY corrections to $m_b(m_b)$ must be of order $-(2-4)\%$ [2]. In addition there are three dominant contributions to these SUSY corrections, a gluino loop contribution $\propto \alpha_3 \mu M_{\tilde{g}} \tan \beta / m_{\tilde{b}_1}^2$, a chargino loop contribution $\propto \lambda_t^2 \mu A_t \tan \beta / m_{\tilde{t}_1}^2$, and a term $\propto \log M_{\text{SUSY}}^2$. When m_{16} increases (with $M_{1/2}$ fixed) the parameter A_t becomes more negative, since $A_0 \approx -2m_{16}$ and $A_t \approx -3M_{1/2} + \epsilon A_0$ where $\epsilon \ll 1$. Also, larger values of m_{16} permit a larger range for the ratio $m_{\tilde{b}_1}/m_{\tilde{t}_1}$. Thus larger values of m_{16} allows more freedom in parameter space for fitting the data at both smaller or larger values of μ , $M_{1/2}$.

In the lower two windows in figure 3 we consider two regions with $m_A = 300 \text{ GeV}$ with $m_{16} = 2.5 \text{ TeV}$ (lower left) and $m_{16} = 3 \text{ TeV}$ (lower right). The blue regions are excluded by the CDF bound $\text{BR}(B_s \rightarrow \mu^+ \mu^-) < 2.6 \times 10^{-6}$ [10]. Note for $m_{16} = 2.5 \text{ TeV}$, the region with $\chi^2 < 2$ does not overlap the region with acceptable dark matter abundance (green shaded). However for $m_{16} = 3 \text{ TeV}$, $m_A = 300 \text{ GeV}$, figure 3 (lower right), we find a small region with acceptable $\chi^2 < 1$ and $\Omega_\chi h^2$. Moreover the branching ratio $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ is now close to the CDF bound. On the other hand for $m_{16} = 5 \text{ TeV}$, $m_A = 500 \text{ GeV}$, in figure 3 (upper right), a new region of parameter space consistent with all the data now opens up with larger μ , $M_{1/2}$. This new region becomes cosmologically allowed due to neutralino annihilation into Higgs boson pairs AA and HH and due to coannihilation with the lighter stau.

Hence we see that increasing m_A has two effects. It suppresses the branching fraction $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$. At the same time it moves the s-channel neutralino annihilation channel to larger values of $M_{1/2}$; hence providing larger regions with $0.095 < \Omega_\chi h^2 < 0.13$ (compare figures 1 and 3 (lower right) or figures 3 (upper and lower left)). In fact, the two (green) branches of the preferred range $0.095 < \Omega_\chi h^2 < 0.13$ correspond to the two sides (except for the upper left window of figure 3 where just one side is evident) of the wide A resonance in the neutralino pair-annihilation. On the other hand, increasing m_A above 1 TeV or so would move the regions of preferred $\Omega_\chi h^2$ too far to the right, in potential conflict with a lower bound on m_h .

Finally, we comment on $\text{BR}(B \rightarrow X_s \gamma)$. The current experimental range [26, 27] is $\text{BR}(B \rightarrow X_s \gamma)_{\text{expt}} = (3.41 \pm 0.36) \times 10^{-4}$, while the SM prediction, including full NLO QCD corrections [28, 29], is $\text{BR}(B \rightarrow X_s \gamma)_{\text{SM}} = (3.70 \pm 0.30) \times 10^{-4}$. In computing the SUSY contribution to $b \rightarrow s \gamma$ we further include full LO and dominant NLO-level $\tan \beta$ -




Figure 4: The upper and lower limits on $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ as a function of m_A in the $\mu, M_{1/2}$ region of parameter space satisfying all the collider constraints, $0.095 < \Omega_\chi h^2 < 0.13$ and $\chi^2 < 2$ for fixed $m_{16} = 3 \text{ TeV}$.

enhanced contributions [30, 31]. Conservatively allowing for the SM+SUSY contribution to be in the range $(3.41 \pm 0.67) \times 10^{-4}$ [32, 27] selects a band $300 \text{ GeV} \lesssim \mu \lesssim 400 \text{ GeV}$ which slowly decreases with increasing $M_{1/2}$. (This remains approximately true for all the cases that we have analyzed except $m_{16} = 5 \text{ TeV}$ and $m_A = 500 \text{ GeV}$ where one finds a narrower range at $\mu \lesssim 200 \text{ GeV}$.) However, $\text{BR}(B \rightarrow X_s \gamma)$ is strongly sensitive to the 2-3 generation down-type squark mixings [27] which are model dependent and which we do not include here. In summary, the process is generally consistent with the most preferred regions of $M_{1/2}$ and μ but we do not use it here as a constraint, since it can be easily relaxed by employing parameters which are less relevant for our analysis.

To summarize, we find that regions satisfying all three constraints exist for $m_{16} \geq 3 \text{ TeV}$ and $m_A \geq 300 \text{ GeV}$. The acceptable range for $\mu, M_{1/2}$ grows with increasing m_A (for fixed $m_{16} = 3 \text{ TeV}$) from approximately $260 \text{ GeV} \leq \mu \leq 295 \text{ GeV}$, $140 \text{ GeV} \leq M_{1/2} \leq 200 \text{ GeV}$ for $m_A = 300 \text{ GeV}$ to $160 \text{ GeV} \leq \mu \leq 520 \text{ GeV}$, $140 \text{ GeV} \leq M_{1/2} \leq 530 \text{ GeV}$ for $m_A = 500 \text{ GeV}$. In addition, the allowed regions grow as m_{16} increases and for $m_A = 500 \text{ GeV}$, $m_{16} = 5 \text{ TeV}$ there is also a large $\mu, M_{1/2}$ solution satisfying $480 \text{ GeV} \leq \mu \leq 820 \text{ GeV}$, $640 \text{ GeV} \leq M_{1/2} \leq 1020 \text{ GeV}$. In the table 1 we present the input and output data from the χ^2 analysis for three points satisfying all the phenomenological constraints.

4. Predictions and summary

In this paper we have analyzed the MSO₁₀SM and found regions of soft SUSY breaking parameter space which fit precision electroweak data, including the top, bottom and

Data points		1	2	3
Input parameters				
α_G^{-1}		24.66	24.92	25.28
$M_G \times 10^{-16}$		3.51	2.83	2.43
ϵ_3		-0.038	-0.034	-0.029
λ		0.66	0.66	0.66
m_{16}		3000	3000	5000
m_{10}/m_{16}		1.30	1.33	1.33
Δm_H^2		0.14	0.15	0.14
$M_{1/2}$		180	400	700
μ		270	350	600
$\tan \beta$		50.9	50.6	50.5
A_0/m_{16}		-1.85	-1.88	-1.91
χ^2 observables	Exp (σ)			
M_Z	91.188 (0.091)	91.18	91.19	91.20
M_W	80.419 (0.080)	80.42	80.42	80.41
$G_\mu \times 10^5$	1.1664 (0.0012)	1.166	1.166	1.166
α_{EM}^{-1}	137.04 (0.14)	137.0	137.0	137.0
$\alpha_s(M_Z)$	0.118 (0.002)	0.1177	0.1176	0.1179
$\rho_{new} \times 10^3$	-0.200 (1.10)	0.427	0.498	0.162
M_t	174.3 (5.1)	173.9	174.7	174.7
$m_b(m_b)$	4.20 (0.20)	4.28	4.28	4.21
M_τ	1.7770 (0.0018)	1.777	1.777	1.777
TOTAL χ^2		0.53	0.61	0.13
h		120	119	117
H		329	556	557
A		299	499	501
H^+		329	540	541
χ_1^0		72	163	293
χ_2^0		133	288	536
χ_1^+		133	287	535
\tilde{g}		474	1032	1768
\tilde{t}_1		300	300	576
\tilde{b}_1		679	736	1262
$\tilde{\tau}_1$		870	721	1180
$a_\mu^{SUSY} \times 10^{10}$	25.6 (16)	2.7	2.8	1.0
$\Omega_\chi h^2$	0.095 - 0.130	0.099	0.130	0.097
$\sigma_p^{SI}(pb) \times 10^7$		1.020	0.158	0.049
$\text{BR}(B_s \rightarrow \mu^+ \mu^-) \times 10^6$	< 2.6	2.58	0.61	0.66
$\text{BR}(B \rightarrow X_s \gamma) \times 10^4$	3.41 (0.67)	5.36	4.34	0.81

Table 1: We present the results of our χ^2 analysis for three different points (labelled 1 - 3), distinguished by the fixed values of μ , $M_{1/2}$ and m_{16} . We give the 8 input parameters, which were varied to minimize χ^2 ; a function of the 9 experimental observables listed in the table. Finally, we show the SUSY spectrum and predictions for several observables.




Figure 5: Predictions for σ_p^{SI} vs. m_χ for different choices of m_{16} and m_A , subject to the collider constraints, $0.095 < \Omega_\chi h^2 < 0.13$ and $\chi^2 < 3$. The green bands are for $m_{16} = 2.5$ TeV, the red for 3 TeV and the blue for 5 TeV. The lighter shading is for $m_A = 300$ GeV, the darker for $m_A = 500$ GeV. In the last case ($m_{16} = 5$ TeV and $m_A = 500$ GeV) there are two branches which correspond to the two cosmologically preferred regions in the upper right panel in figure 3.

tau masses and, in addition, fit the cosmological dark matter abundance for the neutralino LSP and satisfy $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$. Generically, we find solutions to all the constraints with $m_{16} \geq 3$ TeV. The squark and slepton masses have an inverted scalar mass hierarchy with the first and second generation scalar masses of order m_{16} , while the third generation has mass less than 1.3 TeV for $m_{16} = 5$ TeV. This nice feature of the model suppresses SUSY CP and flavor problems. In addition the gaugino masses are typically much lighter, except for the large μ , $M_{1/2}$ region for $m_{16} = 5$ TeV with a gluino mass of order 1.7 TeV (see spectrum in table for selected acceptable points).

Note an immediate consequence of such heavy first and second generation sleptons is the suppression of the SUSY contribution to the anomalous magnetic moment of the muon. We find $a_\mu^{SUSY} \leq 2.8 \times 10^{-10}$ (see table 1). This is consistent with the most recent experimental [33] and theoretical results at 1σ if one uses τ -based analysis [34]. However it is only consistent with an e^+e^- -based analysis at 3σ .

Another interesting result is the enhanced branching ratio for the process $B_s \rightarrow \mu^+ \mu^-$. In figure 4 we show the ranges of values of $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ in the low μ , $M_{1/2}$ region of parameter space satisfying all the phenomenological constraints with $0.095 < \Omega_\chi h^2 < 0.13$ and $\chi^2 < 2$ as a function of m_A for fixed $m_{16} = 3$ TeV. The horizontal red line is the CDF bound. Over a significant region of parameter space $\text{BR}(B_s \rightarrow \mu^+ \mu^-) > 1 \times 10^{-7}$ and may be observable at the Tevatron (Run II) [25].

Finally in figure 5 we present the cross-section for elastic neutralino-proton scattering due to scalar interactions σ_p^{SI} for all regions satisfying the collider constraints, $0.095 <$

$\Omega_\chi h^2 < 0.13$ and $\chi^2 < 2$. The green bands are for $m_{16} = 2.5$ TeV, the red for 3 TeV and the blue for 5 TeV. The lighter shading is for $m_A = 300$ GeV, the darker for $m_A = 500$ GeV. In the last case ($m_{16} = 5$ TeV and $m_A = 500$ GeV) there are two branches which correspond to the two cosmologically preferred regions in the upper right panel in figure 3. Note that lower m_A generally gives larger σ_p^{SI} as expected. For comparison, we also show the bounds from the present dark matter searches and the predictions of the general MSSM [35]. (Other recent studies of σ_p^{SI} in the case of non-universal Higgs mass in a variant of the CMSSM can be found in [36].) Over the next two to five years the experimental sensitivity is expected to gradually improve by some three orders of magnitude. This will cover large parts of the predicted ranges of σ_p^{SI} , especially at lower values of m_A where $B_s \rightarrow \mu^+ \mu^-$ will be accessible at the Tevatron (Run II).

Acknowledgments

We gratefully acknowledge the use of the GUT χ^2 analysis code developed by T. Blažek and a contribution from Y.G. Kim to a routine for computing $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$. L.R. is grateful to A. Melchiorri for helpful comments regarding the recent WMAP analysis. R.D. is supported, in part, by the U.S. Department of Energy, Contract DE-FG03-91ER-40674 and the Davis Institute for High Energy Physics. S.R. received partial support from DOE grant# DOE/ER/01545-841 and from a grant in aid from the Monell Foundation. He is also grateful to the hospitality shown to him by the School of Natural Sciences, Institute for Advanced Study. RRdA and LR are supported in part by the EU Fifth Framework network "Supersymmetry and the Early Universe" (HPRN-CT-2000-00152).

References

- [1] G.L. Kane, C.F. Kolda, L. Roszkowski and J.D. Wells, *Study of constrained minimal supersymmetry*, *Phys. Rev. D* **49** (1994) 6173 [[hep-ph/9312272](#)].
- [2] T. Blazek, R. Dermisek and S. Raby, *Predictions for Higgs and SUSY spectra from SO(10) yukawa unification with $\mu > 0$* , *Phys. Rev. Lett.* **88** (2002) 111804 [[hep-ph/0107097](#)]; *Yukawa unification in SO(10)*, *Phys. Rev. D* **65** (2002) 115004 [[hep-ph/0201081](#)].
- [3] K. Tobe and J.D. Wells, *Revisiting top-bottom-tau Yukawa unification in supersymmetric grand unified theories*, [hep-ph/0301015](#).
- [4] D. Auto et al., *Yukawa coupling unification in supersymmetric models*, [hep-ph/0302155](#).
- [5] M. Olechowski and S. Pokorski, *Electroweak symmetry breaking with nonuniversal scalar soft terms and large $\tan \beta$ solutions*, *Phys. Lett. B* **344** (1995) 201 [[hep-ph/9407404](#)];
D. Matalliotakis and H.P. Nilles, *Implications of nonuniversality of soft terms in supersymmetric grand unified theories*, *Nucl. Phys. B* **435** (1995) 115 [[hep-ph/9407251](#)];
N. Polonsky and A. Pomarol, *Nonuniversal GUT corrections to the soft terms and their implications in supergravity models*, *Phys. Rev. D* **51** (1995) 6532 [[hep-ph/9410231](#)];
H. Murayama, M. Olechowski and S. Pokorski, *Viable t - b - τ yukawa unification in SUSY SO(10)*, *Phys. Lett. B* **371** (1996) 57 [[hep-ph/9510327](#)];
R. Rattazzi and U. Sarid, *The unified minimal supersymmetric model with large Yukawa couplings*, *Phys. Rev. D* **53** (1996) 1553 [[hep-ph/9505428](#)].

- [6] U. Chattopadhyay, A. Corsetti and P. Nath, *Supersymmetric dark matter and Yukawa unification*, *Phys. Rev. D* **66** (2002) 035003 [[hep-ph/0201001](#)].
- [7] C. Pallis and M.E. Gomez, *Yukawa quasi-unification and neutralino relic density*, [hep-ph/0303098](#).
- [8] D.N. Spergel et al., *First year wilkinson microwave anisotropy probe (WMAP) observations: determination of cosmological parameters*, [astro-ph/0302209](#).
- [9] C. Hamzaoui, M. Pospelov and M. Toharia, *Higgs-mediated FCNC in supersymmetric models with large $\tan\beta$* , *Phys. Rev. D* **59** (1999) 095005 [[hep-ph/9807350](#)];
K.S. Babu and C.F. Kolda, *Higgs-mediated $B0 \rightarrow \mu^+\mu^-$ in minimal supersymmetry*, *Phys. Rev. Lett.* **84** (2000) 228 [[hep-ph/9909476](#)];
P.H. Chankowski and L. Slawianowska, *$B0/d, s \rightarrow \mu^-\mu^+$ decay in the MSSM*, *Phys. Rev. D* **63** (2001) 054012 [[hep-ph/0008046](#)];
C. Bobeth, T. Ewerth, F. Kruger and J. Urban, *Analysis of neutral Higgs-boson contributions to the decays $\text{anti-}b/s \rightarrow \ell^+\ell^-$ and $\text{anti-}b \rightarrow k \ell^+\ell^-$* , *Phys. Rev. D* **64** (2001) 074014 [[hep-ph/0104284](#)]; *Enhancement of $b(\text{anti-}b/d \rightarrow \mu^+\mu^-)/b(\text{anti-}b/s \rightarrow \mu^+\mu^-)$ in the MSSM with minimal flavour violation and large $\tan\beta$* , *Phys. Rev. D* **66** (2002) 074021 [[hep-ph/0204225](#)];
A. Dedes, H.K. Dreiner and U. Nierste, *Correlation of $b/s \rightarrow \mu^+\mu^-$ and $(g-2)(\mu)$ in minimal supergravity*, *Phys. Rev. Lett.* **87** (2001) 251804 [[hep-ph/0108037](#)];
G. Isidori and A. Retico, *Scalar flavour-changing neutral currents in the large- $\tan\beta$ limit*, *J. High Energy Phys.* **11** (2001) 001 [[hep-ph/0110121](#)];
A.J. Buras, P.H. Chankowski, J. Rosiek and L. Slawianowska, *Correlation between $\delta m(s)$ and $b/(s,d)0 \rightarrow \mu^+\mu^-$ in supersymmetry at large $\tan\beta$* , *Phys. Lett. B* **546** (2002) 96 [[hep-ph/0207241](#)].
- [10] CDF collaboration, F. Abe et al., *Search for the decays $b/d0 \rightarrow \mu^+\mu^-$ and $b/s0 \rightarrow \mu^+\mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.8 \text{ tev}$* , *Phys. Rev. D* **57** (1998) 3811.
- [11] C. Balazs and R. Dermisek, *Yukawa coupling unification and non-universal gaugino mediation of supersymmetry breaking*, [hep-ph/0303161](#).
- [12] J.A. Bagger, J.L. Feng, N. Polonsky and R.-J. Zhang, *Superheavy supersymmetry from scalar mass a -parameter fixed points*, *Phys. Lett. B* **473** (2000) 264 [[hep-ph/9911255](#)].
- [13] F. Gabbiani, E. Gabrielli, A. Masiero and L. Silvestrini, *A complete analysis of FCNC and CP constraints in general SUSY extensions of the standard model*, *Nucl. Phys. B* **477** (1996) 321 [[hep-ph/9604387](#)];
T. Besmer, C. Greub and T. Hurth, *Bounds on flavor violating parameters in supersymmetry*, *Nucl. Phys. B* **609** (2001) 359 [[hep-ph/0105292](#)].
- [14] E. Kearns, Snowmass 2001.
- [15] R. Dermisek, A. Mafi and S. Raby, *SUSY GUTs under siege: proton decay*, *Phys. Rev. D* **63** (2001) 035001 [[hep-ph/0007213](#)].
- [16] T. Blazek, M. Carena, S. Raby and C.E.M. Wagner, *A global χ^2 analysis of electroweak data in SO(10) SUSY GUTs*, *Phys. Rev. D* **56** (1997) 6919 [[hep-ph/9611217](#)].
- [17] H.E. Haber and R. Hempfling, *The renormalization group improved Higgs sector of the minimal supersymmetric model*, *Phys. Rev. D* **48** (1993) 4280 [[hep-ph/9307201](#)];
M. Carena, J.R. Espinosa, M. Quiros and C.E.M. Wagner, *Analytical expressions for radiatively corrected Higgs masses and couplings in the MSSM*, *Phys. Lett. B* **355** (1995) 209

- [hep-ph/9504316]; *Effective potential methods and the Higgs mass spectrum in the MSSM*, *Nucl. Phys. B* **461** (1996) 407 [hep-ph/9508343].
- [18] D.E. Groom et al., *The review of particle physics*, *Eur. Phys. J. C* **15** (2000) 1.
- [19] M. Beneke and A. Signer, *The bottom \overline{MS} quark mass from sum rules at next-to-next-to-leading order*, *Phys. Lett. B* **471** (1999) 233 [hep-ph/9906475];
G. Corcella and A.H. Hoang, *Uncertainties in the \overline{MS} bottom quark mass from relativistic sum rules*, *Phys. Lett. B* **554** (2003) 133 [hep-ph/0212297].
- [20] P. Langacker, talk at Chicagoland seminar, October 1999.
- [21] T. Nihei, L. Roszkowski and R. Ruiz de Austri, *Towards an accurate calculation of the neutralino relic density*, *J. High Energy Phys.* **05** (2001) 063 [hep-ph/0102308]; *Exact cross sections for the neutralino wimp pair-annihilation*, *J. High Energy Phys.* **03** (2002) 031 [hep-ph/0202009].
- [22] J. Edsjo and P. Gondolo, *Neutralino relic density including coannihilations*, *Phys. Rev. D* **56** (1997) 1879 [hep-ph/9704361].
- [23] T. Nihei, L. Roszkowski and R. Ruiz de Austri, *Exact cross sections for the neutralino slepton coannihilation*, *J. High Energy Phys.* **07** (2002) 024 [hep-ph/0206266].
- [24] P. Gondolo, J. Edsjo, L. Bergstrom, P. Ullio, and T. Baltz,
<http://www.physto.se/edsjo/darksusy/>
- [25] R. Arnowitt, B. Dutta, T. Kamon and M. Tanaka, *Detection of $b/s \rightarrow \mu^+ \mu^-$ at the Tevatron run ii and constraints on the SUSY parameter space*, *Phys. Lett. B* **538** (2002) 121 [hep-ph/0203069].
- [26] CLEO collaboration, S. Chen et al., *Branching fraction and photon energy spectrum for $b \rightarrow s\gamma$* , *Phys. Rev. Lett.* **87** (2001) 251807 [hep-ex/0108032];
ALEPH collaboration, R. Barate et al., *A measurement of the inclusive $b \rightarrow s\gamma$ branching ratio*, *Phys. Lett. B* **429** (1998) 169;
BELLE collaboration, K. Abe et al., *A measurement of the branching fraction for the inclusive $b \rightarrow x/s$ gamma decays with belle*, *Phys. Lett. B* **511** (2001) 151 [hep-ex/0103042];
BABAR collaboration, C. Jessop et al., talk at ICHEP-02, Amsterdam, July 2002.
- [27] K.-i. Okumura and L. Roszkowski, *De-constraining supersymmetry from $b \rightarrow s\gamma$?*,
hep-ph/0208101.
- [28] P. Gambino and M. Misiak, *Quark mass effects in anti- $b \rightarrow x/s$ gamma*, *Nucl. Phys. B* **611** (2001) 338 [hep-ph/0104034].
- [29] A.J. Buras, A. Czarnecki, M. Misiak and J. Urban, *Completing the NLO QCD calculation of anti- $b \rightarrow x/s$ gamma*, *Nucl. Phys. B* **631** (2002) 219 [hep-ph/0203135].
- [30] G. Degrandi, P. Gambino and G.F. Giudice, *$B \rightarrow x/s$ gamma in supersymmetry: large contributions beyond the leading order*, *J. High Energy Phys.* **12** (2000) 009 [hep-ph/0009337].
- [31] M. Carena, D. Garcia, U. Nierste and C.E.M. Wagner, *$b \rightarrow s\gamma$ and supersymmetry with large $\tan\beta$* , *Phys. Lett. B* **499** (2001) 141 [hep-ph/0010003].
- [32] L. Roszkowski, R. Ruiz de Austri and T. Nihei, *New cosmological and experimental constraints on the CMSSM*, *J. High Energy Phys.* **08** (2001) 024 [hep-ph/0106334].

- [33] MUON G-2 collaboration, G.W. Bennett et al., *Measurement of the positive muon anomalous magnetic moment to 0.7 PPM*, *Phys. Rev. Lett.* **89** (2002) 101804 [[hep-ex/0208001](#)], erratum *ibid.* **89** (2002) 129903.
- [34] M. Davier, S. Eidelman, A. Hocker and Z. Zhang, *Confronting spectral functions from e^+e^- annihilation and tau decays: consequences for the muon magnetic moment*, *Eur. Phys. J. C* **27** (2003) 497 [[hep-ph/0208177](#)].
- [35] Y.G. Kim, T. Nihei, L. Roszkowski and R. Ruiz de Austri, *Upper and lower limits on neutralino wimp mass and Spin-independent scattering cross section and impact of new $(g-2)_\mu$ measurement*, *J. High Energy Phys.* **12** (2002) 034 [[hep-ph/0208069](#)].
- [36] J.R. Ellis, A. Ferstl, K.A. Olive and Y. Santoso, *Direct detection of dark matter in the MSSM with non-universal Higgs masses*, [hep-ph/0302032](#);
V. Bertin, E. Nezri and J. Orloff, *Neutralino dark matter beyond CMSSM universality*, *J. High Energy Phys.* **02** (2003) 046 [[hep-ph/0210034](#)].