

New formalism for QCD parton showers

To cite this article: Stefan Gieseke *et al* JHEP12(2003)045

View the [article online](#) for updates and enhancements.

You may also like

- [Jet substructure at the Tevatron and LHC: new results, new tools, new benchmarks](#)
A Altheimer, S Arora, L Asquith et al.
- [PYTHIA 6.4 physics and manual](#)
Torbjörn Sjöstrand, Stephen Mrenna and Peter Skands
- [QCD resummation for hadronic final states](#)
Gionata Luisoni and Simone Marzani

New formalism for QCD parton showers

Stefan Gieseke,^a Philip Stephens^a and Bryan Webber^{ab}

^a*Cavendish Laboratory, University of Cambridge
Madingley Road, Cambridge, CB3 0HE, U.K.*

^b*Theory Division, CERN
1211 Geneva 23, Switzerland*

*E-mail: gieseke@hep.phy.cam.ac.uk, stephens@hep.phy.cam.ac.uk,
webber@hep.phy.cam.ac.uk*

ABSTRACT: We present a new formalism for parton shower simulation of QCD jets, which incorporates the following features: covariant kinematics, improved treatment of heavy quark fragmentation, angular-ordered evolution with soft gluon coherence, more accurate soft gluon angular distributions, and better coverage of phase space. It is implemented in the new **Herwig++** event generator.

KEYWORDS: QCD, Jets, Heavy Quarks Physics.

Contents

1. Introduction	2
2. New variables for parton branching	3
2.1 Final-state quark branching	3
2.1.1 Kinematics	3
2.1.2 Running coupling	4
2.1.3 Evolution variable	4
2.1.4 Branching probability	5
2.2 Gluon splitting	5
2.3 Initial-state branching	6
2.4 Allowed regions and termination of branching	6
2.5 Treatment of colour flows	8
3. Final-final colour connection	8
3.1 Phase space variables	9
3.2 Soft gluon region	10
3.3 Example: $e^+e^- \rightarrow q\bar{q}g$	11
3.3.1 Exact matrix element	12
3.3.2 Soft gluon distribution	13
3.3.3 Dead region contribution	14
4. Initial-initial colour connection	15
4.1 Phase space variables	16
4.2 Example: Drell-Yan process	16
5. Initial-final colour connection	18
5.1 Initial-state branching	18
5.2 Final-state branching	19
5.3 Phase space variables	19
5.4 Example: deep inelastic scattering	20
5.5 Example: $q\bar{q} \rightarrow t\bar{t}$	21
6. Decay colour connection	23
6.1 Initial-state branching	23
6.2 Final-state branching	24
6.3 Phase space variables	24
6.4 Example: top decay	25
7. Conclusions	28

1. Introduction

The parton shower approximation has become a key component of a wide range of comparisons between theory and experiment in particle physics. Calculations of infrared-safe observables, i.e. those that are asymptotically insensitive to soft physics, can be performed in fixed-order perturbation theory, but the resulting final states consist of a few isolated partons, quite unlike the multihadron final states observed experimentally. One can attempt to identify isolated partons with hadronic jets, but then the energy flows within and between jets are not well represented.

At present, the only means of connecting few-parton states with the real world is via parton showers, which generate high-multiplicity partonic final states in an approximation that retains enhanced collinear and soft contributions to all orders. Such multiparton states can be interfaced to a hadronization model which does not require large momentum transfers in order to produce a realistic hadronic final state. Hadronization and detector corrections to the fixed-order predictions can then be computed, and the results have generally been found to be in satisfactory agreement with the data. Infrared-sensitive quantities such as hadron spectra and multiplicities have also been described successfully using parton showers. This increases confidence that similar techniques can be used to predict new physics signals and backgrounds in future experiments.

A crucial ingredient of modern parton showering algorithms¹ is *angular ordering*, which ensures that important aspects of soft gluon coherence are included in an azimuthally-averaged form. The angular shower evolution variable [2] used in the event generator program HERWIG [3] is good for ensuring that angular ordering is built in from the outset, but the phase space is complicated and not invariant under any kind of boosts. Evolution in virtuality looks natural but then angular ordering must be imposed afterwards, as is done in PYTHIA [4].

In the present paper we investigate a new shower evolution formalism, based on an angular variable related to transverse momentum [5]–[9]. The main aim is to retain the direct angular ordering of the shower while improving the Lorentz invariance of the evolution and simplifying the coverage of phase space, especially in the soft region. The new shower variables also permit a better treatment of heavy quark fragmentation, through evolution down to zero transverse momentum and the use of mass-dependent splitting functions, which eliminate the sharply-defined collinear “dead cones” seen in earlier angular-ordered treatments.²

In the following section we define the new shower variables and their associated kinematics and dynamics, including the appropriate argument of the running coupling, the mass-dependent parton branching probability, and the shower evolution cutoff. The variables are defined slightly differently for initial- and final-state parton branching, and depend on the colour connection of the evolving parton, so we consider in subsequent sections the various possible configurations of colour flow between initial and final jets.

¹For a general introduction to the parton shower approximation, see for example Chapter 5 of [1].

²Improvements to heavy quark fragmentation in the framework of the PYTHIA parton shower are presented in ref. [10].

The formalism presented here is implemented in the new Monte Carlo event generator **Herwig++** [11]. Results for e^+e^- annihilation and comparisons with LEP data are presented in a separate publication [12]. The formulae in the present paper could also be used to construct a matching scheme for next-to-leading order (NLO) QCD calculations and **Herwig++** parton showers, similar to that developed for HERWIG showers in [13, 14] and implemented in the MC@NLO event generator [15], or to improve schemes for combining parton showers with fixed-order matrix elements [16].

2. New variables for parton branching

2.1 Final-state quark branching

2.1.1 Kinematics

Consider parton branching in an outgoing quark jet. For improved treatment of quasi-collinear emission from a massive quark, following ref. [8], we define the quark momentum after the i th gluon emission $q_{i-1} \rightarrow q_i + k_i$ (see figure 1) as




Figure 1: Final-state parton branching. The blob represents the hard subprocess.

$$q_i = \alpha_i p + \beta_i n + q_{\perp i} \quad (2.1)$$

where p is the jet’s “parent parton” momentum ($p^2 = m^2$, the *on-shell* quark mass-squared), n is a lightlike “backward” 4-vector ($n^2 = 0$), and $q_{\perp i}$ is the transverse momentum ($q_{\perp i}^2 = -\mathbf{q}_{\perp i}^2$, $q_{\perp i} \cdot p = q_{\perp i} \cdot n = 0$). Then

$$\beta_i = \frac{\mathbf{q}_{\perp i}^2 + q_i^2 - \alpha_i^2 m^2}{2\alpha_i p \cdot n}. \quad (2.2)$$

If the “backward” vector n were not lightlike, eq. (2.2) and the ensuing kinematics would be much more complicated. Ideally, n should point in the direction of the parton that is colour-connected to the quark, forming a colour dipole [8]. If the colour-connected parton is massive, this clearly cannot be the case in all frames of reference. However, as will be illustrated later, we can always define a lightlike n that is close to the required direction by forming an appropriate linear combination of the colour-connected 4-momenta.

The momentum fraction z and relative transverse momentum \mathbf{p}_{\perp} are now defined by

$$z_i = \frac{\alpha_i}{\alpha_{i-1}}, \quad \mathbf{p}_{\perp i} = \mathbf{q}_{\perp i} - z_i \mathbf{q}_{\perp i-1}. \quad (2.3)$$

Then we have

$$q_{i-1}^2 = \frac{q_i^2}{z_i} + \frac{k_i^2}{1 - z_i} + \frac{\mathbf{p}_{\perp i}^2}{z_i(1 - z_i)}. \quad (2.4)$$

2.1.2 Running coupling

To find the optimal argument of α_s , we consider the branching of a quark of virtuality q^2 into an on-shell quark and an off-shell gluon of virtuality k^2 [17]. From eq. (2.4), the quark propagator denominator is

$$q^2 - m^2 = \frac{1-z}{z}m^2 + \frac{k^2}{1-z} + \frac{\mathbf{p}_\perp^2}{z(1-z)} = \frac{1}{1-z} \left\{ k^2 + \frac{1}{z} [\mathbf{p}_\perp^2 + (1-z)^2 m^2] \right\}. \quad (2.5)$$

Thus the scale of gluon virtualities that predominate in the parton cascade, which determine the argument of α_s , is set by $[\mathbf{p}_\perp^2 + (1-z)^2 m^2]/z$. Since there are no soft quark ($z \rightarrow 0$) singularities, we may neglect the factor of $1/z$. Therefore the suggested argument of α_s is $\mathbf{p}_\perp^2 + (1-z)^2 m^2$. This is the natural extension of the usual argument \mathbf{p}_\perp^2 [18] to the massive case: collinear gluon emission from a massive quark remains in the perturbative region provided $1-z \gg \Lambda/m$. In practice we impose a minimum virtuality on light quarks and gluons in the parton shower, and therefore the actual argument used for α_s is slightly more complicated (see below).

2.1.3 Evolution variable

The evolution variable is not simply q^2 since this would ignore angular ordering. For massless parton branching this means the evolution variable should be $\mathbf{p}_\perp^2/[z(1-z)]^2 = q^2/z(1-z)$ [21]. For gluon emission by a massive quark we assume this generalizes to $(q^2 - m^2)/z(1-z)$. To define a resolvable emission we also need to introduce a minimum virtuality Q_g^2 for gluons and light quarks. Therefore from eq. (2.5) the evolution variable is

$$\tilde{q}^2 = \frac{\mathbf{p}_\perp^2}{z^2(1-z)^2} + \frac{\mu^2}{z^2} + \frac{Q_g^2}{z(1-z)^2} \quad (2.6)$$

where $\mu = \max(m, Q_g)$. For the argument of the running coupling we use

$$z^2(1-z)^2\tilde{q}^2 = \mathbf{p}_\perp^2 + (1-z)^2\mu^2 + zQ_g^2. \quad (2.7)$$

Note that for massive quarks this allows us to evolve down to $\mathbf{p}_\perp = 0$, provided the emitted gluon is not too soft. As is well known, gluon emission is dynamically suppressed at $\mathbf{p}_\perp < (1-z)m$, i.e. inside the *dead cone* [19, 20]. In previous angular ordered treatments, such as that in HERWIG, this region was simply cut out of the phase space. As we shall see, the present treatment enables us to describe the distribution of gluon emission inside the dead cone without any sharp angular cutoff.

Angular ordering of the branching $q_i \rightarrow q_{i+1}$ is defined by

$$\tilde{q}_{i+1} < z_i \tilde{q}_i. \quad (2.8)$$

The factor of z_i enters because the angle at each branching is inversely proportional to the momentum fraction of the parent. Similarly for branching on the gluon, $k_i \rightarrow k_{i+1}$, we require

$$\tilde{k}_{i+1} < (1-z_i)\tilde{q}_i. \quad (2.9)$$

2.1.4 Branching probability

For the parton branching probability we use the mass-dependent splitting functions of ref. [7]. These are derived in the *quasi-collinear limit*, in which \mathbf{p}_\perp^2 and m^2 are treated as small (compared to $p \cdot n$) but \mathbf{p}_\perp^2/m^2 is not necessarily small. In this limit the $q \rightarrow qg$ splitting function is

$$P_{qq}(z, \mathbf{p}_\perp^2) = C_F \left[\frac{1+z^2}{1-z} - \frac{2z(1-z)m^2}{\mathbf{p}_\perp^2 + (1-z)^2 m^2} \right]. \quad (2.10)$$

Note that at $\mathbf{p}_\perp = 0$ the factor in square brackets is just $1-z$, i.e. the soft singularity at $z \rightarrow 1$ becomes a zero in the collinear direction. The minimum virtuality Q_g^2 serves only to define a resolvable emission, and therefore we omit it when defining the branching probability in terms of the evolution variable (2.6) as

$$dP(q \rightarrow qg) = \frac{\alpha_s}{2\pi} \frac{d\tilde{q}^2}{\tilde{q}^2} P_{qq} dz = \frac{C_F}{2\pi} \alpha_s [z^2(1-z)^2 \tilde{q}^2] \frac{d\tilde{q}^2}{\tilde{q}^2} \frac{dz}{1-z} \left[1 + z^2 - \frac{2m^2}{z\tilde{q}^2} \right]. \quad (2.11)$$

2.2 Gluon splitting

In the case of a final-state gluon splitting into a pair of heavy quarks of mass m , the quasi-collinear splitting function derived in [7] is

$$P_{gg}(z, \mathbf{p}_\perp^2) = T_R \left[1 - 2z(1-z) \frac{\mathbf{p}_\perp^2}{\mathbf{p}_\perp^2 + m^2} \right]. \quad (2.12)$$

We note that this splitting function is bounded above by its value $T_R = 1/2$ at the phase space boundary $\mathbf{p}_\perp = \mathbf{0}$, and below by $T_R/2$. By analogy with eq. (2.6), in this case the evolution variable \tilde{q} is related to the virtuality of the gluon or the relative transverse momentum of the splitting by

$$\tilde{q}^2 = \frac{q^2}{z(1-z)} = \frac{\mathbf{p}_\perp^2 + m^2}{z^2(1-z)^2}. \quad (2.13)$$

In terms of the variables \tilde{q}, z , the $g \rightarrow q\bar{q}$ branching probability then reads

$$dP(g \rightarrow q\bar{q}) = \frac{T_R}{2\pi} \alpha_s [z^2(1-z)^2 \tilde{q}^2] \frac{d\tilde{q}^2}{\tilde{q}^2} \left[1 - 2z(1-z) + \frac{2m^2}{z(1-z)\tilde{q}^2} \right] dz. \quad (2.14)$$

In the case of gluon splitting into gluons, the branching probability takes the familiar form

$$dP(g \rightarrow gg) = \frac{C_A}{2\pi} \alpha_s [z^2(1-z)^2 \tilde{q}^2] \frac{d\tilde{q}^2}{\tilde{q}^2} \left[\frac{z}{1-z} + \frac{1-z}{z} + z(1-z) \right] dz. \quad (2.15)$$

Since we introduce a minimum virtuality Q_g^2 for gluons, the relationship between the evolution variable and the relative transverse momentum for this splitting is as in eq. (2.13) but with the heavy quark mass m replaced by Q_g . Similarly, for gluon splitting to light quarks we use eq. (2.13) with $\mu = \max(m, Q_g)$ in place of m .

2.3 Initial-state branching

Consider the initial-state (spacelike) branching of a partonic constituent of an incoming hadron that undergoes some hard collisions subprocess such as deep inelastic lepton scattering. The momenta are defined as in eq. (2.1), with the reference vector p along the beam direction. In this case the evolution is performed *backwards* from the hard sub-process to the incoming hadron, as shown in figure 2. Thus we now define in place of eq. (2.3)




Figure 2: Initial-state parton branching. The blob represents the hard subprocess.

$$z_i = \frac{\alpha_{i-1}}{\alpha_i}, \quad \mathbf{p}_{\perp i} = \mathbf{q}_{\perp i-1} - z_i \mathbf{q}_{\perp i}. \quad (2.16)$$

Then

$$q_{i-1}^2 = z_i q_i^2 - \frac{z_i}{1 - z_i} k_i^2 - \frac{\mathbf{p}_{\perp i}^2}{1 - z_i}. \quad (2.17)$$

We assume a massless variable-flavour-number evolution scheme [22, 23] for constituent parton branching, setting $m = 0$ and putting all emitted gluons at the minimum virtuality, $k_i^2 = Q_g^2$. Thus in this case the momentum fraction z corresponds precisely to the lightcone fraction normally used in the definition of parton distribution functions. The angular evolution variable now relates only to the angle of the emitted gluon and therefore we choose

$$\tilde{q}_i^2 = \frac{\mathbf{p}_{\perp i}^2 + z_i Q_g^2}{(1 - z_i)^2}, \quad (2.18)$$

with ordering condition simply $\tilde{q}_{i+1} < \tilde{q}_i$. Correspondingly, for the argument of the running coupling we now use $(1 - z)^2 \tilde{q}^2$.

A different type of initial-state branching occurs in the decay of heavy, quasi-stable coloured objects like the top quark. Here the momentum of the incoming heavy object is fixed and evolution is performed forwards to the hard decay process. In this case we cannot neglect the mass of the parton and eq. (2.18) becomes

$$\tilde{q}_i^2 = \frac{\mathbf{p}_{\perp i}^2 + z_i Q_g^2}{(1 - z_i)^2} + m^2, \quad (2.19)$$

while the branching probability (2.11) is replaced by

$$dP(q \rightarrow qg) = \frac{C_F}{2\pi} \alpha_s [(1 - z)^2 \tilde{q}^2] \frac{d\tilde{q}^2}{\tilde{q}^2} \frac{dz}{1 - z} \left[1 + z^2 - \frac{2zm^2}{\tilde{q}^2} \right]. \quad (2.20)$$

2.4 Allowed regions and termination of branching

The allowed phase space for each branching is given by requiring a real relative transverse momentum, $\mathbf{p}_{\perp}^2 > 0$. In final-state $q \rightarrow qg$ branching, we have from eq. (2.7)

$$z^2(1 - z)^2 \tilde{q}^2 > (1 - z)^2 \mu^2 + z Q_g^2. \quad (2.21)$$

This yields a rather complicated boundary in the (\tilde{q}, z) plane. However, since

$$(1-z)^2\mu^2 + zQ_g^2 > (1-z)^2\mu^2, \quad z^2Q_g^2 \quad (2.22)$$

we see that the phase space lies inside the region

$$\frac{\mu}{\tilde{q}} < z < 1 - \frac{Q_g}{\tilde{q}}, \quad (2.23)$$

and approaches these limits for large values of \tilde{q} . The precise phase space can therefore be filled efficiently by generating values of z between these limits and rejecting those that violate the inequality (2.21). The resulting threshold for \tilde{q} is slightly larger than but of the order of $\mu + Q_g$ (see [12] for a detailed discussion).

In gluon splitting, we obtain the allowed phase space range from eq. (2.13) as

$$z_- < z < z_+, \quad z_{\pm} = \frac{1}{2} \left(1 \pm \sqrt{1 - \frac{4\mu}{\tilde{q}}} \right) \quad \text{and} \quad \tilde{q} > 4\mu \quad (2.24)$$

where $\mu = m$ for splitting into heavy quarks, or $\mu = \max(m, Q_g)$ more generally. Therefore, analogously to eq. (2.23), the phase space lies within the range

$$\frac{\mu}{\tilde{q}} < z < 1 - \frac{\mu}{\tilde{q}}. \quad (2.25)$$

Schematically, the parton shower corresponds to selecting a sequence of (\tilde{q}_i, z_i) values by solving the equations

$$\begin{aligned} R_1 &= \exp \left(- \int_{\tilde{q}_i}^{\tilde{q}_{i-1}} d\tilde{q} \int_{z_-}^{z_+} dz \frac{d^2 P}{d\tilde{q} dz} \right) \\ R_2 &= \frac{\int_{z_-}^{z_i} dz d^2 P / d\tilde{q} dz}{\int_{z_-}^{z_+} dz d^2 P / d\tilde{q} dz} \end{aligned} \quad (2.26)$$

where $R_{1,2} \in [0, 1]$ are uniform pseudorandom numbers. Whenever the algorithm selects a value of \tilde{q} below the relevant threshold, branching of that parton is terminated. The minimum virtuality Q_g thus determines the scale at which soft or collinear parton emission becomes unresolvable. In the absence of such a scale one eventually reaches a region where the perturbative expression for the running coupling is divergent.

After branching has terminated, the outgoing partons are put on mass-shell (or given the virtual mass Q_g if lighter) and the relative transverse momenta of the branchings in the shower are computed. For final-state gluon splitting we have

$$|\mathbf{p}_{\perp}| = \sqrt{z^2(1-z)^2\tilde{q}^2 - \mu^2}, \quad (2.27)$$

or else, if the parent is a quark,

$$|\mathbf{p}_{\perp}| = \sqrt{(1-z)^2(z^2\tilde{q}^2 - \mu^2) - zQ_g^2}. \quad (2.28)$$

The virtualities of the internal lines of the shower can now be computed backwards according to eq. (2.4). Finally, the azimuthal directions of the \mathbf{p}_{\perp} 's can be chosen [24] and the full 4-momenta reconstructed using eqs. (2.1) and (2.2).

In initial-state constituent parton branching the evolution is “guided” by the parton distribution functions (PDFs) of the incoming parent hadron. Since PDFs are often not tabulated below some scale $Q_s > Q_0$, one may wish to terminate branching whenever $\tilde{q} < Q_s$ is selected. In that case the incoming parton is assigned virtuality $q_n^2 \sim -Q_s^2$ and the spacelike virtualities of internal lines are then reconstructed back from q_n^2 to q_0^2 using the transverse momenta deduced from eq. (2.18) inserted in eq. (2.17).

For initial-state branching in the decay of a heavy, quasi-stable coloured object, the branching proceeds in the opposite direction but the reconstruction of momenta is similar, using eq. (2.19) instead of (2.18).

2.5 Treatment of colour flows

The more detailed treatment depend on the choice of the “backward” vector n and on which quantities are to be held fixed during jet evolution. Normally n should be taken along the colour-connected partner of the radiating parton, and the 4-momentum of the colour-connected system should be preserved. The upper limits on the evolution variable \tilde{q} for the colour-connected jets should be chosen so as to cover the phase space in the soft limit, with the best possible approximation to the correct angular distribution. In setting these limits we neglect the minimum virtuality Q_g^2 , which is a good approximation at high energies. We consider separately the four cases that the colour connection is between two final-state jets, two initial-state (beam) jets, a beam jet and a final-state jet, or a decaying heavy parton and a decay-product jet.

3. Final-final colour connection

Consider the process $a \rightarrow b + c$ where a is a colour singlet and b and c are colour-connected. Examples are $e^+e^- \rightarrow q\bar{q}$ and $W \rightarrow q\bar{q}'$. We need to preserve the 4-momentum of a and therefore we work in its rest-frame,

$$p_a = Q(1, \mathbf{0}, 0), \quad p_b = \frac{1}{2}Q(1 + b - c, \mathbf{0}, \lambda), \quad p_c = \frac{1}{2}Q(1 - b + c, \mathbf{0}, -\lambda), \quad (3.1)$$

where $p_a^2 = Q^2$, $b = m_b^2/Q^2$, $c = m_c^2/Q^2$ and

$$\lambda = \lambda(1, b, c) \equiv \sqrt{(1 + b - c)^2 - 4b} = \sqrt{(1 - b + c)^2 - 4c}. \quad (3.2)$$

For emission of a gluon g from b we write

$$q_i = \alpha_i p_b + \beta_i n + q_{\perp i} \quad (3.3)$$

where $\mathbf{q}_{\perp g} = \mathbf{k}_{\perp}$, $\mathbf{q}_{\perp b} = -\mathbf{k}_{\perp}$, $\mathbf{q}_{\perp c} = \mathbf{0}$ and we choose

$$n = \frac{1}{2}Q(\lambda, \mathbf{0}, -\lambda). \quad (3.4)$$

If c is massive, the alignment of n along the colour-connected momentum p_c is exact only in a certain class of Lorentz frames. As mentioned in section 2.1.1, if we try to use

a massive “backward” vector the kinematics become too complicated. However, we can always construct the lightlike vector (3.4) in a covariant way as

$$n = \frac{1}{2} [(1 + b - c + \lambda)p_c - (1 - b + c - \lambda)p_b], \quad (3.5)$$

which is close to p_c as long as b and c are small.

To preserve $p_a = q_b + q_c + q_g$ we require

$$\sum \alpha_i = \sum \beta_i = \frac{2}{1 + b - c + \lambda} \quad (3.6)$$

whereas the mass-shell conditions give

$$\begin{aligned} \beta_b &= \frac{2}{\lambda(1 + b - c + \lambda)} \left(\frac{b + \kappa}{\alpha_b} - b\alpha_b \right) \\ \beta_c &= \frac{2}{\lambda(1 + b - c + \lambda)} \left(\frac{c}{\alpha_c} - b\alpha_c \right) \\ \beta_g &= \frac{2}{\lambda(1 + b - c + \lambda)} \left(\frac{\kappa}{\alpha_g} - b\alpha_g \right) \end{aligned} \quad (3.7)$$

where $\kappa \equiv \mathbf{k}_\perp^2/Q^2$. Our new variables are

$$z = \frac{\alpha_b}{\alpha_b + \alpha_g}, \quad \tilde{\kappa} \equiv \frac{\tilde{q}^2}{Q^2} > b, \quad (3.8)$$

where from eq. (2.6) we have

$$\kappa = (z^2 \tilde{\kappa} - b)(1 - z)^2, \quad (3.9)$$

and so $\sqrt{b/\tilde{\kappa}} < z < 1$. From eqs. (3.6)-(3.9) we find

$$\begin{aligned} \alpha_b &= \frac{z}{1 + b - c + \lambda} \left(1 + b - c + z(1 - z)\tilde{\kappa} + \sqrt{[1 - b + c - z(1 - z)\tilde{\kappa}]^2 - 4b} \right), \\ \alpha_c &= \frac{2}{1 + b - c + \lambda} - \frac{\alpha_b}{z}, \\ \alpha_g &= \frac{1 - z}{z} \alpha_b, \end{aligned} \quad (3.10)$$

with the β_i 's given by eq. (3.7).

3.1 Phase space variables

It is convenient to express the phase space in terms of the Dalitz plot variables

$$x_i = \frac{2p_a \cdot q_i}{Q^2} = (1 + b - c)\alpha_i + \lambda\beta_i. \quad (3.11)$$

Substituting from eqs. (3.7) and (3.10), we find

$$\begin{aligned} x_c &= 1 - b + c - z(1 - z)\tilde{\kappa} \\ x_b &= (2 - x_c)r + (z - r)\sqrt{x_c^2 - 4c} \\ x_g &= (2 - x_c)(1 - r) - (z - r)\sqrt{x_c^2 - 4c} \end{aligned} \quad (3.12)$$

where

$$r = \frac{1}{2} \left(1 + \frac{b}{1+c-x_c} \right). \quad (3.13)$$

The jacobian factor is thus simply

$$\frac{\partial(x_b, x_c)}{\partial(z, \tilde{\kappa})} = z(1-z)\sqrt{x_c^2 - 4c} \quad (3.14)$$

and the quasi-collinear branching probability (2.11) translates to

$$dP(q \rightarrow qg) = C_F \frac{\alpha_s}{2\pi} \frac{dx_b dx_c}{(1-b+c-x_c)\sqrt{x_c^2 - 4c}} \left[\frac{1+z^2}{1-z} - \frac{2b}{1-b+c-x_c} \right] \quad (3.15)$$

where

$$z = r + \frac{x_b - (2-x_c)r}{\sqrt{x_c^2 - 4c}}, \quad (3.16)$$

r being the function of x_c given in eq. (3.13).

For emission from parton c we write

$$q_i = \alpha_i p_c + \beta_i n + q_{\perp i} \quad (3.17)$$

where now we choose

$$n = \frac{1}{2} Q(\lambda, \mathbf{0}, \lambda). \quad (3.18)$$

Clearly, the region covered and the branching probability will be as for emission from parton b , but with x_b and x_c , b and c interchanged.

3.2 Soft gluon region

For emission from parton b in the soft region $1-z = \epsilon \rightarrow 0$ we have

$$x_c \sim 1-b+c-\epsilon\tilde{\kappa}, \quad x_b \sim 1+b-c-\epsilon\tilde{\kappa}' \quad (3.19)$$

where

$$\tilde{\kappa}' = \lambda + \frac{\tilde{\kappa}}{2b}(1-b-c-\lambda). \quad (3.20)$$

Since $\tilde{\kappa}$ is an angular variable, we can express it in terms of the angle θ_{bg} between the directions of the emitting parton b and the emitted gluon in the rest frame of a . In the soft region we find

$$\tilde{\kappa} = \frac{(1+b-c+\lambda)(1+b-c-\lambda \cos \theta_{bg})}{2(1+\cos \theta_{bg})} \quad (3.21)$$

Thus $\tilde{\kappa} = b$ at $\theta_{bg} = 0$ and $\tilde{\kappa} \rightarrow \infty$ as $\theta_{bg} \rightarrow \pi$.

For soft emission from parton c , the roles of x_b and x_c , b and c are interchanged. To cover the whole angular region in the soft limit, we therefore require $\tilde{\kappa} < \tilde{\kappa}_b$ in jet b and $\tilde{\kappa} < \tilde{\kappa}_c$ in jet c , where

$$\frac{\tilde{\kappa}_b}{\tilde{\kappa}_b'} = \frac{\tilde{\kappa}_c'}{\tilde{\kappa}_c} \quad (3.22)$$

and hence

$$(\tilde{\kappa}_b - b)(\tilde{\kappa}_c - c) = \frac{1}{4}(1 - b - c + \lambda)^2. \quad (3.23)$$

In particular, the most symmetric choice is

$$\tilde{\kappa}_b = \frac{1}{2}(1 + b - c + \lambda), \quad \tilde{\kappa}_c = \frac{1}{2}(1 - b + c + \lambda). \quad (3.24)$$

The largest region that can be covered by one jet corresponds to the maximal value of $\tilde{\kappa}$ allowed in eq. (3.10) for real α_b , i.e. for the maximal b jet

$$\tilde{\kappa}_b = 4(1 - 2\sqrt{b} - b + c). \quad (3.25)$$

3.3 Example: $e^+e^- \rightarrow q\bar{q}g$

Here we have $b = c = \rho$, $\lambda = \sqrt{1 - 4\rho} = v$, the quark velocity in the Born process $e^+e^- \rightarrow q\bar{q}$. The phase space and the two jet regions for the symmetrical choice (3.24) are shown in figure 3. The region D, corresponding to hard non-collinear gluon emission, is not included in either jet and must be filled using the $\mathcal{O}(\alpha_s)$ matrix element (see below).

For the maximal quark jet we get from eq. (3.25)

$$\tilde{\kappa}_q = 4(1 - 2\sqrt{\rho}), \quad (3.26)$$

as shown in figure 4 together with the complementary antiquark jet region given by eq. (3.23).




Figure 3: Phase space for $e^+e^- \rightarrow q\bar{q}g$ for $m_q = 5 \text{ GeV}$, $Q^2 = m_Z^2$, with symmetric definition of quark and antiquark jets.




Figure 4: Phase space for $e^+e^- \rightarrow q\bar{q}g$ for $m_q = 5 \text{ GeV}$, $Q^2 = m_Z^2$, with maximal region for the quark jet.

3.3.1 Exact matrix element

The $e^+e^- \rightarrow V \rightarrow q\bar{q}g$ differential cross section, where V represents a vector current such as a virtual photon, is given to first order in α_s by [25, 26]

$$\frac{1}{\sigma_V} \frac{d^2\sigma_V}{dx_q dx_{\bar{q}}} = \frac{\alpha_s}{2\pi} \frac{C_F}{v} \left[\frac{(x_q + 2\rho)^2 + (x_{\bar{q}} + 2\rho)^2 + \zeta_V}{(1 + 2\rho)(1 - x_q)(1 - x_{\bar{q}})} - \frac{2\rho}{(1 - x_q)^2} - \frac{2\rho}{(1 - x_{\bar{q}})^2} \right] \quad (3.27)$$

where

$$\zeta_V = -8\rho(1 + 2\rho) \quad (3.28)$$

and

$$\sigma_V = \sigma_0(1 + 2\rho)v \quad (3.29)$$

is the Born cross section for heavy quark production by a vector current, σ_0 being the massless quark Born cross section.

In the case of the axial current contribution $e^+e^- \rightarrow A \rightarrow q\bar{q}g$, instead of eq. (3.27) we have

$$\frac{1}{\sigma_A} \frac{d^2\sigma_A}{dx_q dx_{\bar{q}}} = \frac{\alpha_s}{2\pi} \frac{C_F}{v} \left[\frac{(x_q + 2\rho)^2 + (x_{\bar{q}} + 2\rho)^2 + \zeta_A}{v^2(1 - x_q)(1 - x_{\bar{q}})} - \frac{2\rho}{(1 - x_q)^2} - \frac{2\rho}{(1 - x_{\bar{q}})^2} \right], \quad (3.30)$$

where

$$\zeta_A = 2\rho[(3 + x_g)^2 - 19 + 4\rho], \quad (3.31)$$

σ_A being the Born cross section for heavy quark production by the axial current:

$$\sigma_A = \sigma_0 v^3. \quad (3.32)$$

3.3.2 Soft gluon distribution

In the soft gluon region $1 - z = \epsilon \rightarrow 0$ the branching probability (3.15) becomes

$$\begin{aligned} \frac{d^2 P}{dx_q dx_{\bar{q}}} &\sim \frac{\alpha_s}{2\pi} \frac{2C_F}{v\epsilon^2} f_s(\tilde{\kappa}) \\ f_s(\tilde{\kappa}) &= \left(\frac{1}{\tilde{\kappa}} - \frac{\rho}{\tilde{\kappa}^2} \right) \Theta(\tilde{\kappa} - \rho). \end{aligned} \quad (3.33)$$

In this limit, the exact vector and axial current matrix elements, eqs. (3.27) and (3.30) respectively, give identical distributions:

$$\begin{aligned} \frac{1}{\sigma_V} \frac{d^2 \sigma_V}{dx_q dx_{\bar{q}}} &\sim \frac{1}{\sigma_A} \frac{d^2 \sigma_A}{dx_q dx_{\bar{q}}} \sim \frac{\alpha_s}{2\pi} \frac{2C_F}{v\epsilon^2} f(\tilde{\kappa}) \\ f(\tilde{\kappa}) &= \left(\frac{1-2\rho}{\tilde{\kappa}\tilde{\kappa}'} - \frac{\rho}{\tilde{\kappa}^2} - \frac{\rho}{\tilde{\kappa}'^2} \right) \Theta(\tilde{\kappa} - \rho) \\ &= f_s(\tilde{\kappa}) \left(\frac{v}{\tilde{\kappa}'} \right)^2. \end{aligned} \quad (3.34)$$

Since from eq. (3.20)

$$\tilde{\kappa}' = v + \tilde{\kappa} \left(\frac{1-v}{1+v} \right) > v, \quad (3.35)$$

the parton shower approximation (3.33) always overestimates the true result in the soft limit, and so correction by the rejection method is straightforward. For small values of ρ we have

$$f(\tilde{\kappa}) = \frac{1}{\tilde{\kappa}} - \frac{\rho}{\tilde{\kappa}^2} + \frac{2\rho^2}{\tilde{\kappa}} - 2\rho + \mathcal{O}(\rho^2). \quad (3.36)$$

Since $\tilde{\kappa} > \rho$ we see that the error in the approximation (3.33) is at most $\mathcal{O}(\rho)$, for any value of $\tilde{\kappa}$ (figure 5)




Figure 5: The function $f(\tilde{\kappa})$ giving the gluon angular distribution in the soft limit, for $m = 5 \text{ GeV}$, $Q^2 = m_Z^2$. The exact result eq. (3.34), solid curve, and shower approximation (3.33), dashed, are not distinguishable on this scale. The dotted curve shows the earlier HERWIG approximation.

The angular-ordered shower variables used previously in HERWIG [19] corresponded to a $1/\tilde{\kappa}$ behaviour at small $\tilde{\kappa}$, cut off sharply at $\tilde{\kappa} = 2\rho$, as indicated by the dotted curve in figure 5. We see that the new treatment greatly improves the description at small $\tilde{\kappa}$, near the collinear direction.

3.3.3 Dead region contribution

The integral over the dead region may be expressed as

$$\frac{1}{\sigma_V} \int_D d^2\sigma_V \equiv \frac{\alpha_s}{2\pi} C_F F_V^D(\tilde{\kappa}_q) \quad (3.37)$$

where $\tilde{\kappa}_q$ parametrizes the boundary of the quark jet. As shown in figure 6, this is actually maximal, but still small, at the symmetric point given by eq. (3.24).

Although the integral in eq. (3.37) is finite, the integrand diverges as one approaches the soft limit $x_q = x_{\bar{q}} = 1$ via the narrow “neck” of the dead region in figure 3 or 4. This could cause problems in generating $q\bar{q}g$ configurations in the dead region in order to apply a matrix element correction [27]. To avoid such problems, one can map the region $x_q, x_{\bar{q}} > 3/4$ into a region whose width vanishes quadratically as $x_q, x_{\bar{q}} \rightarrow 1$, as illustrated in figure 7. The mapping shown is

$$\begin{aligned} x_q &\rightarrow x'_q = 1 - \left[\frac{1}{4} - (1 - x_q) \right] = \frac{7}{4} - x_q, \\ x_{\bar{q}} &\rightarrow x'_{\bar{q}} = 1 - 2(1 - x'_q) \left[\frac{3}{4} - (1 - x_q) \right] = \frac{5}{8} + \frac{1}{2}x_q + \frac{3}{2}x_{\bar{q}} - 2x_q x_{\bar{q}} \end{aligned} \quad (3.38)$$

when $x_q > x_{\bar{q}} > 3/4$. Within the mapped region, the integrand then has an extra weight factor of $2(1 - x'_q)$ which regularizes the soft divergence. When $x_{\bar{q}} > x_q > 3/4$, x_q and $x_{\bar{q}}$ are interchanged in both the mapping and the weight.




Figure 6: The function $F^D(\tilde{\kappa}_q)$ giving the contribution of the dead region to the cross section, for $m = 5 \text{ GeV}$, $Q^2 = m_Z^2$. Solid: vector current. Dashed: axial current.




Figure 7: The soft region, with jet boundaries (solid) and mapped region (dashed), for $m = 5 \text{ GeV}$, $Q^2 = m_Z^2$.

4. Initial-initial colour connection

Here we consider the inverse process $b + c \rightarrow a$ where a is a colour singlet of invariant mass Q and b, c are beam jets. The kinematics are simple because we take beam jets to be massless: in the c.m. frame

$$p_a = Q(1, \mathbf{0}, 0), \quad p_b = \frac{1}{2}Q(1, \mathbf{0}, 1), \quad p_c = \frac{1}{2}Q(1, \mathbf{0}, -1). \quad (4.1)$$

For emission of a gluon g from b we write

$$q_i = \alpha_i p_b + \beta_i p_c + q_{\perp i} \quad (4.2)$$

where $\mathbf{q}_{\perp g} = \mathbf{k}_{\perp}$, $\mathbf{q}_{\perp a} = -\mathbf{k}_{\perp}$, $\mathbf{q}_{\perp b} = \mathbf{q}_{\perp c} = \mathbf{0}$. Notice that in this case the recoil transverse momentum is taken by the colour singlet a so we cannot preserve its 4-momentum. We choose to preserve its mass and rapidity, so that

$$\alpha_a = \beta_a = \sqrt{1 + \kappa}, \quad (4.3)$$

where as before $\kappa \equiv \mathbf{k}_{\perp}^2/Q^2$. Now we have

$$\begin{aligned} \beta_b = \alpha_c = 0, & \quad \alpha_g \beta_g = \kappa, \\ \alpha_a = \alpha_b - \alpha_g, & \quad \beta_a = \beta_c - \beta_g, \end{aligned} \quad (4.4)$$

and our new variables in this case are

$$z = 1 - \frac{\alpha_g}{\alpha_b}, \quad \tilde{\kappa} \equiv \frac{\tilde{q}^2}{Q^2} = \frac{\kappa}{(1-z)^2}. \quad (4.5)$$

Thus we find

$$\begin{aligned}\alpha_a &= \beta_a = \sqrt{1 + (1 - z)^2 \tilde{\kappa}}, \\ \alpha_b &= \frac{1}{z} \sqrt{1 + (1 - z)^2 \tilde{\kappa}}, \\ \beta_c &= \frac{1 + (1 - z) \tilde{\kappa}}{\sqrt{1 + (1 - z)^2 \tilde{\kappa}}}.\end{aligned}\tag{4.6}$$

4.1 Phase space variables

It is convenient to express the kinematics in terms of the “reduced” Mandelstam invariants:

$$\bar{s} = \frac{(q_b + q_c)^2}{Q^2}, \quad \bar{t} = \frac{(q_b - q_g)^2}{Q^2}, \quad \bar{u} = \frac{(q_c - q_g)^2}{Q^2}.\tag{4.7}$$

The phase space limits are

$$1 < \bar{s} < \frac{S}{Q^2}, \quad 1 - \bar{s} < \bar{t} < 0, \quad \bar{u} = 1 - \bar{s} - \bar{t}\tag{4.8}$$

where S is the beam-beam c.m. energy squared. In terms of the shower variables for beam jet b , we have

$$\bar{s} = \alpha_b \beta_c = \frac{1}{z} [1 + (1 - z) \tilde{\kappa}], \quad \bar{t} = -\alpha_b \beta_g = -(1 - z) \tilde{\kappa}, \quad \bar{u} = -(1 - z) \bar{s}.\tag{4.9}$$

Thus curves of constant $\tilde{\kappa}$ in the (\bar{s}, \bar{t}) plane are given by

$$\bar{t} = \frac{\tilde{\kappa}(1 - \bar{s})}{\tilde{\kappa} + \bar{s}}\tag{4.10}$$

and the jacobian factor for conversion of the shower variables to the Mandelstam invariants is

$$\frac{\partial(\bar{s}, \bar{t})}{\partial(z, \tilde{\kappa})} = \frac{1 - z}{z} \bar{s}.\tag{4.11}$$

For the other beam jet c we have $\bar{t} \leftrightarrow \bar{u}$ and thus

$$\bar{t} = \frac{\bar{s}(1 - \bar{s})}{\tilde{\kappa} + \bar{s}}.\tag{4.12}$$

We see that in order for the jet regions to touch without overlapping in the soft limit $\bar{s} \rightarrow 1$, $\bar{t} \rightarrow 0$, we need $\tilde{\kappa} < \tilde{\kappa}_b$ in jet b and $\tilde{\kappa} < \tilde{\kappa}_c$ in jet c , where $\tilde{\kappa}_c = 1/\tilde{\kappa}_b$. The most symmetrical choice is $\tilde{\kappa}_c = \tilde{\kappa}_b = 1$, as shown in figure 8, but we can take $\tilde{\kappa}_b$ or $\tilde{\kappa}_c$ as large as we like.

4.2 Example: Drell-Yan process

Consider radiation from the quark in the Drell-Yan process, $q\bar{q} \rightarrow gZ^0$. In the laboratory frame we have

$$q_q = (Px_1, \mathbf{0}, Px_1), \quad q_{\bar{q}} = (Px_2, \mathbf{0}, -Px_2)\tag{4.13}$$

where $P = \frac{1}{2}\sqrt{S}$ is the beam momentum. If we generated the initial hard process $q\bar{q} \rightarrow Z^0$ with momentum fractions $x_q, x_{\bar{q}}$ and we want to preserve the mass and rapidity of the Z^0 we require

$$x_1 = x_q \alpha_b, \quad x_2 = x_{\bar{q}} \beta_c\tag{4.14}$$




Figure 8: Beam jets (B,C) and dead region (D) in initial-state branching.

where α_b and β_c are given by eqs (4.6).

The branching probability in the parton shower approximation is

$$\frac{d^2 P}{dz d\tilde{\kappa}} = C_F \frac{\alpha_s}{2\pi} \frac{1}{\tilde{\kappa}} \frac{1+z^2}{1-z}, \quad (4.15)$$

which gives a differential cross section ($s = \bar{s}Q^2$, etc.)

$$\frac{1}{\sigma_0} \frac{d^2 \sigma}{ds dt} = \frac{D(x_1)D(x_2)}{D(x_q)D(x_{\bar{q}})} \frac{\alpha_s}{2\pi} C_F \frac{s+u}{s^3 tu} [s^2 + (s+u)^2] \quad (4.16)$$

where σ_0 is the Born cross section. The functions $D(x_1)$ etc. are parton distribution functions in the incoming hadrons; these factors take account of the change of kinematics $x_q, x_{\bar{q}} \rightarrow x_1, x_2$ discussed above.

The exact differential cross section for $q\bar{q} \rightarrow gZ^0$ to order α_s is

$$\frac{1}{\sigma_0} \frac{d^2 \sigma}{ds dt} = \frac{D(x_1)D(x_2)}{D(x_q)D(x_{\bar{q}})} \frac{\alpha_s}{2\pi} C_F \frac{Q^2}{s^3 tu} [(s+t)^2 + (s+u)^2]. \quad (4.17)$$

Since $Q^2 = s+t+u$ and $t \leq 0$, we see that the parton shower approximation (4.16) overestimates the exact expression, becoming exact in the collinear or soft limit $t \rightarrow 0$. Therefore the gluon distribution in the jet regions can be corrected efficiently by the rejection method, and the dead region can be filled using the matrix element, as was done in ref. [28].³ The benefit of the new variables is that the angular distribution of soft gluon emission requires no correction, provided the jet regions touch without overlapping in the soft region. As shown above, this will be the case if the upper limits on $\tilde{\kappa}$ satisfy $\tilde{\kappa}_{\bar{q}} = 1/\tilde{\kappa}_q$.

³An alternative approach, which is adopted in PYTHIA for this process [29], would be to increase the jet regions to cover the whole phase space, and to apply a larger corrections within the jets. In our case this would correspond to choosing $\tilde{\kappa}_b = \tilde{\kappa}_c = S/Q^2$.

5. Initial-final colour connection

Consider the process $a + b \rightarrow c$ where a is a colour singlet and the beam parton b and outgoing parton c are colour-connected. An example is deep inelastic scattering, where a is a (charged or neutral) virtual gauge boson. We need to preserve the 4-momentum of a and therefore we work in the Breit frame:

$$p_a = Q(0, \mathbf{0}, -1), \quad p_b = \frac{1}{2}Q(1+c, \mathbf{0}, 1+c), \quad p_c = \frac{1}{2}Q(1+c, \mathbf{0}, -1+c), \quad (5.1)$$

where $p_a^2 = -Q^2$, $p_b^2 = 0$, and $m_c^2 = cQ^2$. Notice that the beam parton b is always taken to be massless, but the outgoing parton c can be massive (e.g. in $W^+d \rightarrow c$).

5.1 Initial-state branching

For emission of a gluon g from the incoming parton b we write

$$q_i = \alpha_i p_b + \beta_i n + q_{\perp i} \quad (5.2)$$

where $\mathbf{q}_{\perp g} = \mathbf{k}_{\perp}$, $\mathbf{q}_{\perp b} = \mathbf{0}$, $\mathbf{q}_{\perp c} = -\mathbf{k}_{\perp}$ and we choose

$$n = \frac{1}{2}Q(1+c, \mathbf{0}, -1-c). \quad (5.3)$$

To preserve $p_a = q_c + q_g - q_b$ we now require

$$\alpha_b - \alpha_c - \alpha_g = \beta_c + \beta_g - \beta_b = \frac{1}{1+c} \quad (5.4)$$

whereas the mass-shell condition is

$$\alpha_i \beta_i Q^2 (1+c)^2 = \mathbf{q}_{\perp i}^2 + q_i^2 \quad (5.5)$$

which gives

$$1+c = \frac{c}{\alpha_c} + \kappa \left(\frac{1}{\alpha_c} + \frac{1}{\alpha_g} \right). \quad (5.6)$$

The new variables for emission from the beam jet are as in eq. (4.5). Substituting in (5.6), we find

$$\begin{aligned} \alpha_b &= \frac{1}{2z(1+c)} \left(1+c + (1-z)\tilde{\kappa} + \sqrt{[1+c + (1-z)\tilde{\kappa}]^2 - 4z(1-z)\tilde{\kappa}} \right), \\ \alpha_c &= z\alpha_b - \frac{1}{1+c}, \quad \alpha_g = (1-z)\alpha_b, \quad \beta_b = 0, \\ \beta_c &= \frac{1}{1+c} \cdot \frac{c + (1-z)^2\tilde{\kappa}}{z(1+c)\alpha_b - 1}, \quad \beta_g = \frac{(1-z)\tilde{\kappa}}{(1+c)^2\alpha_b}. \end{aligned} \quad (5.7)$$

5.2 Final-state branching

Next consider emission from the outgoing parton c . In this case we write

$$q_i = \alpha_i p_c + \beta_i p_b + q_{\perp i} \quad (5.8)$$

To preserve $p_a = q_c + q_g - q_b$ we require

$$\alpha_c + \alpha_g - \alpha_b = \beta_b - \beta_c - \beta_g = 1 \quad (5.9)$$

whereas the mass-shell condition is now

$$\alpha_i \beta_i (Q^2 + m_c^2) = \mathbf{q}_{\perp i}^2 + q_i^2 - \alpha_i^2 m_c^2. \quad (5.10)$$

The new variables for emission from an outgoing parton are as in eqs. (3.8), (3.9) with b replaced by c :

$$z = \frac{\alpha_c}{\alpha_c + \alpha_g}, \quad \tilde{\kappa} \equiv \frac{\tilde{q}^2}{Q^2} = \frac{1}{z^2} \left[c + \frac{\kappa}{(1-z)^2} \right]. \quad (5.11)$$

Thus in this case we find

$$\begin{aligned} \alpha_b &= 0, & \alpha_c &= z, & \alpha_g &= 1 - z, \\ \beta_b &= \frac{1}{1+c} [1 + c + z(1-z)\tilde{\kappa}], \\ \beta_c &= \frac{1-z}{1+c} [2c + z(1-z)\tilde{\kappa}], \\ \beta_g &= \frac{1-z}{1+c} [z^2\tilde{\kappa} - 2c]. \end{aligned} \quad (5.12)$$

5.3 Phase space variables

In this process the invariant phase space variables are usually taken to be

$$x_p = \frac{Q^2}{2p_a \cdot q_b}, \quad z_p = \frac{q_c \cdot q_b}{p_a \cdot q_b}. \quad (5.13)$$

In terms of the new variables for emission from the beam parton, we have

$$x_p = \frac{1}{(1+c)\alpha_b} = 2z \left(1 + c + (1-z)\tilde{\kappa} + \sqrt{[1 + c + (1-z)\tilde{\kappa}]^2 - 4z(1-z)\tilde{\kappa}} \right)^{-1} \quad (5.14)$$

$$z_p = (1+c)\beta_c = \frac{1}{2} \left(1 - c - (1-z)\tilde{\kappa} + \sqrt{[1 + c + (1-z)\tilde{\kappa}]^2 - 4z(1-z)\tilde{\kappa}} \right), \quad (5.15)$$

with the jacobian

$$\frac{\partial(x_p, z_p)}{\partial(z, \tilde{\kappa})} = \frac{1}{\tilde{\kappa}} \left(\frac{1}{x_p} + \frac{1+c}{1-z_p} - 2 \right)^{-1}. \quad (5.16)$$

In the soft limit $z = 1 - \epsilon$ we therefore find for the beam jet

$$x_p \sim \frac{1}{1+c} \left[1 - \epsilon - \frac{\epsilon c \tilde{\kappa}}{(1+c)^2} \right], \quad z_p \sim 1 - \frac{\epsilon \tilde{\kappa}}{1+c} \quad (5.17)$$




Figure 9: Beam jet (B), outgoing jet (C) and dead region (D) in initial-final state branching: $c = 0.25$, $\tilde{\kappa}_b = 1.25$, $\tilde{\kappa}_c = 1.5$.

and

$$\frac{\partial(x_p, z_p)}{\partial(z, \tilde{\kappa})} \sim \frac{\epsilon}{(1+c)^2}. \quad (5.18)$$

In terms of the variables for emission from the outgoing parton,

$$x_p = \frac{1}{(1+c)\beta_b} = \frac{1}{1+c+z(1-z)\tilde{\kappa}}, \quad z_p = \alpha_c = z, \quad (5.19)$$

so the jacobian is simply

$$\frac{\partial(x_p, z_p)}{\partial(z, \tilde{\kappa})} = z(1-z)x_p^2, \quad (5.20)$$

and in the soft limit

$$x_p \sim \frac{1}{1+c} \left[1 - \frac{\epsilon\tilde{\kappa}}{1+c} \right], \quad z_p \sim 1 - \epsilon, \quad (5.21)$$

with the jacobian again given by eq. (5.18). For full coverage of phase space in the soft limit we require $\tilde{\kappa} < \tilde{\kappa}_b$ in jet b and $\tilde{\kappa} < \tilde{\kappa}_c$ in jet c , where

$$\tilde{\kappa}_b(\tilde{\kappa}_c - c) = (1+c)^2. \quad (5.22)$$

Thus the most symmetrical choice is $\tilde{\kappa}_b = 1+c$, $\tilde{\kappa}_c = 1+2c$, as shown in figure 9. On the other hand, any larger or smaller combination satisfying eq. (5.22) is allowed, as illustrated in figure 10 for $\tilde{\kappa}_b = 10$.

5.4 Example: deep inelastic scattering

Consider deep inelastic scattering on a hadron of momentum P^μ by exchange of a virtual photon of momentum q^μ . By construction, the treatment outlined above preserves q^μ and hence also $Q^2 = -q^2$ and the Bjorken variable $x_B = Q^2/2P \cdot q$. If the contribution to the Born cross section from scattering on a quark of momentum fraction x_B is represented by σ_0 (a function of x_B and Q^2), then the correction due to single gluon emission is given by

$$\frac{1}{\sigma_0} \frac{d^2\sigma}{dx_p dz_p} = \frac{C_F\alpha_s}{2\pi} \frac{D(x_B/x_p)}{D(x_B)} \frac{1 + (x_p + z_p - 1)^2}{x_p(1-x_p)(1-z_p)}. \quad (5.23)$$




Figure 10: Beam jet (B), outgoing jet (C) and dead region (D) in initial-final state branching: $c = 0.25$, $\tilde{\kappa}_b = 10$, $\tilde{\kappa}_c = 0.40625$.

In the soft limit $x_p, z_p \rightarrow 1$ we have, from eqs. (5.17), (5.21) with $c = 0$,

$$(1 - x_p)(1 - z_p) \sim \epsilon^2 \tilde{\kappa} \quad (5.24)$$

and so

$$\frac{1}{\sigma_0} \frac{d^2 \sigma}{dx_p dz_p} \sim \frac{C_F \alpha_s}{\pi} \frac{1}{\epsilon^2 \tilde{\kappa}}, \quad (5.25)$$

whereas the parton shower approximation gives

$$\frac{1}{\sigma_0} \frac{d^2 \sigma}{dz d\tilde{\kappa}} \sim \frac{C_F \alpha_s}{\pi} \frac{1}{\epsilon \tilde{\kappa}}. \quad (5.26)$$

Since the jacobian factor (5.16) or (5.20) in this limit is simply ϵ , the shower approximation is exact in the soft limit.

5.5 Example: $q\bar{q} \rightarrow t\bar{t}$

We denote the momenta in this process by $p_a + p_b \rightarrow p_c + p_d$ and the $2 \rightarrow 2$ invariants by

$$\bar{s} = 2p_a \cdot p_b, \quad \bar{t} = -2p_a \cdot p_c, \quad \bar{u} = -2p_a \cdot p_d, \quad (5.27)$$

so that $\bar{s} + \bar{t} + \bar{u} = 0$. Colour flows from q to t and anticolour from \bar{q} to \bar{t} . Therefore the momentum transfer $q = p_a - p_c = p_d - p_b$ is carried by a colour singlet and we preserve this 4-momentum during showering.

For emission from the incoming light quark or the outgoing top quark, we work in the Breit frame for this system, where

$$q = Q(0, \mathbf{0}, 1), \quad p_a = \frac{1}{2}Q(1 + c, \mathbf{0}, 1 + c), \quad p_c = \frac{1}{2}Q(1 + c, \mathbf{0}, -1 + c) \quad (5.28)$$

with $Q^2 = -\bar{t} - m_t^2$ and $c = m_t^2/Q^2$. Then the treatment of sects. 5.1 and 5.2 can be applied directly, with the substitution $b \rightarrow a$ since the emitting system is now (a, c) rather

than (b, c) . However, the phase space variables are no longer those of section 5.3 since they involve the momenta of the \bar{q} and \bar{t} , which in the frame (5.28) take the general form

$$\begin{aligned} p_b &= \left[\frac{1}{2}Q\sqrt{(1+c)^2+4K}, \mathbf{Q}_\perp, -\frac{1}{2}Q(1+c) \right], \\ p_d &= \left[\frac{1}{2}Q\sqrt{(1+c)^2+4K}, \mathbf{Q}_\perp, \frac{1}{2}Q(1-c) \right], \end{aligned} \quad (5.29)$$

where $K = \mathbf{Q}_\perp^2/Q^2$ is related to the $2 \rightarrow 2$ invariants:

$$\bar{s} = \frac{1}{2}Q^2(1+c)^2 \left[1 + \sqrt{1 + \frac{4K}{(1+c)^2}} \right], \quad \bar{t} = -Q^2(1+c), \quad \bar{u} = -\bar{s} - \bar{t}, \quad (5.30)$$

and so

$$\mathbf{Q}_\perp^2 = -\bar{s} \left[1 + \frac{\bar{s}}{(1+c)\bar{t}} \right]. \quad (5.31)$$

For emission from the incoming light quark we define as in section 5.1

$$q_i = \alpha_i p_a + \beta_i n + q_{\perp i} \quad (5.32)$$

for $i = a, c, g$, where $\mathbf{q}_{\perp a} = \mathbf{0}$, $\mathbf{q}_{\perp c} = -\mathbf{k}_\perp$, $\mathbf{q}_{\perp g} = \mathbf{k}_\perp$, and n is as in eq. (5.3). Then the α_i 's and β_i 's are given by eqs. (5.7) with the substitution $b \rightarrow a$. The light antiquark and the antitop are not affected and therefore $q_b = p_b$, $q_d = p_d$. This allows the complete kinematics of the $2 \rightarrow 3$ process to be reconstructed. The $2 \rightarrow 3$ invariants can be defined as in ref. [14]:

$$s = 2q_a \cdot q_b, \quad t_1 = -2q_a \cdot q_c, \quad t_2 = -2q_b \cdot q_d, \quad u_1 = -2q_a \cdot q_d, \quad u_2 = -2q_b \cdot q_c. \quad (5.33)$$

It is convenient to express $n = p_c - cq$ so that (for $i = a, c, g$)

$$q_i = (\alpha_i - c\beta_i)p_a + (1+c)\beta_i p_c + q_{\perp i}. \quad (5.34)$$

Then we find

$$s = \alpha_a \bar{s}, \quad t_1 = \alpha_a \beta_c (1+c) \bar{t}, \quad t_2 = \bar{t}, \quad u_1 = \alpha_a \bar{u}, \quad u_2 = \beta_c (\bar{u} - c\bar{t}) - \alpha_c \bar{s} - 2\mathbf{k}_\perp \cdot \mathbf{Q}_\perp. \quad (5.35)$$

For emission from the outgoing top we use the results of section 5.2, again with the substitution $b \rightarrow a$. Thus we now have for $i = a, c, g$

$$q_i = \alpha_i p_c + \beta_i p_a + q_{\perp i} \quad (5.36)$$

where the α_i 's and β_i 's are given by eqs. (5.12) with $b \rightarrow a$, and we find that

$$s = \beta_a \bar{s}, \quad t_1 = \alpha_c \beta_a \bar{t}, \quad t_2 = \bar{t}, \quad u_1 = \beta_a \bar{u}, \quad u_2 = \alpha_c \bar{u} - \beta_c \bar{s} - 2\mathbf{k}_\perp \cdot \mathbf{Q}_\perp. \quad (5.37)$$

Similar formulae to eqs. (5.35) and (5.37), with the replacements $a \rightarrow b$ and $c \rightarrow d$, will hold for the case of gluon emission from the colour-connected $(\bar{q}\bar{t})$ system. Using these relations, one can study the distribution of gluon radiation in the parton shower

approximation and compare it with the exact $q\bar{q} \rightarrow t\bar{t}g$ matrix element. Agreement will be good in the soft and/or collinear regions but there will be regions of hard, wide-angle gluon emission in which matrix element corrections should be applied. Alternatively, the above equations can be used to formulate a modified subtraction scheme for combining fixed-order and parton-shower results, as was done in ref. [14] for a different parton-shower algorithm.

6. Decay colour connection

Consider the process $b \rightarrow ca$ where a is a colour singlet and the decaying parton b and outgoing parton c are colour-connected. Examples are bottom quark decay, $b \rightarrow cW^*$, and top decay, $t \rightarrow bW$. Here we have to preserve the 4-momentum of the decaying parton b and therefore we work in its rest frame,

$$p_b = m_b(1, \mathbf{0}, 0), \quad p_c = \frac{1}{2}m_b(1 - a + c, \mathbf{0}, \lambda), \quad p_a = \frac{1}{2}m_b(1 + a - c, \mathbf{0}, -\lambda), \quad (6.1)$$

where $a = m_a^2/m_b^2$, $c = m_c^2/m_b^2$ and now

$$\lambda = \lambda(1, a, c) = \sqrt{(1 + a - c)^2 - 4a} = \sqrt{(1 - a + c)^2 - 4c}. \quad (6.2)$$

6.1 Initial-state branching

For emission of a gluon g from the decaying parton b we write

$$q_i = \alpha_i p_b + \beta_i n + q_{\perp i} \quad (6.3)$$

where $\mathbf{q}_{\perp g} = \mathbf{k}_{\perp}$, $\mathbf{q}_{\perp c} = -\mathbf{k}_{\perp}$, $\mathbf{q}_{\perp b} = \mathbf{0}$ and we choose

$$n = \frac{1}{2}m_b(1, \mathbf{0}, 1), \quad (6.4)$$

i.e. aligned along p_c in the rest frame of b . The mass-shell conditions give

$$\beta_a = \frac{a}{\alpha_a} - \alpha_a, \quad \beta_c = \frac{c + \kappa}{\alpha_c} - \alpha_c, \quad \beta_g = \frac{\kappa}{\alpha_g} - \alpha_g, \quad (6.5)$$

with $\kappa = \mathbf{k}_{\perp}^2/m_b^2$. From momentum conservation

$$\alpha_a + \alpha_c + \alpha_g = \frac{a}{\alpha_a} + \frac{c + \kappa}{\alpha_c} + \frac{\kappa}{\alpha_g} = 1. \quad (6.6)$$

Recall that in initial-state branching of a heavy object our new evolution variable is given by eq. (2.19), so we have

$$\alpha_g = 1 - z, \quad \kappa = (\tilde{\kappa} - 1)(1 - z)^2 \quad (6.7)$$

where $\tilde{\kappa} = \tilde{q}^2/m_b^2 > 1$. Introducing for brevity the notation

$$w = 1 - (1 - z)(\tilde{\kappa} - 1), \quad u = 1 + a - c - (1 - z)\tilde{\kappa}, \quad v = \sqrt{u^2 - 4awz}, \quad (6.8)$$

from eq. (6.6) we find

$$\alpha_a = \frac{u + v}{2w}, \quad \alpha_c = 1 - \alpha_a - \alpha_g = z - \alpha_a. \quad (6.9)$$

6.2 Final-state branching

For radiation from the outgoing parton c we write

$$q_i = \alpha_i p_c + \beta_i n + q_{\perp i} \quad (6.10)$$

where p_c is given by eq. (6.1). Since the colour-connected parton b is at rest in our working frame of reference, the choice of the light-like vector n in this case is somewhat arbitrary. By analogy with the cases treated earlier, we choose it to be opposite to that used for the radiation from b , i.e. along the direction of the colour singlet a :

$$n = \frac{1}{2} m_b (\lambda, \mathbf{0}, -\lambda). \quad (6.11)$$

The kinematics are then identical with those for final-final connection (section 3), with the replacement $b \rightarrow c$, $c \rightarrow a$.

6.3 Phase space variables

As in section 3, it is convenient to use the Dalitz plot variables, which in this case are

$$x_i = \frac{2q_i \cdot p_b}{m_b^2}. \quad (6.12)$$

For emission from the decaying parton b we have $x_i = 2\alpha_i + \beta_i$ and hence, from eq. (6.9),

$$x_a = \frac{u+v}{2w} + \frac{u-v}{2z}, \quad x_c = w + z - x_a, \quad x_g = 2 - w - z = (1-z)\tilde{\kappa}, \quad (6.13)$$

with the jacobian factor

$$\frac{\partial(x_a, x_g)}{\partial(z, \tilde{\kappa})} = (1-z) \left[\frac{u+v}{2w^2} - \frac{u-v}{2z^2} + \frac{a(w-z)^2}{vwz} \right]. \quad (6.14)$$

In the soft limit $z \rightarrow 1 - \epsilon$ we find

$$x_a \sim 1 + a - c - \epsilon \tilde{\kappa}'_b, \quad x_g \sim \epsilon \tilde{\kappa}_b \quad (6.15)$$

where

$$\tilde{\kappa}'_b = \lambda + \frac{\tilde{\kappa}_b}{2} (1 - a + c - \lambda). \quad (6.16)$$

For emission from the outgoing parton c we have, from eq. (3.12) with the replacement $b \rightarrow c$, $c \rightarrow a$:

$$\begin{aligned} x_a &= 1 + a - c - z(1-z)\tilde{\kappa} \\ x_c &= (2 - x_a)r + (z - r)\sqrt{x_a^2 - 4a} \\ x_g &= (2 - x_a)(1 - r) - (z - r)\sqrt{x_a^2 - 4a} \end{aligned} \quad (6.17)$$

where

$$r = \frac{1}{2} \left(1 + \frac{c}{1 + a - x_a} \right). \quad (6.18)$$

In the soft limit we have from eq. (3.20)

$$x_a \sim 1 + a - c - \epsilon \tilde{\kappa}_c, \quad x_g \sim \epsilon \tilde{\kappa}_c' \quad (6.19)$$

where

$$\tilde{\kappa}_c' = \lambda + \frac{\tilde{\kappa}_c}{2c} (1 - a + c - \lambda). \quad (6.20)$$

For full coverage of the soft region we require

$$\frac{\tilde{\kappa}_b}{\tilde{\kappa}_b'} = \frac{\tilde{\kappa}_c'}{\tilde{\kappa}_c} \quad (6.21)$$

which gives in this case

$$(\tilde{\kappa}_b - 1)(\tilde{\kappa}_c - c) = \frac{1}{4}(1 - a + c + \lambda)^2. \quad (6.22)$$

Note that, while there is no upper limit on $\tilde{\kappa}_b$, the largest value that can be chosen for $\tilde{\kappa}_c$ is given by the equivalent of eq. (3.25),

$$\tilde{\kappa}_c < 4(1 + a - 2\sqrt{c} - c). \quad (6.23)$$

6.4 Example: top decay

In the decay $t \rightarrow Wbg$ we have $a = m_W^2/m_t^2 = 0.213$ and $c = m_b^2/m_t^2 = 0.026$, so for simplicity we neglect c . We also neglect the top width Γ_t here, although it can be taken into account through a multi-scale branching procedure [11]. Then for radiation from the top we have from eq. (6.13)

$$x_W = \frac{u+v}{2w} + \frac{u-v}{2z}, \quad x_g = (1-z)\tilde{\kappa}, \quad (6.24)$$

where u, v, w are given by eqs. (6.8) with $c \rightarrow 0$. The phase space is the region

$$0 < x_g < 1 - a, \quad 1 - x_g + \frac{a}{1 - x_g} < x_W < 1 + a. \quad (6.25)$$

Notice that for real x_W we require $u^2 > 4awz$, i.e.

$$1 < \tilde{\kappa} < 1 + a \left[1 - \sqrt{\frac{z(1-a)}{a(1-z)}} \right]^2, \quad (6.26)$$

and

$$1 - \frac{1-a}{\tilde{\kappa} + 2\sqrt{a(\tilde{\kappa}-1)}} < z < 1. \quad (6.27)$$

Thus there is no upper limit on $\tilde{\kappa}$, but the range of z becomes more limited as $\tilde{\kappa}$ increases.

For radiation from the b we have from eq. (6.17)

$$\begin{aligned} x_W &= 1 + a - z(1-z)\tilde{\kappa} \\ x_g &= \frac{1}{2}(2 - x_W) - \left(z - \frac{1}{2}\right) \sqrt{x_W^2 - 4a}. \end{aligned} \quad (6.28)$$




Figure 11: Phase space for decay $t \rightarrow Wbg$, with symmetric choice of emission regions for the b (B) and the t (T1,T2), and the dead region (D).

To cover the soft region we require $\tilde{\kappa} < \tilde{\kappa}_t$ for emission from the top quark and $\tilde{\kappa} < \tilde{\kappa}_b$ for that from the bottom, where eq. (6.22) gives

$$\tilde{\kappa}_b = \frac{(1-a)^2}{\tilde{\kappa}_t - 1}. \quad (6.29)$$

The most symmetrical choice would therefore appear to be $\tilde{\kappa}_b = \tilde{\kappa}_t - 1 = 1 - a = 0.787$, as illustrated in figure 11.

As mentioned above, there is no upper limit on $\tilde{\kappa}_t$. Thus the region covered by gluon emission from the top quark can be as large as we like. However, eq. (6.23) tells us that the upper limit for radiation from the b is

$$\tilde{\kappa}_b < 4(1 - \sqrt{a})^2 = 1.16, \quad (6.30)$$

and correspondingly $\tilde{\kappa}_t > 1 + \frac{1}{4}(1 + \sqrt{a})^2 = 1.53$. Figure 12 shows this maximal region that can be covered by emission from the b , together with the complementary regions of emission from the t .

We note from figures 11 and 12 that, for any value of $\tilde{\kappa}_t$, the region for emission from the top quark consists of two distinct parts that touch at the point $x_g = 1 - \sqrt{a}$, $x_W = 2\sqrt{a}$, where the W boson is at rest: a subregion T1 which includes the soft limit $x_g \rightarrow 0$ and a hard gluon region T2. An important difference from the previous treatment of top decay in HERWIG is that both these regions were absent in the latter, since the decay was performed in the top rest frame and the non-covariant formalism used there did not allow any emission from an object initially at rest [19]. As a consequence, a matrix element correction with an infrared cutoff had to be used to fill these regions [30]. In the new covariant formalism, emission from the top quark is generated consistently in any frame, and the infrared region is fully covered by parton showers from the top and bottom.




Figure 12: Phase space for decay $t \rightarrow Wbg$, with maximal region (B) for emission from the b , together with complementary regions of emission from the t (T1,T2) and the dead region (D).

The exact $t \rightarrow Wbg$ differential decay rate to first order in α_s is given in [30]:

$$\frac{1}{\Gamma_0} \frac{d^2\Gamma}{dx_W dx_g} = \frac{\alpha_s}{\pi} \frac{C_F}{(1+a-x_W)x_g^2} \left\{ x_g - \frac{(1+a-x_W)(1-x_g) + x_g^2}{1-a} + \right. \quad (6.31)$$

$$\left. + x_g \frac{(x_W + x_g - 1 - a)^2}{2(1-a)^2} + \frac{2a(1+a-x_W)x_g^2}{(1-a)^2(1+2a)} \right\}$$

where Γ_0 is the lowest-order decay rate. In the soft region $x_W \rightarrow 1+a$, $x_g \rightarrow 0$ this becomes

$$\frac{1}{\Gamma_0} \frac{d^2\Gamma}{dx_W dx_g} \sim \frac{\alpha_s}{\pi} \frac{C_F}{x_g} \left[\frac{1}{1+a-x_W} - \frac{1}{(1-a)x_g} \right]. \quad (6.32)$$

For soft gluon emission ($1-z=\epsilon \rightarrow 0$) from the top quark we have from eqs. (6.15), (6.16)

$$x_W \sim 1+a-\epsilon(1-a), \quad x_g \sim \epsilon\tilde{\kappa} \quad (6.33)$$

and so the exact form of the soft gluon distribution is

$$\frac{1}{\Gamma_0} \frac{d^2\Gamma}{dx_g dx_W} \sim \frac{\alpha_s}{\pi} \frac{C_F}{\epsilon^2} f_t(\tilde{\kappa}) \quad (6.34)$$

where

$$f_t(\tilde{\kappa}) = \frac{\tilde{\kappa} - 1}{(1-a)\tilde{\kappa}^2}. \quad (6.35)$$

In the same region the parton shower approximation (2.11) gives

$$\frac{d^2P}{dx_g dx_W} \sim \frac{1}{(1-a)\epsilon} \frac{d^2P}{dz d\tilde{\kappa}} \sim \frac{\alpha_s}{\pi} \frac{C_F}{(1-a)\epsilon^2 \tilde{\kappa}} \left(1 - \frac{1}{\tilde{\kappa}} \right) = \frac{\alpha_s}{\pi} \frac{C_F}{\epsilon^2} f_t(\tilde{\kappa}). \quad (6.36)$$

Thus we see that, for emission from the top quark, the shower approximation is exact in the soft limit. At higher gluon energies, inside the region T1 the parton shower overestimates the exact matrix element and can therefore be corrected easily by the rejection method. In the hard gluon region T2, which contributes only a small finite correction to the cross section, the parton shower overestimates the matrix element at lower values of x_g but underestimates it at the highest values. Therefore a combination of rejection and matrix element correction is needed in this region.

For emission from the bottom quark in the soft limit, we use the results of section 3.2 with the substitution $b \rightarrow 0$, $c \rightarrow a$ to obtain

$$x_W \sim 1 + a - \epsilon \tilde{\kappa}, \quad x_g \sim \epsilon \left(1 - a + \frac{\tilde{\kappa}}{1 - a} \right). \quad (6.37)$$

Therefore the exact soft gluon distribution in the b jet should be

$$\frac{1}{\Gamma_0} \frac{d^2 \Gamma}{dx_g dx_W} \sim \frac{\alpha_s}{\pi} \frac{C_F}{\epsilon^2} f_b(\tilde{\kappa}) \quad (6.38)$$

where

$$f_b(\tilde{\kappa}) = \frac{1}{(1 - a)\tilde{\kappa}} \left[1 + \frac{\tilde{\kappa}}{(1 - a)^2} \right]^{-2}. \quad (6.39)$$

On the other hand the parton shower approximation in this case gives simply

$$\frac{d^2 P}{dx_g dx_W} \sim \frac{1}{(1 - a)\epsilon} \frac{d^2 P}{dz d\tilde{\kappa}} \sim \frac{\alpha_s}{\pi} \frac{C_F}{(1 - a)\epsilon^2 \tilde{\kappa}}. \quad (6.40)$$

Thus the soft gluon distribution in the b jet region is overestimated by a factor of

$$\left[1 + \frac{\tilde{\kappa}}{(1 - a)^2} \right]^2, \quad (6.41)$$

which can be corrected by the rejection method. This factor varies from 1 to 5.2 for the symmetric choice of the b jet region $\tilde{\kappa}_b = 1 - a$ depicted in figure 11. For the maximal b jet shown in figure 12, it rises to 8.3. Since the shower approximation is exact in the soft limit for emission from the top, one can reduce the amount of soft correction required by decreasing the b jet region and increasing that for top emission, in accordance with eq. (6.29). However, for large values of $\tilde{\kappa}_t$ the dead region moves near to the collinear singularity at $x_W = 1 + a$ and a large hard matrix element correction becomes necessary.

7. Conclusions

We have presented a new formulation of the parton-shower approximation to QCD matrix elements, which offers a number of advantages over previous ones. Direct angular ordering of the shower ensures a good emulation of important QCD coherence effects, while the connection between the shower variables and the Sudakov-like representation of momenta (2.1) simplifies the kinematics and their relation to phase space invariants. The use of mass-dependent splitting functions with the new variables allows an accurate description

of soft gluon emission from heavy quarks over a wide angular region, including the collinear direction. The separation of showering into contributions from pairs of colour-connected hard partons permits a general treatment of coherence effects, which should be reliable at least to leading order in the number of colours. Since the formulation is slightly different for initial- and final-state showering, we have provided formulae for all colour-connected combinations of incoming and outgoing partons.

As mentioned in the Introduction, this new shower formulation is a key element of the event generator **Herwig++** [11]; detailed results of its implementation for e^+e^- annihilation are presented in ref. [12].

Acknowledgments

We are most grateful to the other **Herwig++** authors, Mike Seymour and Alberto Ribon, for helpful discussions. S.G. and B.W. also acknowledge fruitful conversations with Frank Krauss and Stefano Catani. S.G. and P.S. thank CERN Theory Division for hospitality during part of this work. This research was supported by the UK Particle Physics and Astronomy Research Council.

References

- [1] R.K. Ellis, W.J. Stirling and B.R. Webber, *QCD and collider physics*, *Cambridge Monogr. Part. Phys. Nucl. Phys. Cosmol.* **8** (1996) 1.
- [2] G. Marchesini and B.R. Webber, *Monte Carlo simulation of general hard processes with coherent QCD radiation*, *Nucl. Phys.* **B 310** (1988) 461.
- [3] G. Corcella et al., *HERWIG 6: an event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)*, *J. High Energy Phys.* **01** (2001) 010 [[hep-ph/0011363](#)].
- [4] T. Sjostrand, P. Eden, C. Friberg, L. Lonnblad, G. Miu, S. Mrenna and E. Norrbin, *High-energy-physics event generation with PYTHIA 6.1*, *Comput. Phys. Commun.* **135** (2001) 238 [[hep-ph/0010017](#)].
- [5] S. Catani, B.R. Webber and G. Marchesini, *QCD coherent branching and semiinclusive processes at large x* , *Nucl. Phys.* **B 349** (1991) 635.
- [6] S. Catani, Y.L. Dokshitzer, M. Olsson, G. Turnock and B.R. Webber, *New clustering algorithm for multi-jet cross-sections in e^+e^- annihilation*, *Phys. Lett.* **B 269** (1991) 432.
- [7] S. Catani, S. Dittmaier and Z. Trócsányi, *One-loop singular behaviour of QCD and SUSY QCD amplitudes with massive partons*, *Phys. Lett.* **B 500** (2001) 149 [[hep-ph/0011222](#)].
- [8] S. Catani, S. Dittmaier, M.H. Seymour and Z. Trócsányi, *The dipole formalism for next-to-leading order QCD calculations with massive partons*, *Nucl. Phys.* **B 627** (2002) 189 [[hep-ph/0201036](#)].
- [9] M. Cacciari and S. Catani, *Soft-gluon resummation for the fragmentation of light and heavy quarks at large x* , *Nucl. Phys.* **B 617** (2001) 253 [[hep-ph/0107138](#)].
- [10] E. Norrbin and T. Sjostrand, *QCD radiation off heavy particles*, *Nucl. Phys.* **B 603** (2001) 297 [[hep-ph/0010012](#)].

- [11] S. Gieseke, A. Ribon, M.H. Seymour, P. Stephens and B.R. Webber, *HERWIG++ 1.0, physics and manual*, preprint Cavendish-HEP-03/20, in preparation.
- [12] S. Gieseke, A. Ribon, M.H. Seymour, P. Stephens and B. Webber, *HERWIG++ 1.0: an event generator for e^+e^- annihilation*, hep-ph/0311208.
- [13] S. Frixione and B.R. Webber, *Matching NLO QCD computations and parton shower simulations*, *J. High Energy Phys.* **06** (2002) 029 [hep-ph/0204244].
- [14] S. Frixione, P. Nason and B.R. Webber, *Matching NLO QCD and parton showers in heavy flavour production*, *J. High Energy Phys.* **08** (2003) 007 [hep-ph/0305252].
- [15] S. Frixione and B.R. Webber, *The MC@NLO 2.2 event generator*, hep-ph/0309186.
- [16] S. Catani, F. Krauss, R. Kuhn and B.R. Webber, *QCD matrix elements + parton showers*, *J. High Energy Phys.* **11** (2001) 063 [hep-ph/0109231].
- [17] Y.L. Dokshitzer, G. Marchesini and B.R. Webber, *Dispersive approach to power-behaved contributions in QCD hard processes*, *Nucl. Phys.* **B 469** (1996) 93 [hep-ph/9512336].
- [18] D. Amati, A. Bassetto, M. Ciafaloni, G. Marchesini and G. Veneziano, *A treatment of hard processes sensitive to the infrared structure of QCD*, *Nucl. Phys.* **B 173** (1980) 429.
- [19] G. Marchesini and B.R. Webber, *Simulation of QCD coherence in heavy quark production and decay*, *Nucl. Phys.* **B 330** (1990) 261.
- [20] Y.L. Dokshitzer, V.A. Khoze and S.I. Troian, *On specific QCD properties of heavy quark fragmentation ('dead cone')*, *J. Phys.* **G 17** (1991) 1602.
- [21] G. Marchesini and B.R. Webber, *Simulation of QCD jets including soft gluon interference*, *Nucl. Phys.* **B 238** (1984) 1.
- [22] M.A.G. Aivazis, J.C. Collins, F.I. Olness and W.-K. Tung, *Leptoproduction of heavy quarks, 2. A unified QCD formulation of charged and neutral current processes from fixed target to collider energies*, *Phys. Rev.* **D 50** (1994) 3102 [hep-ph/9312319].
- [23] R.S. Thorne and R.G. Roberts, *An ordered analysis of heavy flavour production in deep inelastic scattering*, *Phys. Rev.* **D 57** (1998) 6871 [hep-ph/9709442].
- [24] P. Richardson, *Spin correlations in Monte Carlo simulations*, *J. High Energy Phys.* **11** (2001) 029 [hep-ph/0110108].
- [25] P. Nason and B.R. Webber, *Scaling violation in e^+e^- fragmentation functions: QCD evolution, hadronization and heavy quark mass effects*, *Nucl. Phys.* **B 421** (1994) 473, erratum *ibid.* **B 480** (1994) 755.
- [26] P. Nason and B.R. Webber, *Non-perturbative corrections to heavy quark fragmentation in e^+e^- annihilation*, *Phys. Lett.* **B 395** (1997) 355 [hep-ph/9612353].
- [27] M.H. Seymour, *A simple prescription for first order corrections to quark scattering and annihilation processes*, *Nucl. Phys.* **B 436** (1995) 443 [hep-ph/9410244].
- [28] G. Corcella and M.H. Seymour, *Initial state radiation in simulations of vector boson production at hadron colliders*, *Nucl. Phys.* **B 565** (2000) 227 [hep-ph/9908388].
- [29] G. Miu and T. Sjostrand, *W production in an improved parton shower approach*, *Phys. Lett.* **B 449** (1999) 313 [hep-ph/9812455].
- [30] G. Corcella and M.H. Seymour, *Matrix element corrections to parton shower simulations of heavy quark decay*, *Phys. Lett.* **B 442** (1998) 417 [hep-ph/9809451].