

Four-loop decoupling relations for the strong coupling

To cite this article: York Schröder and Matthias Steinhauser JHEP01(2006)051

View the [article online](#) for updates and enhancements.

You may also like

- [Four-loop pressure of massless \$O\(N\)\$ scalar field theory](#)
Antti Gynther, Mikko Laine, York Schröder et al.
- [Methods and progress in studying inelastic interactions between positrons and atoms](#)
R D DuBois
- [High-precision epsilon expansions of single-mass-scale four-loop vacuum bubbles](#)
York Schröder and Aleksi Vuorinen

Four-loop decoupling relations for the strong coupling

York Schröder

*Fakultät für Physik, Universität Bielefeld
33501 Bielefeld, Germany
E-mail: yorks@physik.uni-bielefeld.de*

Matthias Steinhauser

*Institut für Theoretische Teilchenphysik, Universität Karlsruhe
76128 Karlsruhe, Germany
E-mail: matthias.steinhauser@uka.de*

ABSTRACT: We compute the matching relation for the strong coupling constant within the framework of QCD up to four-loop order. This allows a consistent five-loop running (once the β function is available to this order) taking into account threshold effects. As a side product we obtain the effective coupling of a Higgs boson to gluons with five-loop accuracy.

KEYWORDS: QCD, NLO Computations.

Contents

1. Introduction	1
2. Theoretical framework	2
3. Running and decoupling for α_s	4
4. Effective coupling between a Higgs boson and gluons	6
5. Conclusions	9
A. Results for ζ_g^{OS}	9

1. Introduction

The strong coupling constant, α_s , constitutes a fundamental parameter in the Standard Model and thus its precise numerical value is very important for many physical predictions. An interesting property of α_s is its scale dependence, in particular its strong rise for low and its small value for high energies which make perturbative calculations within the framework of QCD possible. The scale dependence is governed by the β function. However, in order to relate α_s at two different scales it is also necessary to incorporate threshold effects of heavy quarks which is achieved with the help of the so-called matching or decoupling relations. Thus, when specifying α_s it is necessary to indicate next to the scale also the number of active flavours. In this paper we evaluate the decoupling relations to four-loop accuracy. This makes it possible to perform a consistent running of the strong coupling evaluated at a low scale, like, e.g., the mass of the τ lepton, to a high scale like the Z boson mass — once the five-loop β function is available.

Many different techniques have been developed and applied to various classes of Feynman diagrams. The complexity increases both with the number of legs and the number of loops. As far as the application of multi-loop diagrams to physical processes is concerned the current limit are four-loop single-scale Feynman diagrams, where either all internal particles are massless and one external momentum flows through the diagram (see, e.g., ref. [1] for a recent publication), or all external momenta are zero and besides massless lines there are also particles with a common mass M . The latter case has been developed in refs. [2, 3] and first applications can be found in refs. [4, 5]. In this paper we consider a further very important application: the four-loop contribution to the matching or decoupling relation for the strong coupling.

The paper is organized as follows: In the next section we define the decoupling constants and the theoretical framework of our calculation. In section 3 we present analytical

results and discuss the numerical consequences. In section 4 the connection of the decoupling constant to the coupling of a Higgs boson to two gluons is explained and the corresponding coupling strength is evaluated to five-loop order. Finally, we conclude in section 5. In the appendix we present the result for the decoupling constant parameterized in terms of the on-shell heavy quark mass.

2. Theoretical framework

We consider QCD with n_f active quark flavours. Furthermore it is assumed that n_l quarks are massless and n_h quarks are massive, i.e. we have $n_f = n_l + n_h$. In practice one often has $n_h = 1$, however, it is convenient to keep a generic label for the massive quarks.

The decoupling relations relate quantities in the full and effective theory where the latter is defined through the lagrangian \mathcal{L}' given by

$$\mathcal{L}'(g_s^0, m_q^0, \xi^0; \psi_q^0, G_\mu^{0,a}, c^{0,a}; \zeta_i^0) = \mathcal{L}^{\text{QCD}}(g_s^{0'}, m_q^{0'}, \xi^{0'}; \psi_q^{0'}, G_\mu^{0',a}, c^{0',a}) . \quad (2.1)$$

ψ_q , G_μ^a and c^a are the fermion, gluon and ghost fields, respectively, m_q are the quark masses, ξ is the gauge parameter, and $\alpha_s = g_s^2/(4\pi)$ is the strong coupling constant. \mathcal{L}^{QCD} is the usual QCD Lagrange density and the effective n_l -flavour quantities are marked by a prime. Eq. (2.1) states that the lagrangian in the effective theory has the same form as the original one with rescaled fields, masses and coupling. It is convenient to define the decoupling constants ζ_i in the bare theory through

$$\begin{aligned} g_s^{0'} &= \zeta_g^0 g_s^0, & m_q^{0'} &= \zeta_m^0 m_q^0, & \xi^{0'} - 1 &= \zeta_3^0(\xi^0 - 1), \\ \psi_q^{0'} &= \sqrt{\zeta_2^0} \psi_q^0, & G_\mu^{0',a} &= \sqrt{\zeta_3^0} G_\mu^{0,a}, & c^{0',a} &= \sqrt{\tilde{\zeta}_3^0} c^{0,a}. \end{aligned} \quad (2.2)$$

In a next step the renormalized quantities are obtained by the usual renormalization procedure introduced by the multiplicative renormalization constants through [6]

$$\begin{aligned} g_s^0 &= \mu^\varepsilon Z_g g_s, & m_q^0 &= Z_m m_q, & \xi^0 - 1 &= Z_3(\xi - 1), \\ \psi_q^0 &= \sqrt{Z_2} \psi_q, & G_\mu^{0,a} &= \sqrt{Z_3} G_\mu^a, & c^{0,a} &= \sqrt{\tilde{Z}_3} c^a. \end{aligned} \quad (2.3)$$

Combining eqs. (2.2) and (2.3) leads to renormalized decoupling constants, e.g.

$$\zeta_g = \frac{Z_g}{Z_g'} \zeta_g^0, \quad \zeta_3 = \frac{Z_3}{Z_3'} \zeta_3^0, \quad \tilde{\zeta}_3 = \frac{\tilde{Z}_3}{\tilde{Z}_3'} \tilde{\zeta}_3^0. \quad (2.4)$$

Note that since we are interested in the four-loop results for ζ_i the corresponding renormalization constants have to be known with the same accuracy. In ref. [7] the results up to four-loop order have nicely been summarized (see also refs. [8, 9]).

Due to the well-known Ward identities [6] there are several ways to compute the renormalization constant for the strong coupling, Z_g . A convenient relation, which has the advantage that due to the appearance of renormalization constants involving ghosts less diagrams contribute, is given by

$$Z_g = \frac{\tilde{Z}_1}{\tilde{Z}_3 \sqrt{Z_3}}, \quad (2.5)$$

where \tilde{Z}_1 is the renormalization constant of the ghost-gluon vertex $g_s G\bar{c}c$. The same is true for the corresponding equation for the decoupling constant, such that one can use the relation

$$\zeta_g^0 = \frac{\tilde{\zeta}_1^0}{\tilde{\zeta}_3^0 \sqrt{\zeta_3^0}}, \quad (2.6)$$

where $\tilde{\zeta}_1^0$ denotes the decoupling constant for the ghost-gluon vertex. Alternatively, one can use the renormalized objects ζ_3 , $\tilde{\zeta}_3$ from eq. (2.4) as well as $\tilde{\zeta}_1 = \frac{\tilde{Z}_1}{Z_1'} \tilde{\zeta}_1^0$ and then obtain ζ_g from the renormalized version of eq. (2.6).

In refs. [10, 11] formulae for the bare decoupling constants ζ_i^0 are derived which relate the n -loop decoupling constants to n -loop vacuum integrals. In particular, one has

$$\begin{aligned} \zeta_3^0 &= 1 + \Pi_G^{0h}(0), \\ \tilde{\zeta}_3^0 &= 1 + \Pi_c^{0h}(0), \\ \tilde{\zeta}_1^0 &= 1 + \Gamma_{G\bar{c}c}^{0h}(0,0), \end{aligned} \quad (2.7)$$

where $\Pi_G(p^2)$ and $\Pi_c(p^2)$ are the gluon and ghost vacuum polarizations, respectively, and the superscript h denotes the so-called hard part which survives after setting the external momentum to zero. Specifically, $\Pi_G(p^2)$ and $\Pi_c(p^2)$ are related to the gluon and ghost propagators through

$$\begin{aligned} i \int dx e^{ip \cdot x} \langle T G^{0,a\mu}(x) G^{0,b\nu}(0) \rangle &= \delta^{ab} \left\{ \frac{g^{\mu\nu}}{p^2 [1 + \Pi_G^0(p^2)]} + \text{terms proportional to } p^\mu p^\nu \right\}, \\ i \int dx e^{ip \cdot x} \langle T c^{0,a}(x) \bar{c}^{0,b}(0) \rangle &= - \frac{\delta^{ab}}{p^2 [1 + \Pi_c^0(p^2)]}, \end{aligned} \quad (2.8)$$

respectively, while $\Gamma_{G\bar{c}c}^0(p, k)$ is defined through the one-particle-irreducible (1PI) part of the amputated $G\bar{c}c$ Green function as

$$\begin{aligned} i^2 \int dx dy e^{i(p \cdot x + k \cdot y)} \langle T c^{0,a}(x) \bar{c}^{0,b}(0) G^{0,c\mu}(y) \rangle^{1\text{PI}} \\ = p^\mu g_s^0 \left\{ -i f^{abc} [1 + \Gamma_{G\bar{c}c}^0(p, k)] + \text{other colour structures} \right\}, \end{aligned} \quad (2.9)$$

where p and k are the outgoing four-momenta of c and G , respectively, and f^{abc} are the structure constants of the QCD gauge group. Sample four-loop diagrams for each line of eq. (2.7) are shown in figure 1(a)–(c).

From eqs. (2.6), (2.4) and (2.7) it becomes clear that for the calculation of ζ_g four-loop vacuum diagrams are needed. Currently the only practical method to express an arbitrary four-loop vacuum integral in terms of a small set of master integrals is based on the algorithm developed in ref. [12]. The application to four-loop bubbles has been discussed in ref. [2]. First physical results deal with moments of the photon polarization function [4] and the singlet contribution to the electroweak ρ parameter [5]. The essence of the Laporta algorithm [12] is the generation of large tables containing relations between




Figure 1: Sample diagrams for the gluon (a) and ghost (b) propagator and the ghost-gluon vertex (c). In (d) the lowest-order diagram is shown mediating the Higgs-gluon coupling in the Standard Model and (e) shows an example for a five-loop diagram contributing to the result in eq. (4.4).

arbitrary integrals and the so-called master integrals. For the calculation at hand the tables have a size of about 8 GB and contain 6 million equations.

The master integrals needed for the evaluation of ζ_g have been computed in ref. [13], where, however, some of the higher order coefficients in ϵ could only be determined numerically.

3. Running and decoupling for α_s

Whereas at three-loop level of the order of 1000 diagrams have to be considered, at four loops there are almost 20000 diagrams which contribute to the gluon and ghost propagators and the ghost-gluon vertex. They are generated with the program **QGRAF** [14]. With the help of the packages **q2e** and **exp** [15, 16] the topologies and notation are adopted to the program performing the reduction of the four-loop vacuum diagrams [2]. As an output we obtain the bare four-loop results as a linear combination of several master integrals. All of them have been computed in ref. [13].

Since at four-loop order the renormalization is quite non-trivial, let us in the following briefly describe the procedure necessary to arrive at a finite result. It is convenient to build in a first step the sum of the bare contributions to ζ_3^0 , $\tilde{\zeta}_3^0$ and $\tilde{\zeta}_1^0$ and combine them immediately to ζ_g^0 according to eq. (2.6). Already at this point the gauge parameter, ξ , which for the individual pieces starts to appear at three-loop order, drops out and hence spares us from renormalizing ξ . Let us mention that due to the complexity of the intermediate expressions, the four-loop diagrams have been evaluated for Feynman gauge, whereas the lower-order diagrams were computed for general ξ .

In a next step it is convenient to renormalize the parameters $\alpha_s = g_s^2/(4\pi)$ and m_h applying the usual multiplicative renormalization (cf. eq. (2.3)). The corresponding counterterms have to be known up to the three-loop order. At this point one has to apply eq. (2.4) which requires the ratio Z_g/Z'_g up to four-loop order. In order to evaluate this ratio one has to remember that Z'_g is defined in the effective theory and thus depends on α'_s and n_l whereas Z_g depends on α_s and $(n_l + n_h)$. Thus it is necessary to use ζ_g up to three-loop level in order to transform α'_s to α_s where due to the presence of the divergences in Z'_g also higher-order terms in ϵ of ζ_g have to be taken into account.

Finally one arrives at the following finite result for $(\zeta_g)^2$ which for $N_c = 3$ and $n_h = 1$ is given by

$$\begin{aligned}
 \zeta_g^2 = & 1 + \frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \left(-\frac{1}{6} \ln \frac{\mu^2}{m_h^2} \right) + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^2 \left(\frac{11}{72} - \frac{11}{24} \ln \frac{\mu^2}{m_h^2} + \frac{1}{36} \ln^2 \frac{\mu^2}{m_h^2} \right) \\
 & + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^3 \left[\frac{564731}{124416} - \frac{82043}{27648} \zeta(3) - \frac{955}{576} \ln \frac{\mu^2}{m_h^2} + \frac{53}{576} \ln^2 \frac{\mu^2}{m_h^2} - \frac{1}{216} \ln^3 \frac{\mu^2}{m_h^2} \right. \\
 & + n_l \left(-\frac{2633}{31104} + \frac{67}{576} \ln \frac{\mu^2}{m_h^2} - \frac{1}{36} \ln^2 \frac{\mu^2}{m_h^2} \right) \left. + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^4 \left[\frac{291716893}{6123600} \right. \right. \\
 & + \frac{3031309}{1306368} \ln^4 2 - \frac{121}{4320} \ln^5 2 - \frac{3031309}{217728} \zeta(2) \ln^2 2 + \frac{121}{432} \zeta(2) \ln^3 2 - \frac{2362581983}{87091200} \zeta(3) \\
 & - \frac{76940219}{2177280} \zeta(4) + \frac{2057}{576} \zeta(4) \ln 2 + \frac{1389}{256} \zeta(5) + \frac{3031309}{54432} a_4 + \frac{121}{36} a_5 - \frac{151369}{2177280} X_0 \\
 & + \left(\frac{7391699}{746496} - \frac{2529743}{165888} \zeta(3) \right) \ln \frac{\mu^2}{m_h^2} + \frac{2177}{3456} \ln^2 \frac{\mu^2}{m_h^2} - \frac{1883}{10368} \ln^3 \frac{\mu^2}{m_h^2} + \frac{1}{1296} \ln^4 \frac{\mu^2}{m_h^2} \\
 & + n_l \left(-\frac{4770941}{2239488} + \frac{685}{124416} \ln^4 2 - \frac{685}{20736} \zeta(2) \ln^2 2 + \frac{3645913}{995328} \zeta(3) \right. \\
 & - \frac{541549}{165888} \zeta(4) + \frac{115}{576} \zeta(5) + \frac{685}{5184} a_4 + \left(-\frac{110341}{373248} + \frac{110779}{82944} \zeta(3) \right) \ln \frac{\mu^2}{m_h^2} \\
 & - \frac{1483}{10368} \ln^2 \frac{\mu^2}{m_h^2} - \frac{127}{5184} \ln^3 \frac{\mu^2}{m_h^2} \left. \right) + n_l^2 \left(-\frac{271883}{4478976} + \frac{167}{5184} \zeta(3) + \frac{6865}{186624} \ln \frac{\mu^2}{m_h^2} \right. \\
 & \left. \left. - \frac{77}{20736} \ln^2 \frac{\mu^2}{m_h^2} + \frac{1}{324} \ln^3 \frac{\mu^2}{m_h^2} \right) \right] + \mathcal{O} \left(\left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^5 \right), \tag{3.1}
 \end{aligned}$$

where the heavy quark mass m_h is renormalized in the $\overline{\text{MS}}$ scheme at the scale μ . The corresponding expression for the on-shell mass is given in appendix A. In eq. (3.1), $\zeta(n)$ is Riemann's zeta function and $a_n = \text{Li}_n(1/2) = \sum_{k=1}^{\infty} 1/(2^k k^n)$. The constant X_0 , which is the leading coefficient of a certain finite four-loop master integral, is only known numerically with the value [13]

$$X_0 = +1.808879546208334741426364595086952090. \tag{3.2}$$

Interestingly, in principle the number of numerical coefficients occurring in eq. (3.1) should be three. One relation among them can be established through the separate renormalization of the ghost propagator while a further constant has become available recently in analytical form [17]. Thus one remains with one coefficient which is only known numerically.

Inserting numerical values into eq. (3.1) one obtains

$$\begin{aligned}
 \zeta_g^2 \approx & 1 + 0.1528 \left(\frac{\alpha_s^{(n_l+1)}(m_h)}{\pi} \right)^2 + (0.9721 - 0.0847 n_l) \left(\frac{\alpha_s^{(n_l+1)}(m_h)}{\pi} \right)^3 \\
 & + (5.1703 - 1.0099 n_l - 0.0220 n_l^2) \left(\frac{\alpha_s^{(n_l+1)}(m_h)}{\pi} \right)^4. \tag{3.3}
 \end{aligned}$$

It is interesting to note that the n_l -independent four-loop coefficient is relatively big as compared to the corresponding constants at lower loop-order. However, for the interesting values $n_l = (3, 4, 5)$ one observes a big cancellation leading to a well-defined perturbative series with coefficients $(-0.4288, +0.7790, +1.9428)$ in front of $(\alpha_s/\pi)^4$. Note, that the two-loop result for ζ_g has been computed in refs. [18, 19] and the three-loop terms have been evaluated for the first time in ref. [20].

We are now in a position to study the numerical impact of our result. As an example we consider the evaluation of $\alpha_s^{(5)}(M_Z)$ from $\alpha_s^{(4)}(M_\tau)$, i.e. we apply our formalism to the crossing of the bottom quark threshold with $n_l = 4$. In general one assumes that the value of the scale μ_b , where the matching has to be performed, is of order m_b . However, it is not determined by theory. Thus this uncertainty contributes significantly to the error of physical predictions. On general grounds one expects that while including higher order perturbative corrections the relation between $\alpha_s^{(4)}(M_\tau)$ and $\alpha_s^{(5)}(M_Z)$ becomes insensitive to the choice of the matching scale. This has been demonstrated in refs. [21, 10] for the three- and four-loop evolution, respectively. In the following we want to extend the analysis to five loops.

The procedure is as follows. In a first step we calculate $\alpha_s^{(4)}(\mu_b)$ by exactly integrating the equation

$$\frac{\mu^2 d}{d\mu^2} \frac{\alpha_s^{(n_f)}}{\pi} = \beta^{(n_f)}(\alpha_s^{(n_f)}) = - \sum_{i \geq 0} \beta_i^{(n_f)} \left(\frac{\alpha_s^{(n_f)}}{\pi} \right)^{i+2}, \quad (3.4)$$

with the initial condition $\alpha_s^{(4)}(M_\tau) = 0.36$. Afterwards $\alpha_s^{(5)}(\mu_b)$ is obtained from the renormalized version of the first equation in (2.2) where we use ζ_g parameterized in terms of the on-shell mass (cf. eq. (A.1)) $M_b = 4.7 \text{ GeV}$. Finally, we compute $\alpha_s^{(5)}(M_Z)$ using again eq. (3.4). For consistency, i -loop evolution must be accompanied by $(i-1)$ -loop matching, i.e. if we omit terms of $\mathcal{O}(\alpha_s^{i+2})$ on the right-hand side of eq. (3.4), we need to discard those of $\mathcal{O}(\alpha_s^{i+1})$ in eq. (A.1) at the same time. Since the five-loop coefficient in eq. (3.4) is not yet known we set $\beta_4^{(n_f)}$ to zero in our numerical analysis.

In figure 2 the result for $\alpha_s^{(5)}(M_Z)$ as a functions μ_b is displayed for the one- to five-loop analysis. For illustration, μ_b is varied rather extremely, by almost two orders of magnitude. While the leading-order result exhibits a strong logarithmic behaviour, the analysis is gradually getting more stable as we go to higher orders. The five-loop curve is almost flat for $\mu_b \geq 1 \text{ GeV}$ and demonstrates an even more stable behaviour than the four-loop analysis of ref. [10]. It should be noted that around $\mu_b \approx 1 \text{ GeV}$ both the three-, four- and five-loop curves show a strong variation which can be interpreted as a sign for the breakdown of perturbation theory. Besides the μ_b dependence of $\alpha_s^{(5)}(M_Z)$, also its absolute normalization is significantly affected by the higher orders. At the central matching scale $\mu_b = M_b$, we encounter a rapid convergence behaviour.

4. Effective coupling between a Higgs boson and gluons

In this section we want to discuss the relation between ζ_g and the coupling of a scalar Higgs boson to gluons. Due to the fact that gluons are massless, there is no coupling at




Figure 2: μ_b dependence of $\alpha_s^{(5)}(M_Z)$ calculated from $\alpha_s^{(4)}(M_\tau) = 0.36$ and $M_b = 4.7 \text{ GeV}$. The procedure is described in the text. The dotted, short-dashed, long-dashed and dash-dotted line corresponds to one- to four-loop running. The solid curve includes the effect of the new four-loop matching term.

tree-level. At one-loop order the HGG coupling is mediated via a top-quark loop depicted in figure 1(d).

For an intermediate-mass Higgs boson which formally obeys the relation $M_H \ll m_t$ it is possible to construct an effective lagrangian of the form

$$\mathcal{L}_{\text{eff}} = -\frac{H^0}{v^0} C_1 \mathcal{O}_1, \quad (4.1)$$

with the effective operator

$$\mathcal{O}_1 = (G_{\mu\nu}^a)^2, \quad (4.2)$$

where $G_{\mu\nu}^a$ is the colour field strength. The coefficient function C_1 incorporates the contribution from the top-quark loops. At one-loop order it is easy to see that the contribution from the triangle diagrams can be obtained through the derivative of the one-loop diagram for Π_G^0 with respect to the top-quark mass. However, at higher-loop orders this simple picture does not hold anymore and the relation between the HGG diagrams and derivatives of the two-point functions containing a top-quark loop gets more involved. In ref. [10] an all-order low-energy theorem has been derived which establishes such a relation and which has a surprisingly simple form (for definiteness we specify to the top-quark in this section):

$$C_1 = -\frac{1}{2} \frac{m_t^2}{\partial m_t^2} \ln \zeta_g^2. \quad (4.3)$$

An appealing feature of eq. (4.3) is that at a given order in α_s only the logarithmic contributions of ζ_g are needed for the calculation of C_1 at the same order. Thus, from our calculation we can reconstruct the five-loop logarithms of ζ_g from lower-order terms and the β and γ_m functions governing the running of α_s and the top-quark mass, respectively. This leads to the following result, at $N_c = 3$ and $n_h = 1$,

$$\begin{aligned}
 C_1 = & -\frac{1}{12} \frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \left\{ 1 + \frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \left(\frac{11}{4} - \frac{1}{6} \ln \frac{\mu^2}{m_t^2} \right) \right. \\
 & + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^2 \left[\frac{2821}{288} - \frac{3}{16} \ln \frac{\mu^2}{m_t^2} + \frac{1}{36} \ln^2 \frac{\mu^2}{m_t^2} + n_l \left(-\frac{67}{96} + \frac{1}{3} \ln \frac{\mu^2}{m_t^2} \right) \right] \\
 & + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^3 \left[-\frac{4004351}{62208} + \frac{1305893}{13824} \zeta(3) - \frac{859}{288} \ln \frac{\mu^2}{m_t^2} + \frac{431}{144} \ln^2 \frac{\mu^2}{m_t^2} - \frac{1}{216} \ln^3 \frac{\mu^2}{m_t^2} \right. \\
 & + n_l \left(\frac{115607}{62208} - \frac{110779}{13824} \zeta(3) + \frac{641}{432} \ln \frac{\mu^2}{m_t^2} + \frac{151}{288} \ln^2 \frac{\mu^2}{m_t^2} \right) \\
 & + n_l^2 \left(-\frac{6865}{31104} + \frac{77}{1728} \ln \frac{\mu^2}{m_t^2} - \frac{1}{18} \ln^2 \frac{\mu^2}{m_t^2} \right) \left. \right] \\
 & + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^4 \left[-\frac{69820734619}{27993600} - \frac{39407017}{373248} \ln^4 2 + \frac{11011}{8640} \ln^5 2 + \frac{39407017}{62208} \zeta(2) \ln^2 2 \right. \\
 & - \frac{11011}{864} \zeta(2) \ln^3 2 + \frac{27642438179}{24883200} \zeta(3) + \frac{996205247}{622080} \zeta(4) - \frac{187187}{1152} \zeta(4) \ln 2 - \frac{894391}{4608} \zeta(5) \\
 & - \frac{39407017}{15552} a_4 - \frac{11011}{72} a_5 + \frac{1967797}{622080} X_0 \\
 & - \left(\frac{1276661933}{1492992} - \frac{226222121}{331776} \zeta(3) \right) \ln \frac{\mu^2}{m_t^2} + \frac{33517}{1728} \ln^2 \frac{\mu^2}{m_t^2} + \frac{140357}{20736} \ln^3 \frac{\mu^2}{m_t^2} + \frac{1}{1296} \ln^4 \frac{\mu^2}{m_t^2} \\
 & + n_l \left(\frac{58259821853}{195955200} + \frac{3896297}{580608} \ln^4 2 - \frac{121}{1440} \ln^5 2 - \frac{3896297}{96768} \zeta(2) \ln^2 2 + \frac{121}{144} \zeta(2) \ln^3 2 \right. \\
 & - \frac{74306021071}{348364800} \zeta(3) + \frac{141211087}{3870720} \zeta(4) + \frac{2057}{192} \zeta(4) \ln 2 - \frac{20227}{2304} \zeta(5) + \frac{3896297}{24192} a_4 + \frac{121}{12} a_5 \\
 & - \frac{151369}{725760} X_0 + \left(\frac{23250409}{186624} - \frac{8736121}{82944} \zeta(3) \right) \ln \frac{\mu^2}{m_t^2} + \frac{569}{2304} \ln^2 \frac{\mu^2}{m_t^2} + \frac{2551}{2592} \ln^3 \frac{\mu^2}{m_t^2} \left. \right) \\
 & + n_l^2 \left(-\frac{33014371}{8957952} + \frac{685}{41472} \ln^4 2 - \frac{685}{6912} \zeta(2) \ln^2 2 + \frac{970259}{110592} \zeta(3) - \frac{518509}{55296} \zeta(4) \right. \\
 & + \frac{115}{192} \zeta(5) + \frac{685}{1728} a_4 - \left(\frac{1107181}{186624} - \frac{28297}{9216} \zeta(3) \right) \ln \frac{\mu^2}{m_t^2} - \frac{1729}{13824} \ln^2 \frac{\mu^2}{m_t^2} - \frac{1205}{5184} \ln^3 \frac{\mu^2}{m_t^2} \left. \right) \\
 & + n_l^3 \left(-\frac{255947}{1492992} + \frac{5}{64} \zeta(3) + \frac{481}{5184} \ln \frac{\mu^2}{m_t^2} - \frac{77}{6912} \ln^2 \frac{\mu^2}{m_t^2} + \frac{1}{108} \ln^3 \frac{\mu^2}{m_t^2} \right) \\
 & \left. + 6 \left(\beta_4^{(n_l)} - \beta_4^{(n_l+1)} \right) \right] + \mathcal{O} \left(\left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^5 \right) \left. \right\}, \tag{4.4}
 \end{aligned}$$

with m_t being the $\overline{\text{MS}}$ top-quark mass renormalized at the scale μ . Note the appearance of the flavour-dependent part of β_4 in the five-loop contribution, whereas the corresponding coefficient from the anomalous mass dimension does not appear. We want to stress that the term of order α_s^5 covers the contributions from five-loop diagrams like the one in figure 1(e).

Evaluating eq. (4.4) numerically leads to

$$\begin{aligned}
C_1 \approx & -\frac{1}{12} \frac{\alpha_s^{(n_l+1)}(m_t)}{\pi} \left[1 + 2.7500 \frac{\alpha_s^{(n_l+1)}(m_t)}{\pi} + (9.7951 - 0.6979 n_l) \left(\frac{\alpha_s^{(n_l+1)}(m_t)}{\pi} \right)^2 \right. \\
& + (49.1827 - 7.7743 n_l - 0.2207 n_l^2) \left(\frac{\alpha_s^{(n_l+1)}(m_t)}{\pi} \right)^3 \\
& + \left(-662.5065 + 137.6005 n_l - 2.5367 n_l^2 - 0.0775 n_l^3 + 6 \left(\beta_4^{(n_l)} - \beta_4^{(n_l+1)} \right) \right) \\
& \left. \times \left(\frac{\alpha_s^{(n_l+1)}(m_t)}{\pi} \right)^4 \right]. \tag{4.5}
\end{aligned}$$

Again one observes large cancellations between the n_l^0 and n_l^1 term in the five-loop contribution to C_1 .

Note that the result of eq. (4.4) constitutes a building block for the N⁴LO calculation to the Higgs boson production and decay in the two-gluon channel, for which the complete answer currently is certainly out of range. Still, the five-loop result for C_1 constitutes a high-order result in perturbative QCD which is of theoretical interest by itself.

5. Conclusions

In this paper the decoupling constant of the strong coupling is presented to four-loop order. This constitutes a fundamental quantity of QCD and is one of the very few known to such a high order. The decoupling constant is necessary for performing a consistent running of α_s with five-loop accuracy including important effects from the crossing of quark thresholds. The calculation has been performed analytically, and the main result can be found in eq. (3.1). With the help of a low-energy theorem it is possible to derive the five-loop result for the effective coupling of the Higgs boson to gluons, which constitutes a building block in the corresponding production and decay processes.

We want to mention that the result for ζ_g^2 in eq. (3.1) has been obtained independently in ref. [22]. Except for QGRAF, which is used for the generation of the diagrams, there is no common code. Even the master integrals have meanwhile been computed independently [23] and for the renormalization a different procedure has been chosen.

Acknowledgments

This work was supported by the SFB/TR 9. We would like to thank the authors of ref. [22] for communicating their result before publication.

A. Results for ζ_g^{OS}

Replacing in eq. (3.1) the $\overline{\text{MS}}$ mass m_h by the pole mass M_h using the three-loop approximation [24–26] one gets

$$\begin{aligned}
 (\zeta_g^{\text{OS}})^2 = & 1 + \frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \left(-\frac{1}{6} \ln \frac{\mu^2}{M_h^2} \right) + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^2 \left(-\frac{7}{24} - \frac{19}{24} \ln \frac{\mu^2}{M_h^2} + \frac{1}{36} \ln^2 \frac{\mu^2}{M_h^2} \right) \\
 & + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^3 \left[-\frac{58933}{124416} - \frac{2}{3} \zeta(2) - \frac{2}{9} \zeta(2) \ln 2 - \frac{80507}{27648} \zeta(3) - \frac{8521}{1728} \ln \frac{\mu^2}{M_h^2} \right. \\
 & \left. - \frac{131}{576} \ln^2 \frac{\mu^2}{M_h^2} - \frac{1}{216} \ln^3 \frac{\mu^2}{M_h^2} + n_l \left(\frac{2479}{31104} + \frac{1}{9} \zeta(2) + \frac{409}{1728} \ln \frac{\mu^2}{M_h^2} \right) \right] \\
 & + \left(\frac{\alpha_s^{(n_l+1)}(\mu)}{\pi} \right)^4 \left[-\frac{141841753}{24494400} + \frac{3179149}{1306368} \ln^4 2 - \frac{121}{4320} \ln^5 2 - \frac{697121}{19440} \zeta(2) \right. \\
 & + \frac{1027}{162} \zeta(2) \ln 2 - \frac{2913037}{217728} \zeta(2) \ln^2 2 + \frac{121}{432} \zeta(2) \ln^3 2 - \frac{2408412383}{87091200} \zeta(3) \\
 & + \frac{1439}{216} \zeta(3) \zeta(2) - \frac{71102219}{2177280} \zeta(4) + \frac{2057}{576} \zeta(4) \ln 2 + \frac{49309}{20736} \zeta(5) \\
 & + \frac{3179149}{54432} a_4 + \frac{121}{36} a_5 - \frac{151369}{2177280} X_0 \\
 & - \left(\frac{19696909}{746496} + \frac{29}{9} \zeta(2) + \frac{29}{27} \zeta(2) \ln 2 + \frac{2439119}{165888} \zeta(3) \right) \ln \frac{\mu^2}{M_h^2} - \frac{7693}{1152} \ln^2 \frac{\mu^2}{M_h^2} \\
 & - \frac{8371}{10368} \ln^3 \frac{\mu^2}{M_h^2} + \frac{1}{1296} \ln^4 \frac{\mu^2}{M_h^2} + n_l \left(\frac{1773073}{746496} + \frac{173}{124416} \ln^4 2 + \frac{557}{162} \zeta(2) \right. \\
 & + \frac{22}{81} \zeta(2) \ln 2 - \frac{1709}{20736} \zeta(2) \ln^2 2 + \frac{4756441}{995328} \zeta(3) - \frac{697709}{165888} \zeta(4) + \frac{115}{576} \zeta(5) + \frac{173}{5184} a_4 \\
 & + \left(\frac{1110443}{373248} + \frac{41}{54} \zeta(2) + \frac{2}{27} \zeta(2) \ln 2 + \frac{132283}{82944} \zeta(3) \right) \ln \frac{\mu^2}{M_h^2} + \frac{6661}{10368} \ln^2 \frac{\mu^2}{M_h^2} \\
 & + \frac{107}{1728} \ln^3 \frac{\mu^2}{M_h^2} \Big) + n_l^2 \left(-\frac{140825}{1492992} - \frac{13}{162} \zeta(2) - \frac{19}{1728} \zeta(3) \right. \\
 & \left. - \left(\frac{1679}{186624} + \frac{1}{27} \zeta(2) \right) \ln \frac{\mu^2}{M_h^2} - \frac{493}{20736} \ln^2 \frac{\mu^2}{M_h^2} \right) \Big] \\
 \approx & 1 - 0.2917 \left(\frac{\alpha_s^{(n_l+1)}(M_h)}{\pi} \right)^2 + (-5.3239 + 0.2625 n_l) \left(\frac{\alpha_s^{(n_l+1)}(M_h)}{\pi} \right)^3 \\
 & + (-85.8750 + 9.6923 n_l - 0.2395 n_l^2) \left(\frac{\alpha_s^{(n_l+1)}(M_h)}{\pi} \right)^4. \tag{A.1}
 \end{aligned}$$

References

- [1] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Strange quark mass from τ lepton decays with $\mathcal{O}(\alpha_s^3)$ accuracy*, *Phys. Rev. Lett.* **95** (2005) 012003 [[hep-ph/0412350](#)].
- [2] Y. Schröder, *Automatic reduction of four-loop bubbles*, *Nucl. Phys.* **116** (Proc. Suppl.) (2003) 402 [[hep-ph/0211288](#)].
- [3] C. Sturm, PhD thesis (June 2005), Karlsruhe Univ., Germany (unpublished).
- [4] K.G. Chetyrkin, J.H. Kühn, P. Mastrolia and C. Sturm, *Heavy-quark vacuum polarization: first two moments of the $\mathcal{O}(\alpha_s^3 n_f^2)$ contribution*, *Eur. Phys. J. C* **40** (2005) 361 [[hep-ph/0412055](#)].

- [5] Y. Schröder and M. Steinhauser, *Four-loop singlet contribution to the ρ parameter*, *Phys. Lett. B* **622** (2005) 124 [[hep-ph/0504055](#)].
- [6] T. Muta, *Foundations of quantum chromodynamics*, World Scientific, Singapore, 1987.
- [7] K.G. Chetyrkin, *Four-loop renormalization of QCD: full set of renormalization constants and anomalous dimensions*, *Nucl. Phys. B* **710** (2005) 499 [[hep-ph/0405193](#)].
- [8] T. van Ritbergen, J.A.M. Vermaseren and S.A. Larin, *The four-loop beta function in quantum chromodynamics*, *Phys. Lett. B* **400** (1997) 379 [[hep-ph/9701390](#)].
- [9] M. Czakon, *The four-loop QCD β -function and anomalous dimensions*, *Nucl. Phys. B* **710** (2005) 485 [[hep-ph/0411261](#)].
- [10] K.G. Chetyrkin, B.A. Kniehl and M. Steinhauser, *Decoupling relations to $\mathcal{O}(\alpha_s^3)$ and their connection to low-energy theorems*, *Nucl. Phys. B* **510** (1998) 61 [[hep-ph/9708255](#)].
- [11] M. Steinhauser, *Results and techniques of multi-loop calculations*, *Phys. Rept.* **364** (2002) 247 [[hep-ph/0201075](#)].
- [12] S. Laporta, *High-precision calculation of multi-loop Feynman integrals by difference equations*, *Int. J. Mod. Phys. A* **15** (2000) 5087 [[hep-ph/0102033](#)].
- [13] Y. Schröder and A. Vuorinen, *High-precision epsilon expansions of single-mass-scale four-loop vacuum bubbles*, *JHEP* **06** (2005) 051 [[hep-ph/0503209](#)].
- [14] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279.
- [15] T. Seidensticker, *Automatic application of successive asymptotic expansions of Feynman diagrams*, [hep-ph/9905298](#).
- [16] R. Harlander, T. Seidensticker and M. Steinhauser, *Complete corrections of $\mathcal{O}(\alpha_s)$ to the decay of the Z boson into bottom quarks*, *Phys. Lett. B* **426** (1998) 125 [[hep-ph/9712228](#)].
- [17] Y. Schröder, in preparation.
- [18] W. Bernreuther and W. Wetzel, *Decoupling of heavy quarks in the minimal subtraction scheme*, *Nucl. Phys. B* **197** (1982) 228, erratum *ibid.* **B513** (1998) 758.
- [19] S.A. Larin, T. van Ritbergen and J.A.M. Vermaseren, *The large quark mass expansion of $\Gamma(Z^0 \rightarrow \text{hadrons})$ and $\Gamma(\tau^- \rightarrow \nu_\tau + \text{hadrons})$ in the order α_s^3* , *Nucl. Phys. B* **438** (1995) 278 [[hep-ph/9411260](#)].
- [20] K.G. Chetyrkin, B.A. Kniehl and M. Steinhauser, *Strong coupling constant with flavour thresholds at four loops in the \bar{MS} scheme*, *Phys. Rev. Lett.* **79** (1997) 2184 [[hep-ph/9706430](#)].
- [21] G. Rodrigo, A. Pich and A. Santamaria, *$\alpha_s(m_Z)$ from τ decays with matching conditions at three loops*, *Phys. Lett. B* **424** (1998) 367 [[hep-ph/9707474](#)].
- [22] K.G. Chetyrkin, J.H. Kühn and C. Sturm, *QCD decoupling at four loops*, [hep-ph/0512060](#).
- [23] K.G. Chetyrkin, M. Faisst, C. Sturm and M. Tentukov, in preparation.
- [24] K.G. Chetyrkin and M. Steinhauser, *Short distance mass of a heavy quark at order α_s^3* , *Phys. Rev. Lett.* **83** (1999) 4001 [[hep-ph/9907509](#)].
- [25] K.G. Chetyrkin and M. Steinhauser, *The relation between the \bar{MS} and the on-shell quark mass at order α_s^3* , *Nucl. Phys. B* **573** (2000) 617 [[hep-ph/9911434](#)].
- [26] K. Melnikov and T. v. Ritbergen, *The three-loop relation between the \bar{MS} and the pole quark masses*, *Phys. Lett. B* **482** (2000) 99 [[hep-ph/9912391](#)].