

PAPER • OPEN ACCESS


Radiative cooling of hot C_n^- and C_nH^- molecules

To cite this article: M Kaminska *et al* 2015 *J. Phys.: Conf. Ser.* **635** 112124

View the [article online](#) for updates and enhancements.

You may also like

- [Anion production in high-velocity cluster–atom collisions: the electron capture process revisited](#)
K Béroff, M Chabot, G Martinet et al.
- [Lifetimes of bound excited states of Pt](#)
KC Chartkunchand, M. Kamiska, E. K. Anderson et al.
- [Finite Crystallization and Wulff shape emergence for ionic compounds in the square lattice](#)
Manuel Friedrich and Leonard Kreutz



ECS
The
Electrochemical
Society
Advancing solid state &
electrochemical science & technology

DISCOVER
how sustainability
intersects with
electrochemistry & solid
state science research

Radiative cooling of hot C_n^- and C_nH^- molecules

M. Kaminska^{*†1}, R. F. Nascimento^{*}, M. H. Stockett^{**}, E. K. Anderson^{*}, R. Delaunay[‡],
V. Vizcaino[‡], P. Rousseau[‡], L. Adoui[‡], B. A. Huber[‡], R. D. Thomas^{*}, M. Gatchell^{*},
K. Hansen^{*‡}, H. Zettergren^{*}, H. T. Schmidt^{*}, and H. Cederquist^{*2}

^{*} Department of Physics, Stockholm University, Alba Nova University Centre, SE-106 91 Stockholm, Sweden

[†] Institute of Physics, Jan Kochanowski University, 25-369 Kielce, Poland

^{*} Department of Physics and Astronomy, Aarhus University, Ny Munkegade, DK 8000 Aarhus C, Denmark

[‡] CIMAP, UMR 6252, CEA/CNRS/EINSCAEN/Université de Caen Basse-Normandie, bd Henri Becquerel, BP 5133, F-14070 Caen cedex 05, France

[‡] Department of Physics, University of Gothenburg, SE-412 96 Göteborg, Sweden

Synopsis We have measured the rates of neutrals produced from 10 keV C_n^- or C_nH^- ($n=2, 4, 6, 8$, and 10) ion beams stored in one of DESIREE's 14 K storage rings. For $n=4, 6$, and 8 we observe marked differences between C_n^- and C_nH^- cooling rates as inverse internal conversion [cf. S. Martin *et al* (2013) *Phys. Rev. Lett.* **110**, 063003] processes are effective for the C_n^- ions only. Knowledge of the cooling rates of these ions are important for estimates of their formation and destruction rates in cold interstellar environments.

We investigate the cooling of internally hot C_n^- and C_nH^- ions with $n=2, 4, 6, 8$, and 10 stored in the electrostatic ion storage ring DESIREE [1, 2] operating at 14 K. We record the signal of neutrals detected after a straight section of the ring as a function of time, t , after producing internally hot C_n^- or C_nH^- ions in a sputter ion source. This production method gives wide internal temperature distributions. If the ions do not stabilize radiatively very fast, the hottest ones will decay first giving overall rates of neutrals following t^{-p} power laws [3].

In Fig. 1 we show data for $n=4$ (the results for $n=6$ and $n=8$ are qualitatively similar). While the C_4^- -signal is small and disappears rapidly (sub-milliseconds), the C_4H^- -signal follows a power law for tens of milliseconds. Similar to C_6^- [4, 5], C_4^- and C_8^- are open-shell systems that cool very rapidly as they can convert (part of) their internal vibrational energies to low-lying electronically excited states, which decay through fast radiative transitions. The C_nH^- ions, on the other hand, are closed shell systems and have much higher HOMO-LUMO gaps. This means that inverse internal conversions to electronically excited states become highly unlikely [4, 5] and internal cooling must proceed through much slower vibrational transitions. While C_2H^- and $C_{10}H^-$ have similar decay behaviors as C_4H^- , C_6H^- and C_8H^- , the decays for C_2^- and C_{10}^- are much slower than those for C_4^- , C_6^- and C_8^- . The C_{10}^- decay is also more complex possibly reflecting the presence of linear and ring isomers in the stored beam. For C_n^- and C_nH^- anions, knowledge of their cool-

ing rates are considered to be very important for estimates of their interstellar abundances [6].




Figure 1. Rates of detected neutral particles normalized to the injected ion currents for the two molecular ions C_4^- and C_4H^- . The rapid disappearance of the signal for C_4^- is due to fast cooling through inverse internal conversion [4, 5, 7].

References

- [1] R. D. Thomas *et al* 2011 *Rev. Sci. Instrum.* **82** 065112
- [2] H. T. Schmidt *et al* 2013 *Rev. Sci. Instrum.* **84** 055115
- [3] K. Hansen *et al* 2001 *Phys. Rev. Lett.* **87** 123401
- [4] G. Ito *et al* 2014 *Phys. Rev. Lett.* **112** 183001
- [5] V. Chandrasekaran *et al* 2014 *Phys. Chem. Lett.* **5** 4078
- [6] H. Brünken *et al* 2007 *Astrophysical J.* **664** L43
- [7] S. Martin *et al* 2013 *Phys. Rev. Lett.* **110** 063003

¹E-mail: magdalena.kaminska@fysik.su.se

²E-mail: cederq@fysik.su.se

