PAPER • OPEN ACCESS


Probing the Symmetry Term of the Nuclear Equation of State at High Baryonic Densities

To cite this article: P Russotto et al 2017 J. Phys.: Conf. Ser. 863 012059

View the <u>article online</u> for updates and enhancements.

You may also like

- Fast b-tagging at the high-level trigger of the ATLAS experiment in LHC Run 3
 G. Aad, B. Abbott, K. Abeling et al.
- <u>The ATLAS Fast TracKer system</u> The ATLAS collaboration, G. Aad, B. Abbott et al.
- Operation and performance of the ATLAS semiconductor tracker in LHC Run 2
 The ATLAS collaboration, Georges Aad, Brad Abbott et al.



Probing the Symmetry Term of the Nuclear Equation of State at High Baryonic Densities

```
P Russotto<sup>1</sup>, L Acosta<sup>2,3</sup>, M Adamczyk<sup>4</sup>, A Al-Ajlan<sup>5</sup>, M Al-Garawi<sup>6</sup>,
S Al-Homaidhi<sup>5</sup>, F Amorini<sup>2</sup>, L Auditore<sup>7,8</sup>, T Aumann<sup>9,10</sup>,
Y Ayyad<sup>11</sup>, Z Basrak<sup>12</sup>, J Benlliure<sup>11</sup>, M Boisjoli<sup>13</sup>, K Boretzky<sup>10</sup>,
J Brzychczyk<sup>4</sup>, A Budzanowski<sup>14,35</sup>, C Caesar<sup>9</sup>, G Cardella<sup>1</sup>,
P Cammarata<sup>15</sup>, Z Chajecki<sup>16</sup>, M Chartier<sup>17</sup>, A Chbihi<sup>10</sup>, M Colonna<sup>2</sup>, M D Cozma<sup>18</sup>, B Czech<sup>14</sup>, E De Filippo<sup>1</sup>, M Di Toro<sup>2,19</sup>, M Famiano<sup>20</sup>, S Gannon<sup>17</sup>, I Gašparić<sup>9,12</sup>, L Grassi<sup>12</sup>, C Guazzoni<sup>21,22</sup>,
P Guazzoni<sup>21,23</sup>, M Heil<sup>10</sup>, L Heilborn<sup>15</sup>, R Introzzi<sup>24</sup>, T Isobe<sup>25</sup>, K Kezzar<sup>6</sup>, M Kiš<sup>10</sup>, A Krasznahorkay<sup>26</sup>, S Kupny<sup>4</sup>, N Kurz<sup>10</sup>,
E La Guidara<sup>1</sup>, G Lanzalone<sup>2,27</sup>, P Lasko<sup>4</sup>, A Le Fèvre<sup>10</sup>, Y Leifels<sup>10</sup>,
R C Lemmon<sup>28</sup>, QF Li<sup>29</sup>, I Lombardo<sup>30,31</sup>, J Łukasik<sup>14</sup>, W G Lynch<sup>16</sup>,
P Marini<sup>13,15</sup>, Z Matthews<sup>17</sup>, L May<sup>15</sup>, T Minniti<sup>1</sup>, M Mostazo<sup>11</sup>, A Pagano<sup>1</sup>, E V Pagano<sup>2,19</sup>, M Papa<sup>1</sup>, P Pawłowski<sup>14</sup>, S Pirrone<sup>1</sup>,
G Politi<sup>1,19</sup>, F Porto<sup>2,19</sup>, W Reviol<sup>32</sup>, F Riccio<sup>21,22</sup>, F Rizzo<sup>2,19</sup>,
E Rosato<sup>30,31,35</sup>, D Rossi<sup>9,10</sup>, S Santoro<sup>7,8</sup>, D G Sarantites<sup>32</sup>, H Simon<sup>10</sup>, I Skwirczynska<sup>14</sup>, Z Sosin<sup>4,35</sup>, L Stuhl<sup>26</sup>, W Trautmann<sup>10</sup>, A Trifirò<sup>7,8</sup>,
M Trimarchi<sup>7,8</sup>, M B Tsang<sup>16</sup>, G Verde<sup>1</sup>, M Veselsky<sup>33</sup>, M Vigilante<sup>30,31</sup>, Yongjia Wang<sup>29</sup>, A Wieloch<sup>4</sup>, P Wigg<sup>17</sup>, J Winkelbauer<sup>16</sup>, H H Wolter<sup>34</sup>, P Wu<sup>17</sup>, S Yennello<sup>15</sup>, P Zambon<sup>21,22</sup>, L Zetta<sup>21,23</sup> and M Zoric<sup>12</sup>
 <sup>1</sup>INFN-Sezione di Catania, Catania, Italy
<sup>2</sup>INFN-Laboratori Nazionali del Sud, Catania, Italy
<sup>3</sup>Universidad Nacional Autònoma de Mèxico, Mexico
 <sup>4</sup>Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakòw, Poland
 <sup>5</sup>KACST Riyadh, Riyadh, Saudi Arabia
 <sup>6</sup>King Saud University, Riyadh, Saudi Arabia
 <sup>7</sup>INFN-Gruppo Collegato di Messina, Messina, Italy
 <sup>8</sup>Università di Messina, Messina, Italy
<sup>9</sup>Technische Universität, Darmstadt, Germany
 <sup>10</sup>GSI Helmholtzzentrum, Darmstadt, Germany
 <sup>11</sup>University of Santiago de Compostela, Santiago de Compostela, Spain
 <sup>12</sup>Ruder Boškovič Institute, Zagreb, Croatia
 <sup>13</sup>GANIL, Caen, France
 <sup>14</sup>H. Niewodniczáski Institute of Nuclear Physics, Kraków, Poland
 ^{15}\mathrm{Texas} A&M University, College Station, USA
 ^{16}{\rm NSCL} Michigan State University, East Lansing, USA
 <sup>17</sup>University of Liverpool, Liverpool, UK
```

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

¹⁸IFIN-HH, Magurele-Bucharest, Romania
 ¹⁹Università di Catania, Catania, Italy
 ²⁰Western Michigan University, USA

 $^{^{35}}$ deceased

IOP Conf. Series: Journal of Physics: Conf. Series 863 (2017) 012059

doi:10.1088/1742-6596/863/1/012059

```
<sup>21</sup>INFN-Sezione di Milano, Milano, Ialy
```

E-mail: russotto@lns.infn.it

Abstract. In the ASY-EOS experiment flows of neutrons and light charged particles were measured for $^{197}\mathrm{Au}+^{197}\mathrm{Au}$ collisions at 400 MeV/nucleon, in order to investigate the strength of the symmetry term of the nuclear equation of state at supra-saturation densities. By comparing the experimental data with the UrQMD transport model predictions, we have extracted a new constraint in agreement with the moderately soft to linear density dependence obtained in the former analysis on FOPI-LAND data, but reducing the associated uncertainty by a factor ~ 2 .

1. ASY-EOS experiment results

The neutron-proton elliptic flow ratio has proved to be an effective probe of the high-density behavior of the symmetry energy contribution to nuclear equation of state [1]. By comparing data for the Au+Au system from the FOPI-LAND experiment [2] with calculations performed with the UrQMD transport model [3], an estimate of a moderately soft to linear behavior of the symmetry energy was obtained, although with a considerable statistical uncertainty. The same data set was also compared to calculations performed with the Tübingen QMD model and a constraint compatible with the UrQMD result was obtained [4]. It is worth pointing that in the latter study different parameterizations of the isovector part of the equation of state, the Gogny inspired (momentum dependent) vs. the power law (momentum independent) potential, lead to




Figure 1. Schematic view of the experimental set-up of the ASY-EOS experiment at GSI.

²²Politecnico di Milano, Milano, Italy

²³Università degli Studi di Milano, Milano, Italy

²⁴INFN, Politecnico di Torino, Torino, Italy

²⁵RIKEN, Wako, Japan

²⁶Institute of Nuclear Research, Debrecen, Hungary

²⁷Università Kore, Enna, Italy

²⁸STFC Laboratory, Daresbury, UK

²⁹Huzhou Teachers College, China

³⁰INFN-Sezione di Napoli, Napoli, Italy

³¹Università di Napoli, Napoli, Italy

³²Washington University, St. Louis, USA

³³Institute of Physics, Slovak Academy of Sciences, Bratislava, Slovakia

 $^{^{34}}$ LMU, München, Germany

IOP Conf. Series: Journal of Physics: Conf. Series 863 (2017) 012059

doi:10.1088/1742-6596/863/1/012059




Figure 2. Constraints deduced for the density dependence of the symmetry energy from the ASY-EOS experiment in comparison with the FOPI-LAND result of Ref. [1] as a function of the reduced density ρ/ρ_0 . See Ref. [10] for more details.

very similar results, suggesting the possibility to obtain a model independent constraint.

In order to improve the statistical accuracy, the neutron-charged particles elliptic flow ratio in the symmetric collision system ¹⁹⁷Au+¹⁹⁷Au at 400 MeV/nucleon incident energies was remeasured in the ASY-EOS experiment, carried out at the GSI laboratory in May 2011 (S394 experiment), by coupling the Large Area Neutron Detector (LAND) [5], the Krakòw Triple Telescope Array (KRATTA) [6], the ALADIN Time-of-Flight (AToF) wall [7], 8 rings of the CHIMERA multidetector [8] and 4 rings of the Washington-University Microball array [9], as schematically shown in Fig. 1. More details can be found in Ref. [10].

By comparing the experimental ratio with the UrQMD predictions, a value γ =0.72±0.19 is obtained for the power-law coefficient of the potential part of the symmetry energy parametrization adopted in the UrQMD model, as shown in Fig. 2; the density region probed is $\sim 1.5~\rho_0$. This new result agrees with the older one from the FOPI-LAND data, but reduces by a factor ~ 2 the total uncertainty.

The results of the ASY-EOS experiment confirms the effectiveness of the elliptic flows ratio in probing the symmetry energy at high-densities. We plan to extend such a kind of measurements at GSI to higher beam energies, up to ~ 1 GeV/nucleon, in order to explore higher densities (up to about 2.5 ρ_0), profiting of the unique opportunity offered by the NeuLAND detector, presently under construction as part of the R3B experimental setup [11] and the availability of very asymmetric radioactive ion beams that will be delivered by the FAIR facility.

References

- [1] Russotto P et al 2011 Phys. Lett. B 697 471; Russotto P et al 2014 Eur. Phys. Jour. A 50 38
- [2] Leifels Y et al 1993 Phys. Rev. Lett. 71 963
- [3] Li Q et al 2005 J. Phys G $\bf 31$ 1359
- $[4]\ {\rm Cozma\ M\ D}\ et\ al\ 2013\ Phys.\ Rev.\ C\ {\bf 88}\ 044912$
- [5] Blaich Th et al 1992 Nucl. Instr. and Meth. A 314 136
- [6] Łukasik J et al 2013 Nucl. Instr. and Meth. A 709 120
- [7] Schüttauf A et al 1996 Nucl. Phys. A 607 457
- [8] Pagano A et al 2004 Nucl. Phys. A 734 504
- [9] Sarantites D G et al 1996 Nucl. Instr. and Meth. ${\bf 381}$ 418
- [10] Russotto P et al 2016 Phys. Rev. C 94 034608
- [11] see https://www.gsi.de/r3b.