

LETTER • OPEN ACCESS


Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales

To cite this article: M Zampieri *et al* 2017 *Environ. Res. Lett.* **12** 064008

View the [article online](#) for updates and enhancements.

You may also like

- [Predicting spatial and temporal variability in crop yields: an inter-comparison of machine learning, regression and process-based models](#)
Guoyong Leng and Jim W Hall
- [Reckoning wheat yield trends](#)
M Lin and P Huybers
- [Global gridded crop models underestimate yield responses to droughts and heatwaves](#)
Stefanie Heinicke, Katja Frieler, Jonas Jägermeyr *et al.*



The Breath Biopsy® Guide
Fourth edition

FREE

DOWNLOAD THE FREE E-BOOK

BREATH BIOPSY

OWLSTONE MEDICAL

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED
13 February 2017

REVISED
24 April 2017

ACCEPTED FOR PUBLICATION
10 May 2017

PUBLISHED
5 June 2017

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 3.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the
title of the work, journal
citation and DOI.



Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales

M Zampieri¹, A Ceglar, F Dentener and A Toreti

European Commission, Joint Research Centre, Ispra, Italy

¹ Author to whom any correspondence should be addressed.

E-mail: matteo.zampieri@ec.europa.eu

Keywords: wheat yield, heat waves, drought, water excess, Combined Stress Index (CSI), Heat Magnitude Day (HMD) index

Supplementary material for this article is available [online](#)

Abstract

Heat waves and drought are often considered the most damaging climatic stressors for wheat. In this study, we characterize and attribute the effects of these climate extremes on wheat yield anomalies (at global and national scales) from 1980 to 2010. Using a combination of up-to-date heat wave and drought indexes (the latter capturing both excessively dry and wet conditions), we have developed a composite indicator that is able to capture the spatio-temporal characteristics of the underlying physical processes in the different agro-climatic regions of the world. At the global level, our diagnostic explains a significant portion (more than 40%) of the inter-annual production variability. By quantifying the contribution of national yield anomalies to global fluctuations, we have found that just two concurrent yield anomalies affecting the larger producers of the world could be responsible for more than half of the global annual fluctuations. The relative importance of heat stress and drought in determining the yield anomalies depends on the region. Moreover, in contrast to common perception, water excess affects wheat production more than drought in several countries. We have also performed the same analysis at the subnational level for France, which is the largest wheat producer of the European Union, and home to a range of climatic zones. Large subnational variability of inter-annual wheat yield is mostly captured by the heat and water stress indicators, consistently with the country-level result.

Introduction

Wheat, with about a 2.1 million km² total harvested area, is the most abundant crop in the world: it is the first rain-fed crop after maize and the second irrigated crop after rice (Portmann *et al* 2010). With a total production that surpassed 700 million tons (MTons) in year 2010, it is contributing to about the 20% of the total dietary calories and proteins worldwide (Lobell and Gourdji 2012, Shiferaw *et al* 2013).

Field studies of wheat yield variability have shown that higher than optimal temperatures during the growing season are generally accelerating the progress of the plant phenological stages, affecting photosynthesis and its balance with respiration, and reducing the final yield (Lobell and Gourdji 2012, Rezaei *et al* 2015). Higher temperatures also increase the atmospheric demand for water and reduce the crop water-use efficiency (Ray *et al* 2002). Exposure to extremely

high temperatures (i.e. heat stress) leads to plant damages by inducing perturbations in cellular structures and metabolic processes (Nakamoto and Hiyama 1999). Isolated occurrences of extreme high temperatures around a sensitive stage of crop development, such as flowering and grain-filling, can reduce grain yield considerably (Tashiro and Wardlaw 1990, Ferris *et al* 1998, Porter and Gawith 1999, Luo 2011), while a prolonged period of extreme high temperatures might result in almost total yield loss (Semenov and Shewry 2011).

The detrimental effect of the heat stress on wheat yield may worsen when coinciding with drought (Pradhan *et al* 2012). Irrigation, to alleviate local water stress, can also mitigate the impact of the highest temperatures, by evaporative cooling, even at larger spatial scales (Tack *et al* 2015, Troy *et al* 2015, Mueller *et al* 2016). On the other hand, extreme amounts of precipitation and water excess in the soil can also be

responsible for wheat loss due to proliferation of pests and diseases, leakage of nutrients, inhibition of oxygen uptake by roots, and interference with agronomical practices (e.g. waterlogging during harvest). This may have happened during the 2016 cropping season in France, which showed a huge wheat yield anomaly that has not been officially confirmed yet (MARS 2016, Bloomberg 2016, France24 2016, Arvalis 2016).

Compared to other important crops, the main wheat producing regions are characterized by ‘close-to-average’ yield variability (Ben-Ari and Makowski 2014). However, increasing unfavorable conditions under observed and projected climate change conditions will probably impact wheat production variability (Gourdji *et al* 2013, Deryng *et al* 2014, Siebert and Ewert 2014, Asseng *et al* 2015, Ray *et al* 2015), especially because of the exacerbated effects of heat stress on grain number and grain size (Lobell *et al* 2015).

Using an ensemble of crop model simulations, Lobell *et al* (2015) estimated for the period 1980–2008 a 5.5% reduction of global wheat production due to changes in temperature and precipitation. This reduction occurred together with an increase in wheat yield/production of about 50% over the same period, mostly due to improved management and higher yielding crop varieties. Therefore, considering that the human population increased proportionally from 4.4 billion in 1980 to 6.9 billion in 2010 (UN 2015), negative yield deviations from the mean trend can be a potentially increasingly threat to food security. Indeed, changes in wheat yield variability have already been suggested as one of the primary factors influencing global food prices, market stability and food security, especially in developing countries (FAO 2011, OXFAM 2012, Porter *et al* 2014, Deryng *et al* 2014, GFS 2015). Moreover, the current urbanization trend at the expense of cropland area extent would imply, to be sustainable, a further increase of crop yield (Bren d’Amour *et al* 2016), and presumably an increased sensitivity of food security to the climate induced yield interannual variability.

The supporting evidence for these assertions is still relatively weak, since only few recent studies have addressed the relationships of the observed inter-annual climate and wheat yield fluctuations at the global scale. Lizumi and Ramankutty (2016) found consistent increases in standard deviations of annual yields and of an agro-climatic indicator accounting for solar radiation, heat stress and drought. Ray *et al* (2015), through an automatic spatial selection of several agro-climatic indicators based on temperature and precipitation anomalies (and using high resolution yield data), developed a statistical model accounting for the sign of the fluctuations and explaining about one third of the observed yield variance. This is in line with the composite analysis by Lesk *et al* (2016), based on an *a priori* knowledge of extreme events, showing that droughts and extreme

heat could have reduced national annual yields by 9%–10%. No effects of floods and extreme cold events were identified by Lesk *et al* (2016) on national data.

It is worth noting that none of aforementioned studies explicitly considers the possible impact on wheat yield of water excess events, which are not necessarily only related to floods. Moreover, the indicator used by Lizumi and Ramankutty (2016) accounts more for sub-optimal climatic conditions, rather than for the extremes. On the other hand, the indicators used by Ray *et al* (2015) are computed using the seasonal mean variables, which are not capturing the extremes occurring on a sub-monthly scale as the heat waves. Both of them are based on the exceedance of prescribed thresholds that are not explicitly mentioned. This might not be justified at the global scale, because of the different climates and different agronomical practices (such as irrigation and variety selection).

Here, we aim to model explicitly the effects of temperature and soil moisture anomalies (either positive and negative) on global and regional wheat production anomalies. The proposed approach does not require an *a priori* knowledge of extreme events (as in Lesk *et al* 2016) and it has the clear advantage of disentangling the different contributions of heat and water anomalies, potentially resulting in stress for the crop. Furthermore, it helps to better identify and characterize the role of the major producers in driving global yield anomalies, showing explicitly the effects of the individual climatic anomalies and extremes for each year in the period 1980–2010. Compared to Ray *et al* (2015) and Lizumi and Ramankutty (2016), we use the most up-to-date heat waves and water balance indicators currently available. These are, namely, the Heat Wave Magnitude Index daily (HWMId; Russo *et al* 2015) and the Standardized Precipitation Evaporation Index (SPEI; Vicente-Serrano *et al* 2010, 2013). Compared to previous studies, both indexes have been developed to be generally applicable under a wide range of conditions. They bear the great advantage of allowing the diagnosis and the comparison of extreme events in different regions of the world and time of the year on the basis of globally defined thresholds (Zampieri *et al* 2016, Vicente-Serrano *et al* 2010, 2013, see Data and Methods section).

In this study, the HWMId has been slightly modified in order to account for temperature anomalies and heat waves events during the stages of wheat growth that are sensitive to heat stress (i.e. three months before harvesting, roughly corresponding to the period including anthesis and grain filling). Therefore, the new indicator bears some analogy with the Growing Degree Day (DGG) index, but instead of summing the daily temperature anomaly w.r.t. a base value, it sums up the heat wave magnitude function defined by Russo *et al* 2015 (see equation (1) in the Data and Methods section). We refer to it as *Heat Magnitude Day* (HMD). While the HWMId selects the maximum value of the

summations computed on consecutive periods with maximum daily temperature over a certain threshold (i.e. a heat wave), the HMD integrates all of them similarly to the GDD. Therefore, the HMD is able to capture not only heat stress events, but also sub-optimal conditions.

The SPEI is a multi-temporal-scale index that quantifies persistent anomalies in the soil moisture balance (i.e. precipitation minus potential evaporation) over different time periods (1–24 months). We use the SPEI6 computed during the previous six months before harvesting, and we refer to it simply as SPEI. The SPEI6 provides an estimate of the soil moisture state during the sensitive stage of wheat, as for the heat stress, but also accounting for a preconditioning period of three months before flowering consistently with the time-scale of the root soil moisture dynamics (see e.g. Zampieri *et al* 2009). Large negative values of the SPEI indicate drought, while positive SPEI indicate wetter-than-normal soil conditions.

In order to account for concurrent heat and water stress events, we define a combined index as a simple linear superposition of the standardized HMD and SPEI (*Combined Stress Index*, CSI). We calibrate the CSI by determining the coefficients that maximize the CSI explanatory power with a bilinear ridge regression of the observed yield fluctuations at national level (FAOSTAT data), accounting for the covariance of the two agro-climatic indexes (see the Data and Methods section). Thus, the resulting multiplicative coefficients combining the HMD and the SPEI into the combined index depend on the country. The comparison between their absolute values can be interpreted as our best statistical estimate of the relative contribution of heat and water stresses on the national yield anomalies. Moreover, the sign of the parameter multiplying the SPEI indicates, for each country, larger yield sensitivity either to drought or to water excess.

We test the robustness of our proposed approach by performing a specific case study at the subnational level for France, where wheat is resulting to be more sensitive to over-wet conditions. This country is indeed characterized by pronounced heterogeneity of agro-climatic zones and by the availability of both high-quality yield data (from 1989 to 2014) and meteorological data (updated in near real-time). This allows us to discuss also the possible compensation effects while aggregating the indexes over large countries to compare with the FAOSTAT national yield statistics. Finally, we demonstrate the usefulness of the CSI to characterize the impact of climatic extremes on wheat yield anomaly observed during the 2016 wheat-cropping season in France.

Data and methods

Most physiological processes responsible for the final crop yield are negatively affected by maximum daily

temperatures above 30 °C (Porter *et al* 2014). Above 34 °C–35 °C, wheat reaches its limits for survival (Porter and Gawith 1999, Hatfield *et al* 2011, Rezaei *et al* 2015), the so-called ‘cardinal temperatures’ for wheat (Porter and Gawith 1999).

To provide realistic estimates of fixed based threshold estimators often used to characterize heat stress on wheat, biases in temperatures need to be corrected, especially when gridded temperature data and/or reanalysis are used instead of local field measurements. Furthermore, wheat physiology is sensitive to the temperature of its canopy, which might be higher than the 2 m temperature (the commonly available variable in the observational data sets) in unstably stratified surface layers, especially above arid soils (Siebert *et al* 2014). The thermal vertical structure of the atmospheric surface layer can also be altered by evaporative cooling due to irrigation (Tack *et al* 2015, Troy *et al* 2015, Mueller *et al* 2016). These issues, together with the lack of information about the varieties and the detailed agronomical practices that are implemented in the different regions, raise some concerns about the uniform global applicability of heat stress indicators based on the cardinal temperature thresholds used in previous studies (Lizumi and Ramankutty 2016, Ray *et al* 2015).

Here, we adopt indicators based on statistically determined climate thresholds aiming to account for the physiological effects of heat and soil moisture anomalies (and, in principle, other correlated factors) with respect to the local climate variability.

The heat analysis is inspired by the HWMId (Russo *et al* 2015, Zampieri *et al* 2016). A heat wave is defined as a consecutive period when T_{\max} exceeds the 90th percentile of T_{\max} , computed for all days of the year considering a 2 w time window (T_{90}). Over these periods, the HWMId is then defined as the sum of the difference between T_{\max} and the 25th percentile (T_{25}) normalized by the interquantile temperature range (i.e. the magnitude function M_d , equation (1)), computed in a similar manner to the 90th percentile.

$$M_d(T) = (T - T_{25}) / (T_{75} - T_{25}) \quad \text{if} \\ T > T_{90}; \text{ zero otherwise} \quad (1)$$

This non-parametric approach (i.e. without prior assumption on the statistical distribution of T_{\max}) takes both the length and the intensity of the heat waves into account. Whether a severe or a ‘close-to-normal’ event is diagnosed depends on the numerical value that the HWMId assumes. A great advantage is that the severity of the events can be defined on the basis of global thresholds, no more depending on the region (Russo *et al* 2014, 2015, Zampieri *et al* 2016).

This definition of the HWMId is the same described in Russo *et al* (2015). Here, we slightly modify it to diagnose heat stress and temperature anomalies that are relevant for crops. While the original formulation of the HWMId is based on the

maximum event over the whole year, we adapt it for wheat crop by cumulating the magnitude function over the last three months of the growing season before harvesting similarly to the way the Growing Degree Days (GDD) is defined. Therefore, the modified index accounts for all events occurring during the sensitive stage for the crop. We refer to this modified index as HMD. Figure S9 in the online supplementary information provides a visual representation of the HMD computation, with respect to the T_{90} and the interquartile temperature range computed during 2003 growing season in central France.

Daily T_{\max} data are retrieved from the AgMERRA climate forcing dataset (Ruane *et al* 2015) that was created to support the Agricultural Model Intercomparison and Improvement Project (AgMIP). The AgMERRA dataset is available at a $0.25^\circ \times 0.25^\circ$ resolution from 1980 to 2010 globally. This dataset is based on the NASA Modern-Era Retrospective Analysis for Research and Applications (MERRA, Rienecker *et al* 2011), blending models and observational data through a mathematical procedure called ‘data assimilation’, but it corrects biases in temperature using CRU (Climate Research Unit) data. CRU data are amongst the most widely used climatic datasets. They consist of a gridded datasets derived from quality checked station measurements, excluding time-series that do not pass several quality and homogeneity tests (see Osborn and Jones 2014, and references herein). CRU data are defined on a monthly time-scale, while the day-to-day variability is provided from the MERRA reanalysis. We evaluated the HMD considering also AgCFSR, i.e. the other main agricultural forcings dataset adopted in AgMIP (which is also correcting the bias using CRU data, Ruane *et al* 2015), finding almost equal results (see figure S7 of the supplementary material, available at stacks.iop.org/ERL/12/064008/mmedia).

The soil moisture analysis is tackled with the SPEI (Vicente-Serrano *et al* 2010, 2013). The SPEI is a multi-temporal-scale water balance index integrating the precipitation and potential evapotranspiration budget on the basis of CRU data, available at $0.5^\circ \times 0.5^\circ$ resolution from 1901 to 2014 globally (<http://spei.csic.es/>). It can be used for determining the onset, duration and magnitude of drought conditions with respect to normal conditions in a variety of natural and managed systems such as crops. SPEI anomalies are considered large, or extreme when specific thresholds (globally defined) are surpassed.

To compute the HMD and evaluate the SPEI over the wheat growing season and production areas, we use the global data set of monthly irrigated and rainfed crop areas around the year 2000 (MIRCA2000, Portmann *et al* 2010). MIRCA2000 integrates data from several different sources, with the specific aim to enhance the consistency with subnational statistics collected by countries and by the FAO. It provides both

irrigated and rainfed crop areas of 26 crop classes for each month of the year, and includes information on the occurrence of multiple cropping systems allowing to account for winter and spring sown wheat, or irrigated and rainfed wheat when they coexist in the same grid cell. Portmann *et al* (2010) also provide an estimate of the uncertainties of MIRCA2000 in comparison with other sources. MIRCA2000 is available at various resolutions. Here, we use the 0.5×0.5 degrees resolution, which is compatible with the meteorological data used in this study. MIRCA2000 is consistent with the only other global agricultural calendar data available to our knowledge, as it is mostly based on the same sources (Sacks *et al* 2010).

MIRCA2000 (online supplementary figure S1) provides sowing dates but not the varieties. The majority of the harvested areas correspond to rainfed winter sown wheat (118 million hectares), especially in the middle latitudes (in both hemispheres). Rainfed spring sown wheat (30 million hectares) is dominant mainly in the northwestern US and Canada, and it is present in other regions (Spain, England, Poland and India) as secondary crop. Irrigated winter sown wheat (50 million hectares) is found mainly in India and China, and it is characterized by the largest local concentrations. Irrigated spring sown wheat (8 million hectares) harvested areas are almost negligible with respect to the other categories. Online supplementary figures S2 and S3 show the sowing and harvesting months, respectively, derived from the MIRCA2000 dataset.

Wheat is mostly sensitive to climatic extremes during the three months period before harvesting (Lobell and Tebaldi 2014). We define these periods for each grid cell of the climatic data using MIRCA2000 (see figure S3). Accordingly, we compute the indexes separately for each of the four wheat cropping categories (i.e. irrigated and non-irrigated crops, and double cropping), since the sowing and (most importantly) the harvesting dates can be different. In the grid boxes where different categories coexist, we average the values of the indexes using the harvested area of each wheat category as a weighting factor in order to obtain one value of each index for each year and each grid cell.

The HMD is computed for period 1980–2010 by summing equation (1) over all relevant events during the three months before harvesting. The SPEI, which integrates the precipitation and evaporation anomalies occurred earlier, is evaluated at the harvesting month. The time scale (i.e. the number of months) defining it has to be specified. Assuming that the crop is sensitive to soil moisture anomalies in the same period as for heat, but also accounting for the effects of the previous precipitation and evaporation anomalies in determining the current soil moisture values, we used SPEI6. In this way, we account for the water balance anomalies during the three months sensitive period before harvesting and also during an earlier preconditioning

period of preceding three months, accordingly with the typical time scales of soil moisture dynamics (see e.g. Zampieri *et al* 2009).

Online supplementary figures S4 and S5 show the daily maximum temperature 90th percentile (T_{90}), used as a threshold in the heat stress estimation, corresponding to the beginning and to the end of the three months period from harvesting. Notably, there are regions where the T_{90} is above 30 °C already at the beginning of the growing season, especially for irrigated winter wheat in India, and for rainfed spring wheat (figure S4). On the other hand, there are regions as Canada, for instance, where T_{90} is relatively low (this is confirmed by Jarvis *et al* 2008). T_{90} is above 30 °C in most regions at harvesting time (figure S5).

Global, national and subnational spatial aggregation are computed by averaging grid-point data, using again the MIRCA2000 harvested area data as weighting factors, so that the indexes can be compared with FAOSTAT3 national statistics of wheat production and yield (www.faostat3.fao.org/). The quality of yield data represents a main issue in this kind of studies, especially for countries where there are no alternative estimations available (Desiere *et al* 2016). The subnational analysis for France is based on winter wheat yields from 92 French départements, provided by AGRESTE Ministère de l'Agriculture (AGRESTE, 2015) and covering the period between 1989 and 2014. Subnational crop yields have been quality controlled (Ceglar *et al* 2016). Weather data for the computation of the CSI has been obtained from the high-resolution EOBS gridded weather dataset over Europe (Haylock *et al* 2008).

Since most of the tendencies affecting yield statistics are mostly due to improvement in the agricultural practices, we adopt a non-linear detrending procedure, i.e. the locally weighted scatterplot smoothing (LOESS, Cleveland and Devlin 1988) with the 'lambda' (span) parameter equal to 0.75. This value was optimized by trial-and-error procedure. However, the linear correlation coefficient we obtain are only weakly sensitive to changes of the span parameter around this value. In fact, the final results are robust to changes of the lambda parameter within a reasonable range of values. Moreover, this technique yields comparable results with respect to other analog procedures (see e.g. Ben-Ari *et al* 2016). This processing has been conducted with the R software build-in loess function.

Having removed in this way also the part of the signal that is potentially related to climate, we are compelled to apply the same procedure also to the HMD and SPEI time-series. This is performed only when the time-series exhibit either a significant trend (with the Mann–Kendall test), or a change point (Bai and Perron 1998), using 'Kendall' and 'strucchange' R libraries; otherwise, we just remove the mean. In such a way, we can isolate the effects of climate anomalies and extremes on the year-to-year wheat production

variability. Online supplementary figure S8 shows an example of LOESS detrending (and mean removal) of yield and of the indexes time-series.

Our analysis is based on the anomalies obtained after such procedure. The CSI is then defined as a simple linear combination of the standardized HMD and SPEI (i.e. SPEI6) anomalies as expressed by equation (2):

$$CSI = a \cdot HMD_{\text{detrend, std}} + b \cdot SPEI_{\text{detrend, std}} \quad (2)$$


where the subscripts indicate the fact that the time-series are detrended and standardized. Standardization of the SPEI is advisable for using it in period 1980–2010, consistently with the reference period used to compute the HMD.

In equation (2), the a and b parameters are estimated by ridge linear regression with the country level FAOSTAT data (again after detrending, standardization and inversion [i.e. multiplication by -1]), accounting for the co-linearities between heat stress and drought. Therefore, the CSI is an estimate of the standardized yield anomalies. The numerical values (and sign) of the a and b parameters depend on the country. The sign of a is always resulting positive, as the CSI is defined such that positive values are corresponding to negative yield anomalies, which is the reason of the inversion of yield data. On the other hand, the sign of the b parameter indicates if the national yield is more affected by negative or positive soil moisture anomalies. Since all time-series are standardized, the comparison between the values of the parameters provide a qualitative indication of the relative importance of the two regressors in determining the yield anomalies. A quantitative indication could be provided if no co-linearities between the two regressors were present. In any case, the value of the a parameter provides an estimate of the effects of heat stress alone in determining the yield anomalies (i.e. by imposing no soil moisture anomaly in equation (2)). We refer to the a parameter as the heat sensitivity parameter, and to the b parameter as to the soil moisture sensitivity parameter.

Linear correlations between the individual regressors and national yield data are computed as well, for comparison with the CSI skill and to test the robustness and consistency of the procedure. In all three cases, Pearson linear correlations and tests (10%, two-sided) are applied in order to compute the statistical significance of the correlations coefficients.

Results and discussion

Global wheat production approximately doubled from 1970 to 2015 (figure 1(a)), and increased by about 50% in the period 1980–2010 (the one here considered for the climatic analysis). The largest contributor to the global inter-annual variability is the Russian



rainfed winter wheat production (see online supplementary figure S1), characterized by a standard deviation of the production anomalies of about 12 MTons corresponding to the 25% of the mean national production of 48 MTons per year (2000–2015 average).

Wheat production interannual variabilities of EU (rainfed, mainly winter wheat), China (mainly rainfed and irrigated winter wheat) and USA (rainfed winter wheat) are following in decreasing order. Standard deviations consist of about 7 MTons that correspond to the 5%, 6% and 11% of the averaged national productions of 140, 107 and 57 MTons, respectively. Ukraine (rainfed winter wheat), Australia (rainfed winter wheat), Canada (rainfed spring wheat) and India (all agro-management types) are characterized by decreasing standard deviations (between 5 and 3 MTons) corresponding to the 27%, 19%, 14% and 4% of the averaged productions of 22, 24, 25 and 80 MTons, respectively.

These eight countries and regional aggregations together produced, on average, 495 MTons of wheat, corresponding to the 77% of the global amount of 640 MTons. The associated standard deviation almost equals the global value of 22 MTons, which is 3.4% of the total production (and the 4.4% of the sum of the 8 producers). Quantitative analysis of the fluctuations attributable to the individual regions displayed in figure 1(a) reveals that concurrent anomalies of two of these producers are able to explain more than 50% of the global inter-annual fluctuations in almost all years

considered, if no compensating anomalies of the opposite sign are present.

Figure 1(a) also shows the occurrence of climatic extremes in the wheat cropping regions, as represented by the CSI. During the period 1980–2010, significant negative production departures from the mean global tendency are often captured by the CSI, especially in the years 1987–1988, 1994, 2003, 2006–2007 and 2010, coinciding with the observed spikes in cereal prices (see figures 7-3 in IPCC-WG2). We observed a significant level of consistency also in the years characterized by positive production anomalies and less frequent damaging climatic extreme events, like in 1986, 1992–1993, 1997, 2004–2005 and 2008–2009. The abundant wheat production in 1990, being characterized by moderate climate conditions, represents an outsider in our analysis. That was the last year of intensive fertilization in Russia (Lioubimtseva *et al* 2015), which is the main contribution of the recorded production anomaly.

Overall, most of the largest anomalies that could be related to climate variability are induced by the major producers, while smaller and/or compensating positive and negative anomalies characterize the ‘normal’ years. This is partly expected because the single CSI anomalies are weighted by the national productions in figure 1(a). However, the ranking of the anomalies contributions by region slightly differs from the one of production: the largest contributor to the variance of the CSI is Europe, followed by the US, China, Russia and India.




Figure 2. Left panels: as left panel of figure 1, but for (a) the Heat Magnitude Day (HMD, see equations (1) and (2) in Data and Methods) and (b) the Standardized Precipitation Evaporations Index (SPEI, see Data and Methods). Right panels: linear correlations between (c) national yield data and HMD (negative values indicate the detrimental effect of heat on wheat, ‘HOT’), (d) national yield data and SPEI (positive values indicate larger detrimental effect of drought, while ‘DRY’; negative values indicate larger effect of water excess, ‘WET’), and (e) HMD and SPEI (negative values indicate positive correlations between the occurrence of hot and dry anomalies).

This difference could be a consequence of the spatially variable performance of the CSI in capturing the stress events for wheat, or to the different relative importance of climate compared to other factors in determining the production anomalies in the different regions, or it could be an effect of the spatial aggregations in large heterogeneous regions. Higher resolution yield and production data at the global scale, together with detailed agronomical information, would very likely improve the general understanding and ease the modeling of the yield anomalies.

The overall performance of the CSI is quantified by the linear correlation coefficients with yield anomalies, plotted in figure 1(b). We estimated significant linear correlation coefficients for most countries. Among the largest producers, US, China, India and Argentina display smaller but statistically significant values. As for the US, this result is likely to be caused by the mitigation effect of large-scale irrigation that tends to decouple yield anomalies from climate variability also in regions where it is not applied directly (Tack *et al* 2015, Troy *et al* 2015, Mueller *et al* 2016).

The linear correlation between the total production and the global averaged CSI computed for all wheat cropping areas is -0.65 , which corresponds to the 42% of the total variability, strengthening the result of Ray *et al* (2015).


The effects of heat stress are comparable or larger than the effects of water stress in most countries (see figure 1(c)). Water stress effects tend to be more substantial in the Mediterranean region. Negative WSI can be associated to drought and related to negative yield anomalies, especially in arid and semi-arid

regions. In the Tropics and the middle/high latitudes, our statistical model suggests an opposite relation with positive WSI being associated to negative yield anomalies (see figure 1, bottom-right panel). This is partly consistent with the findings of Vicente-Serrano *et al* (2013) who showed a reduced sensitivity of natural vegetation dynamics to drought in regions characterized by climatological water excess. Our study suggests that wheat may be, on average, more sensitive to water excess extreme events than to drought in these regions, and that dryer-than-normal conditions could have, on the other hand, beneficial effects.

Figure 2 shows the spatial and temporal characteristics of the individual regressors forming the CSI, as well as their direct relationships with the FAOSTAT yield data.

At the global level, heat stress alone (HMD, figure 2(a)) has an explanatory power similar to the CSI, and displays a larger trend. Linear correlation between total wheat production and globally averaged HMD anomalies is -0.57 , slightly lower than the value of -0.65 obtained for the CSI. HMD has the advantage, w.r.t. to the CSI, that it does not need calibration with yields data. However, including the SPEI integrating the water balance in the six months prior to harvesting (i.e. SPEI6, see Data and Methods) is necessary to significantly explain the production variability in India, China and France, and to improve the correlation coefficients in several other countries (compare figure 1(b) and figure 2(c)).

Given the dual role of the SPEI in the CSI (figure 1 (d)), the globally averaged SPEI (figure 2(b)) is not



significantly correlated with the global wheat production anomalies. With respect to the HMD, SPEI6 time-series are less often significantly correlated with the national productions (compare figure 2(c) and figure 2(d)). In most of the countries the sign of the correlation is positive, i.e. negative cumulated water balance is associated with negative yield anomalies. This is correctly accounted for in the combined index. The CSI is always outperforming the individual indexes such that it can be significantly correlated with the national yield anomalies even if the HMD and/or SPEI are not.

It is worth noting that heat and water anomalies are significantly linearly correlated in several countries (figure 2(e)), but there are many regions where heat and soil moisture anomalies and extreme events seem to be more independent. While several authors suggested a tight linkage between these two types of extreme events in water limited regions (see e.g. Zampieri *et al* 2009, Seneviratne *et al* 2010), our analysis suggests that this is not often the case for most of the wheat producing regions that are found to be more sensitive to water excess than to drought (figure 1(d)).

France, with about 37 MTons of rainfed winter wheat produced per year, is the largest producer of the European Union (25%), and consequently it is responsible for a corresponding fraction of the inter-annual standard deviation. Therefore, EU and France production variation are significantly correlated, with a linear correlation coefficient of 0.49. However, European climate is characterized by pronounced spatial gradients and prominent production anomalies

of the individual countries (see figures 1(b)–(d)) that can compensate at the European level. This compensation effect has been observed to moderate relatively large anomalies preferably in the first part of the time series, in particular in 1980, 1985, 1986, 1988. It appears to be reduced towards the end of the period. Consistently, the linear correlation of the French and EU productions computed until the end of the 1980s is reduced to 0.18. This implies that there is a current tendency towards more coherent anomalies, and, therefore, towards an increased threat to food security. This shifting behavior might also be manifested by other large countries and spatial aggregations. Indeed, it is consistent with the emerging global scale heat waves pattern observed recently (Zampieri *et al* 2016).

According to our analysis at the national scale, the linkage between climate and yield variability in France is controversial. Although there is a significant correlation between heat waves and drought (see figure 2(e)), the heat and water stress indexes are not directly correlated with the production anomalies and the French production is more negatively affected by water excess than drought. This motivated us to conduct a deeper analysis for France.

Figure 3 represents the main results of the analysis conducted for France at the subnational level. Yield data are available from 1989 to 2014, while the meteorological data are updated in near real-time, allowing the evaluation of the indexes for more recent years.

The CSI is able to explain yield variations significantly in a considerable part of the country

(figure 3(a)), with a linear correlation coefficient of about -0.5 , especially in the central-northern part where the main wheat producing areas are located. In western France, facing the Atlantic, no significant correlations are found. The analysis conducted for heat (HMD) and water (SPEI) indexes alone (figures 3(b) and (c)) shows interesting spatial variability. The central part of France is significantly affected by heat waves. Mediterranean France is more sensitive to drought while the northern part results to be more sensitive to water excess (consistently with Ceglar *et al* 2016). The CSI computed at the subnational level indicates higher sensitivity to water excess than to drought (i.e. negative ' b ' parameters, see Data and Methods) in the northern and central parts of the country, where most of the wheat is grown. This is consistent with the results obtained at country level.

We have used the CSI as a diagnostic tool to assess the impact of weather events on French wheat yield in 2016, characterized by an extremely negative anomaly. The indexes computed for 2016 (figure 3(e) and (f)) are diagnosing an excess of water in central, northern and eastern parts of France (mainly due to heavy precipitation in May), while drought occurred in the southern part of the country. In addition, a moderate heat wave influenced the western and southwestern regions (towards the end of the growing season). Therefore, the combined index computed for 2016 would have indeed suggested a negative yield anomaly (figure 3(d)).

By using the CSI to estimate the production anomaly at country level in 2016, we obtain a reduction of about 1 MTON lower than the average, corresponding to the 40% of the national production standard deviation. Therefore, the CSI is probably underestimating the actual anomaly. However, it suggests that the adverse weather conditions responsible for the water excess (possibly around the wheat flowering stage) might have played a central role.

It is worth noting, however, that the CSI does not capture the single precipitation events, nor their timing, but it rather accounts for their cumulative effect on the soil water balance. Therefore, while the CSI would have predicted a negative yield anomaly, the precise causes of the 2016 French yield loss still have to be determined.

Conclusions

We have analyzed the historical global wheat production annual anomalies computed from the FAOSTAT data set from 1980 to 2010. During this period, few of the major producing regions were responsible for significant anomalies at the global level. Concurrent anomalies of two of the major producers suffice to explain more than 50% of the global inter-annual fluctuations in almost all years, if no significant

compensating anomalies were present. Russia is the largest contributor to the global annual variability.

The combined stress index (CSI), defined as a linear superposition of the Standardized Precipitation Evaporation Index (SPEI, Vicente-Serrano *et al* 2010) and of the Heat Magnitude Day (HMD, this study), allows explaining the 42% of the variability globally (confirming and strengthening the result of Ray *et al* 2015), and often at country level as well. Therefore, accurate yield forecasts in the major producing countries could provide useful insights in global production, potentially increasing food security. In fact, we are currently working on the application of the CSI in a seasonal forecasts context (Ceglar *et al* submitted).

Our result points to a clear role of climate anomalies and extreme events on wheat yield anomalies and enables identifying the relevant contributors and the associated effects. Heat stress is often the most important predictor, consistently with field studies and future projections, and it is in general as important as drought. As a prominent exception, we have found that in the Mediterranean countries drought carries a larger detrimental effect on wheat yield than heat stress.

Heat stress over wheat cropping regions increased significantly in the period 1980–2010, especially since the mid-1990s. This produced less compensating and more concurrent yield anomalies, motivating the general concern about food security.

Our results also point to the sensitivity of wheat yield to the water excess issue, rather than to drought, especially in tropical regions and in some regions of the mid/high latitudes. While heat stress globally remains the most important factor determining the yield anomalies (compare figure 2(c) and figure 2(d)), our findings show that water stress, and in particular water excess, is essential to explain yield anomalies of important wheat producers such as China and India.

The results obtained at the subnational level for France (the main European wheat producer) support and strengthen our main findings. The combined indicator was, indeed, able to capture sub-regional yield anomalies, and provided consistent results with the national scale analysis. This case study also showed the importance of accounting for water excess to gain some predictive skill for extreme events such as the large wheat production reduction occurred in France of 2016.

Acknowledgments

We are indebted with Maurits van den Berg for his help during the revision process and two anonymous Reviewers that greatly contributed to the improvement of the quality and clearness of the message we are conveying.

References

- Arvalis 2016 (www.arvalis-infos.fr/view-22401-arvarticle.html?region=&culture=3,4,6,10,13,39,42,44,45,48,52,54,57,61,67,73,78,79,83,84,87)
- Asseng S *et al* 2015 Rising temperatures reduce global wheat production *Nat. Clim. Change* **5** 143–7
- Bai J and Perron P 1998 Estimating and testing linear models with multiple structural changes *Econometrica* **66** 47–8
- Ben-Ari T and Makowski D 2014 Decomposing global crop yield variability *Environ. Res. Lett.* **9** 114011
- Ben-Ari T, Adrian J, Klein T, Calanca P, Van der Velde M and Makowski D 2016 Identifying indicators for extreme wheat and maize yield losses *Agric. Forest Meteorol.* **220** 130–40
- Bloomberg 2016 (www.bloomberg.com/news/articles/2016-08-01/france-to-harvest-least-wheat-in-28-year-s-after-rains-oda-says)
- Bren d'Amour C, Reitsma F, Baiocchi G, Barthel S, Güneralp B, Erb K-H, Haberl H, Creutzig F and Seto K C 2016 Future urban land expansion and implications for global croplands *Proc. Natl Acad. Sci.* **201606036**
- Ceglar A, Toreti A, Lecerf R, van der Velde M and Dentener F 2016 Impact of meteorological drivers on regional inter-annual crop yield variability in France *Agr. Forest Meteorol.* **216** 58–67
- Ceglar A, Prodhomme C, Zampieri M, Turco M, Toreti A and Doblas-Reyes F From seasonal climate to crop yield forecast: importance of realistic soil moisture initialization *Geophys. Res. Lett.* submitted
- Cleveland W S and Devlin S J 1988 Locally weighted regression: an approach to regression analysis by local fitting *J. Am. Stat. Assoc.* **83** 596–610
- Deryng D, Conway D, Ramankutty N, Price J and Warren R 2014 Global crop yield response to extreme heat stress under multiple climate change futures *Environ. Res. Lett.* **9** 034011
- Desiere S, Staelens L and D'Haese M 2016 When the data source writes the conclusion: evaluating agricultural policies *J. Dev. Stud.* **52** 1372–87
- FAO 2011 *The State of Food Insecurity in The World: How Does International Price Volatility Affect Domestic Economies and Food Security* (Rome: Food and Agriculture Organization of the United Nations) p 52
- Ferris R, Ellis R, Wheeler T and Hadley P 1998 Effect of high temperature stress at anthesis on grain yield and biomass of field-grown crops of wheat *Ann. Bot.* **82** 631–9
- Foley J A *et al* 2011 Solutions for a cultivated planet *Nature* **478** 337–342
- France24 2016 (www.france24.com/en/20160809-french-wheat-harvest-catastrophic-30-year-low-economy-agriculture)
- Global Food Security 2015 Extreme Weather and Global Food System Resilience (www.foodsecurity.ac.uk/assets/pdfs/extreme-weather-resilience-of-global-food-system.pdf)
- Gourdji S M, Sibley A M and Lobell D B 2013 Global crop exposure to critical high temperatures in the reproductive period: historical trends and future projections *Environ. Res. Lett.* **8** 024041
- Haylock M R, Hofstra N, Klein Tank A M G, Klok E J, Jones P D and New M 2008 A European daily high-resolution gridded dataset of surface temperature and precipitation *J. Geophys. Res.* **113** D20119
- Hatfield J L, Boote K J, Kimball B A, Ziska L H, Izaurre R C, Ort D, Thomson A M and Wolfe D 2011 Climate impacts on agriculture: implications for crop production *Agron. J.* **103** 351–70
- Jarvis C K, Sapirstein H D, Bullock P R, Naeem H A, Angadi S V and Hussain A 2008 Models of growing season weather impacts on bread-making quality of spring wheat from producer fields in western Canada *J. Sci. Food Agric.* **88** 2357–70
- Lesk C, Rowhani P and Ramankutty N 2016 Influence of extreme weather disasters on global crop production *Nature* **529** 84–7
- Lioubimtseva E, Dronin N and Kirilenko A 2015 Grain production trends in the Russian Federation, Ukraine and Kazakhstan in the context of climate change and international trade *Climate Change and Food Systems: Global Assessments and Implications for Food Security and Trade* ed Aziz Elbehri (Rome: Food Agriculture Organization of the United Nations (FAO)) (www.fao.org/3/a-i4332e/i4332e07.pdf)
- Lizumi T and Ramankutty N 2016 Changes in yield variability of major crops for 1981–2010 explained by climate change *Environ. Res. Lett.* **11** 034003
- Lobell D B and Gourdji S M 2012 The influence of climate change on global crop productivity *Plant Physiol.* **160** 1686–97
- Lobell D B and Tebaldi C 2014 Getting caught with our plants down: the risks of a global crop yield slowdown from climate trends in the next two decades *Environ. Res. Lett.* **9** 074003
- Lobell D B, Hammer G L, Chenu K, Zheng B, Mclean G and Chapman S C 2015 The shifting influence of drought and heat stress for crops in northeast Australia *Glob. Change Biol.* **21** 4115–27
- Luo Q 2011 Temperature thresholds and crop production: a review *Clim. Change* **109** 583–98
- MARS 2016 Crop monitoring in Europe: September 2016 *JRC MARS Bulletin* **24** 1–38
- Mueller D N, Butler E E, McKinnon K A, Rhines A, Tingley M, Holbrook N M and Huybers P 2016 Cooling of US Midwest summer temperature extremes from cropland intensification *Nat. Clim. Change* **6** 317–22
- Nakamoto H and Hiyama T 1999 Heat shock proteins and temperature stress *Handbook of Plant and Crop Stress* ed M Pessarakli (New York: Marcel Dekker) pp 399–416
- Osborn T J and Jones P D 2014 The CRUTEM4 land-surface air temperature data set: construction, previous versions and dissemination via Google Earth *Earth System Science Data* **6** 61–9
- OXFAM 2012 Extreme weather, extreme prices, the cost of feeding the warming World (www.oxfam.org/sites/www.oxfam.org/files/20120905-ib-extreme-weather-extreme-prices-en.pdf)
- Porter J R and Gawith M 1999 Temperatures and the growth and development of wheat: a review *Eur. J. Agron.* **10** 23–36
- Porter J R, Xie L, Challinor A J, Cochrane K, Howden S M, Iqbal M M, Lobell D B and Travasso M I 2014 Food security and food production systems *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed C B Field *et al* (Cambridge and New York: Cambridge University Press) pp 485–533
- Portmann F T, Siebert S and Döll P 2010 MIRCA2000—Global monthly irrigated and rainfed crop areas around the year 2000: a new high-resolution data set for agricultural and hydrological modeling *Glob. Biogeochem. Cycles* **24** GB1011
- Pradhan G P, Prasad P V V, Fritz A K, Kirkham M B and Gill B S 2012 Effects of drought and high temperature stress on synthetic hexaploid wheat *Funct. Plant Biol.* **39** 190–8
- Ray J D, Gesch R W, Sinclair T R and Hartwell Allen L 2002 The effect of vapor pressure deficit on maize transpiration response to a drying soil *Plant Soil* **239** 113–21
- Ray D K *et al* 2015 Climate variation explains a third of global crop yield variability *Nat. Commun.* **6** 5989
- Rezaei E E, Webber H, Gaiser T, Naab J and Ewert F 2015 Heat stress in cereals: mechanisms and modelling *Eur. J. Agron.* **64** 98–113
- Rienecker M M *et al* 2011 MERRA: NASA's modern-era retrospective analysis for research and applications *J. Clim.* **24** 3624–48
- Ruane A C, Goldberg R and Chrysanthacopoulos J 2015 AgMIP climate forcing datasets for agricultural modeling: merged products for gap-filling and historical climate series estimation *Agr. Forest Meteorol.* **200** 233–48

- Russo A, Sillmann J and Fischer E M 2015 Top ten European heatwaves since 1950 and their occurrence in the coming decades *Environ. Res. Lett.* **10** 124003
- Sacks W J, Deryng D, Foley J A and Ramankutty N 2010 Crop planting dates: an analysis of global patterns *Glob. Ecol. Biogeogr.* **19** 607–20
- Semenov M and Shewry P R 2011 Modelling predicts that heat stress, not drought, will increase vulnerability of wheat in Europe *Nat. Sci. Rep.* **1** 66
- Seneviratne S I, Corti T, Davin E L, Hirschi M, Jaeger E B, Lehner I, Orlowsky B and Teuling A J 2010 Investigating soil moisture–climate interactions in a changing climate: a review *Earth-Sci. Rev.* **99** 125–61
- Shiferaw B *et al* 2013 Crops that feed the world 10. Past successes and future challenges to the role played by wheat in global food security *Food Sec.* **5** 291
- Siebert S, Ewert F, Rezaei E E, Kage H and Graß R 2014 Impact of heat stress on crop yield—on the importance of considering canopy temperature *Environ. Res. Lett.* **9** 044012
- Siebert S and Ewert F 2014 Future crop production threatened by extreme heat *Environ. Res. Lett.* **9** 041001
- Tack Jesse, Barkley Andrew and Lanier Nalley Lawton 2015 Effect of warming temperatures on US wheat yields *Proc. Natl Acad. Sci.* **112** 6931–6
- Tashiro T and Wardlaw I F 1990 The response to high temperature shock and humidity changes prior to and during the early stages of grain development in wheat *Aust. J. Plant Physiol.* **17** 551–61
- Troy T J, Kipgen C and Pal I 2015 The impact of climate extremes and irrigation on US crop yields *Environ. Res. Lett.* **10** 054013
- United Nations, Department of Economic and Social Affairs, Population Division 2015 World Population Prospects: The 2015 Revision
- Vicente-Serrano S M, Beguería S and López-Moreno J I 2010 A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index *J. Clim.* **23** 1696–718
- Vicente-Serrano S M *et al* 2013 Response of vegetation to drought time-scales across global land biomes *Proc. Natl Acad. Sci. USA* **110** 52–7
- Zampieri M, D’Andrea F, Vautard R, Ciais P, De Noblet-Ducoudré N and Yiou P 2009 Hot European summers and the role of soil moisture in the propagation of mediterranean drought *J. Clim.* **22** 4747–58
- Zampieri M, Russo S, di Sabatino S, Michetti M, Scoccimarro E and Gualdi S 2016 Global assessment of heat wave magnitudes from 1901 to 2010 and implications for the river discharge of the Alps *Sci. Total Environ.* **571** 1330–9