

LETTER • OPEN ACCESS

Changing population dynamics and uneven temperature emergence combine to exacerbate regional exposure to heat extremes under 1.5 °C and 2 °C of warming

To cite this article: Luke J Harrington and Friederike E L Otto 2018 *Environ. Res. Lett.* **13** 034011

View the [article online](#) for updates and enhancements.

You may also like

- [Regional compound humidity-heat extremes in the mid-lower reaches of the Yangtze River: a dynamical systems perspective](#)
Yixuan Guo, Yu Huang and Zuntao Fu
- [Interacting implications of climate change, population dynamics, and urban heat mitigation for future exposure to heat extremes](#)
P Vahmani, Andrew D Jones and Christina M Patricola
- [Reduced impacts of heat extremes from limiting global warming to under 1.5 °C or 2 °C over Mediterranean regions](#)
Hao Yu, Di Wu, Xiuting Piao et al.



The Breath Biopsy® Guide
Fourth edition

FREE

DOWNLOAD THE FREE E-BOOK

BREATH BIOPSY

OWLSTONE MEDICAL

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED
31 October 2017

REVISED
24 January 2018

ACCEPTED FOR PUBLICATION
25 January 2018

PUBLISHED
21 February 2018

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 3.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the
title of the work, journal
citation and DOI.



Changing population dynamics and uneven temperature emergence combine to exacerbate regional exposure to heat extremes under 1.5 °C and 2 °C of warming

Luke J Harrington^{1,2} and Friederike E L Otto¹

¹ Environmental Change Institute, University of Oxford, South Parks Road, Oxford, OX1 3QY, United Kingdom

² Author to whom any correspondence should be addressed.

E-mail: luke.harrington@ouce.ox.ac.uk

Keywords: climate change emergence, heat extremes, population exposure, paris agreement

Supplementary material for this article is available [online](#)

Abstract

Understanding how continuing increases in global mean temperature will exacerbate societal exposure to extreme weather events is a question of profound importance. However, determining population exposure to the impacts of heat extremes at 1.5 °C and 2 °C of global mean warming requires not only (1) a robust understanding of the physical climate system response, but also consideration of (2) projected changes to overall population size, as well as (3) changes to where people will live in the future. This analysis introduces a new framework, adapted from studies of probabilistic event attribution, to disentangle the relative importance of regional climate emergence and changing population dynamics in the exposure to future heat extremes across multiple densely populated regions in Southern Asia and Eastern Africa (SAEA). Our results reveal that, when population is kept at 2015 levels, exposure to heat considered severe in the present decade across SAEA will increase by a factor of 4.1 (2.4–9.6) and 15.8 (5.0–135) under a 1.5°- and 2.0°-warmer world, respectively. Furthermore, projected population changes by the end of the century under an SSP1 and SSP2 scenario can further exacerbate these changes by a factor of 1.2 (1.0–1.3) and 1.5 (1.3–1.7), respectively. However, a large fraction of this additional risk increase is not related to absolute increases in population, but instead attributed to changes in which regions exhibit continued population growth into the future. Further, this added impact of population redistribution will be twice as significant after 2.0 °C of warming, relative to stabilisation at 1.5 °C, due to the non-linearity of increases in heat exposure. Irrespective of the population scenario considered, continued African population expansion will place more people in locations where emergent changes to future heat extremes are exceptionally severe.

1. Introduction

There is significant societal interest in understanding how changes to climate extremes will proliferate in a warming climate (Seneviratne *et al* 2016, Stott 2016, Stott *et al* 2016). Following the successful signing of the Paris Agreement in December 2015 (Schleussner *et al* 2016b, Rogelj and Knutti 2016), a targeted focus has emerged within the scientific community to better understand how changes to the global climate system will evolve in response to specific thresholds of future global mean warming, such as 1.5 °C or 2 °C above

‘pre-industrial levels’ (Mitchell *et al* 2016b, Hawkins *et al* 2017, King *et al* 2017, Kraaijenbrink *et al* 2017).

Multiple recent studies have evaluated the changes to climate extremes expected to occur in response to limiting global mean warming to 1.5 °C and 2 °C, with several different modelling frameworks and complementary methodologies being used to provide these estimates (Fischer and Knutti 2015, Schleussner *et al* 2016a, Ciavarella *et al* 2017, Sanderson *et al* 2017, Schleussner *et al* 2017, Lewis *et al* 2017, Henley and King 2017, Mitchell *et al* 2017, King *et al* 2017, King and Karoly 2017, Perkins-Kirkpatrick and

Gibson 2017, Russo *et al* 2017). This variety in analytical approaches is particularly important, recognising estimates of future risk may differ depending on potentially subjective framing choices (James *et al* 2017). These include, but are not limited to: (1) the treatment of ‘pre-industrial’ baseline periods (Hawkins *et al* 2017, Schurer *et al* 2017); (2) what timescales are used to measure 1.5 °C and 2 °C (Henley and King 2017, Rogelj *et al* 2017); (3) whether or not temporary exceedances in a given temperature target are permitted before stabilisation (Knutti *et al* 2016, Rogelj and Knutti 2016, Schleussner *et al* 2016b); and (4) the specific combination of forcing pathways selected, recognising that different combinations of greenhouse gas and aerosol emissions, for example, can produce stabilisation at the same level of global mean warming (Millar *et al* 2017, Wang *et al* 2017). The analysis presented here only utilises a modelling framework specifically designed to understand changes to high-impact extreme weather events under 1.5 °C and 2 °C of warming (Mitchell *et al* 2016b, 2017), so should therefore be interpreted in the context of other studies using different but complementary methods of analysis.

Recognising that more-frequent and more-intense climate extremes will often lead to substantially higher impacts when they occur in highly populated areas, recent research has focused on the implications of future changes to heat extremes for specific regions of interest (Im *et al* 2017, Mishra *et al* 2017, Russo *et al* 2017, Liu *et al* 2017, Jones *et al* 2015, Pal and Eltahir 2016). However, multiple lines of evidence also demonstrate that different regions of the world can experience substantially different rates of emergent climate change given the same amount of global mean warming (Mahlstein *et al* 2011, Diffenbaugh and Scherer 2011, Hawkins and Sutton 2012, Harrington *et al* 2016, Diffenbaugh and Charland 2016, Davis and Diffenbaugh 2016, Herold *et al* 2017, Frame *et al* 2017, Harrington *et al* 2017, Mora *et al* 2017). Therefore, consideration must be given as to how changes to *where* people live, as well as changes to overall population size, will influence collective exposure to future heat extremes. By combining very large ensembles of high-resolution climate model simulations with spatially explicit projections of future population change (Jones and O'Neill 2016), this study will quantify the relative contributions of (1) emergent temperature change, (2) projected changes in total population size and (3) changes to where people live in exacerbating exposure to future heat extremes.

2. Data and methods

2.1. Model framework

This analysis employs atmosphere-only model simulations using an experimental framework developed under the ‘Half a degree Additional warming,


Prognosis and Projected Impacts’ (HAPPI) project (Mitchell *et al* 2016b). The HAPPI model framework prescribes sea surface temperatures and other boundary conditions (sea ice, greenhouse gases, aerosols) that would be consistent with the current climate, as well as for future worlds under 1.5 °C and 2 °C of global mean warming (see Mitchell *et al* (2017) for further details of the experimental setup). We apply these boundary conditions, via the Weather@Home (W@H) distributed computing framework, to produce thousands of atmosphere-only regional climate model simulations over a spatial domain encompassing a large region of Asia, the Middle East and Eastern Africa (hereafter SAEA region, figure 1). The W@H project utilises the spare computing power of volunteers to run thousands of simulations of the global atmosphere-only model HadAM3P (horizontal resolution of $1.875^\circ \times 1.25^\circ$), with a nested regional model (HadRM3P) operating over SAEA at a horizontal resolution of approximately 50 kilometres (see Guillod *et al* (2017) for further details of the W@H modelling framework). This combination of very large model ensembles (between 1500 and 2500 model years per experiment, see table S1) with high-resolution climate data provides a unique opportunity to accurately characterise population exposure to exceptionally severe heat events under future warming.

2.2. Quantifying changes to extreme heat after 1.5 °C and 2.0 °C of warming

To quantify changes in extreme heat across the SAEA region under future warming scenarios, we first calculate five-day running mean values of daily maximum temperatures at each grid point in the region, and then extract the maximum value found across the period spanning April to September for each model year (hereafter Tx5DX). This yields 2692, 1785 and 1863 estimates of Tx5DX at each grid point for model simulations of the present-decade climate, 1.5 °C and 2 °C worlds, respectively (see supplementary table S1 available at stacks.iop.org/ERL/13/034011/mmedia). The choice of a five-day window is subjective, but is intended to reflect the approximate length of heat events for which severe impacts manifest themselves across the regions of interest.

The threshold of Tx5DX which occurs for one-in-ten model years under each of the future scenarios is then identified for each grid cell, and we then quantify the probability of this same threshold being exceeded in the present-decade runs. We then calculate the ratio of the probability of occurrence in the future (one in ten, by definition) relative to the probability of occurrence today, and denote this term the risk ratio (or RR_{10}). Hence, this metric represents the increase in frequency of future extreme heat, relative to the present-day climate.

While previous studies provide confidence in the capability of HadAM3P to faithfully simulate high-temperature extremes (Uhe *et al* 2016, Guillod *et al*




2017), the decision to present changes between warming levels in a relative framework (rather than in terms of absolute temperatures) ensures raw model output can be used directly, and avoids the need to make subjective decisions about what bias correction techniques might be most appropriate (Sippel *et al* 2016).

Finally, while the risk ratio metric applied here remains consistent with the approach employed by many previous studies (Stott *et al* 2016, NAS 2016), it is noted that our method differs by defining an event threshold with respect to the future world, rather than the present-day world. This choice was made to ensure that estimates of very small probabilities will be

calculated with the experiment which has the largest ensemble size. Further, it is noted that the choice of a one-in-ten year event in the future is an arbitrary threshold, but enables consideration of the most severe heat extremes which could occur in the future.

2.3. Patterns of extreme heat under a 1.5°- and 2°-warmer world

Figures 1(a) and (b) present ensemble-mean increases in Tx5DX across the SAEA region under the 1.5 °C and 2 °C experiments, relative to the present-decade experiment. Both results show regions of faster-than-average warming in higher latitudes and in continental



interiors, while less rapid signals are found in coastal regions and at lower latitudes—such patterns are consistent with expected changes to absolute temperature under future warming (Joshi *et al* 2008, Collins *et al* 2013).

To interpret the relative significance of these future changes in extreme heat, figures 1 (c) and (d) show the same results as for panels a and b, but this time normalised by the variability in Tx5DX experienced in the current climate (taken as the standard deviation across all ‘present-decade’ model runs, calculated separately for each individual grid box). When warming is kept to 1.5°C , changes remain relatively modest across the full region, with most places experiencing changes in Tx5DX of less than 1σ (with the exception of parts of the Middle East). But after 2.0°C of warming, changes exceeding 2 standard deviations can be found across large regions of equatorial Africa, as well as large sections of Indonesia, the Middle East and Iran. Such changes are all the more striking when ensemble-mean estimates of RR_{10} are considered (figures 1(e) and (f)): while under 1.5°C of warming, most regions experience risk ratios less than ten, increases in the frequency of severe local Tx5DX escalate to well beyond 50 f old for many regions in the 2.0°C -warmer world. In fact, the regions in black denote those locations for which the threshold of temperature seen every ten years in the future was never seen to occur across more than 2600 model years in the present-day climate. Such results corroborate previous studies, which found significant changes to the frequency of heat extremes are possible after only 2.0°C of warming, though such dramatic increases could be minimised if global mean warming is kept to 1.5°C (Perkins-Kirkpatrick and Gibson 2017, Russo *et al* 2017, Lewis *et al* 2017).

2.4. Quantifying probabilistic changes to heat exposure

Previous research has shown that the signal of changes to extreme heat emerges first when assessed as an average over large spatial scales (Angélil *et al* 2014), while higher variability at smaller scales may suppress any detectable signal of change (King *et al* 2015). However, multiple studies have also demonstrated how robust changes in extremes can in fact be found at small spatial scales when the results of many individual locations are aggregated together (Fischer *et al* 2013, Westra *et al* 2013, Fischer *et al* 2014, Schleussner *et al* 2017). An adapted version of this framework is applied here to quantify levels of population exposure to future changes in extreme heat, as found in figure 1, using gridded population data for the year 2015 (figure 2, CIESIN 2016).

Following equation (1), the average heat exposure for a given region is calculated across all ensemble members, N , by summing the risk ratio (RR_{10}) estimates across all grid cells (spanning latitudes ϕ_1 and ϕ_2 , and longitudes λ_1 and λ_2) with a weighting proportional to the fraction of people living in that grid cell, P/P_{TOT} . More generally, this approach produces a probability distribution which characterises population exposure to different risk ratios after 1.5°C and 2°C of warming. This framework therefore emphasises the importance of changing extremes where people live (Frame *et al* 2017), and will take into account not only future changes to distributions of temperature, but also changes to the number of people living in various locations.

$$\text{Exposure} = \frac{1}{N} \sum_{i=\phi_1}^{\phi_2} \sum_{j=\lambda_1}^{\lambda_2} \sum_{n=1}^N \left(\text{RR}_{10_{ijn}} \times \frac{P_{ijn}}{P_{\text{TOT}}} \right) \quad (1)$$

2.5. Population changes consistent with a 1.5°- and 2°-warmer world

Of course, when considering changes in population exposure to heat extremes under warming targets consistent with the Paris Agreement, a key determinant concerns the range of population scenarios which would be consistent with such a future. The Shared Socioeconomic Pathways (SSPs) provide a range of plausible future pathways of population and socioeconomic changes, and encompass the full spread of corresponding climate change scenarios from the Representative Concentration Pathway (RCP) database (KC and Lutz 2017, O'Neill *et al* 2016, Jones and O'Neill 2016). In addition to present-day population observations, this study also examines population projections for the year 2090 under an SSP1 and SSP2 scenario: these two were selected as 'baseline' scenarios most closely aligned, respectively, with the RCP2.6 and RCP4.5 climate scenario (O'Neill *et al* 2016). Since linear combinations of model simulations from both these RCPs were also used to produce the 1.5 °C and 2.0 °C HAPPI experiments (Mitchell *et al* 2017), we hereafter consider both SSP scenarios as feasible to occur in a future world where temperatures stabilise at 1.5 °C.

Figure 3 shows changes in absolute population between 2015 and 2090 under the two SSP scenarios considered. Both population data sets were first interpolated to a common $0.25^\circ \times 0.25^\circ$ resolution before calculations were made. Under both scenarios, there are two large-scale patterns of population change which emerge: (1) widespread decreases in population over many mid-latitude regions, including Central Asia and China, as well as large regions of South-east Asia; and (2) robust increases in population over the Middle East, parts of Pakistan and Afghanistan, and most notably over widespread areas of Eastern Africa, including already densely-populated regions in Ethiopia, Kenya, Tanzania and Uganda. Whilst more obvious under SSP2, smaller but more concentrated patterns of growth also exist under SSP1 over parts of East Africa, Turkey and even India—a feature likely indicative of continued urban intensification (Jiang and O'Neill 2017). Interestingly, population projections across India are more divergent than other regions: negligible changes in total population between 2015 and 2090 are found under an SSP1 scenario, while at least another 400 million people will inhabit the region under an SSP2 scenario (KC and Lutz 2017). Such large variations in trajectories between the scenarios for different regions can be primarily linked to (a) differences in present-day population age structures, and (b) the differing prospects of changing education rates for younger adult females (aged 20–39, KC and Lutz 2017).

To better understand how the SAEA-wide changes in exposure to extreme heat will translate down to smaller spatial scales with differing projections of future population, two sub-regions have been selected for further analysis: one encompassing the East African (11°S–15°N, 25°E–40°E; hereafter 'EAF') domain

exhibiting strong growth irrespective of SSP; the other encompassing an area over India, Bangladesh and Sri Lanka (5°N–28°N, 70°E–93°E; hereafter 'IND') which displays scenario-dependent population growth characteristics.

3. Results


Table 1 presents results of population exposure to RR_{10} for the three regions of interest, under the multiple combinations of population and climate scenarios of relevance in this study. To represent the variability in exposure implicit in looking over a large population grouping, we hereafter quantify the ensemble-median level of exposure to RR_{10} , as well as the RR_{10} which will be very likely and very unlikely (90% and 10% likelihood) exceeded by a given person across the SAEA, EAF and IND regions.

3.1. Aggregate changes to future heat exposure

For the full SAEA region, median exposure to severe T_{x5DX} increases by a factor of 4.1 (2.4–9.6) under 1.5 °C of warming and when population is kept fixed to 2015 levels, while this number rises to 15.8 (5.0–135) after 2.0 °C of warming. When population changes are also accounted for, SAEA-wide exposure increases by a further 50% under the SSP2 population scenario, while increases are also seen, though less significant, in the lower-population SSP1 scenario. Interestingly, median RR_{10} increases by a factor of four (2.1–25) if temperatures reach 2.0 °C of warming, instead of stabilising at 1.5 °C, irrespective of the population data set chosen. Such results are even more dramatic than previous-reported estimates of a doubling in extreme heat between 1.5 °C and 2.0 °C—however, those results were found using coarser-resolution models and with respect to the entire globe (Fischer and Knutti 2015).

Results for the sub-region encompassing India, Bangladesh and Sri Lanka (IND) qualitatively follow the same patterns of absolute exposure as for the wider SAEA region—such an outcome is unsurprising, given about one-third of the population across the entire SAEA domain is concentrated within the smaller IND region (table s2).

Across the East African (EAF) sub-region, increases in extreme heat are found to become 5.7 (2.1–12.2) times more likely after 1.5 °C warming if population remains fixed at 2015 levels, while risk ratios reach 9.4 (4.1–22.2) and 13.0 (5.7–30.7) if future population changes follow an SSP1 or SSP2 scenario respectively. Interestingly, there is a robust change in the frequency of extremes between the lower- and higher-warming scenarios: an estimated 5.2 fold (2.5–22) increase in extremes is expected between the 1.5- and 2 degree worlds irrespective of population scenario. These larger increases in RR overall for EAF (relative to the entire SAEA region) are indicative of the faster emergence in extreme temperatures found in figure 1, and also



agree with many other studies focusing on the relative rapidity of changes in heat extremes across the African continent (Mahlstein *et al* 2011, Russo *et al* 2016, Harrington *et al* 2016).

3.2. Quantifying the relative impact of changes in climate versus population dynamics

While absolute changes in the frequency of heat extremes are highly informative for regional stakeholders, the methods chosen in this study enable us to further interrogate the changes in population exposure to RR_{10} , by quantifying the relative importance of three factors: (1) changes attributable to increases in global mean temperatures (RR_{CLIM}); (2) changes attributable

to differences in total population size ($RR_{\Delta POPN}$); and (3) changes attributable to the spatial reorganisation of the population exposed ($RR_{REDISTRIBUTION}$). Combined, we find

$$RR_{10} = RR_{CLIM} \times RR_{\Delta POPN} \times RR_{REDISTRIBUTION} \quad (2)$$

where RR_{CLIM} is found by keeping population levels fixed at 2015 observations (first column of table 1) and $RR_{\Delta POPN}$ simply equals the percentage increase in total population for the region of interest under the different SSP scenarios considered (relative to 2015 observations). To calculate $RR_{REDISTRIBUTION}$, we repeat the

Table 1. Projected changes in population exposure to extreme Tx5DX for three regions under a range of future climate scenarios and population scenarios. Main answers denote the risk ratio which the median (50th percentile) person would experience; the answers in brackets respectively denote the risk ratios experienced by at least 90% and 10% of the relevant population grouping.

(a) SAEA	2015 population	SSP1–2090	SSP2–2090
RR ₁₀ at 1.5 °C	4.1 (2.4/9.6)	4.4 (2.3/10.6)	5.9 (3.0/13.8)
RR ₁₀ at 2.0 °C	15.8 (5.0/135)	18.9 (4.9/264)	26.6 (6.5/346)
RR _{2.0} / RR _{1.5}	3.9 (2.1/14)	4.3 (2.1/25)	4.5 (2.2/25)
(b) EAF			
RR ₁₀ at 1.5 °C	5.7 (2.3/12.2)	9.4 (4.1/22.2)	13.0 (5.7/30.7)
RR ₁₀ at 2.0 °C	29.9 (5.9/269)	48.8 (10.0/488)	67.5 (13.8/675)
RR _{2.0} / RR _{1.5}	5.2 (2.5/22)	5.2 (2.4/22)	5.2 (2.4/22)
(c) IND			
RR ₁₀ at 1.5 °C	4.3 (2.0/10.0)	4.3 (2.0/9.9)	5.7 (2.6/13.1)
RR ₁₀ at 2.0 °C	17.9 (4.1/269)	17.9 (4.0/268)	25.3 (5.4/354)
RR _{2.0} / RR _{1.5}	4.1 (2.0/27)	4.1 (2.0/27)	4.4 (2.0/27)

process of spatially aggregating risk ratios weighted according to where people live, but first normalise the maps of population for all three scenarios, so individual grid cells contain information about what percentage of the region's people live there, rather than what is the total number of people living there. By removing the effect of changes in total population size, the fractional difference in risk ratio between these results using 2015 populations, and those using SSP scenarios, will equal the percentage increase in risk of extreme heat attributable to changes in where people live (and thus $RR_{\text{REDISTRIBUTION}}$).

Table 2 presents the range of estimates of $RR_{\text{REDISTRIBUTION}}$ and $RR_{\Delta\text{POP}}$ for the various regions of interest. Results reveal that statistically significant increases in exposure to heat extremes occurs for the SAEA region as a whole, with best-guess (ensemble median) estimates of $RR_{\text{REDISTRIBUTION}}$ of 1.09 (1.03–1.12) after 1.5 °C of warming, and best-guess estimates exceeding 1.2 for a 2.0 °C-warmer world. This result can be explained by the fact that those locations where future population growth (decline) is most significant (figure 3) also preferentially align with the regions of substantially higher (lower) risk ratios in the future (figures 1(e)–(f)). Estimates of $RR_{\text{REDISTRIBUTION}}$ are also robust irrespective of whether an SSP1 or SSP2 scenario is considered—a result which is surprising given some of the large regional differences in population change. The approximate doubling in $RR_{\text{REDISTRIBUTION}}$ after 2.0 °C of warming relative to the 1.5 °C experiment reflects the non-linear changes in heat extremes which occur at the higher warming level—this acceleration of risk ratios in fast-emerging locations will further exacerbate the relative impact of a higher population fraction living there in the future.

When considering the smaller sub-regions, we find negligible changes in risk ratios over India associated with the redistribution of where people live—this is unsurprising however, since population density is extremely high across the entirety of the region. Instead, the added increases in risk ratio in table 1 under SSP2

are the direct result of increases in total population size ($RR_{\Delta\text{POP}}$). By contrast, there is actually a robust decrease in population exposure to heat extremes associated with changes to where people live for the EAF region, with best-guess estimates of an 8% and 17% reduction in heat exposure after 1.5 °C and 2.0 °C of warming, respectively. This effect therefore helps to mitigate some of the impact of 80% and 150% population increases expected for the region under SSP1 and SSP2 scenarios (respectively). However, as table 1 reveals, the net effect is still that of exposure to higher-than-average risk ratios being observed for the EAF region.

4. Discussion and limitations

While certainly not the only manifestation of climate change under a 1.5 °C and 2.0 °C scenario (Schleussner *et al* 2016a), heat extremes are the events where we see the strongest impact of climate change, are the key determinant in changing rates of heat stress incidence (Pal and Eltahir 2016, Mora *et al* 2017, Li *et al* 2018), and the only extreme where climate change is making previously rare events more likely by orders of magnitude (Christidis and Stott 2013, King and Karoly 2017, King *et al* 2017, Lewis *et al* 2017). Though the patterns of change to heat extremes are not the same worldwide, all regions do show the same first-order response, with monotonic increases in both frequency and severity occurring everywhere under future warming scenarios (figure 1). However, future population changes can be dramatically different depending on the location considered (figure 3): for the densely populated SAEA region, there are widespread decreases in population predicted over China, South-east and Central Asia, while regions across Eastern Africa continue to witness as much as a doubling in population over the same period. These regional patterns of population decrease and increase happen to align, respectively, with regions which experience slower-than-average and faster-than-average changes in the emergence of heat extremes.

As a direct consequence of this redistribution effect, population exposure to worsening heat extremes can increase by as much as 30% under 2.0 °C of global-mean warming, with non-trivial impacts also detectable for smaller sub-regions. The effect of changing population dynamics further compounds the already-significant increases in extreme heat exposure expected due to warming only. Therefore, while restricting global-mean warming to 1.5 °C instead of 2.0 °C can reduce the exposure by a factor of five over some of the most vulnerable regions, regional decision makers would further benefit from policies to ensure future population changes are compatible with an SSP1 scenario. Further research is also needed to understand how these patterns of heat exposure are compounded by the anticipated increases in urbanisation

Table 2. Same as for table 1, but instead of showing absolute risk ratios, answers have been separated to show the increase in risk of extreme heat attributable to changes in where people live ($RR_{\text{REDISTRIBUTION}}$) and to changes in the overall size of the relevant population grouping ($RR_{\Delta\text{POP}}$). See main text for details of the relevant methodology.

	SSP1–2090		SSP2–2090	
(a) SAEA	$RR_{\text{REDISTRIBUTION}}$	$RR_{\Delta\text{POP}}$	$RR_{\text{REDISTRIBUTION}}$	$RR_{\Delta\text{POP}}$
1.5 °C world	1.08 (1.03/1.11)	0.981	1.10 (1.04/1.12)	1.283
2.0 °C world	1.20 (1.06/1.30)		1.25 (1.12/1.34)	
(b) EAF				
1.5 °C world	0.92 (0.87/0.96)	1.814	0.92 (0.89/0.94)	2.509
2.0 °C world	0.83 (0.75/0.96)		0.83 (0.75/0.93)	
(c) IND				
1.5 °C world	1.00 (0.98/1.02)	0.995	1.01 (1.00/1.03)	1.313
2.0 °C world	1.00 (1.00/1.06)		1.06 (1.00/1.10)	

rates worldwide (Jiang and O'Neill 2017), particularly in terms of urban heat island impacts.

This analysis uses a very large ensemble of atmosphere-only model simulations, thereby enabling a more complete characterisation of internal climate variability and hence more accurate estimates of temperature distribution tails. However, there are limitations implicit with employing only a single type of climate model, and the uncertainty bounds presented in this analysis may therefore underestimate the ‘true’ range of uncertainty (Bellprat and Doblas-Reyes 2016). It is worth noting though that the spatial aggregation techniques applied in this study should render such discrepancies to be less relevant to the overall results, particularly when compared to studies on specific regions. Further, compared to other large-ensemble GCM simulations, HadAM3P exhibits a climate response to warming that is on the lower end of the spectrum (e.g. Philip *et al* 2017, van Oldenborgh *et al* 2018) so estimates given here are likely conservative. Nevertheless, future work will benefit from replicating these results using super-ensembles of multiple other high-resolution regional climate models.

5. Summary

Stakeholders and policy makers require information on changing risks today and in the future. Understanding changes in the severity of extreme heat in response to 1.5 °C and 2.0 °C of warming is an important part of future risk assessments, both for global and regional decision makers—however, the hazard is only one of three drivers of risk (Pachauri *et al* 2014). Exposure, and thus the role of future changes to population, often remains overlooked or assessed independently of the hazard on different scales, preventing risk estimation on the scales that decisions are made on.

The results presented in this analysis disaggregates large-scale patterns of change in the climate system from changes in patterns of population growth. This study also improves upon previous methods of quantifying population exposure to future heat

(Jones *et al* 2015, Liu *et al* 2017), by combining techniques previously developed to (1) quantify probabilistic changes to climatic extremes (Stott *et al* 2004, Otto 2016, Stott 2016) and (2) better characterise localised changes across multiple regions, through the use of spatial aggregation (Fischer *et al* 2013). The combination of changes in the hazard (extreme heat) with changes in exposure allow for a more comprehensive assessment of future risks. By understanding what differences in heat exposure may emerge in a 1.5 °C or 2.0 °C world, and the influence of different population scenarios compatible with the Paris Agreement targets, decision makers can begin to resolve the minimum adaptation measures which will be necessary even in the case of successful mitigation policies.

While local adaptation planners might be primarily be interested in how the patterns of heat extremes align with changes in population over their immediate community, it is equally important for decision makers to recognise the broader implications of heat exposure increases driven by future changes in where people live. Such patterns may lead to higher-than-anticipated impacts from extreme heat, including greater food insecurity for vulnerable countries (Lobell *et al* 2011, Liu *et al* 2016, Asseng *et al* 2015, Zhao *et al* 2017, Lobell and Asseng 2017), increases in excess-heat mortality rates (Mitchell *et al* 2016a, Gasparrini *et al* 2015, 2017, Mora *et al* 2017, Watts *et al* 2017a, 2017b), and the potential for higher rates of emigration from the most severely-affected regions (Reuveny 2007, Black *et al* 2011b, 2011a, Gemenne *et al* 2014, Lister 2014, Watts *et al* 2017a, 2017b).

Acknowledgments

We would like to thank our colleagues at the Oxford eResearch Centre: A Bowery, S Li, M Rashid, S Sparrow and D Wallom for their technical expertise. We would like to thank the Met Office Hadley Centre PRECIS team for their technical and scientific support for the development and application of Weather@Home. Finally, we would like to thank all of the volunteers who have donated their computing time to climateprediction.net and Weather@Home.

ORCID iDs

Luke J Harrington  <https://orcid.org/0000-0002-1699-6119>

Friederike E L Otto  <https://orcid.org/0000-0001-8166-5917>

References

- Angélil O, Stone D A, Tadross M, Tummon F, Wehner M and Knutti R 2014 Attribution of extreme weather to anthropogenic greenhouse gas emissions: Sensitivity to spatial and temporal scales *Geophys. Res. Lett.* **41** 2150–5
- Asseng S *et al* 2015 Rising temperatures reduce global wheat production *Nat. Clim. Change* **5** 143–7
- Bellprat O and Doblas-Reyes F 2016 Attribution of extreme weather and climate events overestimated by unreliable climate simulations *Geophys. Res. Lett.* **43** 2015GL067189
- Black R, Adger W N, Arnell N W, Dercon S, Geddes A and Thomas D 2011a The effect of environmental change on human migration *Glob. Environ. Change* **21** S3–11
- Black R, Bennett S R G, Thomas S M and Beddington J R 2011b Climate change: Migration as adaptation *Nature* **478** 447–9
- Center for International Earth Science Information Network—CIESIN—Columbia University 2016 Gridded Population of the World, Version 4 (GPWv4): Population Density (<https://doi.org/10.7927/H4NP22DQ>)
- Christidis N and Stott P A 2013 Change in the odds of warm years and seasons due to anthropogenic influence on the climate *J. Clim.* **27** 2607–21
- Ciavarella A, Stott P and Lowe J 2017 Early benefits of mitigation in risk of regional climate extremes *Nat. Clim. Change* **7** 326–30
- Collins M *et al* 2013 Long-term climate change: projections, commitments and irreversibility climate change 2013: the physical science basis *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed T F Stocker, D Qin, G-K Plattner, M Tignor, S K Allen, J Boschung, A Nauels, Y Xia, V Bex and P M Midgley (Cambridge: Cambridge University Press) pp 1029–136
- Davis S J and Diffenbaugh N 2016 Dislocated interests and climate change *Environ. Res. Lett.* **11** 061001
- Diffenbaugh N S and Charland A 2016 Probability of emergence of novel temperature regimes at different levels of cumulative carbon emissions *Front. Ecol. Environ.* **14** 418–23
- Diffenbaugh N S and Scherer M 2011 Observational and model evidence of global emergence of permanent, unprecedented heat in the 20th and 21st centuries *Clim. Change* **107** 615–24
- Fischer E M, Beyerle U and Knutti R 2013 Robust spatially aggregated projections of climate extremes *Nat. Clim. Change* **3** 1033–8
- Fischer E M and Knutti R 2015 Anthropogenic contribution to global occurrence of heavy-precipitation and high-temperature extremes *Nat. Clim. Change* **5** 560–4
- Fischer E M, Sedláček J, Hawkins E and Knutti R 2014 Models agree on forced response pattern of precipitation and temperature extremes *Geophys. Res. Lett.* **41** 2014GL062018
- Frame D, Joshi M, Hawkins E, Harrington L J and Roiste M 2017 Population-based emergence of unfamiliar climates *Nat. Clim. Change* **7** 407–11
- Gasparrini A *et al* 2015 Mortality risk attributable to high and low ambient temperature: a multicountry observational study *The Lancet* **386** 369–75
- Gasparrini A *et al* 2017 Projections of temperature-related excess mortality under climate change scenarios *Lancet Planet. Health* **1** e360–7
- Gemenne F, Barnett J, Adger W N and Dabelko G D 2014 Climate and security: evidence, emerging risks, and a new agenda *Clim. Change* **123** 1–9
- Guillot B P *et al* 2017 weather@home 2: validation of an improved global–regional climate modelling system *Geosci. Model Dev.* **10** 1849–72
- Harrington L J, Frame D J, Fischer E M, Hawkins E, Joshi M and Jones C D 2016 Poorest countries experience earlier anthropogenic emergence of daily temperature extremes *Environ. Res. Lett.* **11** 055007
- Harrington L J, Frame D J, Hawkins E and Joshi M 2017 Seasonal cycles enhance disparities between low- and high-income countries in exposure to monthly temperature emergence with future warming *Environ. Res. Lett.* **12** 114039
- Hawkins E *et al* 2017 Estimating changes in global temperature since the pre-industrial period *Bull. Am. Meteorol. Soc.*
- Hawkins E and Sutton R 2012 Time of emergence of climate signals *Geophys. Res. Lett.* **39** L01702
- Henley B J and King A D 2017 Trajectories toward the 1.5 °C Paris target: Modulation by the Interdecadal Pacific Oscillation *Geophys. Res. Lett.* **44** 2017GL073480
- Herold N, Alexander L, Green D and Donat M 2017 Greater increases in temperature extremes in low versus high income countries *Environ. Res. Lett.* **12** 034007
- Im E-S, Pal J S and Eltahir E A B 2017 Deadly heat waves projected in the densely populated agricultural regions of South Asia *Sci. Adv.* **3** e1603322
- James R, Washington R, Schleussner C-F, Rogelj J and Conway D 2017 Characterizing half-a-degree difference: a review of methods for identifying regional climate responses to global warming targets *Wiley Interdiscip. Rev. Clim. Change* **8** e457
- Jiang L and O'Neill B C 2017 Global urbanization projections for the Shared Socioeconomic Pathways *Glob. Environ. Change* **42** 193–9
- Jones B and O'Neill B C 2016 Spatially explicit global population scenarios consistent with the Shared Socioeconomic Pathways *Environ. Res. Lett.* **11** 084003
- Jones B, O'Neill B C, McDaniel L, McGinnis S, Mearns L O and Tebaldi C 2015 Future population exposure to US heat extremes *Nat. Clim. Change* **5** 652–5
- Joshi M M, Gregory J M, Webb M J, Sexton D M H and Johns T C 2008 Mechanisms for the land/sea warming contrast exhibited by simulations of climate change *Clim. Dyn.* **30** 455–65
- KC S and Lutz W 2017 The human core of the shared socioeconomic pathways: Population scenarios by age, sex and level of education for all countries to 2100 *Glob. Environ. Change*
- King A D, Donat M G, Fischer E M, Hawkins E, Alexander L V, Karoly D J, Dittus A J, Lewis S C and Perkins S E 2015 The timing of anthropogenic emergence in simulated climate extremes *Environ. Res. Lett.* **10** 094015
- King A D and Karoly D 2017 Climate extremes in Europe at 1.5 and 2 degrees of global warming *Environ. Res. Lett.*
- King A D, Karoly D J and Henley B J 2017 Australian climate extremes at 1.5 °C and 2 °C of global warming *Nat. Clim. Change* **7** 412–6
- Knutti R, Rogelj J, Sedláček J and Fischer E M 2016 A scientific critique of the two-degree climate change target *Nat. Geosci.* **9** 13–8
- Kraaijenbrink P D A, Bierkens M F P, Lutz A F and Immerzeel W W 2017 Impact of a global temperature rise of 1.5 °C on Asia's glaciers *Nature* **549** 257–60
- Lewis S C, King A D and Mitchell D M 2017 Australia's unprecedented future temperature extremes under Paris limits to warming *Geophys. Res. Lett.* **44** 9947–56
- Li J, Chen Y D, Gan T Y and Lau N-C 2018 Elevated increases in human-perceived temperature under climate warming *Nat. Clim. Change* **8** 43
- Lister M 2014 Climate change refugees *Crit. Rev. Int. Soc. Polit. Phil.* **17** 618–34
- Liu B *et al* 2016 Similar estimates of temperature impacts on global wheat yield by three independent methods *Nat. Clim. Change* **6** 1130–6

- Liu Z, Anderson B, Yan K, Dong W, Liao H and Shi P 2017 Global and regional changes in exposure to extreme heat and the relative contributions of climate and population change *Sci. Rep.* **7** srep43909
- Lobell D B and Asseng S 2017 Comparing estimates of climate change impacts from process-based and statistical crop models *Environ. Res. Lett.* **12** 015001
- Lobell D B, Bänziger M, Magorokosho C and Vivek B 2011 Nonlinear heat effects on African maize as evidenced by historical yield trials *Nat. Clim. Change* **1** 42–5
- Mahlstein I, Knutti R, Solomon S and Portmann R W 2011 Early onset of significant local warming in low latitude countries *Environ. Res. Lett.* **6** 034009
- Millar R J, Fuglestedt J S, Friedlingstein P, Rogelj J, Grubb M J, Matthews H D, Skeie R B, Forster P M, Frame D J and Allen M R 2017 Emission budgets and pathways consistent with limiting warming to 1.5 °C *Nat. Geosci.* **10** 741–7
- Mishra V, Mukherjee S, Kumar R and Stone D 2017 Heat wave exposure in India in current, 1.5 °C, and 2.0 °C worlds *Environ. Res. Lett.*
- Mitchell D *et al* 2017 Half a degree additional warming, prognosis and projected impacts (HAPPI): background and experimental design *Geosci. Model Dev.* **10** 571–83
- Mitchell D, Heaviside C, Vardoulakis S, Huntingford C, Masato G, Guillod B P, Frumhoff P, Bowery A, Wallom D and Allen M 2016a Attributing human mortality during extreme heat waves to anthropogenic climate change *Environ. Res. Lett.* **11** 074006
- Mitchell D, James R, Forster P M, Betts R A, Shiogama H and Allen M 2016b Realizing the impacts of a 1.5 °C warmer world *Nat. Clim. Change* **6** 735–7
- Mora C *et al* 2017 Global risk of deadly heat *Nat. Clim. Change* **7** 501–6
- National Academies of Sciences, Engineering, and Medicine, Committee on Extreme Weather Events and Climate Change Attribution, Board on Atmospheric Sciences and Climate and Division on Earth and Life Studies 2016 *Attribution of Extreme Weather Events in the Context of Climate Change* (Washington, DC: National Academies Press)
- van Oldenborgh G J, Philip S, Kew S, van Weele M, Uhe P, Otto F, Singh R, Pai I, Cullen H and AchutaRao K 2018 Extreme heat in India and anthropogenic climate change *Nat. Hazards Earth Syst. Sci.* **18** 365–81
- O'Neill B C *et al* 2016 The scenario model intercomparison project (ScenarioMIP) for CMIP6 *Geosci. Model Dev.* **9** 3461–82
- Otto F E L 2016 Extreme events: the art of attribution *Nat. Clim. Change* **6** 342–3
- Pachauri R K *et al* 2014 *Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Geneva: IPCC)
- Pal J S and Eltahir E A B 2016 Future temperature in southwest Asia projected to exceed a threshold for human adaptability *Nat. Clim. Change* **6** 197–200
- Perkins-Kirkpatrick S E and Gibson P B 2017 Changes in regional heatwave characteristics as a function of increasing global temperature *Sci. Rep.* **7** 12256
- Philip S *et al* 2017 Attribution analysis of the Ethiopian drought of 2015 *J. Clim.*
- Reuveny R 2007 Climate change-induced migration and violent conflict *Polit. Geogr.* **26** 656–73
- Rogelj J and Knutti R 2016 Geosciences after Paris *Nat. Geosci.* **9** 187–9
- Rogelj J, Schleussner C-F and Hare W 2017 Getting it right matters: temperature goal interpretations in geoscience research *Geophys. Res. Lett.* 2017GL075612
- Russo S, Marchese A F, Sillmann J and Immé G 2016 When will unusual heat waves become normal in a warming Africa? *Environ. Res. Lett.* **11** 054016
- Russo S, Sillmann J and Sterl A 2017 Humid heat waves at different warming levels *Sci. Rep.* **7** 7477
- Sanderson B M *et al* 2017 Community climate simulations to assess avoided impacts in 1.5 °C and 2 °C futures *Earth Syst. Dyn.* **8** 827–47
- Schleussner C-F *et al* 2016a Differential climate impacts for policy-relevant limits to global warming: the case of 1.5 °C and 2 °C *Earth Syst. Dyn.* **7** 327–51
- Schleussner C-F, Rogelj J, Schaeffer M, Lissner T, Licker R, Fischer E M, Knutti R, Levermann A, Frieler K and Hare W 2016b Science and policy characteristics of the Paris agreement temperature goal *Nat. Clim. Change* **6** 827–35
- Schleussner C-F, Pfleiderer P and Fischer E M 2017 In the observational record half a degree matters *Nat. Clim. Change* **7** 460–2
- Schurer A P, Mann M E, Hawkins E, Tett S F B and Hegerl G C 2017 Importance of the pre-industrial baseline for likelihood of exceeding Paris goals *Nat. Clim. Change* **7** 563–7
- Seneviratne S I, Donat M G, Pitman A J, Knutti R and Wilby R L 2016 Allowable CO₂ emissions based on regional and impact-related climate targets *Nature* **529** 477–83
- Sippel S, Otto F E L, Forkel M, Allen M R, Guillod B P, Heimann M, Reichstein M, Seneviratne S I, Thonicke K and Mahecha M D 2016 A novel bias correction methodology for climate impact simulations *Earth Syst. Dyn.* **7** 71–88
- Stott P 2016 How climate change affects extreme weather events *Science* **352** 1517–8
- Stott P A *et al* 2016 Attribution of extreme weather and climate-related events *Wiley Interdiscip. Rev. Clim. Change* **7** 23–41
- Stott P A, Stone D A and Allen M R 2004 Human contribution to the European heatwave of 2003 *Nature* **432** 610–4
- Uhe P, Otto F E L, Hausteine K, van Oldenborgh G J, King A D, Wallom D C H, Allen M R and Cullen H 2016 Comparison of methods: attributing the 2014 record European temperatures to human influences *Geophys. Res. Lett.* **43**
- Wang Z, Lin L, Zhang X, Zhang H, Liu L and Xu Y 2017 Scenario dependence of future changes in climate extremes under 1.5 °C and 2 °C global warming *Sci. Rep.* **7** srep46432
- Watts N *et al* 2017a The Lancet countdown: tracking progress on health and climate change *Lancet* **389** 1151–64
- Watts N 2017b The Lancet Countdown on health and climate change: from 25 years of inaction to a global transformation for public health *Lancet*
- Westra S, Alexander L V and Zwiers F W 2013 Global Increasing Trends in Annual Maximum Daily Precipitation *J. Clim.* **26** 3904–18
- Zhao C *et al* 2017 Temperature increase reduces global yields of major crops in four independent estimates *Proc. Natl Acad. Sci.* **114** 9326–31