

**OPEN ACCESS**

## Recent global sea level acceleration started over 200 years ago?

To cite this article: Svetlana Jevrejeva *et al* 2009 *IOP Conf. Ser.: Earth Environ. Sci.* **6** 012026

View the [article online](#) for updates and enhancements.

### You may also like

- [Upper limit for sea level projections by 2100](#)  
S Jevrejeva, A Grinsted and J C Moore
- [The Si vacancy: an example of a pressure-sensitive Jahn-Teller system](#)  
E Tarnow
- [Flood damage costs under the sea level rise with warming of 1.5 °C and 2 °C](#)  
S Jevrejeva, L P Jackson, A Grinsted et al.



**ECS**  
The  
Electrochemical  
Society  
Advancing solid state &  
electrochemical science & technology

**DISCOVER**  
how sustainability  
intersects with  
electrochemistry & solid  
state science research

**P01.09**

**Recent global sea level acceleration started over 200 years ago?**

*Svetlana Jevrejeva(1), J Moore(2), A Grinsted(3), P Woodworth(1)*

*(1) Proudman Oceanographic Laboratory, United Kingdom*

*(2) Arctic Centre, University of Lapland, Finland*

*(3) Niels Bohr Institute, University of Copenhagen, Denmark*

We present a reconstruction of global sea level (GSL) since 1700 calculated from tide gauge records and analyse the evolution of global sea level acceleration during the past 300 years. We provide observational evidence that sea level acceleration up to the present has been about 0.01 mm/yr<sup>2</sup> and appears to have started at the end of the 18th century. Sea level rose by 6 cm during the 19th century and 19 cm in the 20th century. Superimposed on the long-term acceleration are quasi-periodic fluctuations with a period of about 60 years. If the conditions that established the acceleration continue, then sea level will rise 34 cm over the 21st century. Long time constants in oceanic heat content and increased ice sheet melting imply that the latest Intergovernmental Panel on Climate Change (IPCC) estimates of sea level are probably too low.