PAPER

Measuring the self-healing of the spatially inhomogeneous states of polarization of vector Bessel beams

To cite this article: Giovanni Milione et al 2015 J. Opt. 17 035617

View the article online for updates and enhancements.

You may also like

- <u>Entropy production and its large deviations</u> in an active lattice gas Tal Agranov, Michael E Cates and Robert L Jack
- <u>Physics of active emulsions</u> Christoph A Weber, David Zwicker, Frank Jülicher et al.
- <u>Stripe structures in phase separated</u> <u>magnetic oxides</u> O G Udalov and I S Beloborodov

J. Opt. 17 (2015) 035617 (7pp)

Measuring the self-healing of the spatially inhomogeneous states of polarization of vector Bessel beams

Giovanni Milione^{1,2,3,4}, Angela Dudley⁵, Thien An Nguyen^{1,2}, Ougni Chakraborty^{1,2}, Ebrahim Karimi⁶, Andrew Forbes^{5,7} and Robert R Alfano^{1,2,3,4}

¹ Institute for Ultrafast Spectroscopy and Lasers, 160 Convent Ave., New York, NY 10031, USA ² Physics Department, City College of the City University of New York, 160 Convent Ave., New York, NY 10031, USA

³Graduate Center of the City University of New York, 365 Fifth Ave., New York, NY 10016, USA

⁴New York State Center for Complex Light, 160 Convent Ave., New York, NY 10031, USA

⁵CSIR National Laser Centre, PO Box 395, Pretoria 0001, South Africa

⁶ Department of Physics, University of Ottawa, 150 Louis Pasteur, Ottawa, Ontario, K1N 6N5, Canada

⁷ School of Physics, University of the Witwatersrand, Johannesburg 2000, South Africa

E-mail: giomilione@gmail.com and angelajvr@gmail.com

Received 10 November 2014, revised 8 December 2014 Accepted for publication 17 December 2014 Published 20 February 2015



Abstract

We experimentally measured the self-healing of the spatially inhomogeneous states of polarization of vector Bessel beams. Radially and azimuthally polarized vector Bessel beams were experimentally generated via a digital version of Durnin's method, using a spatial light modulator in concert with a liquid crystal *q*-plate. As a proof of principle, their intensities and spatially inhomogeneous states of polarization were experimentally measured using Stokes polarimetry as they propagated through two disparate obstructions. It was found, similar to their intensities, that their spatially inhomogeneous states of polarization self-healed. The self-healing can be understood via geometric optics, i.e., the interference of the unobstructed conical rays in the shadow region of the obstruction, and may have applications in, for example, optical trapping.

Keywords: Bessel beams, vector beams, radial polarization, azimuthal polarization, optical vortex, orbital angular momentum, self-healing

(Some figures may appear in colour only in the online journal)

A Bessel beam is a light beam that is a solution to the Helmholtz wave equation, existing over a limited region of propagation, and experimentally generated by the interference of conical rays [1, 2]. It possesses a property referred to as 'self-healing,' i.e., its intensity reappears after propagation through an obstruction. Self-healing can be simply understood via geometric optics [3]. When a portion of a Bessel beam is obstructed in one plane, the unobstructed conical rays interfere in its shadow region in another plane, as shown in figure 1(e). Due to this property, Bessel beams have been extensively studied and have been used for a number of applications; for comprehensive reviews see [4–

7]. For example, when using a Bessel beam for optical trapping, it is possible to simultaneously trap multiple particles in well separated planes [8], and make particle tractor beams [9-12].

A vector beam is a light beam possessing a spatially inhomogeneous state of polarization such as radial or azimuthal polarization as shown in figures 2(b) and (c), respectively. Vector beams have received significant interest [13, 14], due in great part to their ability to produce stronger longitudinal field components [15, 16], and smaller spot sizes [17], as compared to scalar light beams, upon focusing by high numerical aperture objectives. Also, when using



Figure 1. Experimental setup: (a) computer generated hologram (CGH) of an annular slit additionally superimposed with a linear grating displayed on spatial light modulator (SLM). (b) Linear polarized annular ring of light before *q*-plate. (c) Schematic of q = 1/2 *q*-plate. (d) Radially polarized annular ring of light after *q*-plate. (e) Schematic of vector Bessel beam propagation though an obstruction in the focal region of L_B . The intensity of the vector Bessel beam is shown at a propagation distance of roughly (I) 64 mm (unobstructed), (II) 82 mm (obstructed), (III) 118 mm (semi-healed) and (IV) 136 mm (self-healed) after the Fourier transforming lens LB.

vector beams for optical trapping, it is possible to improve the axial and transverse stiffness of the optical trap via radial and azimuthal polarization, respectively [18–20]. Most recently, vector beams have been used for optical communication [21].

Like any other light beam a Bessel beam can have a *scalar* (spatially homogeneous) or *vector* (spatially inhomogeneous) state of polarization [22, 23]. The great majority of experimental investigations of Bessel beams concern scalar Bessel beams. Yet, a vector Bessel beam possesses the properties of a Bessel beam and a vector beam, as described above, and may be used in comparable applications. For example, it may be possible to improve the axial and transverse stiffness of a tractor beam when using a vector Bessel beam. While there are extensive studies on self-healing of scalar Bessel beams, there are limited studies on self-healing of vector Bessel beams [24–27], particularly with respect to their spatially inhomogeneous states of polarization. Previous work only measured the propagation and self-healing of the *intensities* of vector Bessel beams.

In this work, we experimentally measured the self-healing of the spatially inhomogeneous states of *polarization* of vector Bessel beams. Radially and azimuthally polarized vector Bessel beams were experimentally generated via a digital version of Durnin's method, using a spatial light modulator (SLM) in concert with a liquid crystal *q*-plate. As a proof of principle, their intensities and spatially inhomogeneous states of polarization were experimentally measured using Stokes polarimetry as they propagated through two disparate obstructions.

A schematic of the experimental setup is shown in figure 1. First, a scalar Bessel beam was generated following Durnin's method. In Durnin's method, an annular slit is placed in the back focal plane of a lens and illuminated with a



Figure 2. Experimentally measured intensity of scalar (linearly polarized) and vector Bessel beams. (a) (First row) scalar Bessel beam. (b) (Second row) radial polarization. (c) (Third row) azimuthal polarization. The columns show I_0 (second column), I_{45} (third column), I_{90} (fourth column), I_{135} (fifth column) or each Bessel beam as described in the text.

collimated light beam resulting in an annular ring of light. Each point along the annular ring acts as a point source which the lens transforms into a good approximation to a Bessel beam in its focal region [1, 2]. A digital method of Durnin's method was implemented [28]. An annular slit, additionally superimposed with a linear grating, was created using a computer generated hologram displayed on a reflective phase only SLM (HoloEye) as shown in figure 1(a). A linear polarized HeNe laser beam ($\lambda \sim 633$ nm) was spatially filtered by a single mode optical fiber, expanded, collimated by a lens (L_1) , and then illuminated the SLM. The light at the plane of the SLM was then spatially filtered by an aperture (A) in the first diffraction order of a 4 f imaging system (L_2 and L_3), imaged onto the back focal plane of a 10 cm focal length lens (L_B) , resulting in the linear polarized annular ring of light shown in figure 1(b). A good approximation to a scalar (linearly polarized) Bessel beam was formed in the focal region of lens L_B .

Next, the scalar (linearly polarized) Bessel beam was converted into a vector Bessel beam. There are many methods to generate vector beams including the use of metasurfaces and optical fibers [29–32]. Here, we use a *q*-plate [33]. A *q*-plate is a liquid crystal technology comprising of a thin layer of liquid crystal molecules in-between two thin glass plates. The orientation of the liquid crystal molecules is described by $q\phi$, where ϕ is the azimuthal coordinate, and *q* is a half integer. A q = 1/2 *q*-plate is schematically shown in figure 1(c). Effectively, a *q*-plate is a half wave plate (HWP) with an azimuthally varying fast axis that can be represented

by the Jones matrix [34]:

$$\hat{Q} = \begin{pmatrix} \cos 2q\phi & \sin 2q\phi \\ \sin 2q\phi & -\cos 2q\phi \end{pmatrix}.$$
(1)

Using Jones calculus, it can be easily shown, for a q = 1/2 qplate, horizontal (vertical) polarization can be converted into radial (azimuthal) polarization [35]. In general, a q-plate converts any state of polarization on the Poincare sphere to a higher-order state of polarization on the higher-order Poincare sphere [36, 37]. The q-plate is also 'tunable'; the amount of the incident light's power converted to radial or azimuthal polarization is directly controlled, i.e., tuned, via a voltage over the q-plate. For $\lambda \sim 633$ nm, when the voltage over the q-plate is $V_o \sim 5$ volts, no light will be converted to radial (azimuthal) polarization, i.e., the light remains linear polarized. When the voltage over the q-plate is $V_v \sim 2.3$ volts, all of the incident light's power will be converted to radial (azimuthal) polarization [38, 39].

The *q*-plate was placed close to the back focal plane of lens L_B . A linear polarizer (Pol) was placed just before the *q*-plate to ensure the incident light was completely linear polarized. A HWP was used to rotate the light's polarization horizontal (vertical). A signal generator generating a 1 kHz square wave was used to apply a voltage over the *q*-plate. When the voltage was V_o , a good approximation to a scalar (linearly polarized) Bessel beam was generated in the focal region of lens L_B , as shown in figure 2(a). When the voltage was V_v , and the light's polarization was rotated horizontal (vertical), a good approximation to a radial (azimuthal) polarized vector Bessel beam was generated in the focal

region of lens L_B , as shown in figures 2(b) and (c). A 10X microscope objective was used to image each Bessel beam in the focal region of lens L_B onto a CCD camera.

Stokes polarimetry was used to measure the state of polarization of each Bessel beam [40, 41]. In Stokes polarimetry, the Stokes parameters are measured via intensity measurements and used to calculate the polarization orientation and polarization ellipticity at every spatial point of a light beam. The first three Stokes parameters are given by [40, 41]:

$$S_0(r, \phi) = I_0(r, \phi) + I_{90}(r, \phi), \tag{2}$$

$$S_1(r, \phi) = I_0(r, \phi) - I_{90}(r, \phi), \qquad (3)$$

$$S_2(r,\phi) = I_{45}(r,\phi) - I_{135}(r,\phi), \tag{4}$$

where $I_0(r, \phi)$, $I_{45}(r, \phi)$, $I_{90}(r, \phi)$, and $I_{135}(r, \phi)$ are the intensity of the light beam, at every spatial point, measured after a linear polarizer whose transmission axis is rotated 0°, 45°, 90°, 135°, respectively. (r, ϕ) are cylindrical coordinates. The total intensity of the light beam is given by $S_0(r, \phi)$ and the orientation of the state of polarization is given by [40, 41]:

$$\psi(r, \phi) = \frac{1}{2} \tan^{-1} \left(\frac{S_2(r, \phi)}{S_1(r, \phi)} \right).$$
(5)

A linear polarizer (Pol) was placed just before the CCD camera in the experimental setup described above. The linear polarizer was used to measure $I_0(r, \phi)$, $I_{45}(r, \phi)$, $I_{90}(r, \phi)$, and $I_{135}(r, \phi)$ for each Bessel beam as shown in the respective columns of figure 2. While the third Stokes parameter, S_3 , and therefore the polarization ellipticity, was not measured, as shown in the experimental results, it is qualitatively enough to visualize the self-healing of the spatially inhomogeneous state of polarization of each vector Bessel beam via the polarization orientation.

Finally, each Bessel beam was made to propagate through an obstruction. The results are shown in figure 3. The obstruction was created by placing a pitted glass slide possessing multiple, random, speckled, and opaque obstructions in the path of the Bessel beam. The slide's position was adjusted until an isolated and appropriately sized obstruction was found. The size and position of the obstruction was chosen such that it obstructed approximately a small portion of the Bessel beam near its center. The obstruction is outlined by a dashed white line in figure 3. The intensities and states of polarization of each Bessel beam were measured via Stokes polarimetry, as described above, at four propagation distances as they propagated through the obstruction. The four propagation distances are schematically shown in figure 1(e): (I) unobstructed (II) obstructed (III) semi-healed (IV) self-healed. $S_0(r, \phi)$ is overlaid with $\psi(r, \phi)$ such that any change in intensity and orientation of the state of polarization as the Bessel beam propagates through the obstruction can be visualized simultaneously; $S_0(r, \phi)$ is encoded by relative contrast, i.e., bright to dark, and $\psi(r, \phi)$ is encoded by color.

The self-healing of the intensity and spatially inhomogeneous state of polarization of the scalar (linear polarized) Bessel beam was experimentally measured as it propagated through the obstruction. The results are shown in the first row of figure 3(a). As can be seen, as the scalar Bessel beam propagated through the obstruction from (I) to (IV), its intensity self-heals as is expected [4–7]. Next, the self-healing of the intensities and spatially inhomogeneous states of polarizations of the radially and azimuthally polarized vector Bessel beams were experimentally measured as they propagated through the obstruction. The results are shown in the second and third rows of figure 3(a), respectively. As the radially and azimuthally polarized vector Bessel beam propagated through the obstruction from (I) to (IV), their intensities self-healed similar to the scalar Bessel beam. As can be seen, similar to their intensities, the spatially inhomogeneous states of polarization of the radially and azimuthally polarized vector Bessel beams also self-healed. This is the salient result of this letter. There is qualitative agreement between the spatially inhomogeneous states of polarization of the vector Bessel beams when they are unobstructed at (I) and when they self-heal at (IV).

Also, the self-healing of the intensities and spatially inhomogeneous states of polarization of the radially and azimuthally polarized vector Bessel beams were experimentally measured as they propagated through a disparate obstruction. The results are shown in figure 3(b). As shown in figure 3(b), the size and position of this obstruction was chosen such that it obstructed a larger portion of the vector Bessel beams near their center. The obstruction is outlined by a dashed white line. As can be seen, the intensities and the spatially inhomogeneous states of polarization of the radially and azimuthally polarized vector Bessel beams again self-heal in the presence of the larger obstruction. Similar to the first obstruction, there is qualitative agreement between the spatially inhomogeneous states of polarization of the vector Bessel beams when they are unobstructed at (I) and when they self-heal at (IV).

Similar, to scalar Bessel beams, the self-healing of vector Bessel beams can be understood via geometric optics, i.e., the interference of conical rays in the shadow region of the obstruction, as shown in figure 1(e). It is a well-known that the distance in which a Bessel beam is able to reform is given by [3]:

$$z_{\min} \approx \frac{ak}{2k_z},$$
 (6)

where *a* is the width of the obstruction and *k* and k_z are the wave-vector and longitudinal wave-vector, respectively. Equation (6) illustrates that the distance in which a Bessel beam self-heals is dependent on the size and position of the obstruction as well as the opening angle of the cone on which the wave-vectors of the Bessel beam propagate. Here it is assumed that the input field is larger than the obstruction.

There is a relationship between light's space and polarization degrees of freedom when light is scattered by an obstruction, e.g. Rayleigh or Mie particles [42]. There are comparable relationships in multimode optical fiber [43, 44]. In this respect, a more detailed theoretical analysis of selfhealing of the spatially inhomogeneous states of polarization



Figure 3. Experimentally measured intensities and spatially inhomogeneous states of polarization of scalar (linearly polarized) and vector Bessel beams as they propagated through two different obstructions at four different propagation distances. The obstruction is outlined by a white dashed line. (Column I) unobstructed. (Column II) obstructed. (Column III) semi-healed. (Column IV) self-healed. The Bessel beams' total intensities, $S_0(r, \phi)$, are encoded by relative contrast, and overlaid with the orientation of their states of polarization, $\psi(r, \phi)$, encoded by color. (a) First obstruction (first row) scalar Bessel beam. (Second row) radial polarization (third row) azimuthal polarization. (b) Second obstruction (first row) radial polarization. (Second row) azimuthal polarization.

of radially and azimuthally polarized vector Bessel beams, analogous to that of scalar Bessel beams [3], is the subject of future work. Nonetheless, as vector Bessel beams possess the properties of Bessel beams and a vector beams, they may have applications in, for example, optical trapping, where self-healing and 'vectorness' are both needed, e.g. it may be possible to the improve the axial and transverse stiffness of a tractor beam when using a vector Bessel beam.

In conclusion, we experimentally measured the selfhealing of the spatially inhomogeneous states of polarization of vector Bessel beams. Radially and azimuthally polarized vector Bessel beams were experimentally generated via a digital version of Durnin's method, using an SLM in concert with a liquid crystal *q*-plate. As a proof of principle, their intensities and spatially inhomogeneous states of polarization were measured using Stokes polarimetry as they propagated through two disparate obstructions. It was found, similar to their intensities, that their spatially inhomogeneous states of polarization self-healed. The self-healing can be understood via geometric optics, i.e., the interference of the unobstructed

conical rays in the shadow region of the obstruction, and may have applications in, for example, optical trapping.

While there are extensive studies on self-healing of scalar Bessel beams [4–7], there are limited studies on self-healing of vector Bessel beams [24–27], particularly with respect to their spatially inhomogeneous states of polarization. Previous work only measured the propagation and self-healing of the *intensities* of vector Bessel beams. To our knowledge, this is the first experimental measurement of the self-healing of the spatially inhomogeneous states of *polarization* of radially and azimuthally polarized vector Bessel beams. Future work includes experimentally measuring the self-healing of the spatially inhomogeneous states of *polarization* of other types of vector Bessel beams such as full Poincaré beams [45]. In contrast to radially and azimuthally polarized light beams, full Poincaré beams experience non-trivial dynamics as they propagate [46, 47].

Acknowledgments

We thank S Slussarenko and L Marrucci for the *q*-plates. GM acknowledges helpful discussions with D Grier and D Ruffner, and support from AFOSR Grant. No. 47221-00-01, ARO Grant. No. 52759-PH-H, NSF GRFP Grant. No. 40017-00-04, and Corning, Inc. E K acknowledges the support of the Canada Excellence Research Chairs (CERC) Program.

References

- Durnin J 1987 Exact solutions for nondiffracting beams: I. The scalar theory J. Opt. Soc. Am. A 4 651
- [2] Durnin J, Miceli J J Jr and Eberly J H 1987 Diffraction-free beams Phys. Rev. Lett. 58 1499
- [3] Litvin I A, McLaren M G and Forbes A 2009 A conical wave approach to calculating Bessel–Gauss beam reconstruction after complex obstacles *Opt. Commun.* 282 1078–82
- [4] Dudley A, Lavery M, Padgett M and Forbes A 2013
 Unraveling Bessel beams Opt. Photon. News 24 2229
- [5] Mazilu M, Stevenson J D, Gunn-Moore F and Dholakia K 2010 Light beats the spread: 'non-diffracting' beams *Laser Photon. Rev.* 4 529
- [6] Mcgloin D and Dholakia K 2005 Bessel beams: diffraction in a new light *Contemp. Phys.* 46 15
- [7] Duocastella M and Arnold C B 2012 Bessel and annular beams for materials processing *Laser Photon. Rev.* 6 607–21
- [8] Garcs-Chvez V, McGloin D, Melville H, Sibbett W and Dholakia K 2002 Simultaneous micromanipulation in multiple planes using a self-reconstructing light beam *Nature* 419 145–7
- [9] Sukhov S and Dogariu A 2011 Negative nonconservative forces: optical tractor beams for arbitrary objects *Phys. Rev. Lett.* 107 203602
- [10] Novitsky A, Qiu C-W and Wang H 2011 Single gradientless light beam drags particles as tractor beams *Phys. Rev. Lett.* 107 203601
- [11] Novitsky A, Qiu C-W and Lavrinenko A 2012 Materialindependent and size-independent tractor beams for dipole objects *Phys. Rev. Lett.* **109** 023902
- [12] Ruffner D B and Grier D G 2012 Optical conveyors: a class of active tractor beams *Phys. Rev. Lett.* **1109** 163903

- [13] Zhan Q 2009 Cylindrical vector beams: from mathematical concepts to applications Adv. Opt. Photon. 1 1–57
- [14] Brown T G 2011 Unconventional polarization states: beam propagation, focusing and imaging *Prog. Opt.* 56 81
- [15] Youngworth K and Brown T 2000 Focusing of high numerical aperture cylindrical-vector beams Opt. Express 7 77–87
- [16] Novotny L, Beversluis M R, Youngworth K S and Brown T G 2001 Longitudinal field modes probed by single molecules *Phys. Rev. Lett.* 86 5251
- [17] Dorn R, Quabis S and Leuchs G 2003 Sharper focus for a radially polarized light beam *Phys. Rev. Lett.* 91 233901
- [18] Nieminen T, Heckenberg N and Rubinsztein-Dunlop H 2008 Forces in optical tweezers with radially and azimuthally polarized trapping beams *Opt. Lett.* 33 122–4
- [19] Roxworthy B J and Toussaint K C Jr 2010 Optical trapping with π-phase cylindrical vector beams New J. Phys. 12 073012
- [20] Kozawa Y and Sato S 2010 Optical trapping of micrometersized dielectric particles by cylindrical vector beams Opt. Express 18 10828–33
- [21] Milione G et al 4 × 20 Gbit/s mode division multiplexing over free space using vector modes and a q-plate mode (de) multiplexer arXiv:1412.2717
 - Lavery M P J *et al* 2014 Space division multiplexing in a basis of vector modes *Proc. Ecoc.* 2014 doi:10.1109/ ECOC.2014.6964136
- [22] Bouchal Z and Olivik M 1995 Non-diffractive vector Bessel beams J. Mod. Opt. 42 1555–66
- [23] Ornigotti M and Aiello A 2013 Radially and azimuthally polarized nonparaxial Bessel beams made simple *Opt. Express* 21 15530–7
- [24] Niv A, Biener G, Kleiner V and Hasman E 2004 Propagationinvariant vectorial Bessel beams obtained by use of quantized Pancharatnam–Berry phase optical elements *Opt. Lett.* 29 238–40
- [25] Vyas S, Kozawa Y and Sato S 2011 Self-healing of tightly focused scalar and vector Bessel–Gauss beams at the focal plane J. Opt. Soc. Am. A 28 837–43
- [26] He M, Chen Z, Sun S and Pu J 2013 Propagation properties and self-reconstruction of azimuthally polarized nondiffracting beams *Opt. Commun.* 294 36–42
- [27] Wu G, Wang F and Cai Y 2014 Generation and self-healing of a radially polarized Bessel–Gauss beam *Phys. Rev.* A 89 043807
- [28] Vasilyeu R, Dudley A, Khilo N and Forbes A 2009 Generating superpositions of higher-order Bessel beams Opt. Express 17 23389
- [29] Lin J, Genevet P, Kats M A, Antoniou N and Capasso F 2013 Nanostructured holograms for broadband manipulation of vector beams *Nano. Lett.* 13 4269–74
- [30] Sztul H I, Nolan D A, Milione G, Chen X, Koh J and Alfano R R 2009 Cylindrical vector beam generation from spun fiber *Proc. SPIE* 7227 722704
- [31] Milione G, Sztul H I and Alfano R R 2010 Cylindrical vector beam generation from a multi elliptical core optical fiber *Proc. SPIE* 7613 761305
- [32] Milione G, Sztul H, Nolan D, Kim J, Etienne M, McCarthy J, Wang J and Alfano R 2011 Cylindrical vector beam generation from a multi elliptical core optical fiber *CLEO:2011—Laser Applications to Photonic Applications OSA Technical Digest (CD)* (Optical Society of America) paper CTuB2 doi:10.1364/CLEO_S1.2011.CTuB2
- [33] Dudley A, Li Y, Mhlanga T, Escuti M and Forbes A 2013 Generating and measuring nondiffracting vector Bessel beams *Opt. Lett.* 38 3429
- [34] Marrucci L, Manzo C and Paparo D 2006 Optical spin-toorbital angular momentum conversion in inhomogeneous anisotropic media *Phys. Rev. Lett.* **96** 163905

- [35] Cardano F, Karimi E, Slussarenko S, Marrucci L, de Lisio C and Santamato E 2012 Polarization pattern of vector vortex beams generated by *q*-plates with different topological charges *Appl. Opt.* **51** C1
- [36] Milione G, Sztul H I, Nolan D A and Alfano R R 2011 Higherorder Poincaré sphere, Stokes parameters, and the angular momentum of light *Phys. Rev. Lett.* **107** 053601
- [37] Milione G, Evans S, Nolan D A and Alfano R R 2012 Higher order Pancharatnam–Berry phase and the angular momentum of light *Phys. Rev. Lett.* **108** 190401
- [38] Slussarenko S, Murauski A, Du T, Chigrinov V, Marrucci L and Santamato E 2011 Tunable liquid crystal q-plates with arbitrary topological charge Opt. Express 19 4085
- [39] Rumala Y, Milione G, Nguyen T, Pratavieira S, Hossain Z, Nolan D, Slussarenko S, Karimi E, Marrucci L and Alfano R 2013 Tunable supercontinuum light vector vortex beam generator using a *q*-plate Opt. Lett. 38 5083
- [40] Goldstein D 2003 Polarized Light (Boca Raton, FL: CRC Press)
- [41] Dudley A, Milione G, Alfano R R and Forbes A 2014 Alldigital wavefront sensing for structured light beams *Opt. Express* 22 14031–40

- [42] Shumyatsky P, Milione G and Alfano R R 2014 Optical memory effect from polarized Laguerre Gaussian light beam in light-scattering turbid media *Opt. Commun.* 321 116–23
- [43] Alexeyev C N, Alexeyev A N, Lapin B P, Milione G and Yavorsky M A 2013 Spin–orbit-interaction-induced generation of optical vortices in multihelicoidal fibers *Phys. Rev.* A 88 063814
- [44] Milione G, Nolan D A and Alfano R R 2015 Determining principal modes in a multimode optical fiber using the mode dependent signal delay method J. Opt. Soc. Am. B 32 143–9
- [45] Beckley A, Brown T and Alonso M 2010 Full Poincare beams Opt. Express 18 10777–85
- [46] Beckley A, Alonso M and Brown T 2010 Diffraction free Stokes distributions in a full Poincare beam Frontiers in Optics 2010/Laser Science XXVI OSA Technical Digest (CD) (Optical Society of America) paper FThN2 doi:10.1364/FIO.2010.FThN2
- [47] Philip G M, Kumar V, Milione G and Viswanathan N K 2012 Manifestation of the Gouy phase in vector-vortex beams *Opt. Lett.* **37** 2667