BANYAN. VII. A NEW POPULATION OF YOUNG SUBSTELLAR CANDIDATE MEMBERS OF NEARBY MOVING GROUPS FROM THE BASS SURVEY

JONATHAN GAGNÉ¹, JACQUELINE K. FAHERTY^{2,3,9}, KELLE L. CRUZ^{3,4,5}, DAVID LAFRENIÉRE¹, RENÉ DOYON¹, LISON MALO^{1,6},

Adam J. Burgasser⁷, Marie-Eve Naud¹, Étienne Artigau¹, Sandie Bouchard¹, John E. Gizis⁸, and Loïc Albert¹ Institut de Recherche sur les Exoplanètes (iREx), Université de Montréal, Département de Physique,

C.P. 6128 Succ. Centre-ville, Montréal, QC H3C 3J7, Canada; jonathan.gagne@astro.umontreal.ca
 ² Department of Terrestrial Magnetism, Carnegie Institution of Washington, Washington, DC 20015, USA

³ Department of Astrophysics, American Museum of Natural History, Central Park West at 79th Street, NY 10024, USA

⁴ Department of Physics & Astronomy, Hunter College, City University of New York, 695 Park Avenue, NY 10065, USA

⁵ Department of Physics, Graduate Center, City University of New York, 365 Fifth Avenue, New York, NY 10016, USA ⁶ Canada–France–Hawaii Telescope, 65-1238 Mamalahoa Hwy, Kamuela, HI 96743, USA

⁷ Center for Astrophysics and Space Sciences, University of California, San Diego, 9500 Gilman Dr., Mail Code 0424, La Jolla, CA 92093, USA

Department of Physics and Astronomy, University of Delaware, 104 The Green, Newark, DE 19716, USA

Received 2015 April 2; accepted 2015 June 25; published 2015 August 21

ABSTRACT

We present the results of a near-infrared (NIR) spectroscopic follow-up survey of 182 M4-L7 low-mass stars and brown dwarfs (BDs) from the BANYAN All-Sky Survey (BASS) for candidate members of nearby, young moving groups (YMGs). We confirm signs of low gravity for 42 new BD discoveries with estimated masses between 8 and $75 M_{Jup}$ and identify previously unrecognized signs of low gravity for 24 known BDs. We refine the fraction of low-gravity dwarfs in the high-probability BASS sample to \sim 82%. We use this unique sample of 66 young BDs, supplemented with 22 young BDs from the literature, to construct new empirical NIR absolute magnitude and color sequences for low-gravity BDs. We show that low-resolution NIR spectroscopy alone cannot differentiate between the ages of YMGs younger than ~ 120 Myr, and that the BT-Settl atmosphere models do not reproduce well the dust clouds in field or low-gravity L-type dwarfs. We obtain a spectroscopic confirmation of low gravity for 2MASS J14252798–3650229, which is a new \sim 27 M_{Jup} , L4 γ bona fide member of AB Doradus. We identify a total of 19 new low-gravity candidate members of YMGs with estimated masses below 13 M_{Jup} , 7 of which have kinematically estimated distances within 40 pc. These objects will be valuable benchmarks for a detailed atmospheric characterization of planetary-mass objects with the next generation of instruments. We find 16 strong candidate members of the Tucana-Horologium association with estimated masses between 12.5 and $14 M_{Jup}$, a regime where our study was particularly sensitive. This would indicate that for this association there is at least one isolated object in this mass range for every $17.5^{+6.6}_{-5.0}$ main-sequence stellar member, a number significantly higher than expected based on standard log-normal initial mass function, however, in the absence of radial velocity and parallax measurements for all of them, it is likely that this over-density is caused by a number of young interlopers from other associations.

Key words: brown dwarfs – methods: data analysis – proper motions – stars: kinematics and dynamics – stars: low-mass

Supporting material: data behind figures, machine-readable tables

1. INTRODUCTION

Young moving groups (YMGs) consist of stars that formed recently (≤ 120 Myr) from a molecular cloud and that are too young to have experienced significant gravitational perturbations from their environment. The members of YMGs share similar galactic velocities within a few km s^{-1} . The closest and youngest moving groups include the TW Hydrae association (TWA; 5–15 Myr; de la Reza et al. 1989; Kastner et al. 1997; Zuckerman & Song 2004; Weinberger et al. 2013), β Pictoris (\beta PMG; 20-26 Myr; Zuckerman et al. 2001a; Binks & Jeffries 2014; Malo et al. 2014b), Tucana-Horologium (THA; 20-40 Myr; Torres et al. 2000; Zuckerman & Webb 2000; Zuckerman et al. 2001b; Kraus et al. 2014b), Carina (CAR; 20-40 Myr; Torres et al. 2008), Columba (COL; 20-40 Myr; Torres et al. 2008), Argus (ARG; 30-50 Myr; Makarov & Urban 2000), and AB Doradus (ABDMG; 110-130 Myr; Zuckerman et al. 2004; Luhman et al. 2005;

Barenfeld et al. 2013). YMGs are ideal laboratories to measure fundamental properties of star formation such as the initial mass function (IMF) because their members are coeval. This is of particular interest in the case of very low-mass stars and substellar-mass objects (spectral types ≥M5) since these populations are still poorly characterized. The massive, bright population of YMGs has already been explored, thanks to the Hipparcos survey (Perryman et al. 1997). However, fainter members are hard to identify mainly because of the lack of radial velocity (RV) and trigonometric distance measurements that are necessary to obtain their spacial velocities and galactic positions. Several efforts have been made to identify the very low-mass members of YMGs (Kiss et al. 2011: Faherty et al. 2012, 2013; Schlieder et al. 2012b; Shkolnik et al. 2012; Liu et al. 2013b; Rodriguez et al. 2013; Kraus et al. 2014b; Malo et al. 2014a; Riedel et al. 2014; Murphy & Lawson 2015); however, as of today, it is likely that most of them still remain to be identified.

⁹ Hubble Fellow.

The Bayesian Analysis for Nearby Young AssociatioNs tool¹⁰ (BANYAN; Malo et al. 2013), which is based on naive Bayesian inference, identified promising candidate members of YMGs among a sample of low-mass stars that do not have prior RV or parallax measurements. The BANYAN II tool¹¹ (Gagné et al. 2014c; Paper II hereafter) was subsequently developed to identify substellar candidate members with a similar but improved algorithm. BANYAN II is an expansion on BANYAN I that is focused on very-low mass stars and brown dwarfs (BDs) with spectral types \geq M5. The BANYAN All-Sky Survey (BASS; Gagné et al. 2015; Paper V hereafter) was initiated by our team to search for the elusive late-type ($\geq M5$) members of YMGs, using the BANYAN II tool on an all-sky cross-match of the 2MASS (Skrutskieet al. 2006) with the AllWISE survey (Kirkpatrick et al. 2014). The AllWISE survey is based on a combination of the cryogenic phase of the Widefield Survey Explorer mission (WISE; Wright et al. 2010) and the Near-Earth Object WISE (Mainzeret al. 2011) postcryogenic phase.

We present here the results of a near-infrared (NIR) spectroscopic follow-up survey of substellar candidate members of YMGs identified in BASS. In Section 2, we summarize BASS and the method that we used to build the sample of candidate members from a cross-match of 2MASS and AllWISE. We detail our NIR spectroscopic follow-up and its motivation in Section 3. In Section 4, we present our method to assign a spectral and gravity classification. We present the resulting spectral types and updated YMG membership probability for our sample in Section 5. In Section 6, we use new discoveries presented here and other known low-gravity BDs and low-mass stars to build empirical photometric sequences, and we then investigate the physical properties of young BDs. We summarize and conclude in Section 7.

2. THE BASS SURVEY

BASS is a systematic all-sky search for later-than-M5 candidate members to nearby YMGs that was the focus of an earlier publication (Gagné et al. 2015). In this work, we undertake a spectroscopic follow-up of the BASS sample, which we briefly summarize in this section. We refer the reader to (Gagné et al. 2015) for an extensive description of the BASS survey.

We cross-matched the 2MASS and AllWISE catalogs outside of the galactic plane and crowded regions (>2.5 objects per square arcminute) using a cross-match radius of 25" and applied color, confusion and photometric quality cuts to produce a starting sample of 98 970 targets with NIR colors consistent with ≥M5 spectral types and proper motion measurements larger than 30 mas yr⁻¹ at $\geq 5\sigma$ (see Gagné et al. 2015 for the detailed crossmatch and selection algorithm). Astrometry provided in the 2MASS and AllWISE catalogs as well as the mean epochs of observation for both surveys (JD keyword in 2MASS; WIMJDMEAN keyword in AllWISE) were used to calculate proper motions. We used W1MJDMAX-W1MJDMIN as a conservative measurement error on the AllWISE astrometric epoch, which typically corresponds to ~ 6 months to one year, compared to a ~11 years baseline between 2MASS and AllWISE. This uncertainty as well as those on astrometric measurements themselves were propagated to the proper motion

contaminants; (2) *Low-Priority BASS* (LP-BASS) consists of targets that have NIR colors only slightly redder than field dwarfs; and (3) BASS is the final sample presented in Paper V that contains targets at least 1σ redder than field dwarfs and that

measurement errors. We obtain a typical proper motion precision of $\sim 15 \text{ mas yr}^{-1}$.

We used the BANYAN II tool to select only objects that have a Bayesian probability >10% of belonging to any YMG considered here (this threshold ensures that known bona fide members are recovered; see Paper V). The BANYAN II tool takes the sky position, proper motion and J, H, K_s , W1 and W2 photometry as input quantities. It then uses a naive Bayesian classifier to compare those measurements with spatial and kinematic models (SKMs) of YMGs, as well as with old and young color-magnitude diagram (CMD) sequences in both $M_{W1}(J - K_S)$ and $M_{W1}(H - W2)$ spaces. Those CMD sequences were chosen because they were found as the most efficient independent sequences to distinguish between young and field M6-L4 dwarfs. Probabilities generated from a naive Bayesian classifier can be biased when the input parameters are not independent (which is the case here); however, the relative ranking of hypotheses for a given object overcomes this bias (Hand & Yu 2001).

It is known that there is a large scatter in the NIR colors of young BDs even though they are redder than field dwarfs on average (e.g., Faherty et al. 2012). The inclusion of the CMD sequences described above in BANYAN II will systematically bias our sample toward red NIR colors, and decrease our sample completeness for YMG members that are not especially red. However, this effect is likely less important than the color criteria that were applied in selecting the 98,970 objects that were input to BANYAN II. Furthermore, a total of only two independent photometric observables (corresponding to the CMDs) are used in BANYAN II, compared to four kinematics observables when no RV or parallax is available; the relative weight of kinematics is thus twice that of photometry in the calculation of probabilities. Parallax motion was not accounted for in our proper motion measurements or in the BANYAN II tool; the maximal relative importance of this effect will become as large as our typical 2MASS-AllWISE proper motion precision only for objects closer than $\sim 10 \text{ pc}$ (considering the 11 yr baseline between 2MASS and AllWISE). This correction will properly be accounted for in a future version of the BANYAN II tool.

We performed a Monte Carlo simulation based on the Besançon galactic model (Robin et al. 2012; A. C. Robin et al. 2015, in preparation) and the SKMs of YMGs to obtain a field contamination probability for each individual target in our sample, which allows for a more absolute interpretation in terms of the expected contamination fraction. We used the results of this simulation to reject any candidate member with a >50% probability from this Monte Carlo analysis is not necessarily complementary with the YMG Bayesian probability (see Paper V for more detail). We refer the reader to Paper V for an extensive description of all filters that were used to build the BASS sample (e.g., minimal proper motion, color and quality filters, etc.).

¹⁰ Publicly available at http://www.astro.umontreal.ca/~malo/banyan.php.

¹¹ Publicly available at http://www.astro.umontreal.ca/~gagne/banyanII.php.

There are three samples that are referred to in this Paper: (1) PRE-BASS consists of targets that were initially selected as potential members and followed up with spectroscopy, but that were later rejected as we modified our selection criteria to reject

has a lower fraction of contaminants. As discussed in Paper V, the statistical distance associated with the most probable YMG of a candidate member can be used to place it in two CMDs $(M_{W1} \text{ versus } J - K_S \text{ and } H - W2)$ and compare its position to known field and young BDs and low-mass stars.

All candidate members that were placed blueward of the field sequence in any of the two CMDs were rejected from BASS and LP-BASS. Those that were not at least 1σ redder than both field sequences were grouped into the LP-BASS sample, which is expected to be more contaminated by field objects and young M dwarfs with spectral types earlier than M5. We note that the PRE-BASS sample does not necessarily consist of erroneous YMG candidate members; however, it likely suffers from a higher contamination rate from field interlopers or members of moving groups not considered in BANYAN II.

3. OBSERVATIONS

Because of their recent formation, young, low-mass objects have inflated radii compared to their field counterparts and are warmer for a given mass. As a consequence, they have a lower surface gravity at a given temperature (and spectral type). It is well known that these low-gravity dwarfs display weaker alkali and molecular absorption lines (K1 at 7665 and 7669 Å in the optical and 1.17 and 1.25 μ m in the NIR; Na1 at 8183 and 8195 Å in the optical and 1.14 and 2.21 μ m in the NIR; Rb1 at 7800 and 7948 Å; Cs I at 8521 and 8943 Å; FeH at 8692 Å in the optical and 0.99, 1.20 and 1.55 μ m in the NIR; TiO at 8432 Å; and CrH at 8611 Å). This is due to a lower-pressure in their photosphere, which is a direct consequence of their lower surface gravity. Collision-induced absorption (CIA) of the H₂ molecule is also decreased in this lower pressure environment, causing a flatter K-band plateau at 2.18–2.28 μ m (see the H₂(K) index of Canty et al. 2013), leaving the effect of water vapor to become apparent from the triangular-shaped continuum of the H band (Lucas et al. 2001; Kirkpatrick et al. 2006; Allers et al. 2007; Rice et al. 2010; see the H-cont index of Allers & Liu 2013). Furthermore, VO condensate clouds get thicker in the external layers of low-pressure atmospheres, causing deeper absorption bands at 7300-7550 and 7850-8000 Å in the optical and 1.06 μ m in the NIR (These effects are discussed in more detail by Gorlova et al. 2003; McGovern et al. 2004; Kirkpatrick et al. 2006, 2008; Cruz et al. 2009; Allers & Liu 2013 and Canty et al. 2013). Gravity-sensitive features were initially identified by comparing the optical spectra of M-type giants and M-type dwarfs (Kleinmann & Hall 1986; Joyce et al. 1998), and it was later demonstrated that the same features could be used to identify young, inflated M-type dwarfs by observing members of star-forming regions (Martín et al. 1996; Luhman et al. 1997; Slesnick et al. 2004; Lucas et al. 2001; Allers et al. 2007; Lodieu et al. 2008).

A number of low-gravity features (CIA effects of H₂ on the continuum, weaker FeH absorption and stronger VO absorption) can be measured in low-mass stars and BDs with spectral types later than M6 using low-resolution ($R \sim 75$) NIR spectroscopy, providing an efficient way of identifying field interlopers in a set of YMG candidates. A higher spectral resolution ($R \sim 1000$) allows for a more robust determination of low-gravity features through the measurement of the pseudo-equivalent width (EW) of the atomic lines listed above. We thus obtained low-resolution NIR spectra of 241 candidate YMG members from the BASS, LP-BASS, and PRE-BASS samples. We describe in this section all observations and the

individual instrumental configurations that were used. A description of individual observations is included in Table 1.

3.1. FIRE at Magellan

We obtained NIR spectroscopy for 17 targets with the Folded-port InfraRed Echellette (FIRE: Simcoe et al. 2008, 2013) at the Magellan Telescopes in 2013 April and December, as well as 2014 May, June, August and September and 2015 February. We used both the crossdispersed and high-throughput prism modes to obtain respective resolving powers $R \sim 450$ (prism mode) and $R \sim 6000$ (echelle mode) across the 0.8–2.45 μ m range. Total exposure times ranged from 200 to 1800s, depending on source brightness, instrument configuration and weather conditions. This allowed us to obtain a typical signal-to-noise ratio (S/N) > 100 per resolution element. Science targets were observed in an ABBA pattern along the slit, and a standard A0type star was observed immediately before or after each of them at a similar airmass to ensure a proper telluric correction. We obtained ThAr (prism mode) or NeNeAr (echellette mode) lamp exposures between every science target to perform wavelength calibration, as well as high- and low-illumination flat fields that were combined to obtain a flat-field image with a large S/N across all orders while avoiding saturation. We reduced all data using the Interactive Data Language (IDL) pipeline FIREHOSE, which is based on the MASE (Bochanski et al. 2009) and SpeXTool (Vacca et al. 2003; Cushing et al. 2004) packages. We supplemented the list of Ar atomic lines with those listed in Norlén (1973) to allow a more robust wavelength solution in the K band in the case of prism data.

The six echellette spectra that we obtained here have a sufficient resolution to measure radial velocities down to a precision down to a few km s⁻¹. These measurements will be presented in a future publication along with a significant number of additional FIRE echellette spectra.

3.2. SpeX at IRTF

We obtained NIR spectroscopy with SpeX (Rayner et al. 2003) at the IRTF telescope for 118 targets from 2007 to 2015. We used the cross-dispersed and prism modes with slits of 0".6, 0".8, and 1".0 depending on the seeing to obtain resolving powers ranging from $R \sim 75$ to $R \sim 750$ over the 0.8–2.45 μ m range. We used ABBA nodding patterns along the slit with typical exposure times of 60–250 s which yielded typical S/N > 100 per resolution element. A standard early A-type star was observed immediately before or after every science target at a similar airmass to ensure a proper telluric correction. Several high-S/N quartz lamp and Ar lamp exposures were obtained immediately after every target to ensure a proper wavelength calibration and flat field correction. The data were reduced with the IDL SpeXTool package (Vacca et al. 2003; Cushing et al. 2004).

3.3. Flamingos-2 at Gemini-South

We used Flamingos-2 (Eikenberry et al. 2004) at Gemini-South to obtain NIR spectroscopy for 101 targets from 2013 to 2015. We observed each target with both the *JH* and *HK* low resolution grisms and the 0".72 slit to obtain a resolving power of $R \sim 500$ over 0.9–2.4 μ m. Targets were observed in an ABBA pattern along the slit, with total exposure times ranging from 120 to 3400 s, to obtain S/N > 80 per resolution element.

		Ot	oserving Log					
2MASS		Observing	Telescope	Instrument	Slit	Resolving	Tot. Exp.	Num.
Designation	J	Date (UT)			Width (")	Power	Time (s)	Exposures
		Candidate Membe						
00011217+1535355	15.52	2014 Aug 06	Magellan	FIRE	0.6	450	720	6
00065794-6436542	13.39	2014 Jan 16	Gemini-South	Flamingos-2	0.72	500	840	4
00182834-6703130	15.46	2013 Sep 28	Gemini-South	Flamingos-2	0.72	500	2000	8
00191296-6226005	15.64	2013 Oct 29	Gemini-South	Flamingos-2	0.72	500	6200	20
00192626+4614078	12.60	2012 Oct 26	IRTF	SpeX	0.8	750	1200	8
00274534-0806046	11.57	2013 Oct 02	Gemini-South	Flamingos-2	0.72	500	240	8
00344300-4102266	15.71	2014 Oct 12	Gemini-South	Flamingos-2	0.72	500	8575	35
00381489-6403529	14.52	2014 May 31	Gemini-South	Flamingos-2	0.72	500	2080	8
00390342+1330170 A & B	10.94	2013 Jul 31	IRTF	SpeX	0.8	750	900	6
00413538-5621127	11.96	2013 Nov 25 & 2014 Jan 21	Gemini-South	Flamingos-2	0.72	500	360	12
00464841+0715177	13.89	2013 Aug 11	IRTF	SpeX	0.8	750	1600	8
00514561-6227073	12.58	2014 Jan 16	Gemini-South	Flamingos-2	0.72	500	1680	8
00584253-0651239	14.31	2013 Aug 11	IRTF	SpeX	0.8	750	1600	8
01205114-5200349	15.64	2013 Sep 27	Gemini-South	Flamingos-2	0.72	500	6200	20
01265327–5505506	12.04	2014 Jan 21	Gemini-South	Flamingos-2	0.72	500	240	8
01294256-0823580	10.65	2013 Oct 02	Gemini-South	Flamingos-2	0.72	500	240	8
01344601–5707564	12.07	2013 Nov 25	Gemini-South	Flamingos-2	0.72	500	240	8
01484859–5201158	10.87	2013 Oct 02	Gemini-South	Flamingos-2	0.72	500	240	8
01531463-6744181	16.41	2013 Apr 21	Magellan	FIRE	0.6	6000	1500	2
02103857-3015313	15.07	2013 Dec 16	Gemini-South	Flamingos-2	0.72	500	1680	8
02265658-5327032	15.40	2013 Oct 02	Gemini-South	Flamingos-2	0.72	500	2400	8
02282694+0218331	12.12	2013 Oct 18	Gemini-South	Flamingos-2	0.72	500	240	8
02404759-4253377	12.20	2013 Oct 28	Gemini-South	Flamingos-2	0.72	500	240	8
02410564-5511466	15.39	2014 Dec 10, 12 & 15	Gemini-South	Flamingos-2	0.72	500	10605	42
02441019-3548036	15.34	2013 Oct 19	Gemini-South	Flamingos-2	0.72	500	5280	16
02501167-0151295	12.89	2014 Aug 02 & 03	Gemini-South	Flamingos-2	0.72	500	2920	16
02534448-7959133	11.34	2013 Oct 27	Gemini-South	Flamingos-2	0.72	500	240	8
02002877 2014252	15.91	2013 Dec 12 2012 Oct 10	Magellan Comini South	FIKE	0.6	500	1310	2
03132588 2447246	11.30	2013 Oct 19 2013 Dec 16	Gemini-South	Flamingos-2	0.72	500	240	0
03182507 3708118	12.35	2013 Dec 10 2013 Oct 28	Gemini-South	Flamingos-2	0.72	500	1680	0
03204010 3313400	12.57	2013 Oct 28	Gemini-South	Flamingos-2	0.72	500	240	0
03224919=3515400	12.34	2013 Oct 19	Gemini-South	Flamingos-2	0.72	500	240	8
03264225_2102057	16.13	2013 Oct 17 2007 Nov 13	IRTE	SneX	0.72	150	180	6
03333313_3215181	13.17	2007 Nov 15 2013 Aug 11	IRTE	SpeX	0.5	750	1200	8
03363144_2619578	10.68	2013 Aug 11 2012 Feb 05	IRTE	SpeX	0.8	750	600	4
03370359_1758079	15.62	2012 Feb 05	IRTE	SpeX	0.5	150	1440	8
03370362-3709236	12.75	2011 Dec 00 2013 Aug 11	IRTE	SpeX	0.8	750	1200	8
03390160-2434059	10.90	2012 Feb 05	IRTE	SpeX	0.8	750	600	4
03420931-2904317	15.92	2013 Apr 21	Magellan	FIRE	0.6	6000	1204	2
03550477-1032415	13.08	2013 Oct 20	Gemini-South	Flamingos-2	0.72	500	1680	8
03552337+1133437	14.05	2013 Feb 16	IRTF	SpeX	0.8	750	1440	8
03582255-4116060	15.85	2011 Dec 08	IRTF	SpeX	0.5	150	1440	8
04185879-4507413	16.16	2014 Dec 10 & 15	Gemini-South	Flamingos-2	0.72	500	7820	31
04231498-1533245	12.54	2014 Sep 16	Gemini-South	Flamingos-2	0.72	500	3240	27
04400972-5126544	15.69	2013 Oct 27	Gemini-South	Flamingos-2	0.72	500	5600	20
04433761+0002051	12.51	2012 Feb 05	IRTF	SpeX	0.8	750	600	4
04532647-1751543	15.14	2012 Feb 05	IRTF	SpeX	0.8	750	1800	12
04584239-3002061	13.50	2015 Feb 03	IRTF	SpeX	0.8	90	800	16
05002100+0330501	13.67	2015 Feb 02	IRTF	SpeX	0.8	750	1000	4
05012406-0010452	14.98	2013 Feb 16	IRTF	SpeX	0.8	750	2880	16
05104958-1843548	15.35	2015 Feb 03	IRTF	SpeX	0.8	90	2000	10
05123569-3041067	11.90	2013 Oct 08 & 20	Gemini-South	Flamingos-2	0.72	500	480	16
05181131-3101529	11.88	2012 Feb 05	IRTF	SpeX	0.8	750	600	4
05264316-1824315	12.36	2013 Feb 19	IRTF	SpeX	0.8	750	800	4
05361998-1920396	15.77	2012 Oct 26	IRTF	SpeX	0.8	750	150	1
06022216+6336391	14.27	2008 Jan 08	IRTF	SpeX	0.6	120	540	6
06272161-5308428	16.39	2015 Jan 23	Gemini-South	Flamingos-2	0.72	500	4800	16
07140394+3702459	11.98	2015 Feb 03	IRTF	SpeX	0.8	90	300	8
08095903+4434216	16.44	2008 Jan 12	IRTF	SpeX	0.6	120	1500	10

Table 1

Table 1
(Continued)

2MASS		Observing	Telescope	Instrument	Slit	Resolving	Tot. Exp.	Num.
Designation	J	Date (UT)			Width (")	Power	Time (s)	Exposures
09532126-1014205	13.47	2014 Jan 22	IRTF	SpeX	0.8	750	2000	- 8
10212570-2830427	16.91	2015 Feb 13 & 26	Gemini-South	Flamingos-2	0.72	500	6300	21
10284580-2830374	10.95	2012 May 12	IRTF	SpeX	0.8	750	1200	8
10455263-2819303	12.82	2012 May 12	IRTF	SpeX	0.8	750	1200	8
10513331-1916530	14.69	2014 Jan 22	IRTF	SpeX	0.8	750	4000	16
11064461-3715115	14.49	2012 May 12	IRTF	SpeX	0.8	750	2400	16
11271382-3735076	16.47	2015 Feb 09	Magellan	FIRE	0.6	450	480	4
11480096-2836488	16.11	2015 Feb 13	Gemini-South	Flamingos-2	0.72	500	4800	16
12073346-3932539	12.99	2012 May 12	IRTF	SpeX	0.8	95	1200	8
12074836-3900043	15.49	2013 May 10	IRTF	SpeX	0.8	750	800	4
12214223-4012050	16.47	2015 Feb 08	Magellan	FIRE	0.6	450	1080	9
12310489-3801065	14.68	2015 Feb 08	Magellan	FIRE	0.6	450	240	2
12474428-3816464	14.78	2013 May 10	IRTF	SpeX	0.8	750	720	4
12535039-4211215	16.00	2015 Apr 06	Gemini-South	Flamingos-2	0.72	500	2400	16
12563961-2718455	16.42	2014 May 12	Magellan	FIRE	0.6	6000	1800	2
12574463-3635431	14.57	2014 Aug 03	Gemini-South	Flamingos-2	0.72	500	7595	31
12574941-4111373	13.02	2014 Feb 14	Gemini-South	Flamingos-2	0.72	500	400	8
13262009-2729370	15.85	2009 Jun 30	IRTF	SpeX	0.6	120	1200	8
14252798-3650229	13.75	2010 Jul 07	IRTF	SpeX	0.6	120	720	6
19350976-6200473	16.25	2014 Jul 21 & Aug 03	Gemini-South	Flamingos-2	0.72	500	6060	24
19395435-5216468	14.66	2014 Jun 18	Magellan	FIRE	0.6	450	360	4
20004841-7523070	12.73	2014 Aug 04	Gemini-South	Flamingos-2	0.72	500	1920	16
20113196-5048112	16.42	2014 Jun 18	Magellan	FIRE	0.6	450	480	4
20224803-5645567	11.76	2013 Oct 09	Gemini-South	Flamingos-2	0.72	500	240	8
20282203-5637024	13.84	2014 May 28	Gemini-South	Flamingos-2	0.72	500	800	8
20334473-5635338	15.72	2015 Apr 13	Gemini-South	Flamingos-2	0.72	500	3200	24
20334670-3733443	10.85	2013 Oct 09	Gemini-South	Flamingos-2	0.72	500	240	8
20414283-3506442	14.89	2014 May 28	Gemini-South	Flamingos-2	0.72	500	2480	8
20484222-5127435	15.38	2014 Jun 02	Gemini-South	Flamingos-2	0.72	500	6000	24
20505221-3639552	13.00	2014 Jul 29	Gemini-South	Flamingos-2	0.72	500	7560	42
21121598-8128452	10.67	2013 Oct 20	Gemini-South	Flamingos-2	0.72	500	240	8
21144103-4339531	13.02	2013 Aug 20	IRTF	SpeX	0.8	750	1500	10
21490499-6413039	10.35	2013 Oct 30	Gemini-South	Flamingos-2	0.72	500	240	8
21543454-1055308	16.44	2008 Sep 17	IRTF	SpeX	0.5	120	1800	10
21544859-7459134	14.29	2013 Oct 28	Gemini-South	Flamingos-2	0.72	500	2400	8
22021125-1109461	12.36	2013 Aug 11	IRTF	SpeX	0.8	750	1950	13
22025794-5605087	14.36	2014 Jun 02	Gemini-South	Flamingos-2	0.72	500	1360	8
22064498-4217208	15.56	2013 Aug 01	IRTF	SpeX	0.8	750	450	3
22191486-6828018	13.92	2014 Jul 21 & Aug 02	Gemini-South	Flamingos-2	0.72	500	6240	24
22351658-3844154	15.18	2013 Sep 27	Gemini-South	Flamingos-2	0.72	500	4480	16
22353560-5906306	14.28	2014 Jul 10	Gemini-South	Flamingos-2	0.72	500	640	4
22400144+0532162	11.72	2013 Aug 20	IRTF	SpeX	0.8	750	1500	10
22444835-6650032	11.03	2013 Oct 30	Gemini-South	Flamingos-2	0.72	500	240	8
22511530-6811216	12.10	2013 Oct 28	Gemini-South	Flamingos-2	0.72	500	240	8
23102196-0748531	11.60	2013 Jul 31	IRTF	SpeX	0.8	750	900	6
23130558-6127077	10.93	2013 Oct 21	Gemini-South	Flamingos-2	0.72	500	240	8
23143092-5405313	11.50	2013 Oct 21	Gemini-South	Flamingos-2	0.72	500	240	8
23155665-4747315	16.08	2014 Jun 19	Magellan	FIRE	0.6	450	600	5
23225240-6151114	11.53	2013 Oct 20	Gemini-South	Flamingos-2	0.72	500	240	8
23225299-6151275	15.55	2013 Oct 21	Gemini-South	Flamingos-2	0.72	500	1680	8
23255604-0259508	15.96	2013 Aug 01	IRTF	SpeX	0.8	750	450	3
23270843+3858234	11.74	2013 Jul 31	IRTF	SpeX	0.8	750	1200	8
23290437+0329113	11.11	2012 Oct 26	IRTF	SpeX	0.8	95	800	8
23310161-0406193	12.94	2014 Jul 21 & Aug 02	Gemini-South	Flamingos-2	0.72	500	1920	16
23353085-1908389	11.51	2013 Jul 31	IRTF	SpeX	0.8	750	900	6
23355015-3401477	11.64	2013 Oct 28	Gemini-South	Flamingos-2	0.72	500	240	8
23360735-3541489	14.65	2014 Jul 10	Gemini-South	Flamingos-2	0.72	500	760	4
23433470-3646021	16.57	2014 Jun 19	Magellan	FIRE	0.6	450	480	4
23520507-1100435	12.84	2010 Jul 07	IRTF	SpeX	0.6	120	540	12
23532556–1844402 A & B	11.24	2012 Oct 26	IRTF	SpeX	0.8	750	1040	13

Table 1
(Continued)

				. .		D 11	m , n	
2MASS		Observing	Telescope	Instrument	Slit	Resolving	Tot. Exp.	Num.
Designation	J	Date (UT)	mhare from DDE	DASS	Width (")	Power	Time (s)	Exposures
		Calificate Me	inders nom i KE-	DA33				
00020382+0408129 A & B	10.40	2013 Jul 31	IRTF	SpeX	0.8	750	1200	8
00045753-1709369	11.00	2013 Nov 25	Gemini-South	Flamingos-2	0.72	500	240	8
00171571-3219539	10.64	2013 Nov 25	Gemini-South	Flamingos-2	0.72	500	240	8
00174858-0316334	13.23	2013 Aug 11	IRTF	SpeX	0.8	750	1200	8
00210589-4244433	13.52	2013 Aug 01	IRTF	SpeX	0.8	750	1200	8
00425923+1142104	14.75	2013 Aug 11	IRTF	SpeX	0.8	750	3200	16
00461551+0252004	14.40	2013 Aug 01	IRTF	SpeX	0.8	750	1800	12
01035369–2805518 A & B	11.66	2012 Oct 26	IRTF	SpeX	0.8	750	1200	8
02590146-4232204	12.24	2013 Oct 02	Gemini-South	Flamingos-2	0.72	500	240	8
03005033-5459267	12.42	2013 Nov 25	Gemini-South	Flamingos-2	0.72	500	240	8
03111547+0106307	10.68	2013 Oct 19	Gemini-South	Flamingos-2	0.72	500	240	8
03140344+1603056	12.53	2013 Feb 16	IRTF	Spex	0.8	750	1200	8
03263956-0617586	12.96	2013 Feb 18	IRIF	Spex	0.8	/50	1500	0
03350208+2342356	12.25	2012 Feb 05	IKIF	Spex	0.8	750	600	4
03442859+0/16100 A & B	12.72	2012 Oct 26	IKIF	Spex	0.8	750	1200	8
04044052+2010275 A & B	12.05	2012 Oct 26		Spex	0.8	750	1200	8 12
04070732+1340437	13.46	2012 Feb 03	IKIF Comini South	Spex Eleminade 2	0.8	730 500	240	12
04173830 - 1140230 04281061 + 1820021	11.75	2013 Dec 04 2012 Oct 26	IDTE	Flamingos-2	0.72	500 750	1200	0
04201001+1859021	12.50	2012 Oct 20 2013 Dec 16	IKIT Gemini South	Elemingos 2	0.8	500	480	0
04402383 = 1820414 04403288 ± 1607226	14.05	2013 Dec 10 2012 Eeb 05	IRTE	SpeX	0.72	750	480 600	4
05071137+1430013 A & B	10.57	2012 Feb 16	IRTE	SpeX	0.8	750	600	4
05201794+0511521	13.04	2013 Teb 10 2012 Oct 26	IRTE	SpeX	0.8	750	1200	8
0526179 + 0511521 05243009 + 0640349	11.98	2012 Eet 20 2013 Feb 19	IRTE	SpeX	0.8	750	800	4
05271676+0007526 A & B	12.17	2013 Feb 16	IRTF	SpeX	0.8	750	1200	8
05370704–0623170	15.70	2013 Oct 21	Gemini-South	Flamingos-2	0.72	500	1680	8
05402325-0906326	14.59	2013 Oct 19 & 21	Gemini-South	Flamingos-2	0.72	500	3120	24
05404919-0923192	11.31	2014 Jan 16	Gemini-South	Flamingos-2	0.72	500	240	8
05410983-0737392	13.46	2013 Feb 19	IRTF	SpeX	0.8	750	600	4
05415929-0217020	13.22	2013 Dec 23	Gemini-South	Flamingos-2	0.72	500	240	8
05431887+6422528	13.57	2008 Jan 09	IRTF	SpeX	0.6	120	360	4
05451198-0121021	13.83	2013 Nov 24 & Dec 26	Gemini-South	Flamingos-2	0.72	500	3360	16
05484454-2942551	10.56	2013 Feb 18	IRTF	SpeX	0.8	750	150	1
06021735-1413467	14.34	2013 Feb 19	IRTF	SpeX	0.8	750	1000	4
06353541-6234059	12.42	2013 Oct 20	Gemini-South	Flamingos-2	0.72	500	240	8
06494706-3823284	11.65	2013 Nov 02 & 20	Gemini-South	Flamingos-2	0.72	500	120	4
07083261-4701475	14.16	2013 Oct 29	Gemini-South	Flamingos-2	0.72	500	1680	8
07200325-0846499	10.63	2014 Jan 22	IRTF	SpeX	0.8	750	480	8
07202582–5617224	12.88	2013 Oct 21	Gemini-South	Flamingos-2	0.72	500	1680	8
07525247–7947386	12.83	2013 Oct 28 & 2014 Jan 17	Gemini-South	Flamingos-2	0.72	500	2520	12
07583046+1530004	10.43	2012 Oct 26	IRTF	SpeX	0.8	750	800	4
07583098+1530146 A & B	9.97	2012 Oct 26	IRTF	SpeX	0.8	750	800	4
08034469+0827000	11.83	2013 Feb 15		Spex	0.8	750	900	6
08045433-6346180	9.93	2013 NOV 02 2012 E-h 16	Gemini-South	Flamingos-2	0.72	500	240	8
08055944+2505028 A & B	11.55	2013 Feb 16 2012 Oct 26		Spex	0.8	750	900	0
08141709+0255199	11.52	2012 Oct 20	IKIF Comini South	Spex Eleminade 2	0.8	730 500	240	4
08194309-7401232	14.82	2013 Oct 20 & 25 2014 Jap 17	Gemini-South	Flamingos-2	0.72	500	1680	0
08194551-0450071	14.62	2014 Jan 17 2013 Apr 21	Magellan	Fiamingos-2	0.72	6000	1500	0 2
08204440-7514571	15.39	2013 Apr 21	Magellan	FIDE	0.0	6000	1310	2
08255896±03/0198	10.01	2013 Feb 18	IRTE	SpeX	0.0	750	300	2
08540240-3051366	9.01	2013 Nov 20	Gemini-South	Flamingos-2	0.72	500	240	8
08561384_1342242	13.60	2013 Nov 20 2012 May 12	IRTE	SpeX	0.72	750	1200	8
09104094-7552528	13.62	2012 May 12 2014 Feb 07	Gemini-South	Flamingos-?	0.72	500	560	8
09451445-7753150	13.89	2014 Feb 15	Gemini-South	Flamingos-2	0.72	500	800	8
09510459+3558098	10.58	2013 Feb 18	IRTF	SneX	0.8	750	300	2
10051641+1703264	11.13	2013 Feb 16	IRTF	SpeX	0.8	750	600	4
10130718–1706349 A & B	12.79	2013 Feb 16	IRTF	SpeX	0.8	750	1200	8
10352029–2058382	11.66	2014 Jan 22	IRTF	SpeX	0.8	750	600	4
11014673-7735144	15.97	2014 Feb 16	Gemini-South	Flamingos-2	0.72	500	8320	32

Table 1
(Continued)

2MASS		Observing	Telescope	Instrument	Slit	Resolving	Tot. Exp.	Num.
Designation	J	Date (UT)			Width (")	Power	Time (s)	Exposures
11083081+6830169	13.12	2009 Mar 04	Palomar	TripleSpec	1.0	2700	1200	4
11195251-3917150	13.13	2014 Jan 22	IRTF	SpeX	0.8	95	1200	8
11335700-7807240	13.20	2014 Feb 15	Gemini-South	Flamingos-2	0.72	500	400	8
11532691-3015414	12.31	2014 Jan 22	IRTF	SpeX	0.8	750	900	6
11544223-3400390	14.19	2008 Jan 09	IRTF	SpeX	0.6	120	480	4
11560224-4043248	16.00	2014 Feb 17	Gemini-South	Flamingos-2	0.72	500	8000	32
12002750-3405371	9.61	2014 Mar 17	Gemini-South	Flamingos-2	0.72	500	1600	32
12042529-2806364	16.11	2014 Mar 17 & 20	Gemini-South	Flamingos-2	0.72	500	16000	64
12212770+0257198	13.17	2014 Feb 14	Gemini-South	Flamingos-2	0.72	500	400	8
12265135-3316124	10.69	2012 May 12	IRTF	SpeX	0.8	750	1200	8
12271545-0636458	14.19	2014 Feb 14	Gemini-South	Flamingos-2	0.72	500	1040	8
12492353-2035592	9.32	2014 Feb 16	Gemini-South	Flamingos-2	0.72	500	400	8
12521062-3415091	11.65	2013 May 27	Gemini-North	GNIRS	0.675	800	120	4
13015465-1510223	14.54	2013 Aug 20	IRTF	SpeX	0.8	750	1500	6
13252237+0600290	12.25	2012 May 12	IRTF	SpeX	0.8	750	1200	8
13582164-0046262	10.81	2012 May 12	IRTF	SpeX	0.8	750	1200	8
14112131-2119503	12.44	2012 May 12	IRTF	SpeX	0.8	750	1200	8
15104786-2818174	12.84	2014 Feb 14	Gemini-South	Flamingos-2	0.72	500	360	8
15291017+6312539	11.64	2013 Aug 20	IRTF	SpeX	0.8	750	800	8
15424676-3358082	17.02	2014 Jun 01	Gemini-South	Flamingos-2	0.72	500	16000	64
15470557–1626303 A & B	13.86	2013 Aug 11	IRTF	SpeX	0.8	750	1800	12
16210134-2346554	15.16	2013 Aug 20	IRTF	SpeX	0.8	750	2400	12
16221255-2346418	10.90	2013 Aug 11	IRTF	SpeX	0.8	750	2400	12
16232017-2353248	13.38	2012 May 12	IRTF	SpeX	0.8	750	1200	8
16251377-2358021	13.75	2013 May 17	Gemini-North	GNIRS	0.675	800	360	4
16272178-2411060	13.98	2013 May 17	Gemini-North	GNIRS	0.675	800	360	4
16330142-2425083	16.16	2013 Aug 11	IRTF	SpeX	0.8	95	1200	8
16422788-1942350	15.23	2014 Jun 18	Magellan	FIRE	0.6	450	480	4
17065487-1314396	14.52	2013 Aug 20	IRTF	SpeX	0.8	750	2400	12
18393308+2952164	11.01	2013 Aug 11	IRTF	SpeX	0.8	750	600	4
18460473+5246027 A & B	11.03	2013 Aug 11	IRTF	SpeX	0.8	750	300	2
18462188-5706040	15.06	2014 May 28	Gemini-South	Flamingos-2	0.72	500	1680	8
19033113-3723302	13.41	2013 Sep 28	Gemini-South	Flamingos-2	0.72	500	1680	8
19480544+5944412 A & B	11.49	2013 Aug 20	IRTF	SpeX	0.8	750	600	4
20025265-1316418	14.48	2013 Aug 11	IRTF	SpeX	0.8	750	1600	8
20050639-6258034	11.75	2013 Oct 20	Gemini-South	Flamingos-2	0.72	500	240	8
20385687-4118285	11.66	2013 Nov 26	Gemini-South	Flamingos-2	0.72	500	240	8
20391314-1126531	13.79	2013 Aug 01	IRTF	SpeX	0.8	750	1200	8
20482880-3255434	14.71	2013 Aug 11	IRTF	SpeX	0.8	750	2800	14
21272613-4215183	13.32	2008 Jul 14	IRTF	SpeX	0.6	120	240	4
21342814-1840298	11.04	2013 Oct 09	Gemini-South	Flamingos-2	0.72	500	240	8
21484123-4736506	10.97	2013 Oct 20	Gemini-South	Flamingos-2	0.72	500	240	8
22062157-6116284	16.61	2014 May 15	Magellan	FIRE	0.6	450	240	2
22444905-3045535	14.65	2013 Aug 01	IRTF	SpeX	0.8	750	450	3
22573768-5041516	14.96	2014 Jul 17	Gemini-South	Flamingos-2	0.72	500	5280	16
23231347-0244360	13.58	2008 Nov 03	IRTF	SpeX	0.6	120	720	8
23453903+0055137	13.77	2008 Jul 14	IRTF	SpeX	0.6	120	270	6

Standard A0 to A6-type stars were observed immediately before or after every science target at a similar airmass to ensure a proper telluric correction. Several high-S/N quartz lamp and Ar lamp exposures were obtained immediately after every telluric standard star to ensure a proper wavelength calibration and flat field correction. Dark exposures were obtained at the end of each night, using similar exposure times than all of the science and calibration data to ensure a proper correction of the dark current. A numbers of observations were split between a few nights when observing conditions changed before the required S/N could be obtained. We used a custom IDL pipeline to apply dark current subtraction and flat field calibrations, correct the trace curvature, optimally extract the spectrum (Horne 1986) and perform a wavelength calibration using the Ar lamp observations. A dark current subtraction is usually not needed when data are reduced in A-B pairs, like is the case here; however, we found that applying this correction improved the quality of the data. This is likely due to the large exposure times that were used for some targets, which resulted in a large contribution from the dark current that must be corrected both in the data and flat field exposures before applying the flat field correction.

A low-pass filter was applied to the flat field exposures to avoid contaminating data with scattered light. We observed that the spectral dispersion (and thus wavelength solution) generally varied from one exposure to another; the wavelength solutions obtained from the Ar calibrations are hence only approximate.

To address this problem, we used several telluric absorption features in the raw spectra of the science and telluric observations to refine individual wavelength solutions. The *JH* and *HK* blocking filters also caused significant fringing in the data (up to $\sim 7\%$). We corrected this by adjusting a sinusoid fringing solution to the low frequencies of the raw spectra. We found that a complete fringing solution (which includes finesse as an additional parameter) did not improve the results; we thus chose the simpler sinusoid approach to have a more robust algorithm.

The extracted science and telluric spectra were combined and telluric-corrected using a modified version of the SpeXtool package adapted for Flamingos-2. We observed that the slope of the continuum in the overlapping region of both observing modes (in the *H* band) varied in a systematic way at the edge of the detector. Hence, we removed these regions before combining the spectra. A few objects for which we obtained Flamingos-2 data (e.g., 2MASS 07083261–4701475, 2MASS 20414283-3506442, and 2MASS J12042529-2806364) turned out to be field dwarfs that closely match literature SpeX-prism spectra of other known objects of the same spectral type: this is an indication that the systematics mentioned above were accurately corrected.

3.4. GNIRS at Gemini-North

We used GNIRS at Gemini-North to obtain NIR spectroscopy for three targets in 2013. We used the 321 mm^{-1} grating centered at 1.65 μ m in the cross-dispersed mode with the 0".675 slit to achieve a resolving power of $R \sim 750$ over 0.9–2.45 μ m. We nodded exposures along the slit in ABBA patterns with total exposure times ranging from 120 to 360 s to reach S/N > 100 per resolution element. A0-type telluric standard stars were observed immediately before or after science targets at a similar airmass to ensure a proper telluric correction. Several high-S/N quartz lamp and Ar lamp exposures were obtained immediately after every target to ensure a proper wavelength calibration and flat field correction. The data were reduced with the XDGNIRS IRAF package provided by Gemini.

3.5. TripleSpec at Hale

We used TripleSpec (Herter et al. 2008) at the Palomar Observatory 5 m Hale Telescope to obtain NIR spectroscopy for one target in the cross-dispersed mode with the 1".0 slit, yielding a resolving power $R \sim 3800$ over 1.0–2.45 μ m. We observed the science target in a four-position ABBA nodding pattern along the slit with a total exposure time of 1200 s to reach a S/N > 100 per resolution element. High-S/N quartz lamp and NeAr lamp exposures were obtained to ensure a proper wavelength calibration and flat field correction. We reduced the data using an adapted version of SpeXtool (see Section 3.2).

4. SPECTRAL TYPES AND LOW-GRAVITY CLASSIFICATION

We describe in this section the method that we used to assign spectral types to our new observations. Our typing scheme consists of two distinct dimensions : the first dimension consists of the usual spectral subtypes and is mostly sensitive to $T_{\rm eff}$. The second dimension, introduced by Kirkpatrick (2005) and Kirkpatrick et al. (2006), aims at characterizing the surface gravity with the use of a greek-letter suffix. Field-gravity dwarfs are designated with the α suffix or no suffix, intermediate-gravity dwarfs with the β suffix, and very low-gravity dwarfs with the γ suffix. The δ suffix was also introduced by Kirkpatrick et al. (2006) to designate objects with an even younger age (typically less than a few Myr) and lower surface gravity than those associated to the γ suffix.

Optical spectral standards were used to classify NIR spectra of field K7–M9 spectral types. We used the NIR data of GJ 820 B (K7), Gl 229 A (M1), Gl 411 (M2), Gl 213 (M4), Gl 51 (M5), Gl 406 (M6), GJ 644 C (M7), GJ 752 B (M8), and LHS 2924 (M9) as field-gravity spectral standards for these respective spectral types. These standards were identified from the list maintained by Eric Mamajek¹² (Boeshaar 1976; Kirkpatrick et al. 1991; Pecaut & Mamajek 2013) and their spectra were downloaded from the IRTF spectral library.¹³ We did not use any of the suggested K8, K9, M0, and M3-type standards, since none of them were available in the IRTF spectral library.

While NIR L dwarfs spectral standards have been identified by Kirkpatrick et al. (2010), we have opted to use optically anchored NIR spectral average templates for classifying field L0–L9 dwarfs. Templates are constructed by median-combining all spectra of a given optical spectral type and gravity class. These templates were provided by K. Cruz and their creation will be discussed in detail and be made public as part of a forthcoming paper (K. L. Cruz et al. 2015, in preparation). The spectral morphology of these templates is consistent with the Kirkpatrick et al. (2010) spectral standards but since they are an average of many objects, they also reflect the diversity of spectral morphologies present in each spectral type.

Spectral standards have been determined for low-gravity M and L dwarfs by Allers & Liu (2013), but we opted to use spectral average templates in this case too, for the reasons mentioned above. We generated M6–M9 γ templates with data published in Allers & Liu (2013) and sent to us directly by the authors. These templates are available at the Montreal Spectral Library.¹⁴ The optically anchored L0 β , L1 β , and L0–L4 γ templates were provided by K. Cruz. They will be discussed in detail and be made public as part of a forthcoming paper et al. (Cruz et al. 2015, in preparation).

All template, standard, and target spectra were re-sampled to the same resolution and wavelength grid as SpeX prism observations with the 0".6 slit ($R \sim 120$). Following the method of Cruz & Núñez (2012), the spectra were normalized in three sections in order to minimize the effect of large NIR color variations within a given spectral type. The spectra were broken into three sections: 0.80–1.35 μ m, 1.40–1.80 μ m, and 1.95–2.40 μ m, roughly corresponding to the *zJ*, *H*, and *K* bands.

In a first step to estimating a spectral type, we categorized our 245 new spectra with the spectral template and standard grid described above. There were 11 objects, however, that did not have a good visual match to any standard or template in the grid; this number excludes the early-type contaminants which are discussed later in this work. We collected additional low-

¹² http://www.pas.rochester.edu/~emamajek/spt/

¹³ Maintained by Michael C. Cushing and available at http://irtfweb.ifa. hawaii.edu/~spex/IRTF_Spectral_Library/.

¹⁴ www.astro.umontreal.ca/~gagne/MSL.php

 Table 2

 An Extended Sequence of Low-gravity Dwarfs

Name	Spectral Type
USco J160603.75-221930.0	L0 δ
USco J160727.82-223904.0	L0 δ
USco J160737.99-224247.0	LO δ
USco J160818.43-223225.0	LO δ
USco J160828.47-231510.4	LO δ
USco J160843.44-224516.0	L0: δ
USco J160918.69-222923.7	L0 δ
USco J161228.95-215936.1	LO δ
USco J161441.68-235105.9	L0 δ
USco J163919.15-253409.9	L0 δ
CD-35 2722 B	L3 β
2MASS J01531463-6744181	L3 β
2MASS J17260007+1538190	L3 β
2MASS J00011217+1535355	L4 β
2MASS J05120636-2949540	L5 β
2MASS J23174712-4838501	L5 β
2MASS J03264225-2102057	L5 β/γ
SIMP J21543454-1055308	L5 β/γ
2MASS J03552337+1133437	J0355-type (L3–L6 γ)
2MASS J16154255+4953211	J0355-type (L3–L6 γ)
2MASS J23433470-3646021	J0355-type (L3–L6 γ)
WISEP J004701.06+680352.1	J2244-type (L6–L8 γ)
2MASS J22443167+2043433	J2244-type (L6–L8 γ)
PSO J318.5338-22.8603	J2244-type (L6–L8 γ)

Note. All spectral types are from this work and are based on NIR spectra. A : symbol indicates that the spectral type is based on low signal-to-noise data and is uncertain (± 1), and a :: symbol that it is very uncertain (± 2 subtypes); pec indicates peculiar features; β and γ respectively indicate intermediate gravity and very low gravity.

gravity BD spectra in the literature to identify 19 more objects that do not match our standards.

We performed a visual analysis of all of the unclassifiable spectra and identified enough objects with similar spectral morphologies to create tentative new spectral types and templates for L0 δ , L3 β , L4 β , and L5 β . The objects that were used in the creation of these templates are listed in Table 2. We list the revised spectral types that we obtain for other spectra from the literature in Table 3. We note that our L3 β template includes 2MASS J17260007+1538190, which was suggested by Allers & Liu (2013) as a tentative template for the L3 β spectral type.

We could not build a template for the L2 β spectral type, as the only objects that were confirmed as L2 β from optical data have either very low S/N in the NIR or no NIR data. As we gather more high-S/N spectra of low-gravity L dwarfs, we expect to fill this gap.

The L0 δ template was built from eight candidate members of Upper Scorpius (Lodieu et al. 2008) and one candidate member of β PMG (2MASS 00464841+0715177) that are similar to the L0 γ template except that their *H* band is even more triangular and their *K* band has a redder continuum. It is also notable that the H₂O-dependent slope of the L0 δ at 1.7–1.8 μ m is slightly steeper than what is seen in any other L-type template.

There are two sets of objects with similar spectra, each with three targets, that we identified via our visual analysis; however, we are unable to confidently assign them a spectral type that fits into our grid of templates. For the purposes of this paper, we label these objects as J0355-type and J2244-type. One set is

2MASS J03552337+1133437, composed of 2MASS J16154255+4953211, and 2MASS J23433470-3646021. Their spectra are similar to the L4 γ template except that they have a shallower CO band at 2.3 μ m. The other set is composed of 2MASS J00470038+6803543, PSO J318.5338-22.8603, and 2MASS J22443167+2043433. Their spectra display a significantly redder continuum than our templates, which might be indicative of a later spectral type. We note that two objects have previous classifications based on the index-based scheme of Allers & Liu (2013): 2MASS J00470038+6803543 was classified as an intermediate-gravity L7 dwarf by Gizis et al. (2015) and PSO J318.5338-22.8603 was classified as a very low-gravity L7 dwarf by Liu et al. (2013b). We listed these two sets of objects as well in Table 2.

We adopt a conservative estimate of L3–L6 γ for the spectral type range of the *J0355-type*. The spectral features of the J2244-type are indicative of a spectral type in the range L6–L8 γ range. For both of these new spectral types, we refrain from assigning them a more precise location in the spectral sequence until more data are available at these late low-gravity types. It is unclear at this stage whether J0355-type and J2244-type objects are peculiar or a simple extension of low-gravity BDs at spectral types later than L5. A larger number of late-type, low-gravity L dwarfs will need to be identified before we can assess this. Our set of low-gravity templates is displayed in Figure 1.

We used the index-based classification method of Allers & Liu (2013) to corroborate our visual classification. This method consists of measuring the slope of H₂O continuum features to assign a spectral type, and a combination of several gravitysensitive spectroscopic indices to assign a gravity class. We found that spectral types obtained from the template grid system described above generally agree with index-based spectral types within one subtype (Figure 2). The standard deviation between the two methods for the 163 non-peculiar objects that we categorized is of 0.7 subtypes, with a reduced χ^2 value of 0.8. A reduced $\chi^2 \approx 1$ indicates that measurement errors are representative of the discrepancies. The reduced χ^2 is given by $1/(N-1) \cdot \sum y/\sigma_v$, where N is the number of objects, y is the spectral type discrepancy and σ_{y} is the quadrature sum of the index-based and visual-based spectral type measurement errors. All cases discrepant by more than 1.5 spectral types correspond to low-S/N data, except for 2MASS J21420580–3101162 that gets $L1.5\pm0.3$ from the index-based method and L3 from the visual-based method. This object does not display signs of youth or significantly peculiar features, but it has a slightly redder slope at 1.7–1.8 μ m. It unclear what is the cause of this discrepancy.

We used optical data to assign an adopted spectral type using a template-based visual classification method (Cruz et al. 2009) only for the 4 objects for which no NIR data were available. In all other cases, our adopted spectral types are based on NIR data only. Our NIR spectral types based on a visual comparison with templates show a standard deviation of 0.9 subtype with respect to optical spectral types in the literature, and the reduced χ^2 of the differences is 1.5, hence slightly larger than what would be expected given the uncertainties. If we compare optical spectral types to the index-based spectral types of Allers & Liu (2013), we obtain a slightly larger standard deviation (1.1 subtype) and reduced χ^2 (2.4). This is indicative that our visual-based classification method is more consistent with spectral types based on optical data that were reported in the

Name	Spectral Type ^a											
	Optical	References NIR		References	Adopted							
	Lc	ow-gravity Dwarfs										
2MASS J21324036+1029494			L4.5:	(1)	L4: β/γ							
2MASS J14482563+1031590	L4:	(2)	L3.5	(3)	L5: β							
WISE J174102.78-464225.5			L7::	(4)	L5:–L7: γ							
G196–3B	L3 β	(5)	L3 γ	(6)	L2–L4 γ							
2MASS J00303013-1450333	L7	(7)	L4.5::	(8)	L4–L6 β							
2MASS J20025073-0521524	L6	(9)	L7::	(10)	L5–L7 γ							
	Red Drown Dwarfs	with no Clear Signs of I	Low Gravity									
2MASS J08354256-0819237	L5	(11)	L5	(12)	L4 pec ^b							
2MASS J18212815+1414010	L4.5 pec	(13)	L5 pec	(14)	L4 pec							
2MASS J21512543-2441000	L3	(9)			L4 pec							
2MASS J01033203+1935361	L6 β	(7), (15)	L6 β	(6)	L6 pec							
2MASS J01075242+0041563	L8	(16)	L8 pec	(17)	L7 pec ^b							
2MASS J08251968+2115521	L7.5	(7)	L6	(18)	L7 pec ^b							
2MASS J08575849+5708514	L8	(5)	$L8 \pm 1$	(19)	L8–L9 pec							

 Table 3

 Revised NIR Spectral Types From the Literature

Notes. References to this Table: (1) Chiu et al. (2006), (2) Reid et al. (2008), (3) Wilson et al. (2003), (4) Schneider et al. (2014), (5) Kirkpatrick et al. (2008), (6) Allers & Liu (2013), (7) Kirkpatrick et al. (2000), (8) Burgasser et al. (2010), (9) Cruz et al. (2007), (10) Bardalez Gagliuffi et al. (2014), (11) Cruz et al. (2003), (12) Marocco et al. (2013), (13) Looper et al. (2008b), (14) Kirkpatrick et al. (2010), (15) Faherty et al. (2012), (16) Hawley et al. (2002), (17) Geissler et al. (2011), (18) Knapp et al. (2004), (19) Geballe et al. (2002).

^a All revised spectral types are from this work and are based on NIR spectra.

^b Candidate member of the ~625 Myr-old Hyades association Bannister & Jameson (2007).

literature. This should be expected, as our templates are anchored on optical data. In both cases, we observe no systematic bias (the mean of the differences is smaller than 0.1 subtype). Several objects that deserve further discussion are presented in detail in the appendix.

We note that the index-based field-gravity, intermediategravity and very low-gravity classes defined by Allers & Liu (2013) were built to correspond to the optical α , β and γ classes, which is what we observe in 143/176 (81%) of the cases. Some of the discrepancies arise for objects near the spectral type thresholds where the method of Allers & Liu (2013) stops being applicable ($\leq M6$ or $\geq L6$) or for data with a lower S/N. The δ gravity class does not have an equivalent in the index-based classification of Allers & Liu (2013), but we note that all three of the young dwarfs that we categorized as δ are assigned with the maximal index-based gravity score (2222). It does not seem that this maximal index-based gravity score always translates as a δ visual classification though, as there are four additional objects in our sample that obtained the score 2222 but that we visually categorized as γ (2MASS J00182834–6703130; L0 γ , 2MASS J01205114–5200349; L1 γ ; 2MASS J20113196–5048112; L3 γ ; 2MASS J22351658–3844154; L1.5 γ ; all are THA candidate members). For consistency within this work, we have adopted the visual spectral types in the remaining sections, but we list all visual and index-based spectral types in Table 4. We note that this choice does not affect the conclusions presented in this work.

5. RESULTS

In Figure 3, we present the NIR spectra of several new intermediate (β) and very-low (γ) gravity dwarfs discovered in this work, as well as known dwarfs for which we have obtained new data. Several objects that were uncovered as candidate members of YMGs in BASS had NIR or optical spectroscopy

readily available in the SpeX Prism Spectral Libraries¹⁵ (Burgasser 2014) or the RIZzo Ultracool Spectral Library¹⁶ with no discussion of low gravity in the literature; we included them in our present analysis and the resulting spectral classification is listed in Table 4 along with the new discoveries.

There are some cases where the BANYAN II tool yields ambiguous candidate membership to more than one association (i.e., at least a second moving group shares 10% of the total YMG Bayesian probabilities). In all such cases, we list in Table 4 all plausible YMGs with their relative share of the total YMG probability (i.e., excluding the field probability). An extensive RV and parallax follow-up will be required before more can be said on their YMG membership.

We have identified seven objects (Table 4) that display signs of low gravity, but for which additional information was inconsistent with membership to any of the YMGs presented here (e.g., RV and distance measurements or the effect of interstellar extinction affecting the NIR spectrum which is not consistent with ages older than \sim 5 Myr). It is possible that these objects belong to YMGs or star-forming regions that are not considered here, that their RV or parallax measurements are affected by an unresolved binary companion (see the appendix for a detailed discussion), or that other physical properties such as enhanced dust mimics a lower gravity.

However, Allers & Liu (2013) noted that their index-based classification scheme should not be significantly contaminated by old, dusty BDs, which makes this last hypothesis less likely.

In Figure 4, we show a histogram of all previously known low-gravity dwarfs along with new discoveries or

¹⁵ http://pono.ucsd.edu/~adam/browndwarfs/spexprism

¹⁶ Kirkpatrick et al. (2000, 2008, 2010), Cruz et al. (2003, 2007, 2009), Reid et al. (2008); see http://www.astro.umontreal.ca/~gagne/rizzo.



Figure 1. NIR spectral average of templates of intermediate gravity (Panel (a)) and very low gravity (Panel (b)). All spectra were normalized to their median across the full wavelength range in each band, resampled at the same resolution ($R \sim 120$) and shifted vertically for comparison purposes. The vertical color bars indicate the location of gravity sensitive features.

confirmations of low gravity that are presented here. It might seem surprising that we did not identify any new low-gravity L2 dwarfs, however this is likely the effect of small number statistics and the fact that we still lack a template for the L2 β spectral type, e.g., some low-S/N low-gravity objects presently typed as L1: and L3: might turn out to be L2 dwarfs when more data becomes available. We anticipate our visual-based lowgravity classification scheme to improve as more data is obtained. If we account for the measurement errors on our spectral types using Gaussian probability density functions (PDFs) (which softens the gap at L2) and use Poisson statistics to assess the significance of this lack of L2 dwarfs, we find that the differences between the number of known low-gravity L1, L2, and L3 dwarfs is insignificant (at the level of 0.2σ).

In Figures 5 and 6, we compare all new low-gravity confirmations with the field and low-gravity sequences defined by Allers & Liu (2013). The individual values for these gravity-sensitive spectroscopic indices are listed in Table 5. There are 7 objects in our sample that did not have a discussion of low gravity in the literature and for which optical spectra were



Figure 2. Distribution of the differences between our visual and index-based spectral classifications, for young dwarfs and field objects. Spectral types generally agree within one subtype, with a standard deviation of 0.7 subtypes and a reduced χ^2 value of 0.8. This is indicative that our measurement errors are representative of the observed differences between the two methods. It can also be seen that most of the outliers in the distribution correspond to objects with uncertain spectral types (i.e., measurement errors of one subtype or more.)

available in the Ultracool RIZzo Spectral Library. We used them to revise their spectral types and measure gravitysensitive optical indices defined by Kirkpatrick et al. (1999) and Cruz et al. (2009). These results, based on optical data only, are presented in Table 6. The new spectroscopic observations presented here (95 from BASS, 26 from LP-BASS and 120 from PRE-BASS) allowed us to uncover a total of 108 new M6–L5 low-gravity dwarfs, doubling the number of such known objects (98 before this work).

In addition to several new candidate members of YMGs, we report here that 2MASS J14252798-3650229 (DENIS-P J142527.97-365023.4) is a new low-mass BD bona fide member of ABDMG. This object was identified by Kendall et al. (2004) as an L5 dwarf with an estimated spectrophotometric distance of ~ 10 pc. Blake et al. (2010) measured an RV of 5.37 \pm 0.25 km s⁻¹ and Dieterich et al. (2014) measured a trigonometric distance of 11.57 ± 0.11 pc. Gagné et al. (2015) reported that the galactic position and space velocities of this object are a very good match to ABDMG (Figure 7), suggesting that it would be a new bona fide member if low gravity would be confirmed. They also indicated that its NIR colors are redder than those of field dwarfs of the same spectral type, which hints at low gravity. The low gravity is indeed readily apparent in the new SpeX prism spectrum that we obtained for this object (Figure 8): both a visual comparison and the index-based classification of Allers & Liu (2013) indicate that this object is an L4 γ dwarf. We conclude that 2MASS J14252798-3650229 is a new bona fide member of ABDMG, making it the second latest-type confirmed member of this moving group after the L7 β member WISEP J004701.06+680352.1. At the age of ABDMG, 2MASS J14252798-3650229 has an estimated mass of $26.6^{+0.3}_{-1.0} M_{Jup}$.

5.1. Updated YMG Membership

It is possible to use the spectral type information as well as the youth of candidate members determined from the spectroscopic follow-up presented here as additional inputs in BANYAN II to refine estimates of distance, RV and YMG membership and contamination probabilities. Spectral types are used to assess if the absolute W1 magnitude of a target is consistent with its spectral type at the statistical distance that corresponds to a given YMG membership (using distinct sequences for field and low-gravity dwarfs; see Paper V), whereas prior knowledge of youth reduces the number of potential contaminants from the field and thus improves the probability that the object belongs to a YMG. We reject all objects with spectral types \geq M5 that display no signs of low gravity (17 in BASS, 7 in LP-BASS, and 41 in PRE-BASS), since this implies an age older than the Pleiades (~120 Myr; Cruz et al. 2009; Allers & Liu 2013) and is not consistent with membership to any YMG considered here. These updated results are listed in Table 4, and individual objects of interest are discussed in the appendix.

5.2. X-Ray Luminosity

We followed up several objects that turned out to have spectral types earlier than expected, some of them (\leq M5) to the point where current NIR and optical index-based methods are unable to determine whether they are likely young or field objects. In this section, we take advantage of the *ROSAT* bright and faint source catalogs (Voges et al. 1999, 2000; VizieR catalogs IX/10A and IX/29) to assess whether these objects are young candidate members of YMGs or field interlopers.

Malo et al. (2014a) demonstrated that the distribution of absolute X-ray luminosity for M0–M5 dwarf members of ABDMG and β PMG is significantly distinct from that of field M0–M5 dwarfs. In particular, they showed that β PMG members are ~4 times more X-ray luminous than ABDMG members, a factor that goes up to \gtrsim 40 when instead compared with field dwarfs. We investigated whether any of our M0–M5 candidate members listed in Table 4 display X-ray emission by cross-matching their 2MASS position with the *ROSAT* catalogs with a 15" search radius. We computed the absolute X-ray luminosity for all X-ray sources recovered this way, using trigonometric distances when possible or kinematic distances otherwise.

We have identified ROSAT entries for only three objects: 2MASS J08540240-3051366 (M4 candidate member of β PMG; log $L_X = 28.4 \pm 0.3$) has a low X-ray luminosity compared with M3–M5 members of ABDMG or β PMG (both have $\log L_{\rm X} \approx 28.5 - 29.5$) and could thus be a field interloper (log $L_{\rm X} \approx 27 - 28.5$; see Figures 7 and 8 of Malo et al. 2014a). 2MASS J08194309-7401232 (M4.5 candidate member of COL; $\log L_{\rm X} = 29.3 \pm 0.4$) and 2MASS J21490499-6413039 (M4.5 candidate member of THA; $\log L_{\rm X} = 29.3 \pm 0.3$) both have X-ray luminosities consistent with an age similar or younger than that of ABDMG, making them likely members of their respective moving groups. We note that 2MASS J21490499-6413039 has already been reported as a candidate member of THA by Kraus et al. (2014b), who measured its RV and found it to be consistent with other THA members. Objects that do not have a ROSAT counterpart do not necessarily have a low absolute X-ray luminosity, but might be too distant or located outside of the regions covered by the ROSAT survey.

Using the *ROSAT* bright catalog detection limit of 0.1 ct s⁻¹ in the 0.1–2.4 keV energy band and assuming a hardness ratio HR1 \approx 0, we can only put an upper limit of log $L_X = 28 - 29.8$ on the remaining targets, which is generally not sufficient to reject any more candidate members. Only 3/41 of these targets (2MASS J05484454–2942551, 2MASS J06494706–3823284, and 2MASS J07583098+1530146 AB) have log $L_X < 28.5$,

					1			· · · · · · ·							
2MASS		Spectral Type ^a							ravity ^b		YM	G Membership			Source
Designation	J	Lit. Opt.	References	Lit. NIR	References	H ₂ O Ind.	Adopted	Class	Score	Lit.	References	Updated BASS ^c	$B_P (\%)^d$	$C_P (\%)^{e}$	Catalog
							Bona Fide Me	mbers							
03350208+2342356	12.25	M8.5	29	M7 VL-G	30	M7.3	M7.5 β	INT-G	1n21	BPMG	31	BPMG	84.2	3.9	PRE-BASS
03552337+1133437	14.05	L5 γ	32	L3 VL-G	30	L2.3	L3–L6 γ	VL-G	2122	ABDMG	33	ABDMG	99.5	1.1	BASS
14252798-3650229	13.75	L3:	15	L5	52	L3.1	L4 γ	INT-G	11?1			ABDMG	99.9	0.1	BASS
						Lo	w-gravity Candida	te Members							
00011217+1535355	15.52			L4:	1	L3.7	L4 β	INT-G	1211			ABDMG	97.4	1.1	BASS
00065794-6436542	13.39	L0	4			M7.4	M8 γ	INT-G	1n12	THA	5	THA	>99.9	< 0.1	BASS
00182834-6703130	15.46					L0.4	$L0 \gamma$	VL-G	2222			THA	99.9	< 0.1	BASS
00191296-6226005	15.64					L1.0	L1 γ	VL-G	1212			THA	99.7	< 0.1	BASS
00192626+4614078	12.60	M8	7			M7.4	M8 β	INT-G	1n12	ABDMG	8	ABDMG	92.1	4.0	BASS
00274534-0806046	11.57					M5.7	M5.5 β	INT-G	nn1n			BPMG	85.2	26.3	BASS
00303013-1450333	16.28	L7	7,11	L4.5::	9	L3.2	L4–L6 β	VL-G	2n21			ARG	26.5	2.6	LP-BASS
00344300-4102266	15.71					L1.5	L1: β	VL-G	2121			THA	98.7	< 0.1	BASS
00381489-6403529	14.52					M8.2	M9.5 B	INT-G	1n12			THA	99.9	< 0.1	LP-BASS
00413538-5621127	11.96	M6.5 + M9	13			M7.9	M7.5 β u	VL-G	0n22	THA	14	THA	99.9	< 0.1	BASS
00425923 ± 1142104	14 75					M9.8	M9 ß	INT-G	0n11			ABDMG	19.6	53.1	PRE-BASS
00423723 1142104	14.75					1419.00	mp p	intro	0111			(66); BPMG(33)	17.0	55.1	THE BROS
00464841+0715177	13.89	L0::	15			L0.9	L0 δ	VL-G	2222			BPMG	89.3	25.1	BASS
00514561-6227073	12.58					M7.4	M5.5: β	INT-G	nn1n			THA	>99.9	< 0.1	LP-BASS
00584253-0651239	14.31	L0	11	L1	16	L0.6	L1 B	INT-G	11?1			ABDMG	96.5	0.3	LP-BASS
												(66); BPMG(33)			
01205114-5200349	15.64					L1.4	L1 γ	VL-G	2222			THA	>99.9	< 0.1	BASS
01265327-5505506	12.04					M6.2	M6 γ	FLD-G	0n20			THA	>99.9	< 0.1	BASS
01294256-0823580	10.65	M5	6			M6.1	M7 β	VL-G	2n22			BPMG	95.9	18.9	BASS
01344601-5707564	12.07	M4.5	20			M5.8	M6 β	VL-G	2n20	THA	20	THA	>99.9	< 0.1	BASS
01484859-5201158	10.87					M5.0	M5 β	INT-G	nn1n			THA	>99.9	< 0.1	BASS
01531463-6744181	16.41	L2:	15			L2.9	L3 β	VL-G	2211			THA	>99.9	< 0.1	BASS
02103857-3015313	15.07					M8.4	M9.5 β	INT-G	1n12			THA	99.9	< 0.1	BASS
02265658-5327032	15.40					L0.5	L0: γ	VL-G	2221			THA	>99.9	< 0.1	BASS
02282694+0218331	12.12					M5.3	M5 γ	VL-G	nn2n			THA(68); BPMG(29)	89.9	< 0.1	LP-BASS
02404759-4253377	12.20					M6.1	M6 β	INT-G	1n02			THA	99.9	< 0.1	LP-BASS
02410564-5511466	15.39					L1.6	L1 γ	VL-G	1221			THA	>99.9	< 0.1	BASS
02501167-0151295	12.89					M7.3	M7: β	VL-G	1n22			BPMG	92.9	1.1	BASS
02583123-1520536	15.91					L3.0	L3 B	INT-G	1111			THA	88.9	< 0.1	BASS
02590146-4232204	12.24					M5	M5 γ	VL-G	nn2n	COL	24	COL	8.6	62.5	PRE-BASS
03093877-3014352	11.58	M4.5	20			M6.7	M6.5 γ	VL-G	2n20	THA	20	THA	>99.9	< 0.1	BASS
03111547+0106307	10.68	M5.5	17			M7.6	M6: γ	FLD-G	0n20			BPMG(55); THA(45)	79.0	15.7	PRE-BASS
03164512-2848521	14.58	L0:	7	L1	19	L1.7	L1 β	INT-G	1101			ABDMG	96.9	3.2	BASS
03182597-3708118	13.37					M5.7	M6 : γ	VL-G	2n1n			COL(62); THA(38)	75.5	39.2	LP-BASS ^f
03224622-7940595	12.22					M6.3	M6.5 β	INT-G	2n10			THA	80.9	< 0.1	BASS
03264225-2102057	16.13	L4	28			L4.1	L5 β/γ	FLD-G	0n01			ABDMG	98.8	1.1	BASS
03363144-2619578	10.68	M5.5	24			M5.8	M5.5 β	INT-G	nn1n	THA;COL	24	THA	99.9	< 0.1	BASS
03390160-2434059	10.90	M6	24			M4.8	M5 β	INT-G	nn1n	COL	24	COL(53):	73.9	32.3	BASS
		-					- 1					THA(35); BPMG(12)			
03420931-2904317	15.92					L1.0	L0: β	VL-G	122n			THA	99.7	< 0.1	BASS
03421621-6817321	16.85	L2:	28				L4 γ o			THA	5	THA	99.8	< 0.1	BASS
03550477-1032415	13.08	M8.5	7			M8.7	M8.5 β	INT-G	1n21				93.8	< 0.1	BASS

13

GAGNÉ ET AL.

(Continued) 2MASS Ind. Gravity^b YMG Membership Spectral Type^a Source Updated H₂O Ind. BASS Designation JLit. Opt. References Lit. NIR References Adopted Class Score Lit. References $B_{P}(\%)^{d}$ $C_{P}(\%)^{e}$ Catalog Bona Fide Members THA(76); COL(24) 04185879-4507413 16.16 L2.8 L3 γ VL-G 2211 THA 92.7 < 0.1BASS M8.4 VL-G 1212 86.7 < 0.1 04400972-5126544 15.69 L0: γ THA(73): BASS COL(23) 04402583-1820414 12.65 M6.1 M6 β INT-G 1n20 COL(71); 13.0 PRE-BASS 31.7 BPMG(28) 30 04433761+0002051 12.51 28 L0 VL-G M9.9 M9 γ INT-G 1n21 BPMG 5,12 BPMG 99.8 2.8 BASS M9 γ 04493288+1607226 14.27 M9.0 M9 γ VL-G 2n22 BPMG 1.6 98.2 PRE-BASS 05012406-0010452 14.98 L4 γ 32 L3 VL-G 30 L3.1 VL-G 1212 COL(74); 65.9 2.4 BASS L4 γ ... CAR(26) 12 05071137+1430013 A 10.57 M5.6 M5.5 β INT-G nn1n BPMG BPMG 28.5 84.9 PRE-BASS 05071137+1430013 B 10.57 M5.2 M5.5 β INT-G nn1n BPMG 12 BPMG 28.5 84.9 PRE-BASS L4.5 7,35 19 L3.8 INT-G BPMG 5 57.0 37.9 BASS 05120636-2949540 15.46 L4.5:: L5 β 1n01 BPMG 05123569-3041067 11.90 M6.8 M6.5 γ FLD-G 0n20 COL 96.4 11.5 BASS 11.88 M6.5 2 M7.1 M7 β VL-G 1n22 COL 96.2 8.8 BASS 05181131-3101529 05264316-1824315 12.36 ... M6.2 M7 β VL-G 1n22 COL 93.5 12.8 BASS 05361998-1920396 15.77 L2 γ 28 L2 VL-G 30 L2.8 L2 γ INT-G 2111 COL 5 COL 97.6 7.4 BASS 05402325-0906326 14.59 M8.0 M9 β INT-G 1n11 COL 72.0 16.1 PRE-BASS 1012 87.2 9.1 16.39 L0.1 INT-G CAR BASS 06272161-5308428 L0: β/γ 06322402-5010349 15.02 L3 15 L3 β o ABDMG 5 ABDMG 29.5 76.7 PRE-BASS 06494706-3823284 11.65 M4.2 M5 γ nn2n CAR(85); 38.8 41.1 PRE-BASS COL(15) M8 INT-G 07140394+3702459 11.98 15 M7.5 M7.5 β 1n10 ARG 88.9 0.5 LP-BASS 07202582-5617224 12.88 M5.9 M6 γ INT-G 1n21 BPMG 29.3 73.1 PRE-BASS 07525247-7947386 12.83 M6.1 VL-G CAR 97.1 2.5 PRE-BASS M5: γ nn2n M5.7 INT-G 5.2 08034469+0827000 11.83 M6 β 1n02 ABDMG 91.2 PRE-BASS 08194309-7401232 10.06 M5.0 M4.5 CAR 99.3 1.6 PRE-BASS 08561384-1342242 13.60 M8.6 M8 γ INT-G 1n11 TWA 4.9 < 0.1PRE-BASS 13.89 M8.2 M9 β INT-G 90.4 2.8 09451445-7753150 1n21 CAR PRE-BASS 09532126-1014205 13.47 L0 28 M9.9 M9 β INT-G 1n11 TWA(91): 81.2 < 0.1BASS CAR(9) 16.91 INT-G 1012 92.4 < 0.1BASS 10212570-2830427 L2.5 L4: β/γ TWA 10284580-2830374 10.95 M5 45 M5.7 M6 γ VL-G 2n22 TWA 45 TWA 97.5 < 0.1BASS 10455263-2819303 12.82 M6.1 M6 γ VL-G 1n22 TWA 65.0 < 0.1LP-BASS 11064461-3715115 14.49 M9.4 M9 γ VL-G 2n22 TWA 94.6 < 0.1BASS 11083081+6830169 13.12 L1 28.46 L2.0 $L1 \gamma$ INT-G 1211 CAR 6.0 89.9 PRE-BASS 11271382-3735076 L0.6 $L0 \delta$ VL-G 2222 TWA 92.5 < 0.1 LP-BASS 16 47 M9.7 INT-G < 0.1 0112 TWA 68.9 11480096-2836488 16.11 L1: β BASS 11544223-3400390 14.19 L0 35 L0.5 19 L0.8 $L0 \beta$ INT-G 110? ARG 55.6 46.4 PRE-BASS 49 VL-G 49 < 0.112073346-3932539 12.99 M8 pec M8 VL-G 30 M9.0 M8.5 γ 2n22 TWA TWA 99.9 BASS 2222 12074836-3900043 15.49 $L0 \gamma$ 50 L1 VL-G 50 L1.3 $L1 \delta$ VL-G TWA 50 TWA 99.7 < 0.1BASS 51 M5.7 VL-G 51 TWA 95.3 < 0.1 12265135-3316124 10.69 M5 M5.5 γ nn2n TWA PRE-BASS 12271545-0636458 14.19 M9 7 ... M8.1 M8.5 β INT-G 2n10 TWA 1.5 0.6 PRE-BASS 14.78 M9 VL-G 50 M8.7 M9 γ VL-G 2n22 TWA 50 TWA 46.6 < 0.1 BASS 12474428-3816464 L0.3 VL-G 2n22 TWA 59.3 0.0 12535039-4211215 16.00 M9.5 γ BASS L4.3 VL-G 2021 TWA 15.9 < 0.1BASS 12563961-2718455 16.42 L3: β 12574463-3635431 14.57 M6.6 M6:: γ TWA 25.6 < 0.1LP-BASS 12574941-4111373 13.02 M5.9 M6 γ VL-G 2n20 TWA 67.2 < 0.1BASS 15104786-2818174 12.84 M8 49 M9.3 M9 β INT-G 2n11 ARG 59.1 60.2 PRE-BASS 15291017+6312539 M7.8 M8 β INT-G 1n10 ABDMG 24.6 79.2 PRE-BASS 11 64 15470557-1626303 A 13.86 M9.6 M9 β INT-G 0n11 ABDMG 10.6 63.4 PRE-BASS 13.86 VL-G ABDMG 10.6 63.4 PRE-BASS 15470557-1626303 B M5:: nn2n

Table 4

PPLEMENT SERIES, 219:33 (54pp), 2015 August

The

ASTROP

HYSICAL

JOURNAL

 $S_{\rm C}$

Gagné et

A

							(Continue	d)							
2MASS				Spectr	al Type ^a			Ind. G	ravity ^b		YM	G Membership			Source
Designation	J	Lit. Opt.	References	Lit. NIR	References	H ₂ O Ind.	Adopted	Class	Score	Lit.	References	Updated BASS ^c	$B_P\left(\% ight)^{ m d}$	$C_P (\%)^{\mathrm{e}}$	Catalog
							Bona Fide Mer	nbers							
19350976-6200473	16.25					L1.0	L1 γ	VL-G	2212			THA	20.8	0.2	BASS
20004841-7523070	12.73	M9	25			M9.2	M9 γ	VL-G	2n22	CAS;BPMG	5,54	BPMG(69); ARG(28)	99.1	13.0	BASS
20113196-5048112	16.42					L2.4	L3 γ	VL-G	2222			THA	42.6	< 0.1	BASS
20224803-5645567	11.76	M5.5	2			M5.5	M5.5 β	INT-G	nn1n			THA	86.2	< 0.1	BASS
20282203-5637024	13.84					M8.0	M8.5 γ	VL-G	2n22			THA	44.3	< 0.1	BASS
20334473-5635338	15.72					L1.2	L0 γ	VL-G	1221			THA	93.4	< 0.1	BASS
20334670-3733443	10.85	M5	6			M6.6	M6: β	INT-G	1n10			BPMG	97.4	10.8	BASS
20391314-1126531	13.79	M8	7			M7.4	M7 β	INT-G	0n11	Pleiades	54	ABDMG	2.2	46.6	PRE-BASS
20505221-3639552	13.00					M5.2	M5 β	INT-G	nn1n			ARG(66); BPMG(34)	85.9	47.9	LP-BASS ^f
21121598-8128452	10.67					M5.5	M5.5 β	INT-G	nn1n			THA	44.6	< 0.1	BASS
21324036+1029494	16.59			L4.5:	39	L2.0	L4: β	FLD-G	00?1			ARG	30.8	61.6	PRE-BASS
21490499-6413039	10.35	M4.5	42			M4.1	M4.5			THA	20	THA	99.7	< 0.1	BASS
21543454-1055308	16.44			L4 INT-G	56	L3.7	L5 β/γ	INT-G	0n11	ARG	56	ARG	83.8	25.1	BASS
21544859-7459134	14.29					M9.8	M9.5: β	VL-G	2n2n			THA	99.4	< 0.1	BASS
21572060+8340575	13.97	L0	15				M9 γ o					ABDMG	30.8	62.9	PRE-BASS
22025794-5605087	14.36					M6.2	M9: γ	VL-G	1n22			THA	98.4	< 0.1	BASS
22064498-4217208	15.56	L2	11			L1.9	L3 γ	VL-G	2?12	ABDMG	5	ABDMG	99.2	1.1	BASS
22064498-4217208	15.56	L2	11			L1.0	L3 γ	VL-G	1222	ABDMG	5	ABDMG	99.2	1.1	BASS
22191486-6828018	13.92					M6.0	M6 β	VL-G	1n22			THA	28.3	< 0.1	LP-BASS
22351658-3844154	15.18					L1.4	L1.5 γ	VL-G	2222			THA	96.2	< 0.1	BASS
22353560-5906306	14.28					M8.6	M8.5 β	INT-G	1n11			THA	99.8	< 0.1	BASS
22444835-6650032	11.03	M5	20			M5.1	M5 γ	VL-G	nn2n	THA	20	THA	99.8	< 0.1	BASS
22511530-6811216	12.10					M7.4	M5: γ	VL-G	nn2n			THA	99.9	< 0.1	BASS
23130558-6127077	10.93	M4.5	20			M5.2	M5 β	INT-G	nn1n	THA	20	THA	99.9	< 0.1	BASS
23143092-5405313	11.50					M5.0	M5 β	INT-G	nnln			THA	99.6	< 0.1	LP-BASS
23225240-6151114	11.53	M5	5			M5.2	M5 γ	INT-G	nn1n	THA	5	THA	99.9	< 0.1	BASS
23225299-6151275	15.55	L2 γ	32	L2	57	L1.8	L1 γ	VL-G	1221	THA	5	THA	>99.9	< 0.1	BASS
23231347-0244360	13.58	M8.5	28			M7.7	M8 β	INT-G	1n?1			BPMG	30.6	54.4	PRE-BASS
23255604-0259508	15.96	L3:	9	L3	9	L1.3	L1 γ	INT-G	1111			ABDMG	73.4	12.3	BASS
23255604-0259508	15.96	L3:	9	L3	9	L2.8	L1 γ	INT-G	0121			ABDMG	73.4	12.3	BASS
23353085-1908389	11.51					M5.4	M5 β	INT-G	nnln	•••		ABDMG (86);	84.8	9.2	BASS ^f
												BPMG(13)			
23355015-3401477	11.64					M5.1	M6: γ	VL-G	2n1n			BPMG	76.8	31.1	BASS
23360735-3541489	14.65					M8.6	M9 β	VL-G	1n22			ABDMG (60);	50.8	30.3	BASS
23433470-3646021	16.57					L3.7	L3–L6 γ	VL-G	2012			THA(39) ABDMG (46):BPMG	68.9	4.8	BASS
												(38); THA(16)			
23520507-1100435	12.84	M7	28			M7.6	M8 β	INT-G	1n11			ABDMG	90.6	4.0	BASS
23532556-1844402 A	11.24					M5.8	Μ6.5 γ	VL-G	0n22			THA(46); BPMG(34); ABDMG(20)	61.4	<0.1	LP-BASS
23532556–1844402 B	11.24					M5.2	M4.5 pec					THA(46); BPMG(34);	61.4	< 0.1	LP-BASS
					(andidate Merr	hers with no Cons	traint on Surfac	e Gravity			ADDiviO(20)			
00020382+0408120 *	10.40				,		M3					ARDMG	90.6	0.7	PRF-RASS
00020382+0408129 R	10.40						M3					ABDMG	99.6	0.7	PRE-BASS

Table 4

The Astrophysical Journal Supplement Series, 219:33 (54pp), 2015 August

15

GAGNÉ ET AL.

							Table 4(Continue)	d)							
2MASS				Spectr	al Type ^a			Ind. G	ravity ^b		YM	G Membership			Source
								~				Updated	n (m)d		
Designation	J	Lit. Opt.	References	Lit. NIR	References	H ₂ O Ind.	Adopted	Class	Score	Lit.	References	BASS	$B_P(\%)^{*}$	$C_P(\%)^2$	Catalog
00171571 2010520	10.64	244.5	6			N(4.2	Bona Fide Mer	nders				DDMC(5()	82.0	17.6	
001/15/1-3219539	10.64	M4.5	0			M4.3	M4					ARG(44)	83.0	17.6	PRE-BASS
00390342+1330170 A	10.94					M5.8	M4 pec			ABDMG	12	BPMG(92); ABDMG(8)	91.9	11.2	BASS
00390342+1330170 B	10.94					M6.0	M5 pec			ABDMG	12	BPMG(92); ABDMG(8)	91.9	11.2	BASS
01035369–2805518 A	11.66	M4.5	17			M4.8	M4					ABDMG	83.2	3.5	PRE-BASS
01035369-2805518 B	11.66					M4.9	M4					ABDMG	83.2	3.5	PRE-BASS
03132588–2447246	12.53					M4.8	M5.5: pec					BPMG(47); THA(49)	97.8	<0.1	LP-BASS
03442859+0716100 A	12.72						M4					BPMG	27.5	85.1	PRE-BASS
03442859+0716100 B	12.72					M4.1	M4.5					BPMG	27.5	85.1	PRE-BASS
03582255-4116060	15.85	L5	28			L5.4	L6 pec		nnn?			BPMG	67.1	16.6	BASS
04173836-1140256	11.75						M2:					ARG(76); BPMG(24)	12.4	92.9	PRE-BASS
04231498-1533245	12.54					M4.7	M4: pec					BPMG	95.4	1.3	LP-BASS
05104958-1843548	15.35					L2.5	L2: β ?	INT-G	1?1?			COL	68.6	6.6	LP-BASS
05201794+0511521	13.04						K0					COL(51); BPMG(49)	9.0	62.5	PRE-BASS
05484454–2942551	10.56					M4.2	M4::					COL(56); BPMG(43)	62.4	24.0	PRE-BASS
06021735-1413467	14.34						<K0					COL	81.8	25.3	PRE-BASS
07583046+1530004	10.43	M4.5	38				M4					TWA(45); ARG(39); ABDMG(16)	35.7	<0.1	PRE-BASS
07583098+1530146 A	9.97	M3.5	38			M4.2	M4.5					ARG(69); TWA(31)	47.7	41.2	PRE-BASS
07583098+1530146 B	9.97	M3.5	38			M4.3	M4.5					ARG(69); TWA(31)	47.7	41.2	PRE-BASS
08045433-6346180	9.93						<m2< td=""><td></td><td></td><td></td><td></td><td>CAR</td><td>98.8</td><td>1.8</td><td>PRE-BASS</td></m2<>					CAR	98.8	1.8	PRE-BASS
08095903+4434216	16.44			L6	39	L5.4	L6 pec(red)	INT-G	nnn1			ARG	80.7	27.4	BASS
08540240-3051366	9.01	M4	42				M4					BPMG	89.1	23.6	PRE-BASS
09104094-7552528	13.62						<k0< td=""><td></td><td></td><td></td><td></td><td>ARG(59); CAR(28); PPMC(10)</td><td>69.9</td><td>73.8</td><td>PRE-BASS</td></k0<>					ARG(59); CAR(28); PPMC(10)	69.9	73.8	PRE-BASS
09510459+3558098	10.58	M4 5	44				M5					ABDMG	18.9	31.1	PRF-BASS
11195251-3917150	13.13						M3					TWA	99.8	< 0.1	PRE-BASS
11532691–3015414	12.31					M4.7	M4.5					TWA(84);	6.1	<0.1	PRE-BASS
12002750-3405371	9.61						M4					TWA	97 7	< 0.1	PRE-BASS
12492353-2035592	9.32						M2					TWA	3.6	< 0.1	PRE-BASS
13262009-2729370	15.85	L5	49	L6.5:	19	L5.9	L7		nnnn			ARG	85.5	18.7	BASS
19480544+5944412 A	11.49					M4.2	M4					ARG(86); ABDMG(14)	16.8	69.6	PRE-BASS
19480544+5944412 B	11.49					M4.2	M4.5					ARG(86); ABDMG(14)	16.8	69.6	PRE-BASS
23102196-0748531	11.60					M5.0	M5					BPMG(61); ABDMG(39)	96.9	8.3	LP-BASS
23290437+0329113	11.11					M5.2	M5.5 pec Field Contami	 nants				BPMG	67.5	30.0	BASS ^f
00045753-1709369	11.00	M5.5	2	M5.5	3	M5.6	M6	FLD-G	0n00			Field			PRE-BASS
00193927-3724392	15.52	L3:	9	L3.5:	9	L2.2	L3	INT-G	10?1			Field			PRE-BASS
00210589-4244433	13.52	M9.5	10		-	M9.8	L0.5	FLD-G	000?			Field			PRE-BASS

16

GAGNÉ ET AL.

The Astrophysical Journal Supplement Series, 219:33 (54pp), 2015 August

(Continued) 2MASS Ind. Gravity^b YMG Membership Spectral Type^a Source Updated BASS Designation JLit. Opt. References Lit. NIR References H2O Ind. Adopted Class Score Lit. References $B_{P}(\%)^{d}$ $C_{P}(\%)^{e}$ Catalog Bona Fide Members 14.40 M9.7 FLD-G PRE-BASS 00461551+0252004 L0 pec 1000 Field 18 L4.5 19 L3.0 FLD-G 2010 5 PRE-BASS 01291221+3517580 16.78 L4 L3.5 ARG Field 01550354+0950003 14.82 L5 15 L5: 9 L3.2 L4 INT-G 101? Field 65.3 23.7 PRE-BASS 15.34 FLD-G 02441019-3548036 L0.8 L2 pec 1000 Field BASS 23 02534448-7959133 11.34 M5.5 M7.1 M6 pec FLD-G 0n10 Field BASS 03005033-5459267 12.42 M5.5 FLD-G nn0n Field PRE-BASS M5 12.53 L0 15,25 M9.3 M9 pec FLD-G 0n00 UMA 03140344+1603056 UMA 26 PRE-BASS 12.54 M8.7 FLD-G 03204919-3313400 M5.5: nn0n Field BASS 03263956-0617586 12.96 M5 27 M5.0 M5 FLD-G nn0n Field PRE-BASS 03333313-3215181 M6.3 M6.5 FLD-G LP-BASS 13.17 0n10 Field 03370359-1758079 15.62 L4.5 11 L4.3 L4 FLD-G 1000 Field LP-BASS 03370362-3709236 12.75 M5.8 M5.5 FLD-G nn0n Field LP-BASS 04070752+1546457 15.48 L3.5 15 L3.5 22 L3.0 L3 FLD-G 2100 Field PRE-BASS 04532647-1751543 L2.5 INT-G 8.0 15.14 L3: L3 1110 Field 96.5 BASS 7 04584239-3002061 13.50 M6.1 M6.5 FLD-G 0n00 Field LP-BASS 13.67 L4 15 L4 34 L4.2 L4 pec FLD-G 0000 Field BASS 05002100+0330501 05431887+6422528 13.57 L1 15 L2 36 L1.9 L2 INT-G 1110 Field PRE-BASS 06022216+6336391 14.27 L1: 15 L1.5 19 L1.7 L2 FLD-G 1??0 Field BASS 07083261-4701475 14.16 M8.8 M8.5 FLD-G 0n00 Field PRE-BASS 37 M9: M9.8 FLD-G 1000 Field PRE-BASS 07200325-0846499 10.63 L0 pec 08055944+2505028 A 11.53 M4.3 M4 Field PRE-BASS 08055944+2505028 B 11.53 M5.2 M5 FLD-G nn0n Field PRE-BASS 08141769+0253199 M5.2 M5 FLD-G Field PRE-BASS 11.52 nn0n 08194351-0450071 14.82 L0.7 L1: pec FLD-G 010n Field PRE-BASS 08204440-7514571 16.59 L2.8 L3.5 INT-G 101? Field PRE-BASS 08254335-0029110 15.45 L0.3 L0.5 FLD-G 1010 Field PRE-BASS $08255896 {+} 0340198$ Field 10.01 M3 PRE-BASS 08503593+1057156 16.47 L6+L6 40,41 L6.2 L7 pec(red) u FLD-G nnn0 Field PRE-BASS 08511627+1817302 16.57 L4.5: 39 L4.4 L5: FLD-G 0n00 Field PRE-BASS 15.04 L8 35 43 08575849+5708514 L8: L7.1 L8 pec Field PRE-BASS FLD-G 10051641+1703264 11.13 M4.9 M5 nn0n Field PRE-BASS 12.79 M5.1 M5 pec FLD-G nn0n Field PRE-BASS 10130718-1706349 A 12.79 M5.1 FLD-G 10130718-1706349 B M5 pec nn0n Field PRE-BASS 10352029-2058382 11.66 M5.5 M5.5 FLD-G nn0n Field PRE-BASS 10513331-1916530 14.69 L0.3 M9 pec FLD-G 0n00 Field BASS 10584787-1548172 14.15 L3 18 L3 1 L2.2 L3 FLD-G 0000 Field BASS 47 f 11335700-7807240 13.20 M8 M6.2 M6: pec FLD-G 0n01 Field PRE-BASS 11555389+0559577 15.66 L0 48 L7.5 1 L6.8 L6-L8 pec FLD-G nnn0 Field PRE-BASS M9.7 FLD-G 0010 PRE-BASS 12042529-2806364 16.11 1.05 Field 12212770+0257198 13.17 L0 15 M9.7 M9 pec FLD-G 1n00 Field PRE-BASS 12310489-3801065 14.68 M8.0 M8 pec FLD-G 0n10 Field BASS 12521062-3415091 11.65 M5.2 M5.5 FLD-G nn0n Field PRE-BASS 13015465-1510223 14.54 L1 15 M9.6 L1.5: FLD-G 0001 Field PRE-BASS 13252237+0600290 12.25 < M0Field PRE-BASS ... 17065487-1314396 14.52 L5.0 L5 pec FLD-G 0n00 Field PRE-BASS 11.01 53 M6.9 FLD-G 0n00 Field M6 5 M7 PRE-BASS 18393308+2952164 15.06 L0.2 INT-G 2020 Field PRE-BASS 18462188-5706040 L1: 19395435-5216468 14.66 M9.7 L1 INT-G 1021 Field BASS 20025265-1316418 14.48 M8.9 M8.5 FLD-G 0n00 Field PRE-BASS 20050639-6258034 11.75 M4.3 M5 FLD-G nn0n Field PRE-BASS 20414283-3506442 14.89 L2: 28 L2 19 L0.7 L2 FLD-G 1010 Field BASS 20482880-3255434 14.71 M9.4 M9 FLD-G 0n00 Field PRE-BASS 20484222-5127435 10?0 15.38 L1.9 L2 pec FLD-G Field BASS

Table 4

GAGNÉ ET

Ł

The

ASTROP

HYSICAL

JOURNAL

SUPPLEMENT

SERIES,

219:33

(54pp),

201

5

Augu

IST

							Table 4(Continue)	k d)							
2MASS				Spectra	al Type ^a			Ind. G	ravity ^b		YMG Membership				Source
Designation	J	Lit. Opt.	References	Lit. NIR	References	H ₂ O Ind.	Adopted	Class	Score	Lit.	References	Updated BASS ^c	$B_P (\%)^{\mathrm{d}}$	$C_P\left(\% ight)^{\mathrm{e}}$	Catalog
							Bona Fide Mer	mbers	1				-		
20484222-5127435	15.38					L0.7	L2 pec	FLD-G	1001			Field			BASS
21144103-4339531	13.02					M9.0	M7.5 pec	FLD-G	0n00	CAS	54	Field			LP-BASS
21272613-4215183	13.32	M7.5	15	M8	19	M8.5	M8	FLD-G	0n00	Pleiades	54	Field			PRE-BASS
21342814-1840298	11.04						M3: pec					Field			PRE-BASS
21420580-3101162	15.84	L3	55	L2	9	L1.5	L3	FLD-G	10?0			Field			BASS
21484123-4736506	10.97					M4.4	M5	FLD-G	nn0n			Field			PRE-BASS
22021125-1109461	12.36	M6.5	29			M7.2	M7	FLD-G	0n00			Field			LP-BASS
22062157-6116284	16.61					L1.0	L0: pec	FLD-G	1020			Field			PRE-BASS
22400144+0532162	11.72					M5.7	M6 pec	FLD-G	0n02			Field			BASS
22444905-3045535	14.65					L0.2	M9 pec	FLD-G	0n00			Field			PRE-BASS
23155665-4747315	16.08					L5.2	L3 pec	FLD-G	1010			Field			BASS
23270843+3858234	11.74					M6.3	M5.5 pec	FLD-G	nn0n			Field			BASS
23310161-0406193	12.94	M8 + L3	58		<u>^</u>	M8.4	M8 pec u	FLD-G	1n00			Field			LP-BASS
23392527+3507165	15.36	L3.5	15	L4.5	9	L3.0	L4 pec	FLD-G	1020		_	Field			BASS
23512200+3010540	12.47	L5.5	59	L5 p(red)	59	L3.9	L5 pec	FLD-G	0n01	ARG	5	Field	92.8	5.8	PRE-BASS
							Young Contam	inants							
02530084+1652532	8.39	M6.5	21	M7	22	M7.4	M7.5 β	INT-G	1n12	ARG	5	Field			PRE-BASS
05243009+0640349	11.98					M5.5	M5.5 β	INT-G	nn1n			Field			PRE-BASS
06353541-6234059	12.42					M6.3	M6.5 β	INT-G	0n12			Field			PRE-BASS
13582164-0046262	10.81					M5.5	M5.5 γ	VL-G	nn2n			Field			PRE-BASS
14112131-2119503	12.44	M9	7	M9	52	M8.7	M8.5 β	INT-G	1n10			Field			PRE-BASS
20385687-4118285	11.66					M5.2	M5 β	INT-G	nn1n			Field			PRE-BASS
23453903+0055137	13.77	M9	15			M9.7	M9 β	INT-G	1n11			Field			PRE-BASS
							Reddened Contai	minants							
00174858-0316334	13.23					M8.2	M7 β					Field			PRE-BASS
04044052+2616275 A	12.65						<m0< td=""><td></td><td></td><td></td><td></td><td>TAU?</td><td></td><td></td><td>PRE-BASS</td></m0<>					TAU?			PRE-BASS
04044052+2616275 B	12.65						<m0< td=""><td></td><td></td><td></td><td></td><td>TAU?</td><td></td><td></td><td>PRE-BASS</td></m0<>					TAU?			PRE-BASS
04281061+1839021	13.38						<m3< td=""><td></td><td></td><td></td><td></td><td>TAU?</td><td></td><td></td><td>PRE-BASS</td></m3<>					TAU?			PRE-BASS
05271676+0007526 A	12.17						M0					OMC			PRE-BASS
05271676+0007526 B	12.17						M3					OMC			PRE-BASS
05370704-0623170	15.70						<m0< td=""><td></td><td></td><td></td><td></td><td>OMC?</td><td></td><td></td><td>PRE-BASS</td></m0<>					OMC?			PRE-BASS
05404919-0923192	11.31						<m0< td=""><td></td><td></td><td></td><td></td><td>Field</td><td></td><td></td><td>PRE-BASS</td></m0<>					Field			PRE-BASS
05410983-0737392	13.46						<m0< td=""><td></td><td></td><td></td><td></td><td>Field</td><td></td><td></td><td>PRE-BASS</td></m0<>					Field			PRE-BASS
05415929-0217020	13.22						<m0< td=""><td></td><td></td><td></td><td></td><td>Field</td><td></td><td></td><td>PRE-BASS</td></m0<>					Field			PRE-BASS
05451198-0121021	13.83						<m0< td=""><td></td><td></td><td></td><td></td><td>Field</td><td></td><td></td><td>PRE-BASS</td></m0<>					Field			PRE-BASS
11014673-7735144	15.97					M4.2	<m5:< td=""><td></td><td></td><td></td><td></td><td>CHA?</td><td></td><td></td><td>PRE-BASS</td></m5:<>					CHA?			PRE-BASS
11560224-4043248	16.00						<m5< td=""><td></td><td></td><td></td><td></td><td>Field</td><td></td><td></td><td>PRE-BASS</td></m5<>					Field			PRE-BASS
12214223-4012050	16.47						<m0< td=""><td></td><td></td><td></td><td></td><td>Field</td><td></td><td></td><td>BASS</td></m0<>					Field			BASS
15424676-3358082	17.02						<k0< td=""><td></td><td></td><td></td><td></td><td>Lupus?</td><td></td><td></td><td>PRE-BASS</td></k0<>					Lupus?			PRE-BASS
16210134-2346554	15.16						<m5< td=""><td></td><td></td><td></td><td></td><td>SCC?</td><td></td><td></td><td>PRE-BASS</td></m5<>					SCC?			PRE-BASS
16210134-2346554	15.16						<m5< td=""><td></td><td></td><td></td><td></td><td>SCC?</td><td></td><td></td><td>PRE-BASS</td></m5<>					SCC?			PRE-BASS
16221255-2346418	10.90						<m5< td=""><td></td><td></td><td></td><td></td><td>SCC?</td><td></td><td></td><td>PRE-BASS</td></m5<>					SCC?			PRE-BASS
16232017-2353248	13.38					M7.0	M5 β					SCC?			PRE-BASS
16251377-2358021	13.75						<m0< td=""><td></td><td></td><td></td><td></td><td>ρOPH?</td><td></td><td></td><td>PRE-BASS</td></m0<>					ρ OPH?			PRE-BASS
16272178-2411060	13.98						<m0< td=""><td></td><td></td><td></td><td></td><td>ρOPH?</td><td></td><td></td><td>PRE-BASS</td></m0<>					ρ OPH?			PRE-BASS
16330142-2425083	16.16						<m5< td=""><td></td><td></td><td></td><td></td><td>SCC?</td><td></td><td></td><td>PRE-BASS</td></m5<>					SCC?			PRE-BASS
16422788-1942350	15.23						<m5< td=""><td></td><td></td><td></td><td></td><td>SCC?</td><td></td><td></td><td>PRE-BASS</td></m5<>					SCC?			PRE-BASS
18460473+5246027 A	11.03						K0					Field			PRE-BASS
18460473+5246027 B	11.03						K0					Field			PRE-BASS
19033113-3723302	13.41						<m0< td=""><td></td><td></td><td></td><td></td><td>Field</td><td></td><td></td><td>PRE-BASS</td></m0<>					Field			PRE-BASS

Table	4
Continu	(led

2MASS			Spectral Type ^a					Ind. G	ravity ^b	YMG Membership				Source	
Designation	J	Lit. Opt.	References	Lit. NIR	References	H ₂ O Ind.	Adopted	Class	Score	Lit.	References	Updated BASS ^c	$B_P\left(\% ight)^{\mathrm{d}}$	$C_P (\%)^{\mathrm{e}}$	Catalog
							Bona Fide Mer	nbers			1				
22573768-5041516	14.96						<m5< td=""><td></td><td></td><td></td><td></td><td>Field</td><td></td><td></td><td>PRE-BASS</td></m5<>					Field			PRE-BASS

Notes. References to this Table: (1) Knapp et al. (2004), (2) Crifo et al. (2005), (3) Deshpande et al. (2012), (4) Martín et al. (2010), (5) Gagné et al. (2014c), (6) Reid et al. (2007), (7) Cruz et al. (2003), (8) Schlieder et al. (2012b), (9) Burgasser et al. (2010), (10) Basri (2000), (11) Kirkpatrick et al. (2000), (12) Schlieder et al. (2012a), (13) Reiners & Basri (2010), (14) Reiners & Basri (2009), (15) Reid et al. (2008), (16) Marocco et al. (2013), (17) Bochanski et al. (2005), (18) Kirkpatrick et al. (1999), (19) Bardalez Gagliuffi et al. (2014), (20) Kraus et al. (2014b), (21) Teegarden et al. (2003), (22) Burgasser et al. (2008), (23) Phan-Bao & Bessell (2006), (24) Rodriguez et al. (2013), (25) Schmidt et al. (2007), (26) Seifahrt et al. (2010), (27) West et al. (2008), (28) Cruz et al. (2007), (29) Reid et al. (2002), (30) Allers & Liu (2013), (31) Shkolnik et al. (2012), (32) Cruz et al. (2009), (33) Liu et al. (2013a), (34) Bardalez Gagliuffi et al. (2014), (35) Kirkpatrick et al. (2008), (36) Kirkpatrick et al. (2014), (37) Scholz (2014), (38) Gizis et al. (1997), (39) Chiu et al. (2006), (40) Faherty et al. (2011), (41) Dupuy & Liu (2012), (42) Riaz et al. (2006), (43) Geballe et al. (2002), (44) Shkolnik et al. (2009) (45) Schneider et al. (2012), (46) Gizis et al. (2000), (47) Luhman (2007), (48) Schmidt et al. (2014), (51) Rodriguez et al. (2011), (52) Kendall et al. (2004), (53) Reid et al. (2003), (54) Gálvez-Ortiz et al. (2010), (55) Liebert & Gizis (2006), (56) Gagné et al. (2014b), (57) Manjavacas et al. (2014), (58) Caballero (2007), (59) Kirkpatrick et al. (2010).

^a All spectral types determined in this work (fourth column) are based on NIR spectra, except those with the o suffix, which are based on optical data. A semi-colon in the optical and NIR spectral types indicates that the subtype is uncertain (± 1), and a double semi-colon indicates that the subtype is very uncertain (± 2 subtypes or more); pec indicates peculiar features; β and γ respectively indicate intermediate gravity and very low gravity, determined from a visual classification.

^b The index-based gravity classes and scores are defined by Allers & Liu (2013) and based on the FeH and VO features, alkali lines depth and the *H*-band continuum shape, respectively. We used our adopted spectral subtypes in their calculation. A score value of 0 indicates field gravity, 1 indicates intermediate gravity and 2 indicates a very low gravity. A score of n indicates that no conclusion can be drawn either because the spectroscopic indicator in question is not sensitive to gravity at this spectral type or because the data does not cover the wavelength of this index. A question mark indicates that the quality of the data is not good enough to draw any conclusion (i.e., the measurement is consistent with low gravity, but the measurement error bar overlaps with the field sequence). The final gravity score is taken as the median of these individual gravity scores, ignoring n or ? scores and taking the average of the two central values when an even number of scores is used.

^c Results reported in the BASS survey paper (Paper V), updated using data presented here.

^d Bayesian probability for membership in a YMG.

^e The probability that this object is a field contaminant, based on a Monte Carlo analysis and the Besançon galactic model (see text and Paper II).

^f Low-probability Sample.

(This table is available in its entirety in machine-readable form.)





Figure 3. NIR spectra of all new observations and objects for which spectral types were revised in this work. All spectra were re-sampled to a spectral resolution of $R \sim 120$ and a dispersion relation identical to SpeX observations in the prism mode with the 0.4 slite. We used alternating colors for visibility.



Figure 4. NIR Spectral type histogram of all known low-gravity dwarfs and those presented in this work. Green bars delimited by dashed lines represent the known population prior to BASS, purple bars delimited by dashed–dotted lines represent known dwarfs for which low-gravity features were identified here for the first time, and orange bars delimited by solid lines represent new discoveries from BASS. The BASS survey has contributed significantly in increasing the number of known low-gravity M6–L5 dwarfs.

potentially making them less interesting candidate members. It should be noted however that one of these three objects (2MASS J06494706–3823284) has weak Na I absorption consistent with a very low surface gravity. This demonstrates how the absence from the *ROSAT* catalog is not a strong enough constraint to reject any of our M0–M5 candidate members. Kraus et al. (2014b) has demonstrated that surveys for M-type moving group members based on either X-ray or UV-bright samples are incomplete because of the sky coverage and detection limits of current X-ray and UV catalogs.

5.3. Sources of Contamination

In Paper II, we demonstrated that a fraction of candidate members identified by the BANYAN II tool are expected to be field interlopers, especially if no prior knowledge is available on age. This fraction of contaminants is dependent on the YMG considered: ARG, ABDMG and β PMG are expected to be the most contaminated, mostly due to their proximity and their overlap with the galactic plane. Counting the fraction of lowgravity dwarfs in the spectroscopic follow-up presented here allows us to estimate minimal contamination rates of 18% and 33% in the BASS and LP-BASS samples, respectively. These values are slightly larger than the estimates that we derived in Paper V (12.6% for BASS and 26% for LP-BASS). The most likely explanation is that the kinematic distribution of field BDs is not perfectly reproduced by the Besançon galactic model, on which our previous estimates were based. The reason why these updated estimates correspond to a minimal contamination fraction is that some low-gravity dwarfs in our candidate sample could still be contaminants from associations not considered in BANYAN II, e.g., the Ursa Majoris moving group (UMA; ~300 Myr; Zuckerman & Song 2004), the Hercules-Lyrae moving group (250 Myr; Eisenbeiss et al. 2013), the ϵ Chamaeleontis association (also called Cha-Near; ~10 Myr; Zuckerman & Song 2004), the Octans association (30-40 Myr; Torres et al. 2008, Murphy & Lawson 2015), and the Carina-Near moving group (200 Myr; Zuckerman et al. 2006). Measurements of RV and trigonometric distance will be helpful to identify such contaminants. Besides field-gravity \ge M5 dwarfs, we identified other kinds of contaminants in our sample of candidates, based on our new NIR spectroscopy. We uncovered a number of objects with spectral types earlier than M5 (4 in BASS, 4 in LP-BASS, and 28 in PRE-BASS), for which there is no known reliable lowgravity indicators in the NIR. In addition to those, we uncovered 27 contaminants mostly in the PRE-BASS sample (only one was found in BASS) that correspond to K- and M-type low-mass stars reddened by interstellar dust in the line of sight (Figure 9). A number of these are likely located in starforming regions, such as ρ Ophiucus (ρ OPH), the Scorpius-Centaurus Complex (SCC), and Taurus-Aurigae (TAU; Elias 1978). These objects were all rejected from the BASS sample, mainly because (1) we avoided star-forming regions in the final survey; and (2) the extragalactic WISE color filter defined by Kirkpatrick et al. (2011) and the 2MASS crowding filter defined in Paper V efficiently rejected them.

6. DISCUSSION

6.1. Updated Color–Magnitude Sequences for Young Low-mass Stars and BDs

We complemented the list of all spectroscopically confirmed ≥L0 dwarfs as of 2014 February (Mace 2014) and the DwarfArchives online library¹⁷ with more recent discoveries, measurements of photometry in the literature and additional NIR photometry from a cross-match with 2MASS and WISE, in order to build an up-to-date sequence of field dwarfs. This list currently contains >1800 published \ge L0 low-mass stars and BDs.¹⁸ We compiled a similar list of > 8700 M6–M9 low-mass stars and BDs.¹⁹ These two lists of dwarfs contain photometric data from articles referenced throughout the present work.²⁰ In Figures 10 and 11, we compare our updated population of known young low-mass stars and BDs to the field sequence in various spectral type-color and spectral type-absolute magnitude diagrams. We used data from the two aforementioned lists to build the photometric sequences. In the case of YMG candidate members that do not have a trigonometric distance measurement, we used the statistical distance from BANYAN II, associated with the most probable YMG hypothesis. In each case, we calculated the error-weighted median sequence in bins of 1 subtype and adjusted a polynomial relation by minimizing the χ^2 value. We list in Table 7 the coefficients of these polynomial fits as well as the respective standard deviation of the data with respect to the best fit. We note that our field sequences are slightly redder than those derived from samples based on the Sloan Digital Sky Survey (SDSS; York et al. 2000) such as those presented by West et al. (2008) and Schmidt et al. (2015). This is true because SDSS-based surveys rely directly on spectra and are thus un-biased, whereas

¹⁷ http://dwarfarchives.org

¹⁸ Publicly available at www.astro.umontreal.ca/~gagne/listLTYs.php.

¹⁹ Publicly available at www.astro.umontreal.ca/~gagne/listMs.php.

 $^{^{20}}$ In addition to the following references: Andrei et al. (2011), Artigau et al. (2010), Beichman et al. (2014), Burgasser et al. (2006), Burningham et al. (2008, 2013), Castro et al. (2013), Costa et al. (2005, 2006), Cushing et al. (2011), Deacon et al. (2014), Delorme et al. (2008a, 2008b), Dieterich et al. (2014), Faherty et al. (2013), Hawley et al. (2002), Kendall et al. (2007a, 2007b), Kirkpatrick et al. (2012), Leggett et al. (2000, 2002, 2009, 2010, 2013, 2015), Lodieu et al. (2005, 2007), Looper et al. (2007a), Lucas et al. (2013), Mace et al. (2013b, 2013a), Marocco et al. (2010), Marsh et al. (2013), Monet et al. (1992), Pérez-Garrido et al. (2014), Phan Bao et al. (2008), Pinfield et al. (2003, 2014), van Leeuwen (2007), Warren et al. (2007), Wilson et al. (2003), Zapatero Osorio et al. (2014), and van Altena et al. (1995).



Figure 5. Low-resolution ($R \gtrsim 75$) gravity-sensitive NIR indices defined by Allers & Liu (2013) for all intermediate-gravity (green circles) and very low-gravity (red diamonds) dwarfs from the BASS sample. This sample consists mainly of new discoveries and known dwarfs with a new low-gravity classification. Previously known intermediate-gravity and low-gravity dwarfs from the samples of Allers & Liu (2013) and Manjavacas et al. (2014) are displayed as smaller, open symbols. The thick, blue line and the pale blue region delimited by dashed, purple lines represent the field sequence and its scatter. Random offsets smaller than 0.25 subtypes have been added to the spectral types for clarity. Lower-gravity dwarfs display (1) lower FeH_Z and KI_J indices at spectral types M5.5–L6; (2) larger *H*-cont indices at spectral types M5.5–L6; and (3) larger VO_Z indices at spectral types L0–L4 (see text for more detail). It is readily apparent that low-gravity dwarfs of the same spectral type can display a different set of low-gravity features, which is why a classification based on multiple gravity-sensitive indices is necessary (Allers & Liu 2013). The data used to create this figure are available.

other surveys based on 2MASS and/or *WISE* (e.g., Cruz et al. 2003; Reid et al. 2008; Kirkpatrick et al. 2011) perform a spectroscopic follow-up only on targets that were pre-selected from color cuts, which makes them biased toward detecting red objects more easily. Since 2MASS- and *WISE*-based surveys dominating the population of L dwarfs identified in the literature, our field sequences are consequently redder than those based on SDSS samples. This effect is also demonstrated in Figure 3 of Schmidt et al. (2010).

The radii of young low-mass stars and BDs are inflated compared with old objects of the same spectral type. For this reason, it could be expected that young absolute magnitude sequences fall above the field sequences across all spectral types. However, starting at spectral type ~L0, dust clouds form in the photosphere of BDs. Young BDs have a lower atmospheric pressure, which allows the formation of thick clouds higher in their atmosphere (Kirkpatrick et al. 2006; Looper et al. 2008b). As a result, a fraction of the NIR light at ~0.5–3 μ m gets redirected to longer wavelengths, causing young BDs to display similar absolute *J* magnitudes to those of field BDs around spectral type ~L0, as well as absolute *J* magnitudes even fainter than those of field dwarfs at later spectral types (Faherty et al. 2012, 2013; Liu et al. 2013a; Zapatero Osorio et al. 2014). We could expect that this effect



Figure 6. Moderate-resolution ($R \gtrsim 750$) gravity-sensitive NIR indices defined by Allers & Liu (2013) for all intermediate-gravity and very low-gravity dwarfs from the BASS sample. Symbols and color coding are identical to those of Figure 5. Lower-gravity dwarfs display weaker alkali and FeH absorption features, which results in lower Na 1 and K1 EWs and a lower FeH_J index. The data used to create this figure are available.

will eventually cease around spectral type T, where dust clouds fall below the photosphere. This has yet to be demonstrated, because there is only a very small number of young T dwarfs currently known (e.g., Delorme et al. 2012; Naud et al. 2014). In Figure 12, we show the spectral type at which the young and field sequences cross as a function of spectral band. Horizontal error bars represent the effective width of the photometric filters and vertical error bars are drawn from a 10,000-step Monte Carlo simulation, introducing noise in the data that is representative of photometric uncertainties and repeating the polynomial fit every time. Cases where the sequences do not cross are not included in the calculation of the median and standard deviation of the crossing points. We note that the fraction of Monte Carlo steps where the sequences cross significantly decreases at increasing wavelengths. This is explained by the fact that the photometric sequences become

gradually disjointed in the spectral range considered; it is thus possible that in reality the sequences generally cross at spectral types $\ge L7$ (or not at all) in the W1 and W2 bands. This figure shows a clear correlation which indicates that flux is redistributed out to longer wavelengths in low-gravity dwarfs, a likely effect of the dust clouds (J. K. Faherty et al. 2015, in preparation). This is a known effect which is in part due to the larger opacity from the H₂O, CO, and H₂ molecules at wavelengths larger than ~1 μ m that are masking the effects of clouds (Ackerman & Marley 2001). The BT-Settl isochrones (Baraffe et al. 2003; Allard et al. 2013) do not reproduce this effect, as the young (≤ 100 Myr) and old (≥ 1 Gyr) isochrones do not cross in neither of the *J*, *H*, or *K*_S bands over the range of effective temperatures that correspond to the M and L spectral types (~1300–3000 K; Stephens et al. 2009).

2MASS	Spectral		Eq	uivalent Widths (Å)		Spectral Indices (Allers & Liu 2013)					
Designation	Туре	Na 1 1.138 μm	K 1 1.169 μm	K 1 1.177 μm	K 1 1.244 μm	K 1 1.253 μm	FeHz	VOz	FeH _J	KI_J	H-cont	
					Intermediate-grav	ity (β)						
00011217+1535355	L4 β	7.3 ± 1.0	5.7 ± 1.0	7.0 ± 1.3		2.9 ± 1.4	1.22 ± 0.01	1.17 ± 0.01	1.24 ± 0.01	1.13 ± 0.01	0.92 ± 0.01	
00274534-0806046	M5.5 β	4.1 ± 0.7	1.3 ± 0.7	2.7 ± 0.9	0.6 ± 0.6	1.5 ± 0.6	1.06 ± 0.01	1.01 ± 0.01	1.04 ± 0.01	1.03 ± 0.01	0.97 ± 0.01	
00303013-1450333	L4–L6 β			••••			1.10 ± 0.03	1.03 ± 0.02		1.06 ± 0.02	0.92 ± 0.01	
00344300-4102266	L1: β	7.5 ± 1.5	6.0 ± 1.6	5.5 ± 1.6	2.9 ± 2.1	3.6 ± 1.6	1.18 ± 0.01	1.14 ± 0.01	1.15 ± 0.02	1.09 ± 0.01	0.96 ± 0.01	
00381489-6403529	M9.5 β	8.3 ± 1.4	4.1 ± 1.0	5.2 ± 1.3	3.6 ± 1.9	2.8 ± 1.2	1.12 ± 0.01	1.19 ± 0.01	1.18 ± 0.01	1.07 ± 0.01	1.01 ± 0.01	
00425923+1142104	M9 β	10.2 ± 1.4	4.1 ± 1.3	6.5 ± 1.7	3.0 ± 1.0	3.4 ± 1.3	1.17 ± 0.01	1.10 ± 0.01	1.16 ± 0.02	1.08 ± 0.01	0.95 ± 0.01	
00514561-6227073	M5.5: β	5.1 ± 1.2	2.0 ± 0.9	2.5 ± 0.8	5.1 ± 1.3	2.3 ± 0.7	1.07 ± 0.01	1.03 ± 0.01	1.06 ± 0.01	1.01 ± 0.01	•••	
00584253-0651239	L1 β	12.0 ± 1.4	6.7 ± 1.6	8.7 ± 1.8	2.5 ± 1.5	5.6 ± 1.3	1.23 ± 0.01	1.16 ± 0.01	1.19 ± 0.02	1.10 ± 0.01	0.93 ± 0.01	
01294256-0823580	M7 β	3.7 ± 1.0	0.3 ± 0.5	2.5 ± 0.7	1.6 ± 1.2	1.8 ± 0.6	1.04 ± 0.01	1.00 ± 0.01	1.04 ± 0.01	1.03 ± 0.01	1.03 ± 0.01	
01531463-6744181	L3 β	8.6 ± 0.2	5.6 ± 0.2	6.7 ± 0.2	3.2 ± 0.2	4.5 ± 0.2	1.14 ± 0.01	1.20 ± 0.01	1.19 ± 0.01	1.09 ± 0.01	0.92 ± 0.01	
)2103857-3015313	M9.5 β	7.5 ± 1.9	4.3 ± 1.4	5.7 ± 1.5	2.6 ± 1.9	4.1 ± 1.4	1.15 ± 0.01	1.25 ± 0.01	1.20 ± 0.01	1.08 ± 0.01	1.00 ± 0.01	
)2404759-4253377	M6 β	6.1 ± 0.9	1.2 ± 0.5	3.2 ± 0.9	1.6 ± 1.2	2.6 ± 0.7	1.05 ± 0.01	1.02 ± 0.01	1.07 ± 0.01	1.04 ± 0.01	1.00 ± 0.01	
02501167-0151295	M7 : β	7.2 ± 1.3	0.5 ± 1.1		6.7 ± 1.6		1.06 ± 0.01	0.99 ± 0.01	1.13 ± 0.01	1.05 ± 0.01	1.02 ± 0.01	
02530084+1652532	M7.5 β						1.07 ± 0.01	1.02 ± 0.01		1.05 ± 0.01	1.00 ± 0.01	
)3164512-2848521	L1 β						1.16 ± 0.01	1.18 ± 0.03		1.12 ± 0.02	0.95 ± 0.01	
)3224622-7940595	M6.5 β	5.1 ± 1.0	1.9 ± 0.6	1.3 ± 0.7	0.2 ± 0.8	1.9 ± 0.7	1.06 ± 0.01	0.99 ± 0.01	1.03 ± 0.01	1.04 ± 0.01	0.98 ± 0.01	
)3550477-1032415	M8.5 β	8.8 ± 1.2	3.2 ± 0.9	3.0 ± 1.2	3.7 ± 1.3	3.6 ± 1.0	1.11 ± 0.01	1.07 ± 0.01	1.15 ± 0.01	1.06 ± 0.01	0.97 ± 0.01	
04402583-1820414	M6 β	6.5 ± 1.2	-0.1 ± 0.9	2.0 ± 0.9		1.4 ± 0.8	1.05 ± 0.01	1.02 ± 0.01	1.09 ± 0.01	1.06 ± 0.01	0.99 ± 0.01	
)5071137+1430013 A	M5.5 β	4.9 ± 0.9	1.4 ± 0.7	1.9 ± 0.8		1.3 ± 0.6	1.06 ± 0.01	1.01 ± 0.01	1.04 ± 0.01	1.04 ± 0.01	1.00 ± 0.01	
)5071137+1430013 B	M5.5 B	42 ± 10	1.3 ± 0.7	2.0 ± 0.8	1.3 ± 0.9	18 ± 0.6	1.05 ± 0.01	1.00 ± 0.01	1.02 ± 0.01	1.03 ± 0.01	1.01 ± 0.01	
)5120636-2949540	L5 β			2.0 ± 0.0			1.21 ± 0.04	1.09 ± 0.02		1.10 ± 0.02	0.91 ± 0.02	
)5181131_3101529	M7 β	70 ± 09	1.7 ± 0.7	30 ± 09	1.7 ± 0.7	18 ± 0.6	1.21 ± 0.01 1.08 ± 0.01	1.09 ± 0.02 1.04 ± 0.01	1.08 ± 0.01	1.10 ± 0.02 1.05 ± 0.01	1.00 ± 0.01	
5243009 ± 0640349	M5 5 B	48 ± 0.7	0.9 ± 0.5	2.0 ± 0.5	1.7 ± 0.7 1.2 ± 0.4	1.0 ± 0.0 1.3 ± 0.4	1.00 ± 0.01 1.06 ± 0.01	1.01 ± 0.01 1.00 ± 0.01	1.00 ± 0.01 1.05 ± 0.01	1.03 ± 0.01 1.03 ± 0.01	1.00 ± 0.01 1.00 ± 0.01	
)5264316_1824315	M7 B	7.1 ± 1.0	18 ± 0.6	35 ± 0.0	1.2 ± 0.4 2.0 ± 0.6	1.5 ± 0.4 2.1 + 0.6	1.00 ± 0.01 1.09 ± 0.01	1.00 ± 0.01 1.02 ± 0.01	1.05 ± 0.01 1.08 ± 0.01	1.05 ± 0.01 1.04 ± 0.01	1.00 ± 0.01 1.00 ± 0.01	
)5/02325_0006326	MQ B	10.0 ± 1.0	5.0 ± 1.0	5.3 ± 0.9 5.2 ± 1.7	2.0 ± 0.0 2.5 ± 2.6	2.1 ± 0.0 4.4 ± 1.6	1.05 ± 0.01 1.16 ± 0.01	1.02 ± 0.01 1.11 ± 0.01	1.00 ± 0.01 1.16 ± 0.02	1.04 ± 0.01 1.09 ± 0.01	0.97 ± 0.01	
)63535/1_623/050	M65 B	10.0 ± 1.4 6.3 ± 1.2	3.0 ± 1.4 2.2 ± 0.8	3.2 ± 1.7 28 ± 0.9	2.5 ± 2.0 0.4 ± 1.0	4.4 ± 1.0 2.0 ± 0.8	1.10 ± 0.01 1.07 ± 0.01	1.11 ± 0.01 1.01 ± 0.01	1.10 ± 0.02 1.05 ± 0.01	1.09 ± 0.01 1.05 ± 0.01	1.01 ± 0.01	
08034460 + 0827000	M6.5 p	0.5 ± 1.2 8 8 \pm 1 1	2.2 ± 0.3 2.1 ± 0.7	2.0 ± 0.9 3.3 ± 1.0	0.4 ± 1.0 2.4 ± 0.6	2.0 ± 0.6 2.0 ± 0.6	1.07 ± 0.01 1.06 ± 0.01	1.01 ± 0.01 1.00 ± 0.01	1.05 ± 0.01 1.05 ± 0.01	1.03 ± 0.01 1.04 ± 0.01	1.01 ± 0.01 1.00 ± 0.01	
00054409 ± 0027000	MO B	0.0 ± 1.1 10.5 ± 1.7	2.1 ± 0.7 2.8 ± 1.2	5.5 ± 1.0 4.1 ± 1.6	2.4 ± 0.0 2.6 ± 2.0	2.0 ± 0.0 3.6 ± 1.4	1.00 ± 0.01 1.14 ± 0.01	1.00 ± 0.01 1.06 ± 0.01	1.05 ± 0.01 1.11 ± 0.01	1.04 ± 0.01 1.08 ± 0.01	1.00 ± 0.01 0.07 ± 0.01	
0522126 1014205	MO B	10.3 ± 1.7 8.1 ± 1.0	2.6 ± 1.2	4.1 ± 1.0 6.4 ± 1.5	2.0 ± 2.0 2.8 ± 1.1	3.0 ± 1.4	1.14 ± 0.01 1.12 ± 0.01	1.00 ± 0.01 1.22 ± 0.01	1.11 ± 0.01 1.16 ± 0.01	1.08 ± 0.01 1.08 ± 0.01	0.97 ± 0.01	
1544222 2400200	108	0.1 ± 1.0	4.0 ± 1.2	0.4 ± 1.3	2.0 ± 1.1	4.0 ± 1.1	1.13 ± 0.01 1.16 ± 0.02	1.22 ± 0.01 1.10 \pm 0.02	1.10 ± 0.01	1.08 ± 0.01 1.11 ± 0.02	0.90 ± 0.01	
11344223-3400390		 10.1 + 0.2	 6 9 1 6	···· 8 2 1 2 0	20 1 22	10 1 2 1	1.10 ± 0.02	1.19 ± 0.02	1 10 + 0.01	1.11 ± 0.02	0.94 ± 0.01	
12271345-0050456	12. P	12.1 ± 2.3 12.8 ± 0.1	0.0 ± 1.0	0.2 ± 2.0	2.0 ± 2.2	1.0 ± 2.1	1.04 ± 0.01	0.99 ± 0.01	1.10 ± 0.01	1.11 ± 0.01 1.07 ± 0.01	0.93 ± 0.01	
12303901-2716433	LS: ρ	12.0 ± 0.1	1.5 ± 0.2	5.5 ± 0.1	1.9 ± 0.2	2.7 ± 0.1	1.10 ± 0.01	0.97 ± 0.01	1.30 ± 0.01	1.07 ± 0.01	0.91 ± 0.01	
5104/60-26161/4	M9 p	0.3 ± 1.4	5.4 ± 1.0	4.7 ± 1.4	2.0 ± 1.7	5.0 ± 1.1	1.03 ± 0.01	1.11 ± 0.01	1.11 ± 0.01	1.08 ± 0.01	0.93 ± 0.01	
15291017+0512559	M8 p	8.9 ± 1.2	3.2 ± 0.9	4.7 ± 1.4	2.0 ± 0.8	3.4 ± 0.8	1.11 ± 0.01	1.05 ± 0.01	1.11 ± 0.01	1.06 ± 0.01	0.95 ± 0.01	
154/055/-1626505 A	M9 p	9.5 ± 1.5	4.0 ± 1.5	0.9 ± 1.0	3.0 ± 1.2	4.8 ± 1.2	1.18 ± 0.01	1.11 ± 0.01	1.16 ± 0.01	1.10 ± 0.01	0.95 ± 0.01	
20224805-5045507	M5.5 p	5.5 ± 0.9	0.1 ± 0.6	1.3 ± 1.0	2.2 ± 1.9	1.2 ± 0.7	1.09 ± 0.01	1.02 ± 0.01	1.05 ± 0.01	1.03 ± 0.01	1.01 ± 0.01	
20334670-3733443	M6: <i>β</i>	4.1 ± 0.9	1.7 ± 0.6	2.7 ± 0.8	5.6 ± 1.2	1.7 ± 0.5	1.06 ± 0.01	1.01 ± 0.01	1.10 ± 0.01	1.02 ± 0.01	0.99 ± 0.01	
20391314-1126531	M/β	9.5 ± 1.2	2.9 ± 1.0	4.2 ± 1.3	1.8 ± 0.9	3.3 ± 0.9	1.11 ± 0.01	1.05 ± 0.01	1.10 ± 0.01	1.06 ± 0.01	0.99 ± 0.01	
21121598-8128452	M5.5 β	5.3 ± 1.0	0.7 ± 0.6	1.3 ± 0.8	1.8 ± 1.0	1.7 ± 0.5	1.06 ± 0.01	1.01 ± 0.01	1.05 ± 0.01	1.02 ± 0.01	1.00 ± 0.01	
21324036+1029494	L4: β							1.02 ± 0.05		1.09 ± 0.04	0.92 ± 0.02	
21544859–7459134	M9.5: β	10.0 ± 1.7	3.8 ± 1.2	2.0 ± 1.3			0.98 ± 0.01	1.11 ± 0.01	1.09 ± 0.03			
22191486-6828018	M6 β	6.5 ± 1.3	1.5 ± 0.5	2.1 ± 0.9	1.6 ± 1.2	1.7 ± 0.7	1.05 ± 0.01	1.03 ± 0.01	1.06 ± 0.01	1.05 ± 0.01	1.01 ± 0.01	
22353560-5906306	M8.5 β	7.2 ± 1.4	3.5 ± 1.0	5.6 ± 1.5	2.2 ± 1.9	2.7 ± 1.2	1.14 ± 0.01	1.16 ± 0.01	1.16 ± 0.02	1.06 ± 0.01	0.97 ± 0.01	
23231347-0244360	M8 β	•••	•••	•••		•••	1.09 ± 0.01	1.05 ± 0.01	•••	1.07 ± 0.01	0.98 ± 0.01	
23360735-3541489	M9 β	9.4 ± 1.3	4.1 ± 1.2	6.3 ± 1.8	2.8 ± 2.0	3.0 ± 1.4	1.13 ± 0.01	1.12 ± 0.01	1.17 ± 0.01	1.07 ± 0.01	0.99 ± 0.01	

24

Table 5 Equivalent Widths and Spectral Indices

The Astrophysical Journal Supplement Series, 219:33 (54pp), 2015 August

Т	able	e 5

2MASS	Spectral		Eq	uivalent Widths (Å)			Spectral In	ndices (Allers &	Liu 2013)	
Designation	Туре	Na 1 1.138 μm	K 1 1.169 μm	Κ 1 1.177 μm	K 1 1.244 μm	K 1 1.253 μm	FeHz	VOZ	FeH _J	KI_J	H-cont
					Intermediate-grav	ity (β)					
23453903+0055137 23520507-1100435	M9 β M8 β						$\begin{array}{c} 1.12 \pm 0.01 \\ 1.11 \pm 0.01 \end{array}$	$\begin{array}{c} 1.11 \pm 0.01 \\ 1.05 \pm 0.01 \end{array}$		$\begin{array}{c} 1.07 \pm 0.01 \\ 1.07 \pm 0.01 \end{array}$	$\begin{array}{c} 0.97 \pm 0.01 \\ 0.98 \pm 0.01 \end{array}$
			,	V	ery Low-gravity	$(\gamma \text{ or } \delta)$					
00182834-6703130	L0 γ	6.1 ± 1.0	3.8 ± 1.2	4.0 ± 1.6	1.3 ± 1.3	1.4 ± 1.1	1.06 ± 0.01	1.34 ± 0.02	1.09 ± 0.02	1.03 ± 0.01	1.02 ± 0.01
00191296-6226005	L1 γ	8.8 ± 1.5	5.7 ± 1.4	5.6 ± 1.6	4.2 ± 2.1	3.8 ± 1.6	1.13 ± 0.01	1.29 ± 0.01	1.18 ± 0.02	1.11 ± 0.01	1.00 ± 0.01
00464841+0715177	L0 δ	7.5 ± 1.0	3.3 ± 1.1	3.5 ± 1.3	2.4 ± 1.0	1.7 ± 0.9	1.06 ± 0.01	1.27 ± 0.01	1.12 ± 0.01	1.05 ± 0.01	0.98 ± 0.01
01205114-5200349	L1 γ	5.1 ± 1.1	5.2 ± 1.4	6.5 ± 1.7	4.6 ± 2.1	2.7 ± 1.3	1.13 ± 0.01	1.27 ± 0.01	1.14 ± 0.01	1.08 ± 0.01	1.01 ± 0.01
01265327-5505506	M6 γ	7.3 ± 1.6	-0.4 ± 0.4	0.8 ± 0.6	1.2 ± 0.9	1.1 ± 0.6	1.07 ± 0.01	1.01 ± 0.01	1.05 ± 0.01	1.05 ± 0.01	0.98 ± 0.01
02265658-5327032	L0 γ	6.6 ± 1.4	3.8 ± 1.4	4.8 ± 1.4	3.2 ± 1.5	2.7 ± 1.1	1.07 ± 0.01	1.30 ± 0.01	1.13 ± 0.01	1.06 ± 0.01	0.93 ± 0.01
02410564-5511466	L1 γ	5.1 ± 1.1	5.0 ± 1.3	5.7 ± 1.5	3.5 ± 1.8	3.6 ± 1.4	1.19 ± 0.01	1.26 ± 0.01	1.15 ± 0.01	1.10 ± 0.01	0.97 ± 0.01
03264225-2102057	L5 β/γ						1.28 ± 0.11	1.08 ± 0.04		1.12 ± 0.02	0.94 ± 0.02
04185879-4507413	L3 γ	7.7 ± 2.5	6.6 ± 2.0	8.8 ± 2.1	5.7 ± 3.0	4.5 ± 1.9	1.08 ± 0.01	1.31 ± 0.01	1.18 ± 0.02	1.08 ± 0.01	0.94 ± 0.01
04400972-5126544	$L0 \gamma$	9.2 ± 1.6	4.9 ± 1.1	5.2 ± 1.5	0.5 ± 1.5	2.5 ± 1.2	1.11 ± 0.01	1.30 ± 0.01	1.20 ± 0.01	1.09 ± 0.01	1.05 ± 0.01
04493288+1607226	M9 γ	10.6 ± 1.3	4.2 ± 1.5	3.9 ± 1.4	3.1 ± 1.4	2.9 ± 1.1	1.07 ± 0.01	1.16 ± 0.01	1.10 ± 0.04	1.05 ± 0.01	1.00 ± 0.01
05123569-3041067	M6.5 γ	5.9 ± 1.0	0.8 ± 0.4	1.7 ± 0.8	4.0 ± 1.6	0.4 ± 0.6	1.07 ± 0.01	1.02 ± 0.01	1.05 ± 0.01	1.02 ± 0.01	0.98 ± 0.01
07202582-5617224	M6 γ	5.4 ± 0.8	0.9 ± 0.6	1.8 ± 0.8	2.1 ± 1.1	2.0 ± 0.6	1.05 ± 0.01	1.00 ± 0.01	1.05 ± 0.01	1.04 ± 0.01	0.99 ± 0.01
08561384-1342242	M8 γ	7.6 ± 1.1	1.6 ± 1.0	3.9 ± 1.2	2.5 ± 0.9	2.9 ± 0.9	1.12 ± 0.01	1.12 ± 0.01	1.12 ± 0.01	1.06 ± 0.01	0.98 ± 0.01
10284580-2830374	M6 γ	3.9 ± 0.6	0.3 ± 0.3	1.5 ± 0.4	1.0 ± 0.3	0.6 ± 0.3	1.04 ± 0.01	1.00 ± 0.01	1.04 ± 0.01	1.02 ± 0.01	1.01 ± 0.01
10455263-2819303	M6 γ	4.2 ± 0.7	0.8 ± 0.4	1.5 ± 0.5	1.1 ± 0.3	1.0 ± 0.3	1.05 ± 0.01	1.01 ± 0.01	1.05 ± 0.01	1.03 ± 0.01	1.01 ± 0.01
11064461-3715115	M9 γ	6.3 ± 1.0	0.7 ± 0.7	2.1 ± 0.9	1.7 ± 0.7	1.4 ± 0.7	1.06 ± 0.01	1.14 ± 0.01	1.04 ± 0.01	1.05 ± 0.01	1.02 ± 0.01
11083081+6830169	L1 γ	9.5 ± 1.2	6.2 ± 1.2	7.2 ± 1.4	5.2 ± 1.1	4.8 ± 1.0	1.30 ± 0.01	1.28 ± 0.01	1.14 ± 0.02	1.11 ± 0.01	0.95 ± 0.01
12074836-3900043	L1 δ						1.02 ± 0.02	1.26 ± 0.02		1.05 ± 0.02	1.00 ± 0.02
12474428-3816464	M9 γ						1.05 ± 0.01	1.10 ± 0.01		1.05 ± 0.02	0.99 ± 0.01
12574941-4111373	M6 γ	8.0 ± 1.4	0.0 ± 0.6	1.8 ± 0.8	0.6 ± 0.8	1.9 ± 0.6	1.03 ± 0.01	1.01 ± 0.01	1.03 ± 0.01	1.06 ± 0.01	0.99 ± 0.01
13582164-0046262	M5.5 γ	6.6 ± 0.9	1.0 ± 0.6	2.6 ± 0.8	1.4 ± 0.4	1.4 ± 0.5	1.07 ± 0.01	1.00 ± 0.01	1.05 ± 0.01	1.03 ± 0.01	0.99 ± 0.01
14252798-3650229	L4 γ						1.22 ± 0.02	1.11 ± 0.01		1.11 ± 0.02	0.94 ± 0.01
19350976-6200473	L1 γ	5.9 ± 2.0	4.9 ± 2.0	5.9 ± 1.7	0.8 ± 3.8	4.2 ± 1.9	1.11 ± 0.01	1.35 ± 0.01	1.09 ± 0.01	1.09 ± 0.01	1.01 ± 0.01
20113196-5048112	L3 γ	5.4 ± 1.5	1.3 ± 1.4	0.5 ± 1.5		1.2 ± 1.3	1.15 ± 0.01	1.31 ± 0.01	1.13 ± 0.02	1.05 ± 0.01	0.98 ± 0.01
20282203-5637024	M8.5 γ	4.5 ± 0.7	1.1 ± 0.7	2.8 ± 0.9	2.2 ± 1.0	1.3 ± 0.7	1.02 ± 0.01	1.12 ± 0.01	1.10 ± 0.01	1.00 ± 0.01	1.01 ± 0.01
20334473-5635338	$L0 \gamma$	5.7 ± 1.2	5.1 ± 1.3	5.8 ± 1.4	2.9 ± 2.1	2.7 ± 1.6	1.14 ± 0.01	1.27 ± 0.01	1.16 ± 0.02	1.09 ± 0.01	0.96 ± 0.01
21543454-1055308	L5 β/γ						1.27 ± 0.04	1.04 ± 0.03		1.06 ± 0.02	0.92 ± 0.01
22025794-5605087	M9: γ	9.5 ± 1.6	3.9 ± 1.2	7.0 ± 1.6	2.9 ± 2.4	2.4 ± 1.3	1.12 ± 0.01	1.13 ± 0.01	1.12 ± 0.02	1.05 ± 0.01	1.02 ± 0.01
22064498-4217208	L4 γ						1.11 ± 0.03	1.09 ± 0.04		1.08 ± 0.02	0.97 ± 0.01
22351658-3844154	L1.5 γ	5.5 ± 1.0	4.1 ± 1.3	4.9 ± 1.3	1.1 ± 1.5	2.0 ± 1.0	1.05 ± 0.01	1.31 ± 0.02	1.11 ± 0.01	1.04 ± 0.01	0.97 ± 0.01
23255604-0259508	L1 γ			•••			1.16 ± 0.01	1.16 ± 0.01		1.07 ± 0.02	0.93 ± 0.01
23433470-3646021	L3–L6 γ	8.1 ± 1.4	6.0 ± 1.1	6.6 ± 1.1	0.5 ± 1.6	2.9 ± 1.2	1.24 ± 0.01	1.03 ± 0.01	1.13 ± 0.02	1.07 ± 0.01	0.97 ± 0.01
23532556-1844402 A	M6.5 γ	6.0 ± 1.0	0.9 ± 0.8	1.7 ± 1.0	1.4 ± 1.0	1.8 ± 0.7	1.07 ± 0.01	1.01 ± 0.01	1.07 ± 0.01	1.03 ± 0.01	1.00 ± 0.01

25

					IC VISCU	Optical Spee	tial Types and	a spectral fild	lices					
2MASS	Optical						Spec	tral Indices						Low Gravity ^a
Designation	type	K-a	K-b	Na-a	Na-b	VO-a	Cs-a	Cs-b	TiO-b	Rb-a	Rb-b	FeH-a	CrH-a	
03164512-2848521	L1 β	3.27	2.90	0.975 \pm	$0.988~\pm$	1.244 \pm	1.125	1.071	1.374 \pm	1.214 \pm	1.1089 \pm	1.254 \pm	1.196 \pm	8 (1)
		$\pm 0.36^{b}$	± 0.38	0.053 ^b	0.059 ^b	0.095 ^b	± 0.036	± 0.050	0.061 ^b	0.075 [°]	0.080^{b}	0.043 ^b	0.045 ^b	
03264225-2102057	L5 β/γ	4.23	8.80	1.023 \pm	1.064 \pm	0.993 \pm	1.44	1.249	$1.058~\pm$	1.34	1.83	1.345	1.701 \pm	8 (1)
		$\pm 0.94^{b}$	$\pm 3.04^{b}$	0.037 ^b	0.046 ^b	0.030 ^b	$\pm 0.10^{\circ}$	± 0.071	0.022 ^b	$\pm 0.19^{b}$	± 0.27	± 0.053	0.065 ^b	
03421621–6817321	L4 γ	3.99	3.23	0.946 \pm	1.026 \pm	0.930	1.011 \pm	1.123 \pm	$0.873~\pm$	1.164 \pm	$1.281~\pm$	1.102 \pm	1.284 \pm	10(1)
	,	$\pm 0.61^{b}$	$\pm 0.46^{b}$	0.031 ^b	0.035 ^b	± 0.054	0.028 ^b	0.049 ^c	0.056 ^b	0.088 ^b	0.066 ^b	0.038 ^b	0.047 ^b	
06322402-5010349	L3 β	7.83	5.81	1.034 \pm	1.079 \pm	1.059 \pm	1.276	1.020 \pm	$1.081~\pm$	1.316 \pm	1.328 \pm	1.214 \pm	$1.555~\pm$	8 (3)
		$\pm 1.53^{b}$	$\pm 1.29^{\circ}$	0.022 ^b	0.036 ^b	0.031 ^b	± 0.066	0.033 ^b	0.020 ^b	0.099 ^c	0.078 [°]	0.023 ^b	0.028 ^b	
11083081+6830169	L1 γ	6.1	2.97	$0.909~\pm$	1.04	1.20	$1.091~\pm$	0.970 \pm	1.240 \pm	1.32	1.24	$1.067~\pm$	1.324 \pm	9(1)
	,	$\pm 1.7^{b}$	$\pm 0.72^{b}$	0.065 ^b	$\pm 0.12^{b}$	$\pm 0.10^{b}$	0.085 ^c	0.037 ^b	0.051 ^b	± 0.14	± 0.12	0.064 ^b	0.070 ^b	
11544223-3400390	L0 β	3.48	2.43	$1.098~\pm$	1.120 \pm	1.218 \pm	1.106 \pm	1.072	1.504 \pm	1.113 \pm	1.144 \pm	$1.135~\pm$	1.171	5 (5)
11577225-5400550		$\pm 0.38^{b}$	$\pm 0.25^{b}$	0.048 ^c	0.054 ^b	0.025 ^e	0.024 ^c	± 0.027	0.030 ^b	0.036 ^e	0.030 ^c	0.017 ^b	± 0.011	
21572060+8340575	M9 γ	2.33	1.54	$0.920~\pm$	0.943 \pm	1.33	$1.081~\pm$	1.047 \pm	1.61	1.090	$0.797~\pm$	$0.888~\pm$	$0.990~\pm$	9 (2)
	,	$\pm 0.23^{b}$	$\pm 0.14^{b}$	0.037 ^b	0.048 ^b	$\pm 0.14^{b}$	0.040 ^c	0.041 ^c	$\pm 0.10^{b}$	± 0.085	0.068 ^b	0.035 ^b	0.034 ^b	

 Table 6

 Revised Optical Spectral Types and Spectral Indices

Notes.

26

^a This column indicates the total number of spectral indices (out of 14) that are consistent with a low surface gravity. The number in parentheses indicates the total number of measurements that are consistent with an intermediate surface gravity.

^b This index is consistent with a low gravity at this spectral type.

^c This index is marginally consistent with a low gravity at this spectral type (the measurement is consistent with a low gravity but its error bars overlap with the field population).



Figure 7. Galactic position *XYZ* and space velocity *UVW* of the new AB Doradus bona fide member 2MASS J14252798–3650229 (red point and its projections), compared with other bona fide members of ABDMG (green points and their vertical projections on the *XY* and *UV* planes) and the SKM models of ABDMG (as defined in Paper II; orange ellipsoid and its projections).

In Figure 13, we show a M_J versus J - K CMD in the Mauna Kea Observatories NIR filter system (MKO; Simons & Tokunaga 2002) for low-gravity and field dwarfs. When MKO photometry was not available, we used 2MASS photometry with the conversion relations of Stephens & Leggett (2004; L and T dwarfs) and Leggett et al. (2006; M dwarfs). The combined effects of redder colors due to thicker/higher clouds (Marley et al. 2002) and brighter absolute J magnitude due to inflated radii cause a systematic shift of the low-gravity sequence to the right compared to the field sequence. This Figure brings into evidence the fact that the currently known population of young BDs does not reach a color reversal similar to the L/T transition of field dwarfs (at $J - K \sim 1.8$ and $M_I \sim 14.5$), corresponding to the temperature at which



Figure 8. NIR spectrum of the new L4 γ ABDMG bona fide member 2MASS J14252798–3650229 (thick black line), compared with various field and low-gravity L4 BDs. All spectra were degraded to a resolution of $R \sim 120$ and normalized at their median value in the $\sim 1.27-1.33 \mu m$ range. The *H*-band continuum of 2MASS J14252798–3650229 has a typical triangular shape and its global slope is particularly red, which are both telltale signs of low gravity.



Figure 9. NIR spectra of typical contaminants in the PRE-BASS sample. Resolution was degraded in the same way as described in Figure 1. All spectra were normalized to their median across the full wavelength range and shifted vertically for comparison purposes. The contaminants presented in this figure likely correspond to background K- and M-type stars reddened by interstellar dust. We used alternating colors for visibility.

dust clouds fall below the photosphere (Barman et al. 2011a; Dupuy & Liu 2012; Faherty et al. 2012, 2013; Bonnefoy et al. 2013; Liu et al. 2013a; Males et al. 2014; Zapatero Osorio et al. 2014). We chose this parameter space because a significant amount of data are available in these filters and it is very efficient in displaying this color reversal. It can be expected that a color reversal would eventually be reached for young dwarfs around the T spectral type, corresponding to cooler temperatures than the currently known population. The coolest known directly imaged young exoplanets and low-mass BDs (blue stars in Figure 13) tentatively hint at such a color reversal.

Since J - K and M_J are generally correlated for a given spectral type, the (J - K)-spectral type and M_J -spectral type relations listed in Table 7 are not the best representation for the low-gravity and field sequences in this CMD diagram. In the case of the young sequence, the absence of a color reversal



Figure 10. Absolute magnitude–NIR spectral type and color–NIR spectral type sequences for field (black diamonds) and young dwarfs (red dots when trigonometric distances were used), as well as polynomial sequences (blue and orange lines, respectively) defined in Table 7. We used the kinematic distances obtained from the BANYAN II tool (without photometry as input) to include low-gravity candidate members of YMGs that do not have a trigonometric distance measurement. Young dwarfs are generally brighter because of their inflated radii; however, thicker/higher dust clouds compete with this effect at spectral types L0–L7. Low-gravity L dwarfs are systematically redder than their field counterparts because of thicker/higher dust clouds in their photosphere. The data used to create this figure are available.

allowed us to simply fit a polynomial sequence to the young dwarfs directly in the $M_J - (J - K)$ space; however, the field sequence cannot be represented by a simple polynomial relation across the M6–T9 range. We used a Markov Chain Monte Carlo algorithm to construct a parametrized polynomial sequence that fits the field sequence across its complete spectral range. We started from a parametrized equation obtained from the combination of the J - K and M_J polynomial relations described in Table 7, and allowed the eight coefficients of each dimension to vary such that the sequence minimizes the quadrature sum of the bi-dimensional distance of all individual field dwarf positions in the CMD diagram relative to their error bars. This results in a parametrized sequence that describes J - K and M_J as a function of the parametric variable λ . Larger values of λ correspond to later spectral types on average, but no relation between λ and spectral types can be provided as the field sequence is a parametric equation that does not assign a λ value to individual data points. We obtain:

$$(M_J)_{\text{Young}} = 1.61 \times 10^3 - 8.92 \times 10^2 (J - K) + 2.04 \times 10^2 (J - K)^2 - 2.46 \times 10^1 (J - K)^3 + 1.65 (J - K)^4 - 5.87 \times 10^{-2} (J - K)^5 + 8.59 \times 10^{-4} (J - K)^6$$
(1)



Figure 11. Additional color–NIR spectral type sequences for young and field dwarfs as well as polynomial sequences defined in Table 7. The color scheme is identical to that of Figure 10 except that all young dwarfs are displayed with purple circles. The data used to create this figure are available.

$$(J - K)_{\text{Field}} = 1.98 \times 10^{1} - 1.55 \times 10^{1} \lambda + 5.09 \lambda^{2} - 8.76 \times 10^{-1} \lambda^{3} + 8.68 \times 10^{-2} \lambda^{4} - 5.12 \times 10^{-3} \lambda^{5} + 1.76 \times 10^{-4} \lambda^{6} - 3.27 \times 10^{-6} \lambda^{7} + 2.53 \times 10^{-8} \lambda^{8}$$
(2)

$$(M_J)_{\text{Field}} = -5.61 \times 10^1 + 3.73 \times 10^1 \lambda - 8.45 \lambda^2 + 1.04 \lambda^3 - 7.64 \times 10^{-2} \lambda^4 + 3.56 \times 10^{-3} \lambda^5 - 1.06 \times 10^{-4} \lambda^6 + 1.84 \times 10^{-6} \lambda^7 - 1.43 \times 10^{-8} \lambda^8.$$
 (3)

where the young sequence is valid in the range $8.8 \le M_J \le 14.7$ and the field sequence is valid in the range $4.5 \le \lambda \le 28.5$ (i.e., $J \ge -1.1$ and $9.3 \le M_J \le 19.9$).

We note that the NIR colors of young BDs discovered in the BASS and LP-BASS surveys are likely affected by a form of the confirmation bias, in the sense that we specifically looked for red objects in our survey (see Paper V). Hence, this new photometric data should not be taken as additional evidence that young BDs are redder than field BDs. Reinforcing this result would require looking for signs of low gravity in a sample of BDs that were selected independently of their photometric colors. Directly imaged young planets and BD companions do not suffer from this potential bias however, and their colors seem consistent with those of isolated young BDs (Chauvin et al. 2004, 2005; Barman et al. 2011a; Bonnefoy et al. 2013; Delorme et al. 2013; Bowler et al. 2013; Currie

et al. 2014). This might be an indication that our confirmation bias is not significant.

6.2. An Updated Investigation on the Age Dependence of Spectroscopic Indices

Since they are the only BDs with a well calibrated age, members of YMGs provide the exciting opportunity of creating a spectroscopic age calibration applicable to all young BDs. Using Pleiades members and the fact that known low-gravity BDs were located away from star-forming regions, Kirkpatrick et al. (2008) and Cruz et al. (2009) estimated that the very lowgravity (γ) and intermediate gravity (β) classifications likely correspond to ~ 10 and ~ 100 Myr, respectively. Allers & Liu (2013) extended this investigation by using a restrained sample of 25 M6-L5 dwarf members of young associations. They found that very low-gravity (γ) and intermediate-gravity (β) dwarfs likely correspond to ages of $\sim 10-30$ and ≤ 200 Myr; however, they note that BDs with ages older than ~ 30 Myr, such as the ~ 120 Myr ABDMG member 2MASS J03552337 +1133437, can display very strong signs of low gravity that correspond to the very low-gravity (γ) classification.

We used our updated sample of low-gravity candidate members of YMGs to investigate this further. We inspected various spectroscopic index–spectral type relations of candidate members of different YMGs to identify any systematic correlation with age. We assigned the age of the most probable YMG to our candidates, while rejecting any candidate with ambiguous membership (Table 4). We found that the strongest correlations with age in the ~10–130 Myr range resulted from: (1) the mean value of the EW of the three K I doublets at 1.169,

 Table 7

 Polynomial Coefficients for Spectral Type-magnitude and Spectral Type-color Diagrams

Sequence		Field Sequ	ence			Young Sequ	uence	
Name	<i>c</i> ₀	c_1	<i>c</i> ₂	σ	<i>c</i> ₀	<i>c</i> ₁	<i>c</i> ₂	σ
M_J	$8.53 \pm 3.07e - 1$	$3.08e - 1 \pm 5.66e - 2$	2.56e - 3 $\pm 2.42e - 3$	0.66	$4.97 \pm 5.65e - 1$	$7.96e - 1 \pm 1.02e - 1$	-1.14e - 2 $\pm 4.35e - 3$	1.16
M_H	$8.11 \pm 2.86e - 1$	2.81e - 1 $\pm 5.23e - 2$	1.65e - 3 $\pm 2.24e - 3$	0.63	$4.76 \pm 5.04e - 1$	$7.37e - 1 \pm 8.81e - 2$	-1.20e - 2 $\pm 3.60e - 3$	0.72
M_{KS}	8.12 + 2.74e - 1	2.05e - 1 +5.02e - 2	3.86e - 3 +2.19 $e - 3$	0.60	4.69 + 4.74e - 1	6.88e - 1 +8.16e - 2	-1.22e - 2 +3.28e - 3	0.56
M _{W1}	7.84 + 2.94e - 1	2.40e - 1 +5.07e - 2	-1.92e - 4 +2.10e - 3	0.55	4.30 +4.83 e - 1	7.57e - 1 +8.27e - 2	-1.98e - 2 +3.35e - 3	0.57
M_{W2}	7.43 + 2.96e - 1	2.82e - 1 +5.14e - 2	-2.73e - 3 +2.16e - 3	0.56	3.95 + 5.02e - 1	7.84e - 1 +8.69e - 2	-2.25e - 2 +3.60e - 3	0.42
J - H	3.83e - 1 +2.95e - 2	3.83e - 2 +5.75e - 3	2.29e - 4 +2.57e - 4	0.09	3.39e - 1 $\pm 7.39e - 2$	2.43e - 2 $\pm 1.62e - 2$	2.72e - 3 $\pm 8.33e - 4$	0.09
$H - K_S$	3.92e - 2 +2.43e - 2	6.68e - 2 +4.95e - 3	-1.85e - 3 +2.34e - 4	0.08	-1.83e - 2 +5.34e - 2	6.55e - 2 +1.13e - 2	-4.31e - 4 +5.41e - 4	0.07
$K_S - W1$	2.87e - 1 +3.36e - 2	-2.05e - 2 +6.36e - 3	2.99e - 3 +2.87e - 4	0.08	1.85e - 1 +7.56e - 2	-1.40e - 2 +1.60e - 2	4.27e - 3 +8.09e - 4	0.08
W1 - W2	3.00e - 1 $\pm 2.15e - 2$	-1.91e - 2 $\pm 4.07e - 3$	1.55e - 3 $\pm 1.87e - 4$	0.05	3.32e - 2 $\pm 3.59e - 2$	3.68e - 2 $\pm 7.75e - 3$	-3.77e - 4 $\pm 3.95e - 4$	0.07
$J - K_S$	4.48e - 1 +4.55e - 2	9.91e - 2 +9.02e - 3	-1.34e - 3 +4 14e - 4	0.14	3.35e - 1 +1.04e - 1	8.71e - 2 +2.23e - 2	2.38e - 3 +1.11e - 3	0.13
J - W1	6.75e - 1 +6.73e - 2	9.25e - 2 +1.31e - 2	1.02e - 3 +6.01e - 4	0.19	3.80e - 1 +1.07e - 1	9.97e - 2 $\pm 2.35e - 2$	5.57e - 3 +1.16e - 3	0.21
J - W2	9.85e - 1 +5 99e - 2	7.04e - 2 +1.15e - 2	2.73e - 3 +5.04e - 4	0.22	5.12e - 1 +1.36e - 1	1.18e - 1 +2.84e - 2	6.09e - 3 +1 35e - 3	0.23
H - W1	3.69e - 1 +4 50e - 2	3.81e - 2 +8.90e - 3	1.54e - 3 +4 17e - 4	0.14	8.67e - 2 +9.00e - 2	6.93e - 2 +1.93e - 2	3.00e - 3 +9.62e - 4	0.13
H - W2	6.30e - 1 +4 10e - 2	2.64e - 2 +7.96e - 3	2.76e - 3 +3.61e - 4	0.18	1.68e - 1 +1.30e - 1	9.99e - 2 +2.74e - 2	2.82e - 3 +1.36e - 3	0.15
$K_S - W2$	5.80e - 1 $\pm 3.78e - 2$	-3.93e - 2 $\pm 6.86e - 3$	4.56e - 3 $\pm 2.92e - 4$	0.13	2.41e - 1 $\pm 1.11e - 1$	2.29e - 2 $\pm 2.34e - 2$	3.71e - 3 $\pm 1.18e - 3$	0.11

Note. All abscissa are spectral types S_{type} , expressed in decimal value, where zero is M0, ten is L0, etc. A given sequence respects the equation $y = \sum_{i=0}^{N} c_i S_{\text{type}}^i$. The scatter of the data with respect to a best-fitting sequence is given by σ . All sequences are valid in the M6–L8 range.

1.177, and 1.253 μ m; and (2) the mean gravity score defined by Allers & Liu (2013). The resulting sequences are presented in Figure 14. Even though they do correlate with age on average, the scatter is too large to allow a precise determination of the age of an individual system from spectroscopic indices alone. We find that the $H_2(K)$ index defined by Canty et al. (2013) does not seem to correlate significantly with age in the 1-130 Myr range. Our results seem to be in contradiction with the findings of Canty et al. (2013) that the $H_2(K)$ index is sufficient to differentiate between objects from populations of \sim 1–2, \sim 3–10 Myr and field dwarfs in the M8–L0 range: we observe an overlap of the typical values for $H_2(K)$ in populations of $\sim 1-2$ Myr and $\sim 5-15$ Myr. However, our results are consistent with $H_2(K)$ being a good gravity-sensitive index, as it discriminates between the field population and $\lesssim 100$ Myr dwarfs for spectral types in the M6–L1 range, or $\lesssim 130$ Myr for L2–L6. It is possible that interlopers from other young associations not considered in BANYAN II contaminate our sample, which would introduce noise in these relations. A full RV and parallax follow-up of the candidates presented here will be needed to assess this.

6.3. Model Comparison

We used our sample of 86 new low-gravity M6–L5 dwarfs supplemented with 39 low-gravity and 131 field M6–L9 dwarfs from Allers & Liu (2013) and the SpeX Prism Spectral Libraries to investigate the physical properties of our sample of young dwarfs, using BT-Settl atmosphere models (Baraffe et al. 2003; Allard et al. 2013). In Section 6.3.1, we focus on effective temperatures and surface gravities obtained from a comparison of our NIR spectra with atmosphere models. In Section 6.3.2, we focus on the mass and radii that are obtained from a comparison of our photometry with evolution models.

6.3.1. BT-Settl Atmosphere Models

Manjavacas et al. (2014) used BT-Settl atmosphere models to determine the physical parameters of seven young L dwarfs and found that (1) low-gravity L0–L3 dwarfs fit models with similar temperatures of ~1800 K; (2) the continuum shape of the *H* band is not well reproduced by solar-metallicity models; (3) the 1.1–2.5 μ m range in the *zJ* bands is not well reproduced by models; and (4) the global continuum slope is not well reproduced by atmosphere models for L dwarfs.

We used a method similar to that of Cushing et al. (2008) and Naud et al. (2014) to identify the best fitting solarmetallicity CIFIST2011 BT-Settl atmosphere model for our observed spectra, on a grid of effective temperature and surface gravity ranging from $T_{\rm eff} = 500-5000$ K and $\log g = 3.0-5.5$ dex with a grid spacing of 100 K and



Figure 12. Spectral type at which the young and field absolute magnitude polynomial sequences cross (see Figure 10), as a function of the effective wavelength in which each sequence is defined. Young dwarfs are systematically brighter than their field counterparts because of their inflated radii; however, dust clouds are thicker in the high atmosphere of young L dwarfs, which counter-balances this effect and causes the young sequence to cross the field sequence. The fraction of Monte Carlo steps where the sequences crossed is indicated next to a given data point; see text for more detail. Dust clouds are more opaque in the J band (~1.2 μ m), hence the crossing point for this sequence happens at earlier spectral types. At longer wavelengths (~4.5 μ m), dust clouds do not have as much effect. This causes the sequences to cross less often and when they do, they cross at later spectral types.

0.5 dex, respectively. We computed the goodness-of-fit ($G_{k,j}$; Cushing et al. 2008) in each case.

Cushing et al. (2008) demonstrated that $T_{\rm eff}$ can only be recovered efficiently by performing such a model fitting on a very large spectral range in the case of field L1-L8 dwarfs; however, while fitting a single model spectrum in this way allows recovering a good $T_{\rm eff}$ estimate, it does not reproduce well the general slope and the features in individual spectral bands. Since gravity-sensitive spectral features are generally narrow, this method will not yield good estimates of log g. We have thus performed our model fitting in two different steps: (1) by fitting one single BT-Settl spectrum to the full $0.8-5 \,\mu m$ range (WISE W1 and W2 magnitudes were added as additional data to our spectra in order to do this); and (2) by fitting one BT-Settl spectrum to each one of the zJ, H, and K spectral bands. The first method allowed us to obtain an estimate of $T_{\rm eff}$, while the second one allowed us to obtain an estimate of $\log g$ for each object in our sample.

In order to append the *WISE* photometric data to an observed NIR spectrum, we compute the synthetic J, H, and K_S 2MASS magnitudes of the spectrum and determine the three corresponding normalization factors. We then use the median of these factors to bring back the two *WISE* photometric data points to the same scale as the observed spectrum. The dilution factor is treated as a free parameter in our analysis so that no estimate nor measurement of distance is needed in the model fitting. We thus choose the dilution factor that minimizes $G_{k,j}$ for each fitted model. We do so in an analytical way to decrease computing time. We thus define the goodness-of-fit as:

$$G_{k,j} = \sum_{i=1}^{N} W_{i,j} \left(\frac{F_{\text{obs},i,j} - D_{k,j} F_{k,i,j}}{\sigma_{\text{obs},i,j}} \right)^2, \tag{4}$$

where
$$D_{k,j} = \frac{\sum_{i=1}^{N} W_{i,j} \frac{F_{\text{obs},i,j} F_{k,i,j}}{\sigma_{\text{obs},i,j}^2}}{\sum_{i=1}^{N} W_{i,j} \frac{F_{k,i,j}^2}{\sigma_{\text{obs},i,j}^2}},$$
 (5)
$$\frac{d \ln \lambda}{dr}$$

$$W_{i,j} = \frac{\boxed{dx}}{\sum_{i'=1}^{N} \frac{d\ln\lambda}{dx}} \bigg|_{x=x_{i',j}},$$
(6)

where $x_{i,j}$ is the pixel number (i.e., the spectral position), j is the index of the spectral band (i.e., zJ, H, or K, applicable only when we fit by individual bands), k is the atmosphere model index (each value of k corresponds to a given combination of $T_{\rm eff}$ and log g), N is the total number of pixels in the fitting range, λ is the wavelength (μ m), $W_{i,j}$ are the normalized weight factors, $D_{k,j}$ is the dilution factor that minimizes $G_{k,j}$, $F_{obs,i,j}$ and $\sigma_{\text{obs},i,j}$ are the observed spectrum and its measurement error, and $F_{k,i,j}$ is an atmosphere model. The weights are chosen to ensure that equal wavelength ranges in log space equally contribute to the goodness-of-fit. For example, a broadband photometric measurement or one pixel of a low-dispersion spectroscopic order would be given a larger weight than one pixel of a highdispersion spectroscopic order as it covers a larger wavelength range. Cushing et al. (2008) introduced this weighting method except that it was not done in log space; Naud et al. (2014) noted that the log space provides a more physically meaningful scale (i.e., using the log space prevents a bias that would be caused by working in wavelength space rather than frequency space).

We calculated errors on the adjusted parameters a_l (i.e., T_{eff} and log g) from Wolberg (2006):

$$\sigma_{a_l,k,j} = \sqrt{\frac{N}{N-2}} G_{k,j} C_{l,l,k,j}^{-1} \text{ with}$$

$$\tag{7}$$

$$C_{l,m,k,j} = \sum_{i=1}^{N} W_{i,j} \frac{\partial}{\partial a_l} \left(D_{k,j} F_{k,i,j} \right) \frac{\partial}{\partial a_m} \left(D_{k,j} F_{k,i,j} \right), \tag{8}$$

where $C_{l,m,k,j}$ are elements of the correlation matrix. Equation (7) and the equivalent expression of Wolberg (2006) differ by a factor \sqrt{N} to compensate for our use of normalized weights in Equation (4). These error estimates do not take into account any systematic error in either our observations or the BT-Settl atmosphere models, and are thus only based on the variation of the goodness-of-fit with respect to each parameter. We show a few typical examples of per-band model fitting in Figure 15.

As noted by Manjavacas et al. (2014), we find that the BT-Settl models generally fail to accurately reproduce the *zJ*-bands spectra of L dwarfs, especially at wavelengths smaller than $\sim 1 \mu$ m; the general slope seems to be in agreement, but a highgravity solution is almost always preferred for all L dwarfs. Moreover, the FeH absorption features at $\sim 1.6 \mu$ m are not present at all in the atmosphere models, which could be explained by missing opacity sources in the synthetic models. For this reason, we have only kept results from the *H* and *K* bands to determine log *g*. The adopted log *g* value is thus determined from the weighted mean of the values obtained



Figure 13. NIR CMD for young (red dots when trigonometric distances were used, or purple circles when kinematic distances were used) and field (black diamonds) low-mass stars and BDs. The young and field sequences are displayed with the dashed blue line and the solid orange-red line, respectively (see text for more detail). The young sequence is systematically shifted compared to field dwarfs because of the combined effect of larger radii and thicker/higher clouds. Blue stars indicate the positions of known low-mass BDs and directly imaged exoplanets (Goto et al. 2002; Chauvin et al. 2004, 2005; Luhman et al. 2007; Marois et al. 2008; Luhman et al. 2009; Thalmann et al. 2009; Goldman et al. 2010; Lagrange et al. 2010; Janson et al. 2011; Wahhaj et al. 2011; Delorme et al. 2012, 2013; Skemer et al. 2012; Biller et al. 2013; Bowler et al. 2014; Carson et al. 2013; Kuzuhara et al. 2013; Bonnefoy et al. 2014; Currie et al. 2014; Kraus et al. 2014; Males et al. 2014; Males et al. 2014; Males et al. 2015; Marois et al. 2015; Marois et al. 2015; Marois et al. 2014; Males et al. 2015; Marois et al. 2015; Marois et al. 2014; Males et al. 2014

from the *H*-band and *K*-band fitting, where the weights are set to the total values of $W_{i,j}$ (see Equation (6)) within the fitting range divided by the inverse square of the individual measurement errors. This corresponds to the optimal weights that account both for the measurement error and the wavelength range used in the fitting process. Both the measurements and errors were rounded to the nearest half-integer and to the nearest factor of 100 K in the case of log g and T_{eff} , respectively. We imposed a floor on measurement errors that correspond to the grid size of our BT-Settl models, i.e., 0.5 and 100 K for log g and T_{eff} .

Our adopted T_{eff} and log g values are listed in Table 8 for our complete sample of low-gravity and field dwarfs. In Figure 16(a), we show the spectral type- $T_{\rm eff}$ sequence that we obtain, compared to various sequences from the literature (Golimowski et al. 2004; Stephens et al. 2009; Marocco et al. 2013). We find $T_{\rm eff}$ values that are consistent with the literature across the full range of spectral types, except for low-gravity objects which seem to be systematically cooler. This might be an additional indication that low-gravity BDs have cooler effective temperatures compared with field BDs of the same spectral types, an effect that was previously hypothesized and then demonstrated for the young, directly imaged BD and exoplanet companions HD 203030 B, TWA 27 b, HR 8799 b, and β Pictoris b (Metchev & Hillenbrand 2006; Barman et al. 2011a, 2011b; Males et al. 2014), as well as for young BDs (Faherty et al. 2012; Liu et al. 2013b; Filippazzo et al. 2015).

In Figure 16(b), we show the spectral type-log g sequence that we obtain for low-gravity and field dwarfs. The log g values that we derive for our low-gravity sample are

systematically lower than those of our field sample, as expected. However, we observe a large scatter in the $\log g$ values of low-gravity dwarfs, although they are lower on average. This indicates that the model fitting method that we described above might not be very efficient in recovering lowgravity dwarfs in an ensemble of NIR spectra. Additionally, we derive slightly lower log g values for field dwarfs with spectral types M7 and L3–L6, indicating that the false positive rate might be larger when identifying low-gravity dwarfs based solely on model fitting in this range of spectral types. Our results also tentatively indicate that M7 dwarfs are systematically better fit by low-gravity atmosphere models, however this is based on only three objects and is thus possibly an effect of small number statistics.

We tried to reproduce the results of Cushing et al. (2008) showing that fitting individual bands yield systematically offset $T_{\rm eff}$ values, and to extend this result to our full M6–L9 range as well as to low-gravity dwarfs. In Figure 16(c), we show the spectral type- $T_{\rm eff}$ sequence that we obtain if we combine the zJand H-band measurements in a weighted mean (using similar weights than described above for $\log g$). We show that the systematic offsets in $T_{\rm eff}$ values derived with this method are significant in the M9-L5 range, and independent of surface gravity. In Figure 16(d), we compare the difference of $T_{\rm eff}$ values obtained from zJ- and H-band fitting to those obtained from the K-band fitting only. We show that $T_{\rm eff}$ values derived from the K-band only are systematically warmer in the M9-L5 range. The values of T_{eff} obtained from K-band fitting only are thus closer to those presented in Figure 16(a), except that the scatter is much larger. These results confirm the findings of Cushing et al. (2008), while extending them to earlier spectral



Figure 14. Spectroscopic indices versus NIR spectral type for YMG candidates of distinct ages in our sample, binned by spectral type (see legends for color coding). We find that the mean EW of *J*-band K I and the mean gravity score defined by Allers & Liu (2013) seems to correlate with age. However, we do not see a clear correlation in the case of the $H_2(K)$ index in the 1–130 Myr range.

types (down to M9) and seem to indicate that the zJ and H bands are the most likely cause of the systematic offset in $T_{\rm eff}$.

It will be interesting to investigate whether fixing the T_{eff} value using a large spectral coverage, and subsequently determining the best log g value using wavelength regions significantly smaller than a spectral band that are known to be gravity-sensitive, might provide a better way to determine accurate log g values for L dwarfs. This will be the subject of a future work, along with repeating this analysis with future generations of BT-Settl atmosphere models that include a more realistic treatment of dust clouds (see Manjavacas et al. 2014 for a discussion on this topic).

6.3.2. Evolution Models

We estimated the physical parameters (mass, radius, T_{eff} , log g) of all low-gravity candidate members presented here from a comparison of their absolute 2MASS and *WISE* photometry with isochrones from CIFIST2011 BT-Settl models

using a likelihood analysis. The age range of the most probable host YMG was used in each case, and statistical distances from BANYAN II are used when a trigonometric distance is not available. These models do not account for magnetic fields and assume a hot-start formation (large initial entropy). Both effects could cause a systematic underestimation of mass (Konopacky et al. 2010; Stassun et al. 2012; Malo et al. 2014b; Marleau & Cumming 2014). However, it has been demonstrated that BD masses derived from evolution models are systematically too large when compared to dynamical mass measurements (Lane Bouy 2001; et al. 2004: Dupuy et al. et al. 2009a, 2009b, 2009c, 2010, 2014, 2015). This seems in contradiction with what would be expected from the model limitations described above; instead, it is likely that the cooling rate of BDs is slowed down by atmospheric clouds, an effect that is not taken into account in current evolution models (Dupuy et al. 2015).

The resulting physical parameters are presented in Table 8. This allowed us to compile a total of 25 objects with an



Figure 15. Best-fitting BT-Settl atmosphere models for typical field and low-gravity BDs (thick, black line and gray error bars). The zJ (red line), H (purple line), and K (yellow line) dilution factors were adjusted separately so that the goodness-of-fit is optimized (see text). We observe that BT-Settl models are generally unable to reproduce the zJ bands or the H-band dip at $\sim 1.6 \mu m$ that is due to FeH absorption.

estimated mass in the planetary regime ($<13 M_{Jup}$); they are individually discussed in the Appendix. These objects are all likely located within 10–60 pc and will constitute a sample of choice for a detailed study of the connection between the physical properties of BDs and giant, gaseous exoplanets, e.g., using the *James Webb Space Telescope* (Gardner et al. 2006).

In Figure 17, we compare the masses and radii estimated for the objects in our sample with those of other known exoplanets and young BDs, as well as with BD radii measured by Dieterich et al. (2014). We show that our sample overlaps with the regime of giant, gaseous exoplanets. Our sample displays inflated radii and lower masses than field dwarfs, for given spectral types, which is expected for young, low-gravity lowmass stars and BDs.

6.4. Space Density at the Deuterium-burning Limit

Late-type members of YMGs provide the opportunity to measure the low-mass end of the IMF which is still poorly constrained. The BASS survey is still not complete enough to construct individual IMFs for the YMGs under study, but we can already put constraints on the population of objects near the planetary-mass boundary where our survey is particularly sensitive.

We display in Figure 18(a) histogram of the estimated masses of all objects in our sample. We also display in this

Figure a PDF that represents a continuous analog of the histogram which is independent on the binning and that includes individual measurement errors. This PDF is obtained by normalizing the integral of each individual mass estimation PDF to unity and summing them over the full sample. In the case of absolute *W*1 magnitudes, the PDFs that correspond to individual measurements were taken as normalized Gaussian distributions with a characteristic width that corresponds to the measurement error.

There are 15 objects in our sample of THA candidates that have estimated masses in the $12.5-14 M_{Jup}$ range, which corresponds to the planetary-mass limit. This peak-shaped distribution of estimated masses for the THA candidate members uncovered here is the combined effect of a selection bias (we observed the latest-type objects first) and the distance distribution of THA members (~30-70 pc; Paper II), as 12/15of these objects are likely located within 50 pc. Furthermore, we have identified a larger number of THA candidates compared to other YMGs, because its members are more easily identified in an all-sky search—the slightly larger distance of THA ensures that its members have a narrower distribution in space position and proper motion. The relatively large number of $12.5-14 M_{Jup}$ objects compared to objects in the $5-10 M_{Jup}$ or $15-75 M_{Jup}$ ranges is thus a selection effect.

Since our sample is biased on recovering objects more efficiently in the 12.5–14 M_{Jup} range, it remains useful to assess

			Physical Parameters			
2MASS	Spectral	YMG	Estimated fro	om Isochrones ^a	Estimated from	SED Fitting
Designation	Туре	Age (Myr)	Mass (M _{Jup})	Radius (R _{Jup})	$T_{\rm eff}$ (K) ^b	$\log g^{c}$
			Field Objects			
02535980+3206373	M6	•••				5.5 ± 0.5
07522390+1612157	M6					5.5 ± 0.5
00335534-0908247	M7			•••		4.5 ± 1.0
18393308+2952164	M7				2800 ± 200	5.0 ± 1.0
22021125-1109461	M7				2800 ± 200	5.0 ± 1.0
16553529-0823401	M7	•••		•••	2700 ± 100	5.0 ± 0.5
00115060-1523450	M7		•••			5.0 ± 0.5
21144103-4339531	M7.5 pec					5.5 ± 0.5
23540957-3316220	M8					5.5 ± 0.5
21272613-4215183	M8	•••	•••	•••	2300 ± 200	5.5 ± 0.5
23520481-2208032	M8	•••	•••	•••	•••	5.5 ± 0.5
00552554+4150184	M8		•••			5.5 ± 0.5
$1/304839 \pm 0220420$	M8 M8 B		•••		$$ 2700 \pm 200	5.0 ± 1.0 5.0 ± 1.0
20025265 1316418	M8 5	•••	•••	•••	2700 ± 200 2300 ± 200	5.0 ± 1.0 5.5 ± 0.5
12531308 ± 2728028	M8 5				2300 ± 200	5.5 ± 0.5 5.0 ± 1.0
07083261_4701475	M8 5				2700 ± 200	5.0 ± 1.0 5.5 ± 0.5
22444905-3045535	M9 pec				2700 ± 200 2300 ± 200	5.5 ± 0.5 5.5 ± 0.5
12212770 ± 0257198	M9 pec				2300 ± 200 2300 ± 200	5.5 ± 0.5 5.5 ± 0.5
14284323 + 3310391	M9				2500 ± 200 2500 ± 200	5.0 ± 0.5 5.0 ± 0.5
03140344 + 1603056	M9 pec	\sim 500	$712^{+3.6}$	1.06 ± 0.02	2300 ± 200 2300 ± 200	5.0 ± 0.5 5.5 ± 0.5
10513331 1016530	M0 pec	-500	/1.2_3.7	1.00 ± 0.02	2500 ± 200 2100 ± 300	5.5 ± 0.5 5.5 ± 0.5
10313331 = 1910330 12212770 ± 0257108	M9 pec				2100 ± 300 2600 ± 200	5.5 ± 0.5
20482880_3255434	M9				2000 ± 200 2300 ± 200	5.5 ± 0.5 5.5 ± 0.5
10473109-1815574	10				2300 ± 200	5.5 ± 0.5 55 ± 0.5
17312974 + 2721233	1.0			•••		5.5 ± 0.5 5.5 ± 0.5
02281101 + 2537380	LO				2100 ± 400	5.5 ± 0.5
00461551+0252004	L0 pec				2300 ± 200	5.5 ± 0.5
22062157-6116284	L0: pec				2100 ± 200	5.5 ± 0.5
07200325-0846499	L0 pec				2300 ± 200	5.5 ± 0.5
23515044-2537367	L0.5				2600 ± 200	5.0 ± 0.5
08254335-0029110	L0.5				2100 ± 200	5.5 ± 0.5
21073169-0307337	L0.5				2300 ± 200	5.5 ± 0.5
02441019-3548036	L1 pec				1700 ± 100	4.5 ± 0.5
10484281 + 0111580	L1				2100 ± 300	5.5 ± 0.5
00332386-1521309	L1					5.5 ± 0.5
20343769+0827009	L1				2100 ± 300	5.5 ± 0.5
18071593+5015316	L1				2100 ± 300	5.5 ± 0.5
02081833+2542533	L1				2100 ± 200	5.5 ± 0.5
18071593+5015316	L1	•••		•••	•••	5.5 ± 0.5
17054834–0516462	L1		•••		2300 ± 200	5.5 ± 0.5
03454316+2540233	Ll				2300 ± 200	5.5 ± 0.5
14392836+1929149		•••	•••	•••	2300 ± 100	5.5 ± 0.5
16452211-1319516	L1.5		•••		2100 ± 300	5.5 ± 0.5
10332970 ± 0231304 12015465 1510222	L1.5		•••	•••	1900 ± 200 2100 \pm 200	5.5 ± 0.5
20575400 0252302	L1.5 1.2	•••	•••	•••	2100 ± 300 2100 ± 200	5.5 ± 0.5
20373409 = 0232302 20360316 ± 1051295	12				2100 ± 200 2100 ± 200	5.0 ± 0.5
22425317+2542573	L2 nec				1700 ± 100	5.0 ± 0.5 5.5 ± 0.5
20282035+0052265	L2 pec			•••	2100 ± 200	5.5 ± 0.5 5.5 ± 0.5
02055138-0759253	L2 L2				2100 ± 200 2100 ± 200	5.0 ± 0.5 5.0 ± 0.5
08472872-1532372	L2				2100 ± 200	5.5 ± 0.5
02415367-1241069	 L2				2100 ± 200	5.5 ± 0.5
06022216+6336391	L2				2100 ± 200	5.5 ± 0.5
05431887+6422528	L2				2100 ± 200	5.5 ± 0.5
14313029+1436599	L2					5.5 ± 0.5
20484222-5127435	L2 pec				1900 ± 100	5.5 ± 0.5
20414283-3506442	L2				1700 ± 100	5.5 ± 0.5
21041491-1037369	L2				1900 ± 100	5.5 ± 0.5
11463449+2230527	L2.5				1800 ± 100	5.0 ± 0.5
04532647-1751543	L3				2100 ± 100	5.0 ± 0.5

Table 8

			(Continued)			
2MASS	Spectral	YMG	Estimated fro	om Isochrones ^a	Estimated from	SED Fitting
Designation	Туре	Age (Myr)	Mass (M_{Jup})	Radius (R _{Jup})	$T_{\rm eff} ({\rm K})^{\rm b}$	$\log g^{c}$
			Field Objects			
10584787-1548172	L3	•••			1900 ± 100	5.0 ± 0.5
21420580-3101162	L3				1800 ± 100	5.0 ± 0.5
04070752 + 1546457	L3					5.0 ± 0.5
23155665-4747315	L3 pec	•••			1800 ± 100	5.0 ± 0.5
21420580-3101162	L3				1800 ± 100	5.5 ± 0.5
04070752+1546457	L3				•••	5.0 ± 0.5
10584787-1548172	L3	•••	•••	•••	1900 ± 100	5.0 ± 0.5
13571237+1428398	L3				1700 ± 100	5.5 ± 0.5
11000965+4957470	L3				1700 ± 100	5.0 ± 0.5
08234818+2428577	L3			•••	1700 ± 100	5.5 ± 0.5
08204440-7514571	L3.5			•••	1700 ± 100	4.5 ± 0.5
00165953-4056541	L3.5			•••	1600 ± 100	5.0 ± 0.5
22244381-0158521	L3.5	•••	•••	•••	1600 ± 100	5.0 ± 0.5
14482563+1031590	L3.5	•••			1600 ± 100 1700 + 100	4.5 ± 0.5
00193927 - 3724392 01201221 + 2517580	L3.5:				$1/00 \pm 100$ 1600 ± 100	5.0 ± 0.5
01291221+351/580 18212815+1414010	L3.5				1600 ± 100 1600 ± 100	5.0 ± 0.5
10212013 ± 1414010 01550354 ± 0050003	L4 pec	•••		•••	1000 ± 100 1800 ± 100	4.5 ± 0.5
01330334 ± 0930003 02270250 1758070	L4 I 4	•••			1800 ± 100 1500 ± 100	4.3 ± 0.3
03370339-1736079	L4 L4 pag	•••		•••	1300 ± 100 1800 ± 100	4.3 ± 0.3
23392327 ± 3307103 00361617 ± 1821104	L4 pec	•••	•••	•••	1000 ± 100 1000 ± 100	5.0 ± 0.5
00501017 + 1821104 00511078 - 1544169	L4 I 4				1600 ± 100 1600 ± 100	3.0 ± 0.5 4.5 ± 0.5
02050344 ± 1251422	I 4 nec				1600 ± 100 1600 ± 100	4.5 ± 0.5 50 ± 0.5
02030344 + 1231422 06523073 + 4710348	I 4				1600 ± 100 1600 ± 100	3.0 ± 0.5 45 ± 0.5
08014056 + 4628498	I.4: pec				1500 ± 100 1500 ± 100	50 ± 0.5
08354256-0819237	IA pec				1800 ± 100 1800 ± 100	5.0 ± 0.5 5.0 ± 0.5
11040127+1959217	L4				1900 ± 100	5.0 ± 0.5
12392727+5515371	L4 pec				1600 ± 100	4.5 ± 0.5
15065441+1321060	L4				1800 ± 100	5.5 ± 0.5
21512543-2441000	L4 pec				1600 ± 100	5.0 ± 0.5
00043484-4044058	L4.5				1600 ± 100	5.0 ± 0.5
14283132+5923354	L4.5				1800 ± 100	5.0 ± 0.5
02082363+2737400	L5				1600 ± 100	5.0 ± 0.5
06244595-4521548	L5	•••		•••	1500 ± 100	5.0 ± 0.5
09054654+5623117	L5				1700 ± 100	4.5 ± 0.5
17065487-1314396	L5 pec				1900 ± 100	5.0 ± 0.5
08511627+1817302	L5:				1500 ± 100	4.5 ± 0.5
08350622+1953050	L5	•••	•••	•••	1600 ± 100	5.0 ± 0.5
00282091+2249050	L5				•••	5.0 ± 0.5
15261405+2043414	L5				1600 ± 100	5.0 ± 0.5
01443536-0716142	L5 pec				1600 ± 100	5.0 ± 0.5
02052940-1159296	L5.5	•••		•••	1500 ± 100	5.5 ± 0.5
150/4/69-162/386	L5.5	•••	•••	•••	1900 ± 100	5.0 ± 0.5
1/401199+3034030	L5.5				1900 ± 100 1600 ± 100	5.0 ± 0.5
13202981-0038314	L3.5	•••		•••	1000 ± 100 1800 ± 100	3.3 ± 0.3
17502464 - 0010151 06540564 + 6528051	LJ.J	•••	•••	•••	1800 ± 100 1800 ± 100	4.3 ± 0.3 5.0 ± 0.5
08095903+4434216	L6 pec	30-50	81 ± 0.8	1 31+0.01	1600 ± 100 1600 ± 100	3.0 ± 0.5 4.5 ± 0.5
15150083 4847416	Lo pec	50 50	0.1 ± 0.0	1.51_0.03	1600 ± 100 1600 ± 100	4.5 ± 0.5
21011544 ± 1756586	16				1000 ± 100 1500 ± 100	3.0 ± 0.5 45 ± 0.5
16335933-0640552	1.6				1500 ± 100 1500 + 100	4.5 ± 0.5 50 ± 0.5
01033203+1935361	L6 pec				1600 ± 100 1600 + 100	4.5 ± 0.5
09153413+0422045	L6				1600 ± 100 1600 + 100	5.5 ± 0.5
04390101-2353083	L6 pec				1600 ± 100	5.0 ± 0.5
03582255-4116060	L6 pec	20-26	8.2 ± 0.6	1.37 ± 0.01	1600 ± 100	4.5 ± 0.5
23512200+3010540	L6 pec				1600 ± 100	5.0 ± 0.5
10101480-0406499	L6				1600 ± 100	5.0 ± 0.5
21321145+1341584	L6				1500 ± 100	5.5 ± 0.5
12281523-1547342	L6				1700 ± 100	5.0 ± 0.5
07171626+5705430	L6.5				1700 ± 100	5.0 ± 0.5
10433508+1213149	L7				1500 ± 100	5.5 ± 0.5

Table 8

2MASS	Spectral	YMG	Estimated fro	om Isochrones ^a	Estimated from SED Fitting		
Designation	Туре	Age (Myr)	Mass $(M_{\rm lup})$	Radius (R_{Iup})	$T_{\rm eff}$ (K) ^b	$\log g^{c}$	
	71	6 () /	Field Objects	un an Chuby		- 2 0	
23254530+4251488	L7				1200 ± 100	5.5 ± 0.5	
08503593+1057156	L7 pec u	•••			1600 ± 100	4.5 ± 0.5	
14002320+4338222	L7				1600 ± 100	5.5 ± 0.5	
08251968+2115521	L7 pec				1500 ± 200	5.0 ± 0.5	
16303054+4344032	L7	•••			1500 ± 100	5.0 ± 0.5	
03185403-3421292	L7	•••			1600 ± 100	5.0 ± 0.5	
010/5242 + 0041563 10440042 + 0420276	L/ pec	•••				4.5 ± 0.5	
10440942 + 0429376	L/ 175	•••			1600 ± 100 1500 ± 100	5.5 ± 0.5	
15150607 ± 4436483	L7.5				1500 ± 100 1500 ± 100	5.3 ± 0.3 5.0 ± 0.5	
09293364 + 3429527	L7.5				1500 ± 100 1600 ± 100	3.0 ± 0.5 4.5 ± 0.5	
15232263+3014562	L8				1500 ± 100 1500 ± 100	5.5 ± 0.5	
10365305-3441380	L8			•••	1600 ± 200	5.5 ± 0.5	
10430758+2225236	L8 pec				1600 ± 100	4.5 ± 0.5	
00325937+1410371	L8				1400 ± 100	5.5 ± 0.5	
16322911+1904407	L8	•••			1500 ± 100	5.0 ± 0.5	
12195156+3128497	L8				1200 ± 100	4.5 ± 0.5	
08575849+5708514	L8 pec	•••	••••	•••	1600 ± 200	4.5 ± 0.5	
10071185+1930563	L8	•••			1500 ± 100	5.0 ± 0.5	
15400942+3742316	L9	•••			1600 ± 100	5.0 ± 0.5	
09083803 + 5032088	L9 10	•••			1600 ± 100 1500 + 100	5.5 ± 0.5	
20431709-1331031	L9 10	•••			1300 ± 100 1400 ± 100	3.3 ± 0.3	
02550357 - 4700509	1.9				1400 ± 100 1500 ± 100	4.5 ± 0.5 55 + 05	
02550557 + 700505 03105986 + 1648155	L9				1500 ± 100 1500 ± 100	5.0 ± 0.5 5.0 ± 0.5	
03284265+2302051	L9.5				1400 ± 100 1400 ± 100	5.5 ± 0.5	
08523490+4720359	L9.5				1200 ± 100	5.5 ± 0.5	
08583467+3256275	T1				1400 ± 100	5.0 ± 0.5	
			Low-gravity Objects				
05071137+1430013 B	M5.5 β	20–26	$176.8^{+17.7}_{-17.9}$	$4.00\substack{+0.21\\-0.22}$		4.0 ± 1.5	
03363144-2619578	M5.5 β	20-40	$189.1^{+14.1}_{-17.2}$	3.94 ± 0.14		5.0 ± 1.0	
05071137+1430013 A	M5.5 β	20-26	$176.8^{+17.7}_{-17.9}$	$4.00^{+0.21}_{-0.22}$		5.0 ± 1.5	
22191486-6828018	M6 β	20-40	$32.2_{-9.3}^{+6.2}$	$1.76_{-0.05}^{+0.10}$	3100 ± 300	4.5 ± 1.5	
02404759-4253377	M6 β	20-40	$63.3_{-9.1}^{+7.3}$	$2.22_{-0.06}^{+0.07}$	2900 ± 200	4.0 ± 1.5	
04402583-1820414	M6 β	20-40		••••	3000 ± 200	4.5 ± 1.5	
08034469+0827000	M6 β	110-130	$91.0_{-4.0}^{+4.1}$	1.68 ± 0.03	3000 ± 100	5.5 ± 0.5	
03182597-3708118	M6: γ	20-40			3000 ± 400	4.5 ± 2.0	
10284580-2830374	M6 γ	5-15	$100.6^{+23.9}_{-27.5}$	$4.33_{-0.24}^{+0.29}$	2800 ± 300	4.0 ± 1.5	
10455263-2819303	M6 γ	5-15	$27.6^{+9.6}_{-5.2}$	$2.63^{+0.11}_{-0.09}$	2900 ± 100	5.0 ± 1.0	
07202582-5617224	M6 γ	20-26	$15.7^{+2.3}_{-0.7}$	$1.79\substack{+0.05\\-0.04}$	2900 ± 200	5.0 ± 1.0	
20334670-3733443	M6: γ	20-26	$106.2_{-6.9}^{+8.0}$	3.12 ± 0.08	2800 ± 200	4.5 ± 1.5	
01265327-5505506	M6 γ	20-40	$75.0^{+8.7}_{-10.2}$	2.40 ± 0.06	2900 ± 200	4.0 ± 1.5	
12574941-4111373	M6 γ	5-15	$51.5^{+12.4}_{-17.1}$	$3.11_{-0.14}^{+0.26}$	2900 ± 200	5.0 ± 1.0	
23355015-3401477	M6: γ	20-26	59.0 ± 5.1	$2.37_{-0.06}^{+0.07}$	3000 ± 300	4.0 ± 1.5	
03111547+0106307	M6 γ	20-40		•••	3000 ± 200	3.5 ± 0.5	
06353541-6234059	M6.5 β				2900 ± 300	3.5 ± 1.5	
03093877-3014352	M6.5 γ	20-40	$116.0^{+13.2}_{-14.8}$	$2.93_{-0.11}^{+0.13}$		3.5 ± 1.0	
05123569-3041067	M6.5 γ	•••		•••	2900 ± 200	4.0 ± 2.0	
02501167-0151295	M7: β	20-26	$22.9^{+5.4}_{-4.6}$	$1.91\substack{+0.06\\-0.05}$	2800 ± 200	4.5 ± 1.0	
20391314-1126531	M7 β	110-130	$74.0^{+3.6}_{-3.4}$	1.53 ± 0.03	2600 ± 200	5.0 ± 1.0	
05181131-3101529	M7 β	20-40	$97.5^{+14.3}_{-13.6}$	$2.73_{-0.12}^{+0.14}$	2800 ± 100	3.5 ± 1.5	
05264316-1824315	M7 β	20-40	$82.5^{+11.4}_{-11.5}$	$2.52_{-0.11}^{+0.12}$	2800 ± 100	5.0 ± 1.0	
01294256-0823580	M7 β	20-26	$96.3^{+7.3}_{-7.1}$	$2.99\substack{+0.09\\-0.10}$	2700 ± 300	3.5 ± 1.5	
03350208+2342356	M7.5 β	20-26	$60.9^{+4.0}_{-4.4}$	2.40 ± 0.04		5.0 ± 1.0	
00413538-5621127	M7.5 γ u	20-40			2700 ± 200	5.5 ± 0.5	
23231347-0244360	M8 β	20-26	$16.7^{+4.0}_{-0.8}$	1.85 ± 0.03	2600 ± 200	4.5 ± 1.0	
00192626+4614078	M8 β	110-130	$103.4_{-6.5}^{+7.4}$	1.76 ± 0.04	2700 ± 200	5.0 ± 1.0	

Table 8(Continued)

$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	2MASS	Spectral	YMG	Estimated from Isochrones ^a		Estimated from SED Fitting	
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	Designation	Туре	Age (Myr)	Mass $(M_{\rm Jup})$	Radius (R_{Jup})	$T_{\rm eff}$ (K) ^b	$\log g^{c}$
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		• 1		Field Objects	(sup)		
	23520507-1100435	M8 β	110-130	95.4 ± 4.3	1.71 ± 0.03	2600 ± 200	5.0 ± 0.5
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	15291017+6312539	M8 β	110-130	$93.2^{+7.9}_{-7.2}$	1.69 ± 0.06	2700 ± 200	5.0 ± 1.0
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	08561384-1342242	M8 γ	5-15	$14.4_{-1.4}^{+0.8}$	1.86 ± 0.04	2300 ± 200	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	00065794-6436542	M8 γ	20-40	$31.9^{+8.7}_{-9.5}$	$1.82_{-0.05}^{+0.07}$	2100 ± 300	5.5 ± 0.5
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	22353560-5906306	M8.5 β	20-40	$17.7^{+6.0}_{-2.4}$	1.76 ± 0.02	2300 ± 200	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	03550477-1032415	M8.5 β	20-40			2700 ± 200	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	14112131-2119503	M8.5 β				2600 ± 200	5.0 ± 1.0
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	12073346-3932539	M8.5 γ	5-15	$23.4_{-1.0}^{+2.5}$	$2.73_{-0.04}^{+0.03}$	2300 ± 200	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	20282203-5637024	M8.5 γ	20-40	$36.4_{-9.7}^{+6.9}$	$1.83\substack{+0.07\\-0.05}$	2300 ± 200	5.5 ± 0.5
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	05402325-0906326	M9 β	20-40	$19.2^{+5.1}_{-3.6}$	$1.74\substack{+0.03\\-0.06}$	2600 ± 200	5.0 ± 1.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	15104786-2818174	M9 β	30–50	$37.0_{-4.8}^{+4.4}$	$1.66\substack{+0.05\\-0.04}$	2600 ± 200	5.5 ± 0.5
	15474719-2423493	M9 β	30-50	12.9 ± 0.3	1.06 ± 0.01		5.5 ± 0.5
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	15470557–1626303 A	M9 β	110-130			2300 ± 100	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	23360735-3541489	M9 β	20-130			2300 ± 200	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	09451445-7753150	M9 β	20-40	$37.9^{+0.5}_{-8.8}$	$1.84^{+0.07}_{-0.05}$	2600 ± 200	4.5 ± 1.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	09532126-1014205	M9 β	5-40	•••	•••	1800 ± 100	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	00425923+1142104	M9 β	20–130	•••		2300 ± 100	5.5 ± 0.5
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	23453903+0055137	M9 β		10 4+5.5	1 05+0.05	2300 ± 200	5.5 ± 0.5
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	04493288+1607226	M9 γ	20-26	$18.4_{-2.2}$	$1.85_{-0.04}$	2300 ± 200	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	22025794-5605087	M9: γ	20-40	19.9-3.9	$1.79_{-0.16}^{+0.02}$	2600 ± 200	5.5 ± 0.5
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	002/419/+050341/	M9 γ		10 2+1.4	0.00+0.02		3.0 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	11395113-3159214	M9 γ	5-15	$19.3_{-1.0}^{+0.8}$	$0.99_{-0.01}^{+0.01}$	2300 ± 100	3.0 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	124/4428-3816464	M9 γ	5-15	$17.4_{-0.9}^{+0.0}$	2.15 ± 0.06	2100 ± 300	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	20004841-7523070	M9 γ	20-50	 15 6+07	1.06+0.06	2300 ± 200	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	11004401-3/15115	M9 γ	5-15	15.0-1.1	$1.90_{-0.05}$	2500 ± 100	3.0 ± 0.5
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	19355595-2846343	M9 γ		20 (+5.9	1.06+0.06		3.0 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	04433701 ± 0002051	M9 γ	20-26	$20.6_{-3.8}$	$1.80_{-0.05}$	1800 ± 100	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	00381489-0403529	M9.5 p	20-40	$14.9_{-0.4}^{+0.5}$	1.70 ± 0.02	1700 ± 100	5.5 ± 0.5
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	21544859-7459134	M9.5: B	20-40	$20.4_{-4.5}^{+0.5}$	$1.80_{-0.16}^{+0.02}$	2300 ± 300	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	02103857-3015313	M9.5 β	20-40	$14.0_{-0.3}^{+0.1}$	1.62 ± 0.02	1700 ± 100	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	12535039-4211215	M9.5 γ	5-15	$12.5^{+1.2}_{-2.0}$	$1.71^{+0.04}_{-0.03}$	•••	4.5 ± 0.5
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	15525906+2948485	$L0 \beta$			 1 < 0 ⁺⁰ 02		3.0 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	11544223-3400390		30-50	18.2-3.6	$1.60_{-0.03}^{+0.03}$	2100 ± 200	5.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	00325584-4405058		20-26	11.7 ± 0.6	1.00 ± 0.01		5.5 ± 0.5
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	05420951-2904517	$L0: \beta$	20-40	12.7 ± 0.4 g g+1.0	1.48 ± 0.02 1.35 ± 0.03	2100 ± 200 1500 ± 100	3.0 ± 0.5 5.5 ± 0.5
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	12451416 4420077	$L0. p/\gamma$	20-40	$0.0_{-1.1}$	1.35 ± 0.05 1.24 ± 0.03	1300 ± 100	3.5 ± 0.5
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	12431410-4429077	$L0 \gamma$	3-13	$18.4_{-1.3}$	$1.24_{-0.02}$	2300 ± 200	5.0 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	00182834-0703130	LOY	20-40	$15.7_{-0.3}$	1.58 ± 0.02	1700 ± 100 1700 ± 100	3.0 ± 0.3
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	04400972-5120544	$L0 \gamma$	20-40	12 6+0.3		1700 ± 100 2100 \pm 200	4.5 ± 0.5 3.0 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	20334473-3033338	LON	20-40	$13.0_{-0.4}$ 14 7+5.7	1.50 ± 0.02 1.16 ± 0.01	2100 ± 200 1700 ± 100	5.0 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	01413825-4055574	$L0 \gamma$	20-40	$14.7_{-0.4}$	1.10 ± 0.01	1700 ± 100	3.0 ± 0.3
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	02292794-0033282	LOγ	20-40	$13 1^{+0.3}$	1.04 ± 0.02		5.3 ± 0.3 5.0 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	06101201 5803156 b		20-40	$13.1_{-0.4}$ 12 8 ^{+0.4}	1.04 ± 0.02 1.03 ± 0.01		3.0 ± 0.3 3.0 ± 1.0
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	00191291-3803130 0	LOY	20-40	$12.0_{-0.3}$	1.03 ± 0.01	•••	5.0 ± 1.0
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	22134491 - 2130079 00464841 + 0715177	$L0 \gamma$	20-20	15.5 ± 0.5 $15.4^{+0.5}$	1.09 ± 0.01 1 70 ^{+0.03}	$$ 2100 \pm 100	3.0 ± 0.3 5.5 ± 0.5
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	06085283 2753583		20-20	10.8+3.9	$1.79_{-0.02}$ 1.20 ± 0.02	2100 ± 100	3.0 ± 0.5
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	20135152 2806020		20-40	$15.3_{-4.2}$	1.20 ± 0.02 1.10 ± 0.02		3.0 ± 0.5 3.0 ± 0.5
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	02265658_5327032	108	20-20	13.7 ± 0.3	1.10 ± 0.02 1.50 ± 0.02	1700 ± 100	3.0 ± 0.3 4.5 ± 0.5
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	11271382_3735076	108	20 -4 0 5_15	0.7 ± 0.3 $0.2^{+1.4}$	1.59 ± 0.02 1 56 ^{+0.05}	1700 ± 100 2100 ± 200	4.5 ± 0.5 4.5 ± 0.5
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	003//300_/102266	I 1. R	20_40	12.8 ± 0.4	$1.30_{-0.03}$ 1.48 ± 0.01	1700 ± 100	-7.5 ± 0.5 55 ± 0.5
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	11480096-2836488	$L1 \cdot \beta$	20-40 5-15	12.0 ± 0.4 8 5 ^{+1.4}	$1.55^{+0.04}$	1800 ± 100 1800 + 100	3.5 ± 0.5 3.5 ± 0.5
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	03164512_2848521	I1 R	110-130	$340^{+1.6}$	$1.00_{-0.03}$ 1.24 ± 0.01	2100 ± 200	5.5 ± 0.5 55 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	00584253_0651230		20_130	J-1.5	1.27 ± 0.01	2100 ± 200 2100 + 200	5.5 ± 0.5 55 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	19350976-6200473	$L1 \gamma$	20-150	13 5+0.3	1.54 ± 0.02	2100 ± 200 2100 + 200	3.5 ± 0.5 4.5 ± 0.5
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	23225299_6151275	$L1 \sim$	20-40	$13.5_{-0.4}$ 13.6 ± 0.3	1.56 ± 0.02	1700 ± 100	50 ± 0.5
25255004-0259508 L1 γ 110-130 31.0 \pm 17 1.22 \pm 0.01 1800 \pm 100 5.0 \pm 0.5	23255604-0259508	$L1 \gamma$	110-130	$31.0^{+1.4}_{-1.2}$	1.22 ± 0.02	1800 ± 100 1800 ± 100	5.0 ± 0.5 5.0 ± 0.5

Table 8(Continued)

2MASS Designation	Spectral Type	YMG Age (Myr)	Estimated from Isochrones ^a		Estimated from SED Fitting	
			Mass (M _{Jup})	Radius (R _{Jup})	$T_{\rm eff} ({\rm K})^{\rm b}$	$\log g^{c}$
			Field Objects			
02410564-5511466	L1 γ	20-40	13.6 ± 0.3	1.57 ± 0.02	1700 ± 100	5.0 ± 0.5
01205114-5200349	L1 γ	20-40	13.3 ± 0.3	1.54 ± 0.02	1700 ± 100	5.0 ± 0.5
01174748-3403258	L1 γ	20-40	13.6 ± 0.3	1.09 ± 0.01	1800 ± 100	5.0 ± 0.5
04062677-3812102	L1: γ	20-40	$11.7^{+0.7}_{-1.1}$	$0.98^{+0.03}_{-0.02}$	•••	4.5 ± 0.5
05184616-2756457	L1 γ	20-40	$15.3_{-1.5}^{+6.7}$	1.11 ± 0.05		5.0 ± 0.5
00191296-6226005	L1 γ	20-40	13.8 ± 0.3	1.59 ± 0.02	1600 ± 100	5.0 ± 0.5
11083081+6830169	L1 γ	20-40	13.5 ± 0.3	1.55 ± 0.01	1700 ± 100	5.0 ± 0.5
12074836-3900043	L1 δ	5-15	$12.1^{+1.4}_{-2.0}$	1.69 ± 0.04	2100 ± 100	5.0 ± 0.5
22351658-3844154	L1.5 γ	20-40	$14.0^{+0.4}_{-0.3}$	1.62 ± 0.02	1600 ± 100	5.0 ± 0.5
06023045+3910592	L2 B					5.0 ± 0.5
00452143+1634446	$L2\gamma$	30-50	14.7 ± 0.3	1.14 ± 0.01		5.0 ± 0.5
05361998-1920396	L2 γ	20-40	13.0 ± 0.9	$1.50^{+0.07}_{-0.06}$	1600 ± 200	4.5 ± 0.5
02583123-1520536	L3 B	20-40	$13.0^{+0.3}_{-0.4}$	$1.50^{+0.02}_{-0.01}$	1700 ± 100	5.0 ± 0.5
12563961-2718455	L3: β	5-15	$7.7^{+1.4}_{-1.5}$	$1.53^{+0.04}_{-0.02}$	1600 ± 100	4.5 ± 0.5
01531463-6744181	$L3\beta$	20-40	$129^{+0.3}$	149 ± 0.02	1600 ± 100	50 ± 0.5
17260007 ± 1538190	L3 α	20 10	12.7_0.5	1.17 ± 0.02	1000 ± 100	3.0 ± 0.5
0/185870_/507/13	$L3\gamma$	20-40	$120^{+0.3}$	1.49 ± 0.02	1600 ± 100	4.5 ± 0.5 5.0 ± 0.5
22081363 + 2021215		20-40	$12.9_{-0.4}$ 12.0 $^{+0.3}$	1.49 ± 0.02 1.14 ± 0.01	1000 ± 100	3.0 ± 0.5
22081303-2921213	L2 v	20-20	$12.9_{-0.1}$	1.14 ± 0.01		4.5 ± 0.5
20113190-3046112	$L_{2}\gamma$	20-40	12.9 ± 0.4	1.49 ± 0.02	•••	3.0 ± 0.3
10042000 + 3022390 15515227 + 0041148	L3: γ	20-40	12.3 ± 0.4	1.34 ± 0.07	•••	4.0 ± 0.3
13313237 ± 0941140 00011217 ± 1535355	$L_3.\gamma$	110, 130	25.3 ± 1.0	120 ± 0.01	$\frac{1600 \pm 100}{1600}$	3.5 ± 0.5
21324036 ± 1029494	$\mathbf{L} + \boldsymbol{\beta}$ $\mathbf{L} + \boldsymbol{\beta}$	30-50	25.5 ± 1.0 11.4 ± 0.4	1.20 ± 0.01 1.36 ± 0.01	1600 ± 100 1600 ± 100	4.0 ± 0.5
10212570_2830427	$\mathbf{I} 4 \cdot \boldsymbol{\beta} / \boldsymbol{\gamma}$	5-15	$65^{+1.3}$	1.50 ± 0.01 $1.51^{+0.04}$	1600 ± 100 1600 ± 200	50 ± 10
22064408 4217208		110 130	26.1 ± 1.0	$1.51_{-0.03}$ 1.21 ± 0.01	1000 ± 200 1800 ± 100	5.0 ± 0.5
05012406_0010452	L4 Y	20.40	20.1 ± 1.0 10 2 ^{+0.8}	1.21 ± 0.01 1.36 ± 0.02	1600 ± 100 1600 ± 100	3.0 ± 0.3
22405245 + 0044046	L4 Y	20-40	$10.2_{-1.0}$	1.50 ± 0.02	1000 ± 100 1700 ± 100	4.0 ± 0.5
22493343 ± 0044040 14252708 3650220	L4 Y	110, 130	26 6+0.3	120 ± 0.01	1700 ± 100 1600 ± 100	4.3 ± 0.3 5.0 ± 0.5
1+252770-5050227		110-130	$20.0_{-1.0}$	1.20 ± 0.01	1500 ± 100	5.0 ± 0.5
05552557+1155457	$L_{3} = L_{0} \gamma$	20, 120	$22.4_{-1.0}$	1.20 ± 0.01	1500 ± 100	4.3 ± 0.3
23433470-3040021	$L_3 - L_0 \gamma$	20-130	•••		1500 ± 100	4.0 ± 0.5
10134233+4933211	L3-L0 γ	20.26	 6 7 ^{+1.0}	1.26 ± 0.01	$$ 1600 \pm 100	4.0 ± 0.5
03120030-2949340		20-20	$0.7_{-0.9}$	1.30 ± 0.01	1000 ± 100	4.0 ± 0.3
231/4/12-4838501			10.9+0.4			4.5 ± 0.5
00505015-1450555	L4-L0 p	30-50	$10.8_{-0.6}$	1.33 ± 0.01	1500 ± 100	5.0 ± 0.5
21543454-1055308	LS β / γ	30-50	$10.3_{-0.7}^{+0.0}$	1.32 ± 0.01	1600 ± 100	4.0 ± 0.5
03264225-2102057	LS β / γ	110–130	$22.4^{+1.0}_{-1.1}$	1.20 ± 0.01	1600 ± 100	
20025073-0521524	L5–L7 γ	•••		•••	1700 ± 100	4.5 ± 0.5
17410280-4642218	L5:–L7: γ		•••			3.5 ± 0.5
00470038+6803543	L6–L8 γ					4.0 ± 0.5
22443167+2043433	L6–L8 γ	110-130	12.0 ± 0.1	0.86 ± 0.01		4.0 ± 0.5
21140802-2251358	Lo–L8 γ	20-26	9.0 ± 0.3	1.00 ± 0.01	•••	4.0 ± 0.5

Table 8

Notes.

^a Estimated masses and radii were derived from a comparison of the trigonometric or statistical distances, the age of the most probable YMG membership and the 2MASS and *WISE* photometry with AMES-Cond evolutionary models (see the text for a detailed explanation).

^b Values obtained from the modified model fitting that uses WISE W1, W2 photometry as well as NIR spectra in the J, H, and K bands.

^c Values obtained from a weighted mean of *H*-band and *K*-band $\log g$ (see text for more information).

(This table is available in its entirety in machine-readable form.)

the space density of such objects. We will concentrate on the THA candidate members for this as they provide a larger sample. Assuming that we have uncovered all of the 12.5–14 M_{Jup} candidate members of THA in BASS within 50 pc (accounting for 65.6% of the expected population according to our SKM model for THA) and correcting for the expected completeness of BASS for this association (90%; Gagné et al. 2015), we can expect that there are a total number of

 $20.3^{+6.8}_{-5.1}$ objects in THA that lie within this range of masses. The error was estimated assuming that the objects were drawn from a Poisson distribution, and they thus account for small number statistics.

Assuming that the population of $1.00-1.26 M_{\odot}$ stars is complete in THA ($N = 14^{+4.3}_{-3.3}$ using Poisson statistics, see Figure 8 of Kraus et al. 2014b) and adjusting a fiducial lognormal IMF peaking at $0.25 M_{\odot}$ with a width $\sigma = 0.5$ dex



Figure 16. Panel (a): adopted effective temperature (T_{eff}) derived by simultaneously fitting atmosphere models to full *JHK* spectra and *WISE* photometry as a function of spectral type for our sample of field and low-gravity dwarfs, binned by spectral type. The number of data points that were included in each bin is displayed above each symbol. Field dwarfs are represented with black circles, intermediate gravity dwarfs (β) with green diamonds and very low-gravity dwarfs (γ) with purple downside triangles. We added small systematic offsets in the spectral types of very low-gravity and field dwarfs for visibility. The solid orange line, green dashed line and fuchsia dashed–dotted lines represent T_{eff} –spectral type relations from Marocco et al. (2013), Golimowski et al. (2004), Looper et al. (2008a) and Stephens et al. (2009), respectively. We derive systematically cooler temperatures for young BDs. Panel (b): adopted surface gravity (log g) as a function of spectral type for our sample of field and low-gravity dwarfs, binned by spectral type. The color coding is similar to that of Panel (a). The derived log g values for dwarfs with spectroscopic confirmation of low gravity are systematically lower than those of field dwarfs; however, the large scatter hints that model fitting alone is not an efficient way of identifying low-gravity dwarfs. Panel (c): effective temperature, derived by fitting atmosphere models to individual *zJ*- and *H*-bands only without *WISE* photometry, as a function of spectral type for our sample of field and low-gravity dwarfs, a likely effect of dust clouds not being properly reproduced by BT-SettI models. Panel (d): difference in the derived effective temperature from the *K*-band model fitting from that obtained by individual *zJ*- and *H*-bands model fitting (all without using *WISE* photometry), binned by spectral type. Color-coding is identical to Panel (a) and the red dot-dashed lines marks ΔT_{eff} =0. We derive systematically warmer temperatures when fitting

(Jeffries 2012), we can expect a total of 356^{+61}_{-47} main-sequence stars in THA (>75 M_{Jup}) and only $0.56^{+0.17}_{-0.13}$ objects in the 12.5–14 M_{Jup} range (the ratio of 12.5–14 M_{Jup} to 1.00–1.26 M_{\odot} objects derived from that IMF is 0.04). We thus seem to be uncovering at least $36.4^{+16.6}_{-12.5}$ times too many objects in this mass range, compared to the predictions of a typical log-normal IMF anchored on the 1.00–1.26 M_{\odot} population of THA.

It is possible that this is a consequence of a fault in the evolution models rather than a true over-population. For example, one could argue that the models fail to reproduce the effects of clouds which have a larger impact on the spectra of less massive, cooler objects. This could lead us to misinterpret the masses of our 15 low-gravity THA candidates, assigning them 12.5–14 M_{Jup} while their true masses span a larger range. If this effect alone is to explain the over-population, the true range of masses for our 15 objects would have to be extended by 190% in log space, which would mean that their true masses would span 4.5–39 M_{Jup} . This effect is thus unlikely to be the lone explanation of this over-population. It is also possible that the current age estimate of THA is wrong—e.g., β PMG, Upper



Figure 17. Panel (a): radius as a function of mass for intermediate-gravity (green diamonds) and very low-gravity (purple downside triangles) candidate members of YMGs, derived from the BT-Settl models, compared with exoplanets and BD companions that benefit from transit and RV data (orange circles). Isochrones of various ages (gray lines; 10 Myr to 8 Gyr) were added for comparison. It can be seen that our sample overlaps with the planetary regime, and follow isochrones that correspond to the ages of YMGs considered here. Very low-gravity objects have a younger age on average compared with intermediate-gravity objects, which explains that they have larger radius for a given mass. Transit and RV data were obtained from Stassun et al. (2006), Sahu et al. (2006), Deleuil et al. (2008), Winn et al. (2008), Buchhave et al. (2010), Southworth (2010), Bouchy et al. (2011a, 2011b), Tingley et al. (2011), Bakos et al. (2011), Buchhave et al. (2011), Deleuil et al. (2012), Siverd et al. (2012), Cappetta et al. (2012), Triaud et al. (2013), Díaz et al. (2013), Hébrard et al. (2013), Moutou et al. (2013), Blecic et al. (2013), Parviainen et al. (2014), Díaz et al. (2014), Shporer et al. (2014), Littlefair et al. (2014), Montet et al. (2014), and Quinn et al. (2014). Panel (b): mass as a function of spectral type for intermediate-gravity (green diamonds) and very low-gravity (purple downside triangles) candidate members of YMGs, compared with BT-Settl isochrones (gray lines;10 Myr-8 Gyr) from which the masses were derived. The isochrones were mapped on the spectral type dimension by converting effective temperatures to spectral types using the polynomial relation of Stephens et al. (2009). Panel (c): radius as a function of spectral type for intermediate-gravity (green diamonds) and very low-gravity (purple downside triangles) candidate members of YMGs, compared with radii measurements from Dieterich et al. (2014; orange circles). BT-Settl isochrones of various ages (gray lines; 10 Myr-8 Gyr), which were used to derive our radii, were added for comparison. They were mapped on the spectral type dimension by converting effective temperatures to spectral types using the polynomial relation of Stephens et al. (2009). The radii measurements of Dieterich et al. (2014) are all based on a comparison of the synthetic colors of atmosphere models with photometric measurements combined to trigonometric parallaxes. Their data consists in a majority of field objects. It can be seen that we determine radii that are consistent with the young ages of our objects when comparing to isochrones; however, the radii measurements of Dieterich et al. (2014) for dwarfs later than ~L2 yield larger values than those predicted by field-age evolution models alone (see Dieterich et al. 2014 for more detail). The data used to create this figure are available.

Scorpius, AB Doradus and the Pleiades have recently been found to be slightly older than previously thought (Luhman et al. 2005; Pecaut et al. 2012; Binks & Jeffries 2014; Malo

et al. 2014b; Mamajek & Bell 2014). If it turns out that this is also the case for THA, our estimated masses would need to be shifted to larger values. As an example, doubling the age of



Figure 18. Panel (a): histogram of estimated masses (green bars) for low-gravity dwarfs in the BASS sample, obtained from a comparison of NIR photometry and trigonometric or kinematic distances with BT-Settl–CIFIST2011 synthetic models. The continuous PDFs of different subsets of the candidates are indicated with different lines (see legend). They were obtained by combining the individual mass estimation PDFs directly, and they thus provide a histogram-like continuous distribution that include measurement errors and are independent of the binning. Panel (b): histogram of absolute *WISE M*_{W1} magnitude of low-gravity dwarfs in the BASS sample (purple bars), obtained from trigonometric or kinematic distances. The thick aqua distribution is a continuous distribution that does not include binning and takes account of measurement uncertainties, and was built in the same way as that of Panel (a). We find a larger number of objects as M_{W1} gets fainter, up to $M_{W1} \sim 11$ where we stop being sensitive. The most limiting aspect of our survey is the inclusion in the 2MASS catalog, with a limiting magnitude around $J \sim 16-17$. The absence of a strong over-density is not in contradiction with Panel (a), because our sample is composed of objects at different ages ($\sim 12-120$ Myr), hence a given mass can correspond to a different temperature and absolute magnitude. The green, yellow and red vertical dashed lines correspond to the absolute W1 magnitudes of a 10, 40, and 120 Myr object, respectively.

THA would shift the estimated mass of a member from ~13 to ~20 M_{Jup} . This effect alone would thus be insufficient to explain the large number of 12.5–14 M_{Jup} THA candidates that we found. A similar shift of our estimated masses could be caused by systematics in evolution models (see our discussion in Section 6.3.2), although it is difficult to estimate the magnitude of this effect at this time. Dupuy et al. (2015) has shown that masses from evolution model are likely underestimated for dusty BDs at the L/T transition; it could be expected that the same effect is important in young L dwarfs. This would further accentuate the discrepancy between our observations and the predictions from a typical IMF.

Bowler et al. (2013) noted that the age-absolute luminosity model sequences of ~ 13 and $\sim 25 M_{Jup}$ objects at different young ages overlap; such a pile-up in the isochrones could cause a degeneracy in our estimated masses and cause our method to mis-interpret true $\sim 25 M_{Jup}$ objects as planetarymass objects. However, there are several observations that make this explanation unlikely: (1) the likelihood method with which we estimate masses not only generates a measurement and error bars, but it also provides a continuous PDF for each individual mass estimate. If this effect is important, we would thus be able to observe double-peaked individual measurement PDFs, as well as a peak at $\sim 25 M_{Jup}$ in the PDF displayed in Figure 18. Note that even if present, this effect would not introduce a second peak in the histogram, since it was constructed from the most probable values of the estimated masses only. (2) While the young age-absolute luminosity isochrones overlap at different masses, this effect is much more subtle in the individual J, H, K_S , W1, and W2 age-absolute magnitude isochrones. Furthermore, the slight overlap happens at slightly different ages and masses in the different filters, and allows to lift the degeneracy between ~ 13 and $\sim 25 M_{Jup}$ objects. This likely explains why we do not observe dualvalued mass estimate PDFs. (3) Performing a Monte Carlo analysis in which 20 and 40 Myr isochrones are used to estimate the masses of a population of 20,000 synthetic objects with true masses uniformly distributed between 4 and 80 M_{Jup} produces no over-density of estimated masses in the 12–14.5 M_{Jup} range. The absolute *J*, *H*, K_S , *W*1, and *W*2 magnitudes of these synthetic objects are obtained from the model isochrones themselves, hence this Monte Carlo analysis cannot be used to investigate systematics in the model cooling tracks. Instead, it only addresses the potential problem of overlapping isochrones that could produce degenerate mass estimates.

As a consequence of these observations, it does not appear that overlapping isochrones are the cause of the large population of 12–14.5 M_{Jup} THA candidates in our sample. We note that it is, however, possible that a fraction of these THA candidate members are contaminants in our analysis (i.e., young interlopers from other moving groups or associations, considered in BANYAN II or not) despite their high Bayesian probability and the low expected contamination rate in this particular YMG. It will be necessary to measure the RVs and parallaxes for all 12 objects discussed here to assess this, but at this stage it seems that this effect would be the most likely explanation for this over-density. For example, only 5/12 of these objects would need to be interlopers in order for the overdensity to become a 1σ result.

If we assume that the over-density is real, it would mean that there is at least one isolated dwarf in the 12.5–14 M_{Jup} range for every $17.5^{+6.6}_{-5.0}$ main-sequence star in THA. Comparing with the space density of main-sequence stars in the solar neighborhood $(9.3 \times 10^{-2} \text{ stars pc}^{-3}; \text{ Chabrier 2005})$ and assuming that the ratio we observed in THA is valid in the field, this would amount to a field density of $5.3^{+3.8}_{-2.9} \times 10^{-3}$ dwarfs pc⁻³ in the 12.5–14 M_{Jup} range in THA. At ages older than 2.5 Gyr, they will all have temperatures below 450 K that correspond to spectral types later than Y0, and will thus be hard to locate due to their extreme faintness (Cushing et al. 2011; Kirkpatrick et al. 2011; Beamín et al. 2014; Luhman 2014). This is significantly larger than the lower limit measured by

Kirkpatrick et al. (2011) that corresponds to at least one \ge Y0 dwarf for every 78 main-sequence star (or 1.2×10^{-3} dwarfs pc⁻³), especially when considering that the population of field \ge Y0 dwarfs is also probably composed of objects that span a large range of ages and thus masses. Kirkpatrick et al. (2011) noted that their measurement is only a gross underestimation on the space density of Y-type dwarfs due to several biases. We note however that the IMF of YMGs might be different than that of the field, which could be yet another cause for this difference.

A less likely scenario is that our results could be an indication that we are approaching an up-turn in the IMF of isolated objects in THA with masses below the deuteriumburning limit: such an up-turn has already been hinted at by micro-lensing surveys in the galactic plane that measure $1.8^{+1.7}_{-0.8}$ Jupiter-mass object for every main-sequence star (corresponding to space density of $1.7^{+1.6}_{-0.7} \times 10^{-1}$ objects pc⁻³; Sumi et al. 2011). Measurements of RV and distance for the complete set of YMG candidates in BASS will be crucial to assess whether the observed over-density holds, and discovering YMG candidate members at even lower masses will provide a strong constraint on whether there is an up-turn in the IMF of YMGs.

7. SUMMARY AND CONCLUSIONS

We presented a NIR spectroscopic follow-up of 241 candidate members of YMGs identified through the BASS, LP-BASS and PRE-BASS samples. This allowed us to identify 108 new low-gravity M5-L5 candidate members of YMGs with estimated masses spanning the range of $7-189 M_{Jup}$. Thirty-seven of these objects were previously known in the literature, but no signs of low gravity had been reported for them before this work. We complemented this unique sample with 22 low-gravity dwarfs from the literature to (1) build color-spectral type and absolute magnitude-spectral type sequences for field and young dwarfs; (2) show that some gravity-sensitive indices correlate with age in the 10-200 Myr regime, albeit with a large scatter, such that low-resolution NIR spectroscopy does not allow a strong constraint on the age of an individual object; (3) we discuss some limitations of the current BT-Settl models, mainly their improper treatment of dust clouds in L-type dwarfs of all ages; and (4) show that we find an unexpectedly large number of isolated objects with estimated planetary masses in the Tucana-Horologium association, which might be caused by young interlopers from other moving groups. This study represents one of the first steps toward bridging the gap in our knowledge of the the space density of the lowest-mass BDs ($\sim 13 M_{Jup}$; Kirkpatrick et al. 2011) and potential isolated giant planets that were ejected from their stellar system (~1 M_{Jup} ; Sumi et al. 2011). Additional figures, data and information on this work can be found on the website www.astro.umontreal.ca/~gagneand in the Montreal Spectral Library, which is located at www.astro. umontreal.ca/~gagne/MSL.php.

The authors would like to thank the anonymous referee who suggested to improve BANYAN II with the inclusion of parallax motion and significantly helped to improve the quality of this paper, as well as make it more concise and clear. We would like to thank Robert Simcoe, Philippe Delorme, Michael C. Cushing, Rebecca Oppenheimer, Amélie Simon, Gilles Fontaine, Sergio B. Dieterich, Benjamin M. Zuckerman, André-Nicolas Chené Sarah Jane Schmidt, Simon Coudé Daniella C. Bardalez Gagliuffi, and Jonathan B. Foster for useful comments and discussions. We thank Katelvn N. Allers. Michael C. Liu, Federico Marocco, and Brendan P. Bowler for sharing data. We also thank all observatory staff and observers who helped us in this quest-Bernard Malenfant, Ghislain Turcotte, Pierre-Luc Lévesque, Alberto Pastén, Rachel Mason, Stuart Ryder, Rubén Díaz, Stéphanie Côté John P. Blakeslee, Mischa Schirmer, Andrew McNichols, Dave Griep, Brian Cabreira, Tony Matulonis, German Gimeno, Steve Margheim, Percy L. Gomez, René Rutten, Bernadette Rodgers, Tim J. Davidge, Jaehyon Rhee Jay, Inger Jørgensen, Thomas L. Hayward, Andrew Cardwell, Blair C. Conn, Eleazar Rodrigo Carrasco Damele, David A. Krogsrud, Eduardo Marin, Erich Wenderoth, Fredrik T. Rantakyro, Joanna E. Thomas-Osip, Pablo Patricio Candia, Pascale Hibon, Cláudia Winge, Benoit Neichel, Peter Pessev, Matthew B. Bayliss, and Anne Sweet. This work was supported in part through grants from the Fond de Recherche Québécois-Nature et Technologie and the Natural Science and Engineering Research Council of Canada. This research has benefited from the SpeX Prism Spectral Libraries, maintained by Adam Burgasser at http://pono.ucsd. edu/~adam/browndwarfs/spexprism, as well as the M, L, T, and Y dwarf compendium housed at http://DwarfArchives.org and maintained by Chris Gelino, Davy Kirkpatrick, and Adam Burgasser, whose server was funded by a NASA Small Research Grant, administered by the American Astronomical Society. This research made use of the SIMBAD database and VizieR catalog access tool, operated at the Centre de Données astronomiques de Strasbourg, France (Ochsenbein et al. 2000); data products from 2MASS Skrutskie et al. 2006), which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center (IPAC)/California Institute of Technology (Caltech), funded by the National Aeronautics and Space Administration (NASA) and the National Science Foundation (Skrutskie et al. 2006); the Extrasolar Planets Encyclopaedia (exoplanet.eu), which was developed and is maintained by the exoplanet TEAM; data products from the Wide-field Infrared Survey Explorer (WISE; Wright et al. 2010), which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory (JPL)/Caltech, funded by NASA; the NASA/IPAC Infrared Science Archive (IRSA), which is operated by JPL, Caltech, under contract with NASA; the Infrared Telescope Facility (IRTF), which is operated by the University of Hawaii under Cooperative Agreement NNX-08AE38A with NASA, Science Mission Directorate, Planetary Astronomy Program; the Database of Ultracool Parallaxes maintained by Trent Dupuy (Dupuy & Liu 2012); the Hale 5 m telescope at Palomar Observatory, which received funding from the Rockefeller Foundation; and of tools provided by Astrometry.net. This paper includes data gathered with the 6.5 m Magellan Telescopes located at Las Campanas Observatory, Chile (CNTAC program CN2013A-135). Based on observations obtained at the Gemini Observatory through programs number GN-2013A-Q-118, GS-2013B-Q-79, GS-2014A-Q-55, GS-2014B-Q-72, GS-2014B-Q-47, and GS-2015A-Q-60. The Gemini Observatory is operated by the Association of Universities for Research in Astronomy, Inc., under a cooperative agreement with the National Science Foundation (NSF) on behalf of the Gemini partnership: the NSF (United States), the National Research Council (Canada), CONICYT

Potential Isolated Planetary-mass Objects								
2MASS		Spectral	Source	Moving	Estimated	Distance ^a		
Designation	References	Туре	Sample	Group	Mass (M _{Jup})	(pc)		
00303013-1450333	(1)	L4–L6 β	LP-BASS	ARG	$10.8_{-0.6}^{+0.4}$	26.72 ± 3.21^{b}		
00344300-4102266	(2)	L1: β	BASS	THA	12.8 ± 0.4	41.2 ± 2.4		
01531463-6744181	(3)	L3 β	BASS	THA	$12.9_{-0.5}^{+0.3}$	47.0 ± 3.2		
03420931-2904317	(2)	L0: β	BASS	THA	12.7 ± 0.4	$48.2^{+3.6}_{-3.2}$		
03421621-6817321	(4)	L4 γ	BASS	THA	12.4 ± 0.5	48.6 ± 3.6		
04185879-4507413	(2)	L3 γ	BASS	THA	$12.9_{-0.4}^{+0.3}$	$49.8_{-3.6}^{+4.0}$		
05012406-0010452	(3)	L4 γ	BASS	COL,CAR	$10.2\substack{+0.8\\-1.0}$	14.7 ± 2.8^{b}		
05120636-2949540	(5)	L5 β	BASS	BPMG	$6.7^{+1.0}_{-0.9}$	$10.9_{-4.0}^{+4.4}$		
06272161-5308428	(2)	L0: β/γ	BASS	CAR	$8.8^{+1.0}_{-1.1}$	$24.1^{+2.4}_{-2.0}$		
06322402-5010349	(3)	L3 β	PRE-BASS	ABDMG	$11.0\substack{+0.5\\-0.6}$	$7.7^{+3.2}_{-2.8}$		
10212570-2830427	(2)	L4: β/γ	BASS	TWA	$6.5^{+1.3}_{-1.2}$	42.6 ± 5.6		
11271382-3735076	(2)	L0 δ	LP-BASS	TWA	$9.2^{+1.4}_{-1.7}$	$62.2\substack{+8.8\\-8.0}$		
11480096-2836488	(2)	L1: β	BASS	TWA	$8.5^{+1.4}_{-1.6}$	47.8 ± 5.6		
12074836-3900043	(6)	L1 δ	BASS	TWA	$12.1^{+1.4}_{-2.0}$	$58.2_{-6.4}^{+6.8}$		
12271545-0636458	(5)	M8.5 β	PRE-BASS	TWA	$11.6^{+1.4}_{-1.9}$	32.5 ± 3.2		
12535039-4211215	(2)	M9.5 γ	BASS	TWA	$12.5^{+1.2}_{-2.0}$	$81.0^{+7.6}_{-7.2}$		
12563961-2718455	(2)	L3: β	BASS	TWA	$7.7^{+1.4}_{-1.5}$	44.6 ± 5.2		
20113196-5048112	(2)	L3 γ	BASS	THA	12.9 ± 0.4	$53.4_{-3.6}^{+4.0}$		
21324036+1029494	(7)	L4: β	PRE-BASS	ARG	11.4 ± 0.4	34.2 ± 4.8		
21543454-1055308	(8)	L5 β/γ	BASS	ARG	$10.3_{-0.7}^{+0.5}$	22.5 ± 2.8		

Table 9

Notes. References to this Table: (1) Kirkpatrick et al. (2000), (2) This paper, (3) Reid et al. (2008), (4) Cruz et al. (2007), (5) Cruz et al. (2003), (6) Gagné et al. (2014a), (7) Chiu et al. (2006), (8) Gagné et al. (2014b).

^a Kinematic distances estimated from moving group membership unless noted.

^b Trigonometric distance.

(Chile), the Australian Research Council (Australia), Ministério da Ciência, Tecnologia e Inovação (Brazil), and Ministerio de Ciencia, Tecnología e Innovación Productiva (Argentina). All data were acquired through the Canadian Astronomy Data Center and part of it was processed using the Gemini IRAF package. This material is based upon work supported by AURA through the National Science Foundation under AURA Cooperative Agreement AST 0132798 as amended. This publication uses observations obtained at IRTF through programs number 2007B023, 2007B070, 2008A050. 2008B054, 2009A055, 2010A045, 2011B071, 2012A097, 2012B015, 2013A040, 2013A055, 2013B025, 2014B026, and 2015A026. The authors recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

Facilities: IRTF (SpeX), *Magellan*:Baade (FIRE), Gemini: South (Flamingos-2), Hale (TripleSpec).

APPENDIX A DISCUSSIONS ON INDIVIDUAL OBJECTS

Several objects presented here deserve a detailed discussion, either because they display peculiar features, or were reported in the literature as candidate members of other YMGs. Additionally, optical spectra were available in the literature for some objects discussed here, and can serve as an independent assessment of low surface gravity.

A.1 Potential Planetary-mass Low-gravity Candidate Members of YMGs

We list in Table 9 20 potential isolated planetary-mass objects in our sample, 10 of which were discovered as part of this work. A few of these objects deserving further discussion are listed below.

2MASS J05012406-0010452 was discovered by Reid et al. (2008) as an L4 dwarf in the optical, and was categorized as a low-gravity L4 γ by Cruz et al. (2009), using its optical spectrum. Allers & Liu (2013) categorized it as a very-low gravity L3 dwarf in the NIR, whereas we categorize it as an L4 γ dwarf. Faherty et al. (2012) measured a trigonometric distance of 13.1 ± 0.8 pc. We recovered this object in BASS as an ambiguous candidate member of Columba or Carina with respective Bayesian probabilities of 49% and 17%, taking the trigonometric distance measurement of Faherty et al. (2012) into account. If this object is a member of either COL or CAR (both YMGs are coeval at 20-40 Myr), it has an estimated mass of $10.2^{+0.8}_{-1.0}$ M_{Jup}. Zapatero Osorio et al. (2014) independently measured a trigonometric distance of 19.6 ± 1.4 pc, which is discrepant with that of Faherty et al. (2012) at the 5σ level. The reason for this large discrepancy is unclear; the measurement of Faherty et al. (2012) used a smaller number of epochs (11 versus 21); however, they were spread across a larger temporal coverage (3 versus 2 year). If we adopt the distance measurement of Zapatero Osorio et al. (2014), the CAR membership probability becomes negligible and that of COL becomes considerably smaller (7.2%), although we also calculate a low field contamination probability (1.3%). It will be necessary to better constrain the distance of this object to assess whether it is

a viable candidate member of COL or CAR. Obtaining an RV measurement would also be useful for this.

2MASS J05120636–2949540 has been identified as an L4.5 dwarf in the optical by Cruz et al. (2003), Kirkpatrick et al. (2008), and Bardalez Gagliuffi et al. (2014) obtained a NIR spectrum to categorize it as an L4.5 ± 2 dwarf. In Paper II, we determined that this object is a low-probability candidate member of β PMG. We used the NIR spectrum of Bardalez Gagliuffi et al. (2014) to revisit its spectral classification: we find that this object is a very good match to our L5 β template; however, the method of Allers & Liu (2013) assigns it an intermediate gravity. We note that the VO_Z index is significantly larger than that of field L5 dwarfs, but Allers & Liu (2013) only use this index within the L0–L4 spectral types, as later-type low-gravity dwarfs in their sample displayed similar VO absorption than that of field dwarfs of the same spectral types. However, only one low-gravity L5 dwarf was available at the time, hence it is possible that the VO_Z index remains useful to discriminate low-gravity L5 dwarfs. For this reason, we adopt the L5 β spectral type. Due to its low-gravity features, this object is preserved as a candidate member of β PMG. This object has one of the lowest estimated masses among the YMG candidates presented here, with $6.7^{+1.0}_{-0.9} M_{Jup}$. Its statistical distance associated with membership to β PMG is $10.9^{+4.4}_{-4.0}$ pc, which makes it a valuable benchmark to study the atmosphere of planetary-mass objects.

2MASS J12074836-3900043 (2MASS J1207-3900) was discovered as a candidate member of TWA in BASS. Its discovery and NIR spectroscopic follow-up have been presented in Gagné et al. (2014a). They reported an optical spectral type L0 γ and a NIR spectral type L1 γ . Here we used the spectra of several low-gravity candidate members of Upper Scorpius obtained by Lodieu et al. (2008) to define tentative templates for the spectral type L0 δ , which likely correspond to objects younger than \sim 15–20 Myr and have an even more triangular *H*-band continuum than the L0 γ type. Given that both the optical and NIR spectra of 2MASS J1207-3900 are peculiar even in comparison to the best template matches (L0 γ and L1 γ , respectively) and that its H band continuum is more triangular than any β or γ template, we revised its spectral classification by comparing it to Upper Scorpius candidate members. We find that the best match is the L0 δ template; however, 2MASS J1207-3900 displays features that are attributable to a later spectral type (redder slopes at 1.2–1.35 μ m and 1.5–1.6 μ m). We thus suggest a tentative spectral type of L1 δ for this object, but identifying other similar objects will be necessary to confirm this. If it is a member of TWA (5–15 Myr), this object has an estimated mass of $12.1_{-2.0}^{+1.4} M_{Jup}$ and a statistical distance of $58.2_{-6.4}^{+6.8}$ pc.

2MASS J12271545–0636458 was identified as an M9 dwarf by Cruz et al. (2003) using optical spectroscopy. We identified it as a candidate member of TWA in PRE-BASS, and NIR spectroscopy allowed us to categorize it as a low-gravity M8.5 β dwarf. It was initially rejected from the BASS sample because of its low Bayesian probability, which is in part due to the fact that its kinematic distance of 32.5 \pm 3.2 pc if it is a member of TWA does not match its spectrophotometric distance (63.2 \pm 11.4 pc). The latter estimate would place 2MASS J12271545–0636458 at the far-end of the TWA members (~40–62 pc; Paper II; Weinberger et al. 2013; Ducourant et al. 2014). This is reminiscent of 2MASS J12474428–3816464, TWA 29, and TWA 31, which are young and seem to be located between TWA and SCC in terms of distance (Song et al. 2003; Schneider et al. 2012; Gagné et al. 2014a; Paper V). Measurements of distance and RV will be useful to assess whether this is a true member of TWA despite its small Bayesian probability. If it is a true member of TWA (5–15 Myr) located at its statistical distance, this object has an estimated mass of $11.6^{+1.4}_{-1.9} M_{Jup}$.

2MASS J12563961-2718455 was identified in PRE-BASS as a low-probability candidate member of TWA. NIR spectroscopy revealed that this object is a low-gravity L3 $\pm 1 \beta$ dwarf. The probability that this object belongs to TWA is lower than 20%, but the field contamination probability is also very low at <0.1%. This usually points out to either an incomplete SKM for the YMG or to contamination from a source not taken into account in BANYAN II. The most likely a priori explanation would be that this object is a contaminant from SCC (located at \sim 100–150 pc; Sartori et al. 2003); however, the spectrophotometric distance of 2MASS J12563961-2718455 $(43.1 \pm 3 \text{ pc})$ is not consistent with this hypothesis, even when its low gravity is taken into account. Using its 2MASS and WISE photometry and comparing it with other known lowgravity L4 dwarfs, we can rule out a distance larger than 48.5 pc at a 95% confidence level, assuming this object is not an unresolved multiple system. We can hence conclude that as long as this object is not extremely peculiar for a low-gravity L4 dwarf or a multiple system composed of four equalluminosity components, it cannot be a member of SCC. The statistical distance from BANYAN II which is associated to the TWA hypothesis $(46.2^{+4.8}_{-4.4} \text{ pc})$ is similar to those of bona fide members of TWA (~40-62 pc; Paper II; Weinberger et al. 2013; Ducourant et al. 2014), hence this case is different from those of 2MASS J12271545-0636458, 2MASS J12474428-3816464, TWA 29, and TWA 31, which are young and seem to be located between TWA and SCC in terms of distance (Song et al. 2003; Schneider et al. 2012; Gagné et al. 2014a; Paper V). Obtaining a distance measurement for this object will be helpful to assess whether it is a member of TWA. Assuming an age of 5-15 Myr and comparing its statistical distance from BANYAN II with BT-Settl 2MASS and WISE isochrones, the estimated mass of this object is $7.7^{+1.4}_{-1.5}$ M_{Jup}, among the lowest of all candidate YMG members reported here. Its statistical distance associated with membership to TWA is 44.6 ± 5.2 pc.

2MASS J21324036+1029494 was discovered as an L4.5±1 dwarf by Chiu et al. (2006) using low-S/N NIR spectroscopy. We identified it as a candidate member of ARG from PRE-BASS. The NIR spectrum obtained by Chiu et al. (2006) is available in the SpeX PRISM Spectral Libraries, we thus retrieved it to assess whether it is a low-gravity dwarf. We categorize this object as an L4: β dwarf. Its *H*-cont index (Allers & Liu 2013) is consistent with low-gravity objects; however, the quality of the data is not sufficient to assess whether its FeH_Z and KI_J indices are consistent with this. Obtaining a better-quality and higher-resolution NIR spectrum will be useful to confirm the spectral type of this object. If it is a member of ARG (30–50 Myr), this object has an estimated mass of 11.4 ± 0.4 M_{Jup} and a statistical distance of 34.2 ± 4.8 pc.

A.2 Low-gravity Candidate Members of YMGs

2MASS J00413538–5621127 (DENIS-P J00041353–562112) has been identified as a candidate nearby, red dwarf by Phan-

Bao et al. (2001), and spectroscopically confirmed by Schmidt et al. (2007) as an active M8 dwarf. Using high resolution optical spectroscopy, Reiners (2009) revised its spectral type to M7.5 and showed evidence that it displays Li and signatures of active accretion, which indicates that it is a young, $\sim 10 \text{ Myr BD}$. Based on its position, proper motion, and RV, they suggest that it could be a member of THA, or an ejected member of β PMG, which would make it the first accreting BD discovered in either of these associations. Liu et al. (2010) reported that it is a binary with estimated spectral types of M6.5 \pm 1 and M8 from photometry. In Paper II, we corroborated that it is a highprobability candidate member of THA, with estimated masses of 14–41 M_{Jup} and 18–41 M_{Jup} for the individual components. This object was retrieved in BASS as a high-probability candidate of THA. We obtained NIR spectroscopy for the unresolved system, and categorize it as a very low-gravity M7.5 γ BD system, which is consistent with its young age.

2MASS J02590146-4232204 was identified by Rodriguez et al. (2013) as a candidate member of COL with infrared excess indicative of the presence of a circumstellar disk host. We independently identified this object as a candidate member of COL in PRE-BASS; however, it was subsequently rejected from BASS because of its low membership probability and the fact that its WISE colors did not survive the extragalactic filter defined by Kirkpatrick et al. (2011), which is likely a consequence of its infrared excess. NIR spectroscopy allowed us to categorize it as an M5 γ dwarf. This is consistent with the results of Rodriguez et al. (2013), who reported that this object displays weak Na I absorption that is indicative of a low surface gravity. Including the RV measurement of 15.3 \pm 1.5 km s^{-1} from Rodriguez et al. (2013), we find that this object is a low-probability candidate member of COL: this conclusion differs from that of Rodriguez et al. (2013), which found that 2MASS J02590146-4232204 is a candidate member of THA. Obtaining a trigonometric distance will be useful to assess whether this object is a member of COL or THA.

2MASS J03264225-2102057 has been identified as an L4 dwarf with Li absorption by Cruz et al. (2007). Using the DUSTY evolution models (Chabrier et al. 2000), the presence of Li and the spectrophotometric absolute magnitude of this object, they determined that it should be younger than 500 Myr and less massive than $50 M_{Jup}$. We identified this object as a highly probable L5 β/γ candidate member of ABDMG in PRE-BASS. Dahn et al. (2002) measured a trigonometric distance of 32.3 ± 1.6 pc that is consistent with membership to ABDMG. The presence of low-gravity feature in its optical and NIR and optical spectra puts a slightly stronger constraint on the age of 2MASS J03264225-2102057, since it is expected that gravity-sensitive spectral indices remain useful only up to \sim 200 Myr (Cruz et al. 2009; Allers & Liu 2013). We therefore categorize this object as a low-gravity L5 β/γ dwarf. An RV measurement is needed before it can be assessed whether this object is a bona fide member of ABDMG.

2MASS J04493288+1607226 was identified in PRE-BASS as a candidate member of β PMG, but was rejected from the BASS sample because of its proximity with TAU. NIR spectroscopy revealed that it is a low-gravity M9 γ dwarf. We estimate a distance of 54.9 \pm 10.0 pc for this object by comparison with other low-gravity dwarfs. A distance larger than 82 pc can be excluded at a 99% confidence level, which is incompatible with membership to TAU (140 \pm 20 pc; Torres et al. 2007) unless it is an unresolved multiple with at least 3 individual equal-luminosity components. This scenario is unlikely, especially considering that the NIR spectrum of 2MASS J04493288+1607226 is not reddened. We thus preserve this object as a candidate member of β PMG.

2MASS J11083081+6830169 has been discovered by Gizis et al. (2000) as an L1 dwarf in the optical with H α emission. We recovered this object in BASS as a candidate member of ABDMG, and Gizis (2002) identified it as a candidate member of TWA. The RV of -9.8 ± 0.1 km s⁻¹ measured by Blake et al. (2010) does not match the predicted RV of -18.9 ± 1.5 km s⁻¹ for membership to ABDMG. It closely matches that of the CAR hypothesis (-9.7 ± 0.8), but it still obtains a very low Bayesian probability of being a member of CAR. Its statistical distance (15.3 \pm 0.8 pc) places it right into the locus of known young L dwarfs in both an M_{W1} versus $J - K_S$ and M_{W1} versus H - W2 CMDs. This distance places it at only 0.27 km s^{-1} of the CAR bona fide member HIP 33737 in UVW space, and at 17.2 pc of the CAR bona fide member GJ 2079 in XYZ space. We show in Figure 19 its XYZUVW position at its most probable distance: it seems that this object has a most probable position that is consistent with bona fide members of CAR, but our SKM fails to represent this. It can be expected that our SKM of CAR is not accurate because it was derived from a small number of bona fide members. Furthermore, both the NIR spectrum that we obtained and the optical spectrum from the RIZzo spectral library display clear signs of lowgravity and allowed us to categorize it as an L1 γ dwarf, which is consistent with membership to a YMG. A measurement of this object's trigonometric distance will be useful to assess whether or not it is a member of CAR, but we note that it is likely a member despite its low Bayesian membership probability.

2MASS J12265135-3316124 (TWA 32) has been identified by Shkolnik et al. (2011) as an UV-bright M6.5 low-mass star. They measured strong H α emission and Li absorption, as well as an RV of 7.15 \pm 0.26 km s⁻¹. They used this information as well as a photometric distance $(53 \pm 5 \text{ pc})$ to identify it as a new member of TWA, and they noted that it is a 656.1 \pm 0.4 mas visual binary with near-equal luminosity. Rodriguez et al. (2011) independently discovered this object and measured strong H α and HeI emission at 5876 and 6678 Å, as well as strong Li absorption. They argued that the $H\alpha$ full width at 10% of 270 km s^{-1} is consistent with this object being a classical T Tauri star. They measured an RV of 14.8 ± 3 km s⁻¹ and note that its UVW space velocity is consistent with TWA and SCC. We recovered this object in PRE-BASS as a candidate member of TWA, and obtained NIR spectroscopy that allowed us to assign it a spectral type of M5.5 γ . At this spectral type, only the weaker Na1 absorption is a useful low-gravity indicator. We adopted the RV measurement of Shkolnik et al. (2011) which is more precise, and, like them, found that this object is a strong candidate member of TWA. The BANYAN II statistical distance corresponding to the TWA hypothesis is $61.8^{+6.4}_{-6.0}$ pc, which is consistent with the photometric estimate of Shkolnik et al. (2011) that takes its binary nature into account. The SKMs of BANYAN II do not take SCC into account, which includes the Lower Centaurus Crux (LCC) and the Upper Centaurus Lupus (UCL) regions, hence our result does not preclude membership to SCC. The space velocity UVW for this object is (-8.6 ± 1.4) , -15.7 ± 1.1 , -3.4 ± 1.1) km s⁻¹(Shkolnik et al. 2011), at $4.6 \,\mathrm{km \, s^{-1}}$ from the kinematic center of TWA (Paper II), 4.6 km s^{-1} from that of UCL and 4.2 km s^{-1} from that of LCC



Figure 19. Predicted galactic position XYZ and space velocity UVW of the CAR candidate member 2MASS J11083081+6830169 (red point and its projections) using its statistical distance from BANYAN II, compared with bona fide members of CAR (green points and their vertical projections on the XY and UV planes) and the SKM models of CAR (as defined in Paper II; orange ellipsoid and its projections).

(Sartori et al. 2003). Its kinematics are thus consistent with SCC and TWA; however, its photometric distance is not consistent with the distance of this complex (\sim 100–150 pc; Sartori et al. 2003), whereas it is consistent with that of TWA members (\sim 40–62 pc; Paper II; Weinberger et al. 2013; Ducourant et al. 2014). We conclude that TWA 32 is a likely member of TWA, unless it is a multiple system composed of at least three equal-luminosity components. A trigonometric distance measurement will be useful to assess this.

2MASS J20391314–1126531 was discovered as an M9 dwarf by Cruz et al. (2003) using optical spectroscopy. Gálvez-Ortiz et al. (2010) reported that it is a candidate member of the Pleiades stream. Famaey et al. (2005) demonstrated that the Pleiades stream is not a moving group but rather a dynamical stream of stars without a common origin. We identified 2MASS J20391314–1126531 as a candidate member of ABDMG as part of PRE-BASS and obtained NIR spectroscopy which revealed that this is a low-gravity M7 β dwarf. The RV of -18.0 ± 2 km s⁻¹ that was measured by Gálvez-Ortiz et al. (2010) is consistent with a membership to ABDMG, and the fact that it has a low gravity indicates that it might not be a contaminant from the Pleiades stream. A measurement of its distance will be needed to assess this.

A.3 Candidate Members of YMGs with no Age Constraint

2MASS J03582255–4116060 has been discovered by Cruz et al. (2007) as an L5 BD in the optical. We identified it as a low-probability candidate member of β PMG as part of BASS. $R \sim 75$ NIR spectroscopy allowed us to categorize it as a peculiar L6 dwarf. Its continuum is redder and its H band is slightly more triangular than our field L6 template, however it is unclear at this time if these effects are due to a low gravity or not. Obtaining a higher-resolution spectrum would be useful to assess this. If we assume an age of 20–26 Myr and the BANYAN II statistical distance associated with the β PMG hypothesis (18.1 ± 3.2 pc) and compare its 2MASS and WISE photometry with the BT-Settl isochrones, we find that this object has one of the lowest estimated mass of all candidate YMG members reported here, with 8.2 \pm 0.6 M_{Jup} .

2MASS J08095903+4434216 was identified by Knapp et al. (2004) and confirmed by Chiu et al. (2006) as an L6 dwarf. In Paper V, we identified it as a candidate member of ARG as part of BASS. We used its NIR spectrum to revise its spectral type to L6 pec (red) from a visual comparison with field and lowgravity templates. This object has a red continuum and red NIR colors for its spectral type, with $J - K_S = 2.02$ and J - W2 = 3.63, compared with median values of $J - K_S = 1.7 \pm 0.3$ and $J - W^2 = 2.9 \pm 0.4$ for field L6 dwarfs (Figure 11). The low-resolution gravity classification scheme of Allers & Liu (2013) categorizes it as an intermediate-gravity L5.4 dwarf due to its H-cont, KI₁ and FeH_Z indices. However, it is visually a better match to the field L6 template than the field L5 template, albeit it displays a slightly redder continuum. Adopting a spectral type of L6, only the H-cont index remains useful and categorizes it as an intermediate-gravity dwarf, but this index alone does not reject the possibility that this object is a dusty dwarf in the field. Schneider et al. (2014) demonstrated that the $H_2(K)$ index defined by Canty et al. (2013) seems to be gravity-sensitive up to at least L8; we obtain a value of $H_2(K) = 1.056 \pm 0.008$ for 2MASS J08095903+4434216, which is slightly lower than the typical values for field L6 dwarfs (1.06 \pm 0.01; see Figure 14(d) of this work and Figure 10 of Schneider et al. 2014). It is unclear at this time if this object is a lowgravity L6 dwarf; a higher resolution ($R \gtrsim 750$) NIR spectrum will be useful to confirm if this object is a very low-mass, very late-type candidate member of ARG, or more massive and dusty field interloper. At the age of ARG (30-50 Myr), this object would have one of the lowest estimated masses among all YMG candidates presented here, with 8.1 \pm 0.8 M_{Jup} . Its statistical distance associated with membership to ARG is 15.3 ± 2.0 pc.

2MASS J23512200+3010540 was discovered by Kirkpatrick et al. (2010) as L5.5 dwarf in the optical, and as an unusually red L5.5 dwarf in the NIR. In Paper II, we identified



Figure 20. NIR spectrum of the peculiar L0 dwarf 2MASS J00461551 +0252004 that was recovered as a candidate member of ABDMG in PRE-BASS. The *H*-band bump at 1.57 μ m could be a hint of an unresolved T-type companion.

this object as a candidate member of ARG, and it was recovered as such in PRE-BASS. We used its NIR spectrum to categorize it as a peculiar L5 dwarf. Only the *H*-cont index is indicative of a possible young gravity; it is thus a likely scenario that this object is a dusty field interloper. Obtaining a higher-resolution spectrum would be useful to assess this. If it is a member of ARG (30–50 Myr), this object has an estimated mass of $10.0^{+0.6}_{-0.7}$ M_{Jup} and a statistical distance of 20.1 ± 1.8 pc.

A.4 Interlopers from the Field or Other Regions

2MASS J00174858-0316334 was identified as a candidate member of ABDMG as part of PRE-BASS. NIR spectroscopy revealed that this is a reddened low-gravity M7 β dwarf. We de-reddened its spectrum using the fm_unred.pro IDL routine based on the extinction law of Fitzpatrick (1999) and visually compared it with our M7 β template to determine that its total extinction is A(V) = 2.5. We used the parametrization of Fitzpatrick (1999) with a total-to-selective extinction of R(V) = 3.1. This reddening is unlikely caused by interstellar dust, since this object is far from the galactic plane $(b = -64^{\circ}.8)$ and has a spectro-photometric distance of only 65.1 ± 11.5 pc (low gravity was considered in this estimate). It can be expected that this object is still embedded in its formation material, which indicates that it is not a member of ABDMG, but rather a member of another young star-forming region that might not be known. It would be interesting to investigate whether other very young objects can be found in its vicinity.

2MASS J00461551+0252004 was identified as a candidate member of ABDMG in PRE-BASS. NIR spectroscopy revealed that it is a peculiar L0 dwarf with no indication of low gravity. The *H*-band bump at 1.57 μ m is significantly stronger than that of field dwarfs, which could hint at an unresolved T-type component (Figure 20); however, the spectral indices constructed by Bardalez Gagliuffi et al. (2014) do not categorize it as a likely L-type + T-type binary. The cause of its peculiar properties is thus unclear.

2MASS J02441019–3548036 was discovered as candidate member of THA in BASS. NIR spectroscopy ($R \sim 750$)



Figure 21. NIR spectrum of the peculiar L1 dwarf 2MASS J02441019–3548036 that was recovered as a candidate member of THA in BASS. Its NIR continuum is redder than usual and the shape of its H band is peculiar. This could be explained by a dusty atmosphere; however, this object is likely older than the YMGs considered here.

reveals that it is an L2 dwarf that lacks the weaker alkali lines or stronger VO absorption that are typical of low-gravity dwarfs. However, its continuum is unusually red for an L2 dwarf and the shape of its H band is unusual (Figure 21). This could be explained by an unusually dusty atmosphere or an unresolved later-type companion; however, the classification of Bardalez Gagliuffi et al. (2014) based on various spectral indices does not categorize it as a candidate binary. We reject it as a candidate member of THA, as the weaker-than-usual alkali lines are not consistent with a young age even if this object is dusty or multiple.

2MASS J02530084+1652532 (Teegarden's star) was discovered by Teegarden et al. (2003) as a nearby (2.43 \pm 0.54 pc) M6.5 dwarf; Henry et al. (2006) refined its distance measurement to 3.85 ± 0.01 pc. Witte et al. (2011) identified this object as a potential low-gravity dwarf from atmosphere model fitting, and we identified it as a candidate member of ARG in Paper II, but its distance is not consistent with this possibility. We obtained the NIR spectrum of Burgasser et al. (2012) from the SpeX PRISM spectral library and categorized it as an M7.5 β dwarf. The classification scheme of Allers & Liu (2013) assigns it an intermediate gravity, which is consistent with our visual comparison. This is due to a low FeH_Z index (1.0699 \pm 0.0040), a low KI_J index (1.0511 \pm 0.0065) and a high *H*-cont index (0.9974 \pm 0.0049) compared with field M7.5 dwarfs. This object has a relatively blue 2MASS $J - K_S$ color (0.809 \pm 0.053) for an M7.5 dwarf (Figure 11(a)), which is not expected for a low-gravity dwarf. We used the NIR spectrum to measure its synthetic NIR colors and obtained $J - K_S = 0.869 \pm 0.077$ (assuming a 5%) uncertainty in the 2MASS photometric zero points), which places it closer to the locus of low-gravity and field M7 dwarfs, albeit still on the blue end. Its WISE W1 - W2 color (0.265 \pm 0.034) is consistent with field and low-gravity M7.5 dwarfs (Figure 10(i)). Obtaining a higher-resolution NIR spectrum will be useful to assess whether alkali lines are weaker than usual, which would confirm if this object has a low gravity or not. Another explanation could be that this object is an unresolved binary. If Teegarden's star is young, it could be a member of a

YMG that is not considered here, and it would thus be interesting to measure its RV. It is worthwhile mentioning that this object would be the nearest low-gravity dwarf if this is confirmed, a record currently held by LP 944–20 (an L0 β at 6.41 ± 0.04 pc that is a candidate member of the Castor stream; Barrado Y Navascués 1998; Leggett et al. 2001; Allers & Liu 2013; Dieterich et al. 2014).

2MASS J03140344+1603056 was identified by Schmidt et al. (2007) as an L0 dwarf with H α emission. Seifahrt et al. (2010) measured its RV and used its kinematics to assign it as a candidate member of UMA. We initially identified this object as a low-probability candidate member of β PMG in PRE-BASS, but was later rejected because of its large contamination probability, as well as its position on a M_{W1} versus H - W2diagram that is not consistent with young BDs at the most probable statistical distances obtained from BANYAN II. A NIR follow-up allowed us to categorize it as a peculiar M9 dwarf with no apparent sign of low gravity from the classification scheme of Allers & Liu (2013) or a visual comparison with spectroscopic standards. However, it is unclear at what exact age signs of low gravity stop being apparent in moderate-resolution NIR spectra, and Allers & Liu (2013) suggest that this might take place around \sim 200 Myr. We thus reject any possible membership with the younger moving groups considered here, but our data is insufficient to corroborate its possible membership to UMA.

2MASS J04070752+1546457 has been identified as an L3.5 dwarf by Reid et al. (2008) from optical spectroscopy. We identified this BD as an ambiguous candidate member of β PMG and COL as part of PRE-BASS, but we subsequently rejected it because of its alignment with TAU. NIR spectroscopy allowed us to categorize it as field L3 BD. It displays marginal signs of low gravity (weaker FeH and slightly weaker alkali line widths); however, all other features as well as a visual comparison with spectroscopic standards are consistent with a field L3 BD. It could be interesting to investigate whether this object has a peculiar metallicity or a slightly young age (~200 to a few hundred Myrs), but it is most probably not a member of any YMG considered here.

2MASS J05243009+0640349 has been identified as a potential member of β PMG in PRE-BASS. It has been subsequently excluded from BASS because of its low Bayesian probability, but its NIR spectrum allowed us to categorize it as a low-gravity M5.5 β . The only useful sign of low gravity for this spectral type is the weaker Na I absorption. This object has a low galactic latitude (b = -15.92) and is located within the Orion II super bubble (Gatley et al. 1974) at only 10'6 of the Ori C 11 core (see Figure 12(b) of Wood et al. 1994; the B1950 coordinates of this object are $05^{h}21^{m}48.67$, $+06^{\circ}37.54.5$). These clouds are located at significantly larger distances (~400-500 pc; Schlafly et al. 2014) compared to the YMGs considered here. Using the absolute 2MASS and WISE photometry of known young M5.5 dwarfs, we estimate a spectrophotometric distance of 42.0 \pm 7.7 pc for 2MASS J05243009+0640349 and exclude a distance larger than 63 pc at a 99% confidence level, assuming it is not a multiple system. This discrepancy, supplemented with the fact that its spectrum does not seem reddened by interstellar dust, makes it unlikely that this object is a member of the Orion Molecular Complex (OMC) even though it is clearly young. Even when its youth is taken into account, this object has a very low probability of being a member of β PMG. It will be useful to obtain a distance

and RV measurement to investigate whether this object is a member of another YMG that is not considered here.

2MASS J05271676+0007526 has been identified as a potential member of β PMG in PRE-BASS. Its low Bayesian probability as well as color filters $(H - K_S > 0.269)$ and $VR - J \ge 2.63$; Paper II) excluded it from the BASS sample. This object has a low galactic latitude ($b = -18^{\circ}.57$) and is located in the vicinity of the OMC (Gatlev et al. 1974: Schlafly et al. 2014). Acquisition images obtained with SpeX revealed that this is a 2".4 visual binary. We obtained resolved NIR spectroscopy and determined that both components are reddened early M dwarfs. We de-reddened both spectra using the fm unred.pro IDL routine based on the extinction law of Fitzpatrick (1999) and visually compared the results with NIR spectroscopic standards to determine the best matching spectral types and total extinction A(V), using the parametrization of Fitzpatrick (1999) with a total-to-selective extinction of R(V) = 3.1. We find a best match of A(V) = 0.93 with spectral types M0 + M3. We note that the *H*-band continuum of both objects has a rounded triangular shape, which is only seen at those spectral types for very young (≤ 5 Myr) objects. This system is thus likely very young and still embedded in its formation material. Using the 2MASS J magnitude of the unresolved system with the young absolute magnitude-spectral type sequences of Malo et al. (2013), we estimate a distance of \sim 450 pc. We conclude that this system is a probable very young low-mass star member of OMC.

2MASS J08503593+1057156 (2MASS J0850+1057) was first identified from the 2MASS survey as an L6 BD by Kirkpatrick et al. (1999). Subsequently, Reid et al. (2001) and Bouy et al. (2008) identified and confirmed that this object is a 0".16 binary system, and Burgasser et al. (2011) used a template fitting method constrained by the flux ratio of its individual components to assign them spectral types of L7 and L6. They noted the surprising fact that the brighter primary component is the one that gets assigned a later spectral type. They argue that this could be explained either by youth or the latest-type component being an unresolved binary. Faherty et al. (2011) subsequently identified the NLTT 20346 M5+M6 binary system as a very wide (~7700 AU) co-moving companion to 2MASS J0850+1057. They assign an age estimate of 250-450 Myr for NLTT 20346 based on X-ray luminosity, but they note that this estimate is discrepant with that based on H α emission (6.3 \pm 1.0 Gyr and 6.5 \pm 1.0 Gyr for its respective components). They measure a systemic RV of $26 \pm 9 \text{ km s}^{-1}$ for NLTT 20346, and a trigonometric distance of 29 \pm 7 pc for 2MASS J0850+1057. They note that this latter measurement is not precise enough to discriminate between two previous inconsistent measurements in the literature (38 \pm 6 pc from Vrba et al. 2004 and 25.6 \pm 2.3 pc from Dahn et al. 2002). Using their proper motion measurement of $\mu_{\alpha} = -144 \pm 6$ mas yr⁻¹ and $\mu_{\delta} = -38 \pm 6$ mas yr⁻¹, they argued that a faint background contaminant was blended at the epochs used for previous distance measurements, which could explain the discrepancy.

Dupuy & Liu (2012) independently measured a proper motion of -144.2 ± 0.6 mas yr⁻¹, $\mu_{\delta} = -12.6 \pm 0.6$ mas yr⁻¹ and a distance of 33.2 ± 0.9 pc for 2MASS J0850+1057. They also refined the photometry of its resolved components, and used these new measurements to draw different conclusions than those outlined above. First, they used a similar analysis to that of Burgasser et al. (2011) with their updated photometry to argue



Figure 22. NIR spectrum of the peculiar L7 dwarf 2MASS J0850+1057 that was recovered as a candidate member of ARG in BASS. This system is a binary with resolved spectral types of $L6.5\pm1$ and $L8.5\pm1$ estimated from photometry. The NIR continuum of 2MASS J0850+1057 is redder than field L7 dwarfs which is likely an effect of its binary nature, and its kinematics are not consistent with those of ARG.

that the spectral types of the components are rather L6.5±1 and L8.5±1, with the fainter component now associated with the later spectral type. This conclusion does away with the need to invoke youth or any additional component, which was previously based on flux reversal (Burgasser et al. 2011). They thus argued that 2MASS J0850+1057 is a BD system displaying no notable peculiarity. Furthermore, they use their new proper motion measurement at 6.7σ from that of NLTT 20346 with the criterion of Lépine and Bongiorno (2007) to argue that the two systems are likely random alignments, and thus not gravitationally linked.

We measure a proper motion of -141.1 ± 7.7 mas yr⁻¹ and $\mu_{\delta} = -13.1 \pm 9.5 \text{ mas yr}^{-1}$, based on 2MASS and ALLWISE. The μ_{α} component is consistent with both measurements, whereas the μ_{δ} component is at 2.2 σ and 0.05 σ respectively from the measurements of Faherty et al. (2011) and Dupuy & Liu (2012). Our measurement thus favors the later one, but our precision is ~ 16 times lower. We find that even if both components seem to display no peculiarity in their relative fluxes, the unresolved system seems to be unusually red for its absolute magnitude (Figure 22). This could be explained either by additional unresolved later-type components, or the presence of thicker/higher clouds in their atmosphere compared to field BDs. We used the $R \sim 120$ NIR spectrum for the unresolved BD system to assign a spectral type of L7. For such a late spectral type, the only features known to be gravitysensitive in a low-resolution NIR spectrum are the H-cont index of Allers & Liu (2013) and the $H_2(K)$ index of Canty et al. (2013). We find values of H-cont = 0.872 ± 0.025 and $H_2(K) = 1.055 \pm 0.014$, both being only marginally consistent with a low surface gravity. The parallax and proper motion measurements of Dupuy & Liu (2012) preclude a possible membership to ARG; however, obtaining a higher-resolution NIR spectrum for this system would be interesting to assess whether it displays signs of low surface gravity. If this system is younger than $\sim 200 \text{ Myr}$, the individual mass of each component would be well below the deuterium burning limit,



Figure 23. NIR spectrum of the peculiar L8 dwarf 2MASS J08575849 +5708514 that was recovered as a candidate member of ARG in BASS. Its red NIR continuum could be an indication of a low surface gravity, but an RV measurement that was obtained from low-S/N data is not consistent with membership to ARG.

which would make it a remarkable benchmark system to understand the properties of planetary-mass objects.

2MASS J08575849+5708514 has been discovered by Geballe et al. (2002) as an L8 \pm 1 BD. Stephens et al. (2009) used atmosphere model fitting to determine that this object is unusually cloudy and seems to have a low surface gravity $(\log g = 4.5)$. Using our visual comparison with spectral templates, we categorize this object as a peculiar L8 dwarf (Figure 23). No indices from Allers & Liu (2013) are gravitysensitive for such a late spectral type, but the $H_2(K)$ index defined by Canty et al. (2013) seem to remain useful (Schneider et al. 2014). We find a weaker $H_2(K)$ value (1.102 ± 0.006) compared with typical field L8 dwarfs (1.12 \pm 0.02), which could be an indication of a lower surface gravity. We identified this object as a highly probable candidate member of ARG in BASS, with an estimated mass of 8.5 \pm 0.8 $M_{\rm Jup}$ and an estimated distance of 8.9 \pm 0.8 pc. However, Schmidt et al. (2010) measured an RV of -123.5 ± 20.0 km s⁻¹, which is not consistent with membership to ARG or even with the kinematics of any young BD in the solar neighborhood. This large RV thus seems contradictory with its unusually red colors and tentative indications of a lower surface gravity, but it was measured from a low-signal optical spectrum. It is likely that this object is not a young member of ARG but rather an interloping cloudy object from the field. However, obtaining an RV measurement from higher-S/N data will be useful to assess this.

2MASS J11335700–7807240 was identified by Luhman (2007) in the optical as an M8 dwarf in a search for new members of the Chamaeleon I (CHA) star-forming region. They rejected it because it lacks low-gravity indications in its optical spectrum. We independently recovered this object in PRE-BASS as a candidate member of CAR and obtained a NIR spectrum. We categorize this object as a peculiar M6±1 dwarf; its *H* band is more triangular and the slope of its *K* band is bluer. This could be indicative of a low surface gravity, but it lacks all the other usual signatures: only the 1.253 μ m K I line is slightly weaker than that of field M6 dwarfs. All other K I lines, the Na I doublet and FeH absorption are all consistent

with a field M6 dwarf : the classification scheme of Allers & Liu (2013) thus categorizes this object as a field-gravity M6 dwarf. It is unclear what is the source of the peculiar features in this object's NIR spectrum. It is possible that the triangular-shaped *H* band could be caused by dust in its photosphere (Allers & Liu 2013), but this would be unusual at such an early spectral type, and neither its $J - K_S$ color (1.01 \pm 0.04) or its J - W2 color (1.53 \pm 0.04) are redder than those of field M6 dwarfs, which would be unexpected for a dusty object. We thus categorize this object as a peculiar M6 dwarf and reject it as a candidate member of CAR.

2MASS J11555389+0559577 was discovered by Knapp et al. (2004) as an L7.5 dwarf using NIR spectroscopy. We recovered this object in PRE-BASS as a candidate member of ARG. Faherty et al. (2012) measured a trigonometric distance of 17.27 ± 3.04 pc and Schmidt et al. (2010) used a low-quality optical spectrum from SDSS to categorize it as an L0 dwarf and measure an RV of 136.8 ± 20.0 km s⁻¹. If we include only the trigonometric measurement, it remains a modest candidate of ARG; however, the RV measurement is not consistent with this, nor with the kinematics of nearby, young dwarfs (Faherty et al. 2009), much like the case of 2MASS J08575849+ 5708514. We retrieved the NIR spectrum of this object from the SpeX PRISM Libraries and categorize it as a peculiar L6-L8 dwarf. It lacks the triangular-shaped H-band continuum that would be expected for a young object (H-cont = 0.8230 ± 0.0087), and its NIR colors are consistent with those of field dwarfs. We measured the gravity-sensitive $H_2(K)$ index defined by Canty et al. (2013) and find a value of 1.0862 \pm 0.0085, which is consistent with field L6-L7 dwarfs (Schneider et al. 2014). Higher-resolution NIR spectroscopy as well as an RV measurement derived from a high signal-to-noise spectrum would be needed to completely rule out low gravity, but it is very likely that this object is a regular BD; we thus reject it as a candidate member of ARG.

2MASS J20484222–5127435 was identified as a candidate member of THA as part of BASS. NIR spectroscopy revealed that its J and H bands are similar to a field L2 dwarf, but its K band is significantly different, and similar to the K band of field L5 dwarfs (Figure 24). This object is thus unlikely young and we reject it as a candidate member of THA; however, it is unclear what is the cause of its peculiar K band. It would be worthwhile investigating whether this is an early-L/mid-L binary from high-resolution imaging or an RV follow-up.

2MASS J22062157–6116284 has been identified as a candidate member of THA in PRE-BASS. We obtained NIR spectroscopy and categorized it as a peculiar L0±1 dwarf because its *H*-band flux at ~1.57 μ m is stronger than usual (Figure 25). This could indicate the presence of an unresolved T-type component; however, the index-based scheme of Bardalez Gagliuffi et al. (2014) indicates that this scenario is unlikely.

2MASS J23155665–4747315 has been identified as a candidate member of THA in BASS. We obtained NIR spectroscopy and categorized it as a peculiar L3 dwarf; its J band is similar to an L3 dwarf albeit with stronger FeH absorption, and its H and K bands are similar to our field L5 template (Figure 26).

2MASS J23310161–0406193 (Koenigstuhl 3 BC) was discovered by Gizis et al. (2000) as an M9 dwarf in the optical, and Gizis et al. (2003) demonstrated that it is an M8 + L3, 0["].58 binary system. Caballero (2007) discovered that this



Figure 24. NIR spectrum of the peculiar L2 dwarf 2MASS J20484222–5127435 that was recovered as a candidate member of THA in BASS. It lacks indications of a low surface gravity and is thus not a likely member of THA. Its K band is peculiar, as it resembles those of later-type L dwarfs.



Figure 25. NIR spectrum of the peculiar L2 dwarf 2MASS J22062157–6116284 that was recovered as a candidate member of THA in PRE-BASS. It lacks indications of a low surface gravity and is thus not a likely member of THA. Furthermore, its *H*-band bump at \sim 1.57 μ m is stronger than usual.

system is a very wide 451".8 comoving system to the F8 star HR 8931 (HD 221356). Koenigstuhl 3 BC was identified as a candidate member of ABDMG in LP-BASS, but measurements of RV and distance for the co-moving star HR 8931 (-12.86 \pm 0.09 km s⁻¹ and 26.12 \pm 0.37 pc; Nidever et al. 2002; van Leeuwen 2007) preclude a possible membership to all YMGs considered here. We categorize its unresolved spectrum as a peculiar M8 dwarf; our best-matching NIR template is M8 γ , however it presents several differences with it and lacks several low-gravity indications such as weaker alkali lines. We thus conclude that the peculiar nature of this spectrum is likely related to its binary nature, which further rules out a possible membership to ABDMG.

2MASS J23392527+3507165 has been discovered as an L3.5 BD in the optical by Reid et al. (2008), and Burgasser



Figure 26. NIR spectrum of the peculiar L3 dwarf 2MASS J23155665–4747315 that was recovered as a candidate member of THA in BASS. It lacks indications of a low surface gravity and is thus not a likely member of THA. Its J band is similar to a field L3 dwarf with stronger FeH absorption, and its H and K bands are consistent with a later spectral type.

et al. (2010) categorized it as an L4.5 BD in the NIR. We recovered this object as a candidate member of β PMG in BASS. We used its NIR spectral type to categorize it as a peculiar L4 BD that has a stronger *H*-band peak at ~1.57 μ m (Figure 27). However, it is unlikely that this object is young as it lacks the usual low-gravity indications. We thus reject it as a candidate member of β PMG. The peculiar *H*-band feature described above can be an effect of an unresolved T-type companion; however, the index-based scheme of Bardalez Gagliuffi et al. (2014) indicates that this scenario is unlikely.

REFERENCES

- Ackerman, A. S., & Marley, M. S. 2001, ApJ, 556, 872
- Allard, F., Homeier, D., Freytag, B., Schaffenberger, A., & Rajpurohit, A. S. 2013, MmSAI, 24, 128
- Allers, K. N., Jaffe, D. T., Luhman, K. L., et al. 2007, ApJ, 657, 511
- Allers, K. N., & Liu, M. C. 2013, ApJ, 772, 79
- Andrei, A. H., Smart, R. L., Penna, J. L., et al. 2011, AJ, 141, 54
- Artigau, É, Gagné, J., Faherty, J. K., et al. 2015, ApJ, 806, 254
- Artigau, É, Radigan, J., Folkes, S., et al. 2010, ApJL, 718, L38
- Bakos, G. Á, Hartman, J., Torres, G., et al. 2011, ApJ, 742, 116
- Bannister, N. P., & Jameson, R. F. 2007, MNRAS, 378, L24
- Baraffe, I., Chabrier, G., Barman, T. S., Allard, F., & Hauschildt, P. H. 2003, A&A, 402, 701
- Bardalez Gagliuffi, D. C., Burgasser, A. J., Gelino, C. R., et al. 2014, ApJ, 794, 143
- Barenfeld, S. A., Bubar, E. J., Mamajek, E. E., & Young, P. A. 2013, ApJ, 766, 6
- Barman, T. S., Macintosh, B., Konopacky, Q. M., & Marois, C. 2011a, ApJ, 733, 65
- Barman, T. S., Macintosh, B., Konopacky, Q. M., & Marois, C. 2011b, ApJL, 735, L39
- Barrado Y Navascués, D. 1998, A&A, 339, 831
- Basri, G. 2000, ARA&A, 38, 485
- Beamín, J. C., Ivanov, V. D., Bayo, A., et al. 2014, A&A, 570, L8
- Beichman, C., Gelino, C. R., Kirkpatrick, J. D., et al. 2014, ApJ, 783, 68
- Biller, B. A., Liu, M. C., Wahhaj, Z., et al. 2013, ApJ, 777, 160
- Binks, A. S., & Jeffries, R. D. 2014, MNRAS, 438, L11
- Blake, C. H., Charbonneau, D., & White, R. J. 2010, ApJ, 723, 684
- Blecic, J., Harrington, J., Madhusudhan, N., et al. 2013, ApJ, 779, 5
- Bochanski, J. J., Gizis, J. E., Hawley, S. L., et al. 2005, AJ, 130, 1871
- Bochanski, J. J., Hennawi, J. F., Simcoe, R. A., et al. 2009, PASP, 121, 1409
- Boeshaar, P. C. 1976, PhD thesis, Ohio State Univ.
- Bonnefoy, M., Boccaletti, A., Lagrange, A.-M., et al. 2013, A&A, 555, A107



Figure 27. NIR spectrum of the peculiar L4 dwarf 2MASS J23392527 +3507165 that was recovered as a candidate member of β PMG in BASS. It lacks indications of a low surface gravity and is thus not a likely member of β PMG. Furthermore, its *H*-band bump at ~1.57 μ m is stronger than usual.

- Bonnefoy, M., Marleau, G. D., Galicher, R., et al. 2014, A&A, 567, L9
- Bouchy, F., Bonomo, A. S., Santerne, A., et al. 2011a, A&A, 533, A83
- Bouchy, F., Deleuil, M., Guillot, T., et al. 2011b, A&A, 525, A68
- Bouy, H., Duchêne, G., Köhler, R., et al. 2004, A&A, 423, 341
- Bouy, H., Martín, E. L., Brandner, W., et al. 2008, A&A, 481, 757
- Bowler, B. P., Liu, M. C., Kraus, A. L., & Mann, A. W. 2014, ApJ, 784, 65
- Bowler, B. P., Liu, M. C., Shkolnik, E. L., & Dupuy, T. J. 2013, ApJ, 774, 55
- Buchhave, L. A., Bakos, G. Á, Hartman, J. D., et al. 2010, ApJ, 720, 1118
- Buchhave, L. A., Latham, D. W., Carter, J. A., et al. 2011, ApJS, 197, 3
- Burgasser, A. J. 2014, in ASI Conf. Ser. Vol. 11, Int. Workshop on Stellar Spectral Libraries, ed. H. P. Singh, P. Prugniel & I. Vauglin (Pune: India), 7 Burgasser, A. J., Cruz, K. L., Cushing, M., et al. 2010, ApJ, 710, 1142
- Burgasser, A. J., Geballe, T. R., Leggett, S. K., Kirkpatrick, J. D., & Golimowski, D. A. 2006, ApJ, 637, 1067
- Burgasser, A. J., Gelino, C. R., Cushing, M. C., & Kirkpatrick, D. J. 2012, ApJ, 745, 26
- Burgasser, A. J., Gizis, J. E., & Bardalez-Gagliuffi, D. C. 2011, AJ, 141, 70 Burgasser, A. J., Liu, M. C., Ireland, M. J., Cruz, K. L., & Dupuy, T. J. 2008,
- ApJ, 681, 579 Burningham, B., Cardoso, C. V., Smith, L., et al. 2013, MNRAS, 433, 457
- Burningham, B., Pinfield, D. J., Leggett, S. K., et al. 2008, MNRAS, 391, 320
- Caballero, J. A. 2007, ApJ, 667, 520
- Canty, J. I., Lucas, P. W., Roche, P. F., & Pinfield, D. J. 2013, MNRAS, 435, 2650
- Cappetta, M., Saglia, R. P., Birkby, J. L., et al. 2012, MNRAS, 427, 1877
- Carson, J., Thalmann, C., Janson, M., et al. 2013, ApJL, 763, L32
- Castro, P. J., Gizis, J. E., Harris, H. C., et al. 2013, ApJ, 776, 126
- Chabrier, G. 2005, in Astrophysics and Space Science Library Volume 327, The Initial Mass Function 50 Years Later, ed. E. Corbelli & F. Palle (Berlin: Springer), 41
- Chabrier, G., Baraffe, I., Allard, F., & Hauschildt, P. 2000, ApJ, 542, 464
- Chauvin, G., Lagrange, A.-M., Dumas, C., et al. 2004, A&A, 425, L29
- Chauvin, G., Lagrange, A.-M., Zuckerman, B., et al. 2005, A&A, 438, L29
- Chiu, K., Fan, X., Leggett, S. K., et al. 2006, AJ, 131, 2722
- Costa, E., Méndez, R. A., Jao, W. C., et al. 2005, AJ, 130, 337
- Costa, E., Méndez, R. A., Jao, W. C., et al. 2006, AJ, 132, 1234
- Crifo, F., Phan Bao, N., Delfosse, X., et al. 2005, A&A, 441, 653
- Cruz, K. L., Kirkpatrick, J. D., & Burgasser, A. J. 2009, AJ, 137, 3345
- Cruz, K. L., & Núñez, A. 2012, in 17th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun (New York: Wiley)
- Cruz, K. L., Reid, I. N., Kirkpatrick, J. D., et al. 2007, AJ, 133, 439
- Cruz, K. L., Reid, I. N., Liebert, J., Kirkpatrick, J. D., & Lowrance, P. J. 2003, AJ, 126, 2421
- Currie, T., Daemgen, S., Debes, J., et al. 2014, ApJL, 780, L30
- Cushing, M. C., Kirkpatrick, J. D., Gelino, C. R., et al. 2011, ApJ, 743, 50
- Cushing, M. C., Marley, M. S., Saumon, D., et al. 2008, ApJ, 678, 1372
- Cushing, M. C., Vacca, W. D., & Rayner, J. T. 2004, PASP, 116, 362
- Dahn, C. C., Harris, H. C., Vrba, F. J., et al. 2002, AJ, 124, 1170

- de la Reza, R., Torres, C. A. O., Quast, G., Castilho, B. V., & Vieira, G. L. 1989, ApJL, 343, L61
- Deacon, N. R., Liu, M. C., Magnier, E. A., et al. 2014, ApJ, 792, 119
- Deleuil, M., Bonomo, A. S., Ferraz-Mello, S., et al. 2012, A&A, 538, A145
- Deleuil, M., Deeg, H. J., Alonso, R., et al. 2008, A&A, 491, 889
- Delorme, P., Delfosse, X., Albert, L., et al. 2008a, A&A, 482, 961
- Delorme, P., Gagné, J., Girard, J. H., et al. 2013, A&A, 553, L5
- Delorme, P., Gagné, J., Malo, L., et al. 2012, A&A, 548, A26
- Delorme, P., Willott, C. J., Forveille, T., et al. 2008b, A&A, 484, 469
- Deshpande, R., Martín, E. L., Montgomery, M. M., et al. 2012, AJ, 144, 99
- Díaz, R. F., Damiani, C., Deleuil, M., et al. 2013, A&A, 551, L9
- Díaz, R. F., Montagnier, G., Leconte, J., et al. 2014, A&A, 572, A109 Dieterich, S. B., Henry, T. J., Jao, W.-C., et al. 2014, AJ, 147, 94
- Ducourant, C., Teixeira, R., Galli, P. A. B., et al. 2014, A&A, 563, A121
- Dupuy, T. J., & Liu, M. C. 2012, ApJS, 201, 19
- Dupuy, T. J., Liu, M. C., & Bowler, B. P. 2009a, ApJ, 706, 328
- Dupuy, T. J., Liu, M. C., Bowler, B. P., et al. 2010, ApJ, 721, 1725
- Dupuy, T. J., Liu, M. C., & Ireland, M. J. 2009b, ApJ, 692, 729
- Dupuy, T. J., Liu, M. C., & Ireland, M. J. 2009c, ApJ, 699, 168
- Dupuy, T. J., Liu, M. C., & Ireland, M. J. 2014, ApJ, 790, 133
- Dupuy, T. J., Liu, M. C., Leggett, S. K., et al. 2015, ApJ, 805, 56
- Eikenberry, S. S., Elston, R., Raines, S. N., et al. 2004, Proc. SPIE, 5492, 1196
- Eisenbeiss, T., Ammler-von Eiff, M., Roell, T., et al. 2013, A&A, 556, A53
- Elias, J. H. 1978, ApJ, 224, 857
- Faherty, J. K., Burgasser, A. J., Bochanski, J. J., et al. 2011, AJ, 141, 71
- Faherty, J. K., Burgasser, A. J., Cruz, K. L., et al. 2009, AJ, 137, 1
- Faherty, J. K., Burgasser, A. J., Walter, F. M., et al. 2012, ApJ, 752, 56
- Faherty, J. K., Rice, E. L., Cruz, K. L., Mamajek, E. E., & Núñez, A. 2013, AJ, 145.2
- Famaey, B., Jorissen, A., Luri, X., et al. 2005, A&A, 430, 165
- Filippazzo, J. C., et al. 2015, ApJ, submittded
- Fitzpatrick, E. L. 1999, PASP, 111, 63
- Gagné, J., Faherty, J. K., Cruz, K. L., et al. 2014a, The ApJL, 785, L14
- Gagné, J., Lafrenière, D., Doyon, R., et al. 2014b, ApJL, 792, L17
- Gagné, J., Lafrenière, D., Doyon, R., Malo, L., & Artigau, É 2014c, ApJ, 783, 121
- Gagné, J., Lafrenière, D., Doyon, R., Malo, L., & Artigau, É 2015, ApJ, 798.73
- Gálvez-Ortiz, M. C., Clarke, J. R. A., Pinfield, D. J., et al. 2010, MNRAS, 409. 552
- Gardner, J. P., Mather, J. C., Clampin, M., et al. 2006, SSRv, 123, 485
- Gatley, I., Becklin, E. E., Mattews, K., et al. 1974, ApJL, 191, L121
- Geballe, T. R., Knapp, G. R., Leggett, S. K., et al. 2002, ApJ, 564, 466
- Geissler, K., Metchev, S., Kirkpatrick, J. D., Berriman, G. B., & Looper, D. 2011, ApJ, 732, 56
- Gizis, J. E. 2002, ApJ, 575, 484
- Gizis, J. E., Allers, K. N., Liu, M. C., et al. 2015, ApJ, 799, 203
- Gizis, J. E., Hawley, S. L., & Reid, N. I. 1997, AJ, 113, 1458
- Gizis, J. E., Monet, D. G., Reid, I. N., et al. 2000, AJ, 120, 1085
- Gizis, J. E., Reid, I. N., Knapp, G. R., et al. 2003, AJ, 125, 3302
- Goldman, B., Marsat, S., Henning, T., Clemens, C., & Greiner, J. 2010, MNRAS, 405, 1140
- Golimowski, D. A., Leggett, S. K., Marley, M. S., et al. 2004, AJ, 127, 3516
- Gorlova, N. I., Meyer, M. R., Rieke, G. H., & Liebert, J. 2003, ApJ, 593, 1074
- Goto, M., Kobayashi, N., Terada, H., et al. 2002, ApJL, 567, L59
- Hand, D. J., & Yu, K. 2001, Int. Stat. Rev., 69, 385
- Hawley, S. L., Covey, K. R., Knapp, G. R., et al. 2002, AJ, 123, 3409
- Hébrard, G., Almenara, J. M., Santerne, A., et al. 2013, A&A, 554, A114
- Henry, T. J., Jao, W.-C., Subasavage, J. P., et al. 2006, AJ, 132, 2360
- Herter, T. L., Henderson, C. P., Wilson, J. C., et al. 2008, Proc. SIPE, 7017, 70140X
- Horne, K. 1986, PASP, 98, 609
- Janson, M., Carson, J., Thalmann, C., et al. 2011, ApJ, 728, 85
- Jeffries, R. D. 2012, in EAS Publications Series, Vol. 57, Low-Mass Stars and the Transition Stars/Brown Dwarfs, Vol. 57, ed. C. Reylé, C. Charbonnel & M. Schultheis (Cambridge: Cambridge Univ. Press), 45
- Joyce, R. R., Hinkle, K. H., Wallace, L., Dulick, M., & Lambert, D. L. 1998, AJ. 116. 2520
- Kastner, J. H., Zuckerman, B., Weintraub, D. A., & Forveille, T. 1997, Sci, 277.67
- Kendall, T. R., Delfosse, X., Martín, E. L., & Forveille, T. 2004, A&A, 416. L17
- Kendall, T. R., Jones, H. R. A., Pinfield, D. J., et al. 2007a, MNRAS, 374, 445
- Kendall, T. R., Tamura, M., Tinney, C. G., et al. 2007b, A&A, 466, 1059
- Kirkpatrick, J. D. 2005, ARA&A, 43, 195
- Kirkpatrick, J. D., Barman, T. S., Burgasser, A. J., et al. 2006, ApJ, 639, 1120

- Kirkpatrick, J. D., Cruz, K. L., Barman, T. S., et al. 2008, ApJ, 689, 1295
- Kirkpatrick, J. D., Cushing, M. C., Gelino, C. R., et al. 2011, ApJS, 197, 19
- Kirkpatrick, J. D., Gelino, C. R., Cushing, M. C., et al. 2012, ApJ, 753, 156
- Kirkpatrick, J. D., Henry, T. J., & McCarthy, D. W. J. 1991, ApJS, 77, 417
- Kirkpatrick, J. D., Looper, D. L., Burgasser, A. J., et al. 2010, ApJS, 190, 100
- Kirkpatrick, J. D., Reid, I. N., Liebert, J., et al. 1999, ApJ, 519, 802
- Kirkpatrick, J. D., Reid, I. N., Liebert, J., et al. 2000, AJ, 120, 447
- Kirkpatrick, J. D., Schneider, A., Fajardo-Acosta, S., et al. 2014, ApJ, 783, 122
- Kiss, L. L., Moór, A., Szalai, T., et al. 2011, MNRAS, 411, 117
- Kleinmann, S. G., & Hall, D. N. B. 1986, ApJS, 62, 501
- Knapp, G. R., Leggett, S. K., Fan, X., et al. 2004, AJ, 127, 3553
- Konopacky, Q. M., Ghez, A. M., Barman, T. S., et al. 2010, ApJ, 711, 1087
- Kraus, A. L., Ireland, M. J., Cieza, L. A., et al. 2014a, ApJ, 781, 20
- Kraus, A. L., Shkolnik, E. L., Allers, K. N., & Liu, M. C. 2014b, AJ, 147, 146 Kuzuhara, M., Tamura, M., Kudo, T., et al. 2013, ApJ, 774, 11
- Lagrange, A.-M., Bonnefoy, M., Chauvin, G., et al. 2010, Sci, 329, 57 Lane, B. F., Zapatero Osorio, M. R., Britton, M. C., Martín, E. L., &
- Kulkarni, S. R. 2001, ApJ, 560, 390
- Leggett, S. K., Allard, F., Geballe, T. R., Hauschildt, P. H., & Schweitzer, A. 2001, ApJ, 548, 908
- Leggett, S. K., Currie, M. J., Varricatt, W. P., et al. 2006, MNRAS, 373, 781
- Leggett, S. K., Cushing, M. C., Saumon, D., et al. 2009, ApJ, 695, 1517
- Leggett, S. K., Geballe, T. R., Fan, X., et al. 2000, ApJL, 536, L35
- Leggett, S. K., Golimowski, D. A., Fan, X., et al. 2002, ApJ, 564, 452
- Leggett, S. K., Morley, C. V., Marley, M. S., et al. 2013, ApJ, 763, 130
- Leggett, S. K., Morley, C. V., Marley, M. S., & Saumon, D. 2015, ApJ, 799, 37
- Leggett, S. K., Saumon, D., Burningham, B., et al. 2010, ApJ, 720, 252
- Lépine, S., & Bongiorno, B. 2007, AJ, 133, 889
- Liebert, J., & Gizis, J. E. 2006, PASP, 118, 659

326, 695

703. 399

53

Luhman, K. L. 2007, ApJS, 173, 104

Luhman, K. L. 2014, ApJL, 786, L18

Mace, G. N. 2014, PhD thesis, Univ. California

Mainzer, A., Grav, T., Bauer, J., et al. 2011, ApJ, 743, 156 Makarov, V. V., & Urban, S. 2000, MNRAS, 317, 289

Malo, L., Artigau, É, Doyon, R., et al. 2014a, ApJ, 788, 81 Malo, L., Doyon, R., Feiden, G. A., et al. 2014b, ApJ, 792, 37

Malo, L., Doyon, R., Lafrenière, D., et al. 2013, ApJ, 762, 88 Mamajek, E. E., & Bell, C. P. M. 2014, MNRAS, 445, 2169

Marleau, G. D., & Cumming, A. 2014, MNRAS, 437, 1378

Metchev, S. A., & Hillenbrand, L. A. 2006, ApJ, 651, 1166

Monet, D. G., Dahn, C. C., Vrba, F. J., et al. 1992, AJ, 103, 638

Marley, M. S., Seager, S., Saumon, D., et al. 2002, ApJ, 568, 335

Marocco, F., Andrei, A. H., Smart, R. L., et al. 2013, AJ, 146, 161 Marocco, F., Day-Jones, A. C., Lucas, P. W., et al. 2014, MNRAS, 439, 372

Marocco, F., Smart, R. L., Jones, H. R. A., et al. 2010, A&A, 524, A38 Marois, C., Macintosh, B., Barman, T., et al. 2008, Sci, 322, 1348

Marsh, K. A., Wright, E. L., Kirkpatrick, J. D., et al. 2013, ApJ, 762, 119 Martín, E. L., Phan Bao, N., Bessell, M., et al. 2010, A&A, 517, A53

Martín, E. L., Rebolo, R., & Zapatero Osorio, M. R. 1996, ApJ, 469, 706

McGovern, M. R., Kirkpatrick, D. J., McLean, I. S., et al. 2004, ApJ, 600, 1020

- Littlefair, S. P., Casewell, S. L., Parsons, S. G., et al. 2014, MNRAS, 445, 2106
- Liu, M. C., Dupuy, T. J., & Allers, K. N. 2013a, AN, 334, 85
- Liu, M. C., Dupuy, T. J., & Leggett, S. K. 2010, ApJ, 722, 311
- Liu, M. C., Magnier, E. A., Deacon, N. R., et al. 2013b, ApJL, 777, L20
- Lodieu, N., Hambly, N. C., Jameson, R. F., & Hodgkin, S. T. 2008, MNRAS, 383, 1385
- Lodieu, N., Pinfield, D. J., Leggett, S. K., et al. 2007, MNRAS, 379, 1423
- Lodieu, N., Scholz, R. D., McCaughrean, M. J., et al. 2005, A&A, 440, 1061
- Looper, D. L., Burgasser, A. J., Kirkpatrick, J. D., & Swift, B. J. 2007a, ApJL, 669, L97
- Looper, D. L., Gelino, C. R., Burgasser, A. J., & Kirkpatrick, J. D. 2008a, ApJ, 685, 1183 Looper, D. L., Kirkpatrick, J. D., & Burgasser, A. J. 2007b, AJ, 134, 1162

Looper, D. L., Kirkpatrick, J. D., Cutri, R. M., et al. 2008b, ApJ, 686, 528

Lucas, P. W., Roche, P. F., Allard, F., & Hauschildt, P. H. 2001, MNRAS,

Lucas, P. W., Tinney, C. G., Burningham, B., et al. 2010, MNRAS, 408, L56

Luhman, K. L., Mamajek, E. E., Allen, P. R., & Cruz, K. L. 2009, ApJ,

Luhman, K. L., Liebert, J., & Rieke, G. H. 1997, ApJL, 489, L165

Luhman, K. L., Patten, B. M., Marengo, M., et al. 2007, ApJ, 654, 570

Luhman, K. L., Stauffer, J. R., & Mamajek, E. E. 2005, ApJL, 628, L69

Mace, G. N., Kirkpatrick, J. D., Cushing, M. C., et al. 2013a, ApJS, 205, 6

Mace, G. N., Kirkpatrick, J. D., Cushing, M. C., et al. 2013b, ApJ, 777, 36

Manjavacas, E., Bonnefoy, M., Schlieder, J. E., et al. 2014, A&A, 564, A55

Males, J. R., Close, L. M., Morzinski, K. M., et al. 2014, ApJ, 786, 32

- Montet, B. T., Johnson, J. A., Muirhead, P. S., et al. 2014, BASS, 223, #334.03
- Moutou, C., Bonomo, A. S., Bruno, G., et al. 2013, A&A, 558, L6
- Murphy, S. J., & Lawson, W. A. 2015, MNRAS, 447, 1267
- Naud, M.-È, Artigau, É, Malo, L., et al. 2014, ApJ, 787, 5
- Nidever, D. L., Marcy, G. W., Butler, R. P., Fischer, D. A., & Vogt, S. S. 2002, ApJS, 141, 503
- Norlén, G. 1973, PhyS, 8, 249
- Ochsenbein, F., Bauer, P., & Marcout, J. 2000, A&AS, 143, 23
- Parviainen, H., Gandolfi, D., Deleuil, M., et al. 2014, A&A, 562, A140
- Pecaut, M. J., & Mamajek, E. E. 2013, ApJS, 208, 9
- Pecaut, M. J., Mamajek, E. E., & Bubar, E. J. 2012, ApJ, 746, 154
- Pérez-Garrido, A., Lodieu, N., Béjar, V. J. S., et al. 2014, A&A, 567, A6
- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., et al. 1997, A&A, 323, L49
- Phan-Bao, N., & Bessell, M. S. 2006, A&A, 446, 515
- Phan Bao, N., Bessell, M. S., Martín, E. L., et al. 2008, MNRAS, 383, 831
- Phan-Bao, N., Guibert, J., Crifo, F., et al. 2001, A&A, 380, 590
- Pinfield, D. J., Burningham, B., Tamura, M., et al. 2008, MNRAS, 390, 304
- Quinn, S. N., White, T. R., Latham, D. W., et al. 2014, ApJ, 787, 27
- Rayner, J. T., Toomey, D. W., Onaka, P. M., et al. 2003, PASP, 115, 362
- Reid, N. I., Cruz, K. K., & Allen, P. R. 2007, AJ, 133, 2825
- Reid, I. N., Cruz, K. L., Kirkpatrick, J. D., et al. 2008, AJ, 136, 1290
- Reid, N. I., Cruz, K. K., Lowrance, P., et al. 2003, AJ, 126, 3007
- Reid, N. I., Gizis, J. E., Kirkpatrick, D. J., & Koerner, D. W. 2001, AJ, 121, 489
- Reid, N. I., Gizis, J. E., Kirkpatrick, D. J., et al. 2002, AJ, 124, 519
- Reiners, A. 2009, ApJL, 702, L119
- Reiners, A., & Basri, G. 2009, ApJ, 705, 1416
- Reiners, A., & Basri, G. 2010, ApJ, 710, 924
- Riaz, B., Gizis, J. E., & Harvin, J. 2006, AJ, 132, 866
- Rice, E. L., Faherty, J. K., & Cruz, K. L. 2010, ApJL, 715, L165
- Riedel, A. R., Finch, C. T., Henry, T. J., et al. 2014, AJ, 147, 85
- Robin, A. C., Marshall, D. J., Schultheis, M., & Reylé, C. 2012, A&A, 538, 106
- Rodriguez, D. R., Bessell, M. S., Zuckerman, B., & Kastner, J. H. 2011, ApJ, 727, 62
- Rodriguez, D. R., Zuckerman, B., Kastner, J. H., et al. 2013, ApJ, 774, 101
- Sahu, K. C., Casertano, S., Bond, H. E., et al. 2006, Natur, 443, 534
- Sartori, M. J., Lépine, J. R. D., & Dias, W. S. 2003, A&A, 404, 913
- Schlafly, E. F., Green, G., Finkbeiner, D. P., et al. 2014, ApJ, 786, 29
- Schlieder, J. E., Lépine, S., & Simon, M. 2012a, AJ, 143, 80
- Schlieder, J. E., Lépine, S., & Simon, M. 2012b, AJ, 144, 109
- Schmidt, S. J., Cruz, K. K., Bongiorno, B. J., Liebert, J., & Reid, N. I. 2007, AJ, 133, 2258
- Schmidt, S. J., Hawley, S. L., West, A. A., et al. 2015, AJ, 149, 158
- Schmidt, S. J., West, A. A., Hawley, S. L., & Pineda, J. S. 2010, AJ, 139, 1808
- Schneider, A. C., Cushing, M. C., Kirkpatrick, J. D., et al. 2014, AJ, 147, 34
- Schneider, A., Melis, C., & Song, I. 2012, ApJ, 754, 39
- Scholz, R. D. 2014, A&A, 561, A113
- Seifahrt, A., Reiners, A., Almaghrbi, K. A. M., & Basri, G. 2010, A&A, 512, 37
- Shkolnik, E. L., Anglada-Escudé, G., Liu, M. C., et al. 2012, ApJ, 758, 56
- Shkolnik, E. L., Liu, M. C., Reid, I. N., Dupuy, T., & Weinberger, A. J. 2011, ApJ, 727, 6
- Shkolnik, E. L., Liu, M. C., & Reid, N. I. 2009, ApJ, 699, 649
- Shporer, A., O'Rourke, J. G., Knutson, H. A., et al. 2014, ApJ, 788, 92
- Simcoe, R. A., Burgasser, A. J., Bernstein, R. A., et al. 2008, Proc. SPIE, 7014,
- 70140U
- Simcoe, R. A., Burgasser, A. J., Schechter, P. L., et al. 2013, PASP, 125, 270

- Simons, D. A., & Tokunaga, A. 2002, PASP, 114, 169
- Siverd, R. J., Beatty, T. G., Pepper, J., et al. 2012, ApJ, 761, 123
- Skemer, A. J., Hinz, P. M., Esposito, S., et al. 2012, ApJ, 753, 14
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163
- Slesnick, C. L., Hillenbrand, L. A., & Carpenter, J. M. 2004, ApJ, 610, 1045 Song, I., Zuckerman, B., & Bessell, M. S. 2003, ApJ, 599, 342
- Southworth, J. 2010, MNRAS, 408, 1689
- Stassun, K. G., Kratter, K. M., Scholz, A., & Dupuy, T. J. 2012, ApJ, 756, 47
- Stassun, K. G., Mathieu, R. D., & Valenti, J. A. 2006, Natur, 440, 311
- Stephens, D. C., & Leggett, S. K. 2004, PASP, 116, 9
- Stephens, D. C., Leggett, S. K., Cushing, M. C., et al. 2009, ApJ, 702, 154
- Strauss, M. A., Fan, X., Gunn, J. E., et al. 1999, ApJL, 522, L61
- Sumi, T., Kamiya, K., Bennett, D. P., et al. 2011, Natur, 473, 349
- Teegarden, B. J., Pravdo, S. H., Hicks, M., et al. 2003, ApJL, 589, L51
- Thalmann, C., Carson, J., Janson, M., et al. 2009, ApJL, 707, L123
- Thompson, M. A., Kirkpatrick, J. D., Mace, G. N., et al. 2013, PASP, 125, 809
- Tingley, B., Endl, M., Gazzano, J. C., et al. 2011, A&A, 528, A97
- Tinney, C. G., Burgasser, A. J., & Kirkpatrick, J. D. 2003, AJ, 126, 975
- Tinney, C. G., Faherty, J. K., Kirkpatrick, J. D., et al. 2014, ApJ, 796, 39
- Torres, C. A. O., da Silva, L., Quast, G. R., de La Reza, R., & Jilinski, E. 2000, AJ, 120, 1410
- Torres, R. M., Loinard, L., Mioduszewski, A. J., & Rodríguez, L. F. 2007, ApJ, 671, 1813
- Torres, C. A. O., Quast, G. R., Melo, C. H. F., & Sterzik, M. F. 2008, in ASP Monograph Publications, Young Nearby Loose Associations, Vol. I, ed. B. Reipurth (San Francisco, CA: ASP), 757
- Triaud, A. H. M. J., Hebb, L., Anderson, D. R., et al. 2013, A&A, 549, A18
- Vacca, W. D., Cushing, M. C., & Rayner, J. T. 2003, PASP, 115, 389
- van Altena, W. F., Lee, J. T., & Hoffleit, E. D. 1995, The General Catalogue of Trigonometric [Stellar] Parallaxes (4th ed.; New Haven, CT: Yale University Observatory)
- van Leeuwen, F. 2007, A&A, 474, 653
- Voges, W., Aschenbach, B., Boller, T., et al. 1999, A&A, 349, 389
- Voges, W., Aschenbach, B., Boller, T., et al. 2000, IAUC, 7432, 1
- Vrba, F. J., Henden, A. A., Luginbuhl, C. B., et al. 2004, AJ, 127, 2948
- Wahhaj, Z., Liu, M. C., Biller, B. A., et al. 2011, ApJ, 729, 139
- Warren, S. J., Mortlock, D. J., Leggett, S. K., et al. 2007, MNRAS, 381, 1400
- Weinberger, A. J., Anglada-Escudé, G., & Boss, A. P. 2013, ApJ, 762, 118
- West, A. A., Hawley, S. L., Bochanski, J. J., et al. 2008, AJ, 135, 785
- Wilson, J. C., Miller, N. A., Gizis, J. E., et al. 2003, in Proc. of IAU Symposium #211, Brown Dwarfs, ed. E. Martín (San Francisco, CA: ASP), 197
- Winn, J. N., Holman, M. J., Torres, G., et al. 2008, ApJ, 683, 1076
- Witte, S., Helling, C., Barman, T., Heidrich, N., & Hauschildt, P. H. 2011, A&A, 529, A44
- Wolberg, J. 2006, Data Analysis Using the Method of Least Squares Extracting the Most Information from Experiments (Berlin: Springer)
- Wood, D. O. S., Myers, P. C., & Daugherty, D. A. 1994, ApJS, 95, 457
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, AJ, 140, 1868
 - York, D. G., Adelman, J., Anderson, J. E. J., et al. 2000, AJ, 120, 1579
- Zapatero Osorio, M. R., Béjar, V. J. S., Miles-Páez, P. A., et al. 2014, A&A, 568, A6
- Zuckerman, B., Bessell, M. S., Song, I., & Kim, S. 2006, ApJL, 649, L115
- Zuckerman, B., & Song, I. 2004, ARA&A, 42, 685
- Zuckerman, B., Song, I., & Bessell, M. S. 2004, ApJL, 613, L65
- Zuckerman, B., Song, I., Bessell, M. S., & Webb, R. A. 2001a, ApJL, 562, L87
- Zuckerman, B., Song, I., & Webb, R. A. 2001b, ApJ, 559, 388
- Zuckerman, B., & Webb, R. A. 2000, ApJ, 535, 959