CORRIGENDUM

Kinetics of highly excited states in Ar¹⁷⁺ charge exchange recombination fusion plasma spectroscopy

To cite this article: O Marchuk et al 2009 J. Phys. B: At. Mol. Opt. Phys. 42 189801

View the article online for updates and enhancements.

You may also like

- Effects of breathing and oblong mode phonons on transport properties in a single-electron transistor Norihiko Nishiguchi and Martin N Wybourne
- Principles of Plasma Spectroscopy A.L. Osterheld
- High piezoelectric properties and domain configuration in BaTiO₃ ceramics obtained through solid-state reaction route Shoufu Shao, Jialiang Zhang, Zhong Zhang et al.

J. Phys. B: At. Mol. Opt. Phys. 42 (2009) 189801 (1pp)

Corrigendum

Kinetics of highly excited states in Ar¹⁷⁺ charge exchange recombination fusion plasma spectroscopy O Marchuk, Yu Ralchenko, R K Janev, G Bertschinger and W Biel *J. Phys. B: At. Mol. Opt. Phys.* **42** 165701

The equation on page 6 for the rate coefficient of charge exchange contains a misprint in the integration limits and should be replaced by

$$\alpha_{nl}^{\text{cx}} = \frac{1}{\sqrt{\pi}u_1u_2} \int_0^\infty u^2 \sigma_{nl}(u) \left[\exp\left(-\left(\frac{u-u_1}{u_2}\right)^2\right) - \exp\left(-\left(\frac{u+u_1}{u_2}\right)^2\right) \right] du$$

1

This error does not affect any results in the paper as the calculations were performed with the correct limits.