



Selection pressure transforms the nature of social dilemmas in adaptive networks

To cite this article: Sven Van Segbroeck *et al* 2011 *New J. Phys.* **13** 013007

View the [article online](#) for updates and enhancements.

Related content

- [How selection pressure changes the nature of social dilemmas in structured populations](#)
Flávio L Pinheiro, Francisco C Santos and Jorge M Pacheco
- [Large fluctuations in anti-coordination games on scale-free graphs](#)
Daniel Sabsovich, Mauro Mobilia and Michael Assaf
- [Evolutionary dynamics under interactive diversity](#)
Qi Su, Aming Li and Long Wang

Recent citations

- [Rationality alters the rank between peer punishment and social exclusion](#)
Xiukai Sui *et al*
- [Evolutionary advantages of adaptive rewarding](#)
Attila Szolnoki and Matjaž Perc
- [How selection pressure changes the nature of social dilemmas in structured populations](#)
Flávio L Pinheiro *et al*

Selection pressure transforms the nature of social dilemmas in adaptive networks

Sven Van Segbroeck^{1,2,7}, Francisco C Santos^{3,4},
Tom Lenaerts^{1,5} and Jorge M Pacheco^{4,6}

¹ MLG, Université Libre de Bruxelles, Boulevard du Triomphe—CP 212, 1050 Brussels, Belgium

² COMO, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium

³ CENTRIA, Departamento de Informática, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

⁴ ATP-Group, CMAF, Complexo Interdisciplinar, P-1649-003 Lisboa Codex, Portugal

⁵ Department of Computer Science, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium

⁶ Departamento de Matemática e Aplicações, Universidade do Minho, Campus de Gualtar, 4710-057 Braga, Portugal

E-mail: svsegbro@ulb.ac.be, fcsantos@fct.unl.pt, tlenaert@ulb.ac.be and jmpacheco@math.uminho.pt

New Journal of Physics **13** (2011) 013007 (13pp)

Received 8 September 2010

Published 11 January 2011

Online at <http://www.njp.org/>

doi:10.1088/1367-2630/13/1/013007

Abstract. We have studied the evolution of cooperation in structured populations whose topology coevolves with the game strategies of the individuals. Strategy evolution proceeds according to an update rule with a free parameter, which measures the selection pressure. We explore how this parameter affects the interplay between network dynamics and strategy dynamics. A dynamical network topology can influence the strategy dynamics in two ways: (i) by modifying the expected payoff associated with each strategy and (ii) by reshaping the imitation network that underlies the evolutionary process. We show here that the selection pressure tunes the relative contribution of each of these two forces to the final outcome of strategy evolution. The dynamics of the imitation network plays only a minor role under strong selection, but becomes the dominant force under weak selection. We demonstrate how these findings constitute a mechanism supporting cooperative behavior.

⁷ Author to whom any correspondence should be addressed.

Contents

1. Introduction	2
2. A minimal model	3
3. Results and discussion	5
4. Conclusion	9
Acknowledgments	9
Appendix A. The strong selection limit	10
Appendix B. General intensities of selection	11
References	11

1. Introduction

Many biological systems, and especially human societies, show persistent cooperative patterns. The evolution of such patterns is often studied using evolutionary game theory (EGT) [1]–[4]. This framework models interactions between the individuals of a population in terms of simple games. Successful behavior—success being measured in terms of game payoff—will be imitated and spreads in the population. The prototypical game to represent dilemmas of cooperation is the conventional prisoner’s dilemma [5]. In this game, a cooperative act goes at a cost c to the cooperator while conferring a benefit b to another individual (assuming $b > c$). Defectors receive the benefits without spending any costs and are therefore expected to have an evolutionary advantage over cooperators.

However, given the omnipresence of prosocial behavior, certain conditions exist under which cooperation becomes viable [4], [6]–[12]. It has, for instance, been recognized that the topology of the network along which individuals interact and reproduce/imitate affects drastically the evolutionary chances of cooperators [13]–[42] (see Szabó and Fáth for a review [43]). The individuals also shape and reshape their social environment themselves and are, at least partially, responsible for the specific features that characterize their social network [44]–[47]. The process of network reshaping is often coupled with the strategy dynamics: the behavior of an individual influences his social position and vice versa [48]. Networks exhibiting such feedback loops provide sophisticated examples of adaptive networks [49]. Several authors have studied the functioning of such networks in the context of cooperation [50]–[65] (see [66] for a recent review). In general, cooperation has been shown to emerge more easily when interactions that benefit both partners last longer than interactions where one partner is exploited by the other [55, 56, 60].

In structured populations, the outcome of the evolutionary process depends also on the update rule, which dictates how strategies evolve from one generation to the next [67]. The selection pressure enters this update rule, and in EGT this is no exception, also being called intensity of selection [68]. A maximal selection strength implies that individuals only imitate those with a higher game payoff. Reducing the selection strength increases the amount of noise in the imitation process, until, ultimately, evolution becomes a purely random process (neutral selection or random drift). The impact of the selection pressure on the evolutionary dynamics of a finite, well-mixed population is well understood by now [68]–[71]. Additional effects may, however, arise in adaptive networks, where the social environment of an individual changes

according to his behavior. Such a coevolutionary interplay between strategy evolution and graph evolution may give rise to correlations between the strategies of connected individuals. The strategy of an individual will therefore influence not only the payoff he acquires, but also the probability that he will be considered as a potential role model by someone else. When some strategies are more popular than others, the rate of strategy update may effectively change, depending on the intensity of selection.

We study this issue in the framework of active linking (AL) dynamics [55, 56, 62, 65, 72], a model that allows us to study cooperation in adaptive networks analytically in certain limits. Section 2 introduces both our model and the framework in which we will make our analysis. In section 3, we analyze the gradient of selection as a function of the intensity of selection, assuming that the network topology evolves much faster than the individual behavior. Section 4 presents the conclusions.

2. A minimal model

Consider a finite population of N individuals interacting in symmetric, one-shot, two-player games of cooperation defined by the payoff matrix

$$M = \begin{matrix} & \begin{matrix} C & D \end{matrix} \\ \begin{matrix} C \\ D \end{matrix} & \begin{pmatrix} R & S \\ T & P \end{pmatrix} \end{matrix}. \quad (1)$$

We distinguish two possible game strategies: cooperate unconditionally (C) and defect unconditionally (D). Payoff matrix 1 shows that individuals receive a reward R upon mutual cooperation and a punishment P upon mutual defection. When a C meets a D , the C receives the sucker's payoff S , whereas the D acquires the temptation to defect T .

Individuals do not interact with everyone in the population. Instead, a network indicates who meets whom. The structure of this network is dynamic, in the sense that edges appear and disappear over time. Simultaneously, individuals may reconsider their game play. The network dynamics proceed on a characteristic time scale τ_a , the strategy dynamics on another time scale τ_s . Below, we define each of these two dynamical processes separately.

We use the AL model, developed earlier by Pacheco *et al* [55, 56], to define the network's evolution. Each individual has a propensity α to engage in new interactions, such that new edges are formed at a rate α^2 . The lifetime of existing edges depends on the behavior of the individuals connected by this link. Specifically, the rate at which CC -links, CD -links and DD -links disappear is given by γ_{CC} , γ_{CD} and γ_{DD} , respectively.

Consider a network with k C s and $N - k$ D s. The number of CC -links, CD -links and DD -links in such a network can never exceed $N_{CC} \equiv \frac{1}{2}k(k - 1)$, $N_{CD} \equiv k(N - k)$ and $N_{DD} \equiv \frac{1}{2}(N - k)(N - k - 1)$, respectively. Under the assumption that the individuals stick to their game behavior, for a large number of links, we can describe the time evolution of the number of links of each type using the following ordinary differential equations,

$$\begin{aligned} \dot{L}_{CC}(t) &= \alpha^2 [N_{CC} - L_{CC}(t)] - \gamma_{CC} L_{CC}(t), \\ \dot{L}_{CD}(t) &= \alpha^2 [N_{CD} - L_{CD}(t)] - \gamma_{CD} L_{CD}(t), \\ \dot{L}_{DD}(t) &= \alpha^2 [N_{DD} - L_{DD}(t)] - \gamma_{DD} L_{DD}(t), \end{aligned} \quad (2)$$

where $L_{ij}(t)$ denotes the actual number of links at time t between individuals adopting strategy i and individuals adopting strategy j ($i, j \in \{C, D\}$). In the steady state, the number of links of each type is given by

$$\begin{aligned} L_{CC}^* &= \phi_{CC} N_{CC}, \\ L_{CD}^* &= \phi_{CD} N_{CD}, \\ L_{DD}^* &= \phi_{DD} N_{DD}, \end{aligned} \quad (3)$$

where $\phi_{ij} = \alpha^2(\alpha^2 + \gamma_{ij})^{-1}$ denotes the fraction of active ij -links ($i, j \in \{C, D\}$). Note that the stationary configuration of the network depends on the actual strategy configuration of the population, illustrating the interplay between network evolution and strategy evolution.

The second dynamical process in our model, the strategy dynamics in finite populations, is defined by the pairwise-comparison rule [18, 69]. At every strategy update event, a randomly selected individual X imitates a random neighbor Y with probability

$$p = [1 + e^{\beta/N(\Pi_X - \Pi_Y)}]^{-1}. \quad (4)$$

The individual Y can be regarded as the role model of X . Π_X (Π_Y) denotes the total payoff X (Y) receives after interacting once with every neighbor. The parameter β (≥ 0) controls the intensity of selection. When β is large, the imitation process is driven mainly by the payoff values that the individuals acquire. The game becomes progressively less important for decreasing β .

In the following, we assume that the network dynamics proceed much faster than the strategy dynamics ($\tau_a \ll \tau_s$). In this limit, the network always reaches a stationary configuration before a strategy update event occurs. The expected payoff of C s and D s during strategy update events is therefore given by

$$\begin{aligned} \Pi_C^* &= R \phi_{CC}(k-1) + S \phi_{CD}(N-k), \\ \Pi_D^* &= T \phi_{CD}k + P \phi_{DD}(N-k-1). \end{aligned} \quad (5)$$

These payoff values correspond to those obtained in a well-mixed population (complete network) with the same strategy configuration, but using the following rescaled payoff matrix [55, 56],

$$M' = \begin{array}{c} C \quad D \\ C \quad D \end{array} \begin{pmatrix} R' & S' \\ T' & P' \end{pmatrix} = \begin{array}{c} C \quad D \\ C \quad D \end{array} \begin{pmatrix} R\phi_{CC} & S\phi_{CD} \\ T\phi_{CD} & P\phi_{DD} \end{pmatrix}. \quad (6)$$

The network dynamics not only affect the payoffs individuals acquire but also influence the imitation process. Some individuals will act more often as a role model than others, depending on their strategy. In section 3, we show that such differences can have a profound effect on the strategy dynamics.

If $\tau_a \ll \tau_s$, a D will select a C as his role model with probability

$$\frac{\phi_{CD}k}{\phi_{CD}k + \phi_{DD}(N-k-1)}. \quad (7)$$

The term in the numerator corresponds to the average number of CD -links of each D , whereas the term in the denominator reflects the average total number of links of each D . The overall probability that the number of C s will increase during a strategy update event equals

$$T_k^+ = \frac{N-k}{N} \frac{\phi_{CD}k}{\phi_{CD}k + \phi_{DD}(N-k-1)} [1 + e^{\beta/N(\Pi_D^* - \Pi_C^*)}]^{-1}. \quad (8)$$

Similarly, the number of Cs will decrease with probability

$$T_k^- = \frac{k}{N} \frac{\phi_{CD}(N-k)}{\phi_{CD}(N-k) + \phi_{CC}(k-1)} [1 + e^{-\beta/N(\Pi_D^* - \Pi_C^*)}]^{-1}. \quad (9)$$

In well-mixed populations ($\phi_{CC} = \phi_{CD} = \phi_{DD} = 1$), the difference

$$g(k) = T_k^+ - T_k^- = \frac{k}{Z} \frac{Z-k}{Z} \tanh\left(\frac{\beta}{N}(\Pi_D^* - \Pi_C^*)\right) \quad (10)$$

($k \in]0, N[$) [69] can be regarded as a finite population analogue of the gradient of selection associated with the replicator equation in infinite, well-mixed populations [2], which is defined as $\dot{x} = x(1-x)(\Pi_C - \Pi_D)$, where $x \in [0, 1]$ stands for the fraction of cooperators. In both cases, Cs are favored over Ds when $g(k) > 0$ ($\dot{x} > 0$), whereas the opposite is true whenever $g(k) < 0$ ($\dot{x} < 0$).

It is noteworthy that we consider the evolutionary dynamics as a discrete stochastic system while assuming that the number of links is sufficiently large, so that the linking dynamics can be described by a set of ordinary differential equations (see equation (2)). Stochastic effects could be included at the level of the linking dynamics as well, for instance by adopting a discrete version of the AL model like the one proposed recently by Wu *et al* [65]. Nevertheless, the continuous approximation we use here does not weaken the robustness of our conclusions, as shown in the following section by means of computer simulations (see figure 2(d)).

3. Results and discussion

We investigate how the gradient of selection depends on the selection pressure. To do so, one can study the shape of $g(k)$ or, alternatively, that of the ratio $h(k) = T_k^+ / T_k^-$, which is given by

$$h(k) = \frac{\phi_{CD}(N-k) + \phi_{CC}(k-1)}{\phi_{CD}k + \phi_{DD}(N-k-1)} e^{-\beta/N(\Pi_D^* - \Pi_C^*)}. \quad (11)$$

As the solutions of $g(x) = 0$ are sometimes more conveniently obtained solving $h(x) = 1$, we use both interchangeably. For large N , $h(x)$ can be approximated by

$$h(x) \approx \frac{(1-a)x + a}{(a-b)x + b} e^{\beta(u'x + v')}, \quad (12)$$

where $x = \frac{k}{N}$, $a = \frac{\phi_{CD}}{\phi_{CC}}$, $b = \frac{\phi_{DD}}{\phi_{CC}}$, $u' = R' - S' - T' + P'$ and $v' = S' - P'$.

Let us start by studying the two limiting cases: the weak selection limit ($\beta \rightarrow 0$) and the strong selection limit ($\beta \rightarrow \infty$).

In the limit of strong selection, the direction of the gradient of selection depends solely—apart from finite size effects—on the sign of $u'x + v'$ (see appendix A). $g(x)$ will therefore exhibit one of four possible shapes (see figure 1), depending on the ordering of the payoff values of the transformed game M' . When $R' > T'$ and $S' > P'$, $g(x)$ is positive for all $x \in]0, 1[$. This scenario is known as *C-dominance*. Selection will always favor Cs over Ds, irrespective of the strategy configuration of the population. When $R' < T'$ and $S' < P'$, we obtain the opposite scenario: *D-dominance*. The gradient of selection now satisfies $g(x) < 0$ for all $x \in]0, 1[$, implying that Ds are always favored over Cs. When $R' > T'$ and $S' < P'$, $g(x)$ has a root in

$$x^* = \frac{P' - S'}{R' - T' - S' + P'}. \quad (13)$$




Figure 1. The four possible dynamical scenarios defined by two-person symmetric games in well-mixed populations. In each of the four panels, the curve represents a typical shape of the fitness difference $g(x)$ between Cs and Ds for a given fraction x of Cs. The roots of $g(x)$ are the fixed points of the evolutionary dynamics. Stable fixed points are depicted using solid circles and unstable fixed points using open circles. Arrows indicate the expected direction of evolution. Cs (Ds) are favored over Ds (Cs) when the arrow points to the right (left). The particular shape of $g(x)$ can be inferred from the sign of the derivative of g in 0 and in 1.

Furthermore, $g(x) < 0$ for $x < x^*$ and $g(x) > 0$ for $x > x^*$. When the initial fraction of Cs is smaller than x^* , evolution favors Cs. Otherwise, Ds will be favored. This is an example of coordination or bistability. Finally, when $R' < T'$ and $S' > P'$, evolution favors a mixture of Cs and Ds. The gradient $g(x)$ has a root in x^* again, but unlike before, selection favors Cs for $x < x^*$ and Ds for $x > x^*$. Note that one has to correct for self-interactions when computing the finite population analogue of the equilibrium point x^* . The resulting equilibrium k^* is given by

$$\frac{k^*}{N} = x^* + \frac{1}{N} \frac{P - R}{T - S - R + P}. \quad (14)$$

In the other limit, that of weak selection, we have

$$\lim_{\beta \rightarrow 0} h(x) = \frac{(1-a)x + a}{(a-b)x + b}. \quad (15)$$

Because this function is monotonic in x , $h(x) = 1$ can have at most one solution and consequently the gradient $g(x)$ can have at most one root in $]0, 1[$. Hence, $g(x)$ exhibits one of the four shapes shown in figure 1. The particular type of shape that occurs depends on the parameters a and b .

In the following, we assume that links between Cs satisfy both individuals and therefore last longer than other links. Two interacting Ds, on the other hand, are both dissatisfied and prefer to break their connection quickly. Following the same reasoning, Cs would like to break CD-links quickly, whereas Ds prefer to keep such links as long as possible. Hence, on

average, CD -links will have a longer time span than DD -links, but a shorter one than CC -links. Altogether, we obtain the following ordering for the fractions of active links of each type,

$$\phi_{CC} > \phi_{CD} > \phi_{DD}, \quad (16)$$

which is equivalent to

$$1 > a > b. \quad (17)$$

This condition ensures that $h(x) > 1$ for all $x \in]0, 1[$. Hence, C s are dominant in the limit of weak selection, irrespective of the game being played.

Having considered the two limiting cases, we now study the gradient of selection for general values of β . In appendix B, we show that $g(x)$ can have either zero, one or two internal equilibria, depending on the parameter settings. First we discuss examples where the specific intensity of selection leads to a scenario with at most one internal equilibrium. Next we will see that some parameter combinations lead to a scenario with two internal equilibria, which can never occur in the limits of either weak or strong selection.

We reduce the complexity of the game space by normalizing the difference between R and P to 1, taking $R = 2$ and $P = 1$. The contours in figure 2 depict the sign of $g(k)$ for three different D -dominance games, using $\alpha = 0.8$, $\gamma_{CC} = 0.4$, $\gamma_{CD} = 0.5$ and $\gamma_{DD} = 0.6$. In the weak selection limit, these birth/death rates of links lead to dominance of C s, irrespective of the original payoff matrix. The strong selection limit leads to different scenarios, depending on the original game. For instance, $S = 0.5$ and $T = 2.1$ results in bistability (see figure 2(a), $S = 0.9$ and $T = 2.5$ in coexistence of C s and D s (see figure 2(b)), and $S = 0.5$ and $T = 2.5$ in D -dominance (see figure 2(c)). Hence, there exist critical values of β at which the number of internal equilibria changes. We can compute these critical values using the derivative of $g(x)$ in 0 and in 1 (see also figure 1). These derivatives are given by

$$\dot{g}(0) = \frac{\phi_{CD}}{\phi_{DD}} [1 + e^{\beta(P'-S')}]^{-1} - [1 + e^{-\beta(P'-S')}]^{-1}, \quad (18)$$

$$\dot{g}(1) = \frac{\phi_{CD}}{\phi_{CC}} [1 + e^{\beta(R'-T')}]^{-1} - [1 + e^{-\beta(R'-T')}]^{-1}. \quad (19)$$

Assuming at most one internal equilibrium, we obtain C -dominance (D -dominance) when both $\dot{g}(0)$ and $\dot{g}(1)$ are larger (smaller) than zero. Bistability occurs when $\dot{g}(0) < 0$ and $\dot{g}(1) > 0$, and coexistence when $\dot{g}(0) > 0$ and $\dot{g}(1) < 0$. From equation (18), it follows that $\dot{g}(0) > 0$ if and only if

$$\ln\left(\frac{\phi_{CD}}{\phi_{DD}}\right) > \beta (P' - S'), \quad (20)$$

and that $\dot{g}(1) > 0$ if and only if

$$\ln\left(\frac{\phi_{CD}}{\phi_{CC}}\right) > \beta (R' - T'). \quad (21)$$

This means that $\dot{g}(0)$ will be positive for β smaller than

$$\beta_0 \equiv \ln\left(\frac{\phi_{CD}}{\phi_{DD}}\right) (P' - S')^{-1}, \quad (22)$$




Figure 2. The intensity of selection determines the shape of the gradient of selection. (a–c) Each of the contour plots shows the sign of the fitness difference $g(k)$, k being the number of Cs in a network of size N , as a function of β . Regions where $g(k) > 0$ ($g(k) < 0$) are indicated in gray (black). Solid white lines indicate where $g(k) = 0$. In each of the three panels, the individuals engage in a different D -dominance game: panel (a) shows $g(k)$ for $S = 0.5$ and $T = 2.1$, panel (b) for $S = 0.9$ and $T = 2.5$ and, finally, panel (c) for $S = 0.5$ and $T = 2.5$. The other two payoff values are normalized to $R = 2$ and $P = 1$. The network dynamics transform the game so that for $\beta \rightarrow \infty$, we obtain an unstable equilibrium k^* in panel (a), a stable equilibrium k^* in panel (b) and a D -dominance situation in panel (c). Reducing β eventually makes cooperation dominant, irrespective of the original game being played. The dotted vertical lines, indicated by β_0 and β_1 , show the analytical predictions for β at which the sign changes of $\dot{g}(0)$ and $\dot{g}(1)$, respectively. (d) Circles (lines) show the probability, obtained in simulation (analytically), to reach 100% cooperation, starting from a population with 50% C's. We use $\tau = 10^{-3}$ as the relative time scale for network dynamics. Individuals engage in the same game as in panel (c). In all four panels, we use $\gamma_{CC} = 0.4$, $\gamma_{CD} = 0.5$, $\gamma_{DD} = 0.6$, $\alpha = 0.8$ and $N = 100$.

and negative otherwise. Similarly, $\dot{g}(1)$ is positive for β smaller than

$$\beta_1 \equiv \ln\left(\frac{\phi_{CD}}{\phi_{CC}}\right) (R' - T')^{-1}, \quad (23)$$

and negative otherwise. Both β_0 and β_1 are indicated by vertical dotted lines in figure 2. They correspond clearly to the values of β at which each of the game transitions takes place.

We further confirm the validity of the analytical results by means of computer simulations. Figure 2(d) shows the probability, as a function of β , that a population with initially 50% cooperators evolves to full cooperation. Circles, representing simulation results, fit nicely with

the analytical predictions, which are indicated by a solid line. Each simulation starts with a complete network of size $N = 100$, 50% of them being Cs, and runs until the population fixates into either full cooperation or full defection. Strategy and network structure evolve simultaneously under asynchronous updating. Strategy update events take place with probability $(1 + \tau)^{-1}$, where $\tau = \frac{\tau_a}{\tau_s} = 10^{-3}$; network update events occur otherwise. We run 10^4 simulations for each value of β and plot the fraction of runs that end in full cooperation. The analytical predictions are calculated using the following formula [69],

$$\rho_i = \frac{1 + \sum_{k=1}^{i-1} \prod_{j=1}^k \frac{T_j^-}{T_j^+}}{1 + \sum_{k=1}^{N-1} \prod_{j=1}^k \frac{T_j^-}{T_j^+}}, \quad (24)$$

with i being equal to 50. The expressions for T_j^+ and T_j^- are given by equations (8) and (9).

Earlier in this section, we showed that there can be at most one internal equilibrium for the limiting cases $\beta \rightarrow 0$ and $\beta \rightarrow \infty$. At intermediate intensities of selection, two internal equilibria can, however, occur simultaneously (see also appendix B). Figure 3 shows two examples where this is the case. The upper two panels correspond to a scenario where Cs are expected to disappear from the population in case they are rare. If a sufficiently large fraction of Cs is present, however, a mixture of Cs and Ds is favored. The lower two panels illustrate an example of the opposite scenario, in which Ds are expected to go extinct when they are rare.

4. Conclusion

In this paper, we have explored, both analytically and numerically, the evolution of cooperation in dynamical networks that evolve side by side with individuals' behavior, the evolution of the network structure being affected by the dynamics of the individuals and vice versa. We indicate that the adaptive nature of the network of contacts affects both the average payoff associated with each game strategy and the likelihood for each of the different strategies to serve as a role model. The intensity of selection, which controls the contribution of game payoff to fitness, regulates the importance of each of these two effects in the final outcome of strategy evolution. When the intensity of selection is strong, the payoff transformation resulting from the network dynamics provides the dominating contribution to evolution. Weakening the intensity of selection enhances the effect of the adaptive imitation network. By doing so, one is able to effectively transform a D -dominance dilemma into any of the conventional 2×2 symmetric games. We derive analytical conditions, valid under the assumption that there is only one internal equilibrium, that predict which range of intensities of selection leads to which game scenario. We show also that certain parameter combinations may lead to the occurrence of two internal equilibria at intermediate intensities of selection. Our results clearly demonstrate the relevance of the intensity of selection when evolution proceeds on an adaptive structured population. Furthermore, we indicate the conditions under which this coupled dynamics work as an efficient mechanism for the promotion of cooperative behavior.

Acknowledgments

SVS and TL acknowledge financial support from F.R.S.-FNRS (Belgium). JMP and FCS acknowledge financial support from FCT (Portugal).




Figure 3. Two internal equilibria at intermediate intensities of selection. The contour plots show the sign of the fitness difference $g(k)$, like in figure 2. In (a), we use the parameters $R = 2$, $S = -9$, $T = 40$, $P = 1$, $\alpha = 0.2$, $\gamma_{CC} = 0.04$, $\gamma_{CD} = 0.46$, $\gamma_{DD} = 0.85$ and $N = 100$. Two internal equilibria take place at intermediate intensities of selection. The equilibrium closest to full defection is unstable and the other one is stable. This is emphasized in (b), where we zoom in on the actual gradient $g(k)$ for $\beta = 1.0$. Panels (c) and (d) show the same plots, but using $R = 2$, $S = -8$, $T = 3$, $P = 1$, $\alpha = 0.02$, $\gamma_{CC} = 10^{-5}$, $\gamma_{CD} = 6 \times 10^{-4}$, $\gamma_{DD} = 0.9$ and $N = 100$. The equilibrium closest to full defection is now stable, while the other one is unstable.

Appendix A. The strong selection limit

The following inequality holds,

$$\lim_{\beta \rightarrow \infty} h(x) = \lim_{\beta \rightarrow \infty} \frac{(1-a)x + a}{(a-b)x + b} e^{\beta(u'x + v')} \quad (\text{A.1})$$

$$\leq K \lim_{\beta \rightarrow \infty} e^{\beta(u'x + v')}, \quad (\text{A.2})$$

where $K > 0$ is a constant, whose specific value depends on the parameters a and b . Therefore,

$$\lim_{\beta \rightarrow \infty} h(x) = \begin{cases} \infty & (u'x + v' > 0), \\ 0 & (u'x + v' < 0). \end{cases} \quad (\text{A.3})$$

Hence, the direction of the gradient of selection depends solely on the sign of $u'x + v'$. C s are favored over D s if and only if $u'x + v' > 0$.

Appendix B. General intensities of selection

Let us designate the rational factor in equation (12) by $h_1(x)$ and the exponential factor by $h_2(x)$. Both $h_1(x)$ and $h_2(x)$ are monotonic in x :

- The derivative $\frac{d}{dx}h_1(x)$ is given by $\frac{b-a^2}{((a-b)x+b)^2}$. Hence, $h_1(x)$ increases monotonically if $b > a^2$ and decreases monotonically if $b < a^2$.
- The derivative $\frac{d}{dx}h_2(x)$ is given by $u' e^{u'x+v'}$. Hence, $h_2(x)$ increases monotonically if $u' > 0$ and decreases monotonically if $u' < 0$.

$h(x)$ increases (decreases) monotonically in case both $h_1(x)$ and $h_2(x)$ increase (decrease) monotonically. Hence, if $a^2 < b$ and $u' > 0$ (or $a^2 > b$ and $u' < 0$), then $g(x)$ can have at most one root in $]0, 1[$. If $h_1(x)$ increases (decreases) while $h_2(x)$ decreases (increases), then $g(x)$ can have at most two roots in $]0, 1[$.

References

- [1] Maynard Smith J 1982 *Evolution and the Theory of Games* (Cambridge: Cambridge University Press)
- [2] Hofbauer J and Sigmund K 1998 *Evolutionary Games and Population Dynamics* (Cambridge: Cambridge University Press)
- [3] Gintis H 2000 *Game Theory Evolving* (Princeton, NJ: Princeton University Press)
- [4] Nowak M A 2006 *Evolutionary Dynamics: Exploring the Equations of Life* (Cambridge, MA: Belknap Press of Harvard University Press)
- [5] Rapoport A and Chammah A M 1965 *Prisoner's Dilemma* (Ann Arbor, MI: University of Michigan Press)
- [6] Trivers R 1971 The evolution of reciprocal altruism *Q. Rev. Biol.* **46** 35–57
- [7] Axelrod R 1984 *The Evolution of Cooperation* (New York: Basic Books)
- [8] Michod R E 1999 *Darwinian Dynamics: Evolutionary Transitions in Fitness and Individuality* (Princeton, NJ: Princeton University Press)
- [9] Keller L 1999 *Levels of Selection* (Princeton, NJ: Princeton University Press)
- [10] Nowak M A and Sigmund K 2005 Evolution of indirect reciprocity *Nature* **437** 1291–8
- [11] Nowak M A 2006 Five rules for the evolution of cooperation *Science* **314** 1560–3
- [12] Taylor C and Nowak M A 2007 Transforming the dilemma *Evolution* **61** 2281–92
- [13] Nowak M A and May R M 1992 Evolutionary games and spatial chaos *Nature* **359** 826–9
- [14] Nowak M A, Bonhoeffer S and May R M 1994 Spatial games and the maintenance of cooperation *Proc. Natl Acad. Sci. USA* **91** 4877–81
- [15] Herz A V M 1994 Collective phenomena in spatially extended evolutionary games *J. Theor. Biol.* **169** 65–87
- [16] Lindgren K and Nordahl M G 1994 Evolutionary dynamics of spatial games *Physica D* **75** 292–309
- [17] Nakamaru M, Matsuda H and Iwasa Y 1997 The evolution of cooperation in a lattice-structured population *J. Theor. Biol.* **184** 65–81
- [18] Szabó G and Tóke C 1998 Evolutionary Prisoner's Dilemma game on a square lattice *Phys. Rev. E* **58** 69
- [19] Abramson G and Kuperman M 2001 Social games in a social network *Phys. Rev. E* **63** 030901
- [20] Vainstein M H and Arenzon J J 2001 Disordered environments in spatial games *Phys. Rev. E* **64** 051905
- [21] Ebel H and Bornholdt S 2002 Coevolutionary games on networks *Phys. Rev. E* **66** 056118
- [22] Kim B J, Trusina A, Holme P, Minnhagen P, Chung J S and Choi M Y 2002 Dynamic instabilities by asymmetric influence: Prisoners' Dilemma game in small-world networks *Phys. Rev. E* **66** 021907

- [23] Holme P, Trusina A, Kim B J and Minnhagen P 2003 Prisoner's Dilemma in real-world acquaintance networks: spikes and quasiequilibria induced by the interplay between structure and dynamics *Phys. Rev. E* **68** 030901
- [24] Szabó G and Vukov J 2004 Cooperation for volunteering and partially random partnerships *Phys. Rev. E* **69** 036107
- [25] Lieberman E, Hauert C and Nowak M A 2005 Evolutionary dynamics on graphs *Nature* **433** 312–6
- [26] Santos F C and Pacheco J M 2005 Scale-free networks provide a unifying framework for the emergence of cooperation *Phys. Rev. Lett.* **95** 098104
- [27] Szabó G, Vukov J and Szolnoki A 2005 Phase diagrams for an evolutionary prisoner's dilemma game on two-dimensional lattices *Phys. Rev. E* **72** 047107
- [28] Santos F C, Rodrigues J and Pacheco J M 2005 Epidemic spreading and cooperation dynamics on homogeneous small-world networks *Phys. Rev. E* **72** 056128
- [29] Ohtsuki H, Hauert C, Lieberman E and Nowak M A 2006 A simple rule for the evolution of cooperation on graphs *Nature* **441** 502–5
- [30] Santos F C and Pacheco J M 2006 A new route to the evolution of cooperation *J. Evol. Biol.* **19** 726–33
- [31] Santos F C, Pacheco J M and Lenaerts T 2006 Evolutionary dynamics of social dilemmas in structured heterogeneous populations *Proc. Natl Acad. Sci. USA* **103** 3490–4
- [32] Tomassini M, Luthi L and Giacobini M 2006 Hawks and doves on small-world networks *Phys. Rev. E* **73** 016132
- [33] Fu F, Liu L-H and Wang L 2007 Evolutionary prisoner's dilemma on heterogeneous Newman–Watts small-world network *Eur. Phys. J. B* **56** 367–72
- [34] Poncela J, Gómez-Gardeñes J, Floría L M and Moreno Y 2007 Robustness of cooperation in the evolutionary prisoner's dilemma on complex networks *New J. Phys.* **9** 184
- [35] Masuda N 2007 Participation costs dismiss the advantage of heterogeneous networks in the evolution of cooperation *Proc. Biol. Sci.* **274** 1815–21
- [36] Gómez-Gardeñes J, Campillo M, Floría L M and Moreno Y 2007 Dynamical organization of cooperation in complex topologies *Phys. Rev. Lett.* **98** 108103
- [37] Lozano S, Arenas A and Sánchez A 2008 Mesoscopic structure conditions the emergence of cooperation on social networks *PLoS One* **3** e1892
- [38] Szolnoki A, Perc M and Danku Z 2008 Towards effective payoffs in the prisoner's dilemma game on scale-free networks *Physica A* **387** 2075–82
- [39] Assenza S, Gómez-Gardeñes J and Latora V 2008 Enhancement of cooperation in highly clustered scale-free networks *Phys. Rev. E* **78** 017101
- [40] Vukov J, Szabó G and Szolnoki A 2008 Evolutionary prisoner's dilemma game on Newman–Watts networks *Phys. Rev. E* **77** 026109
- [41] Santos F C, Santos M D and Pacheco J M 2008 Social diversity promotes the emergence of cooperation in public goods games *Nature* **454** 214–6
- [42] Pacheco J M, Pinheiro F and Santos F C 2009 Population structure induces a symmetry breaking favoring the emergence of cooperation *PLoS Comput. Biol.* **5** e1000596
- [43] Szabó G and Fáth G 2007 Evolutionary games on graphs *Phys. Rep.* **446** 97–216
- [44] Albert R and Barabási A-L 2002 Statistical mechanics of complex networks *Rev. Mod. Phys.* **74** 47–97
- [45] Newman M E J 2003 The structure and function of complex networks *SIAM Rev.* **45** 167–256
- [46] Dorogotsev S N and Mendes J F F 2003 *Evolution of Networks: From Biological Nets to the Internet and WWW* (Oxford: Oxford University Press)
- [47] Kossinets G and Watts D J 2006 Empirical analysis of an evolving social network *Science* **311** 88–90
- [48] Skyrms B 2004 *The Stag-Hunt Game and the Evolution of Social Structure* (Cambridge: Cambridge University Press)
- [49] Gross T and Blasius B 2008 Adaptive coevolutionary networks—a review *Interface* **5** 259–71
- [50] Bala V and Goyal S 2000 A noncooperative model of network formation *Econometrica* **68** 1181–229

- [51] Skyrms B and Pemantle R 2000 A dynamical model of social network formation *Proc. Natl Acad. Sci. USA* **97** 9340–6
- [52] Zimmermann M G, Eguíluz V M, Cella-Conde C J and San Miguel M 2004 Coevolution of dynamical states and interactions in dynamic networks *Phys. Rev. E* **69** 065102
- [53] Zimmermann M G, Eguíluz V M and San Miguel M 2005 Cooperation and emergence of role differentiation in the dynamics of social networks *Am. J. Soc.* **110** 977
- [54] Santos F C, Pacheco J M and Lenaerts T 2006 Cooperation prevails when individuals adjust their social ties *PLoS Comput. Biol.* **2** 1284–91
- [55] Pacheco J M, Traulsen A and Nowak M A 2006 Active linking in evolutionary games *J. Theor. Biol.* **243** 437–43
- [56] Pacheco J M, Traulsen A and Nowak M A 2006 Co-evolution of strategy and structure in complex networks with dynamical linking *Phys. Rev. Lett.* **97** 258103
- [57] Biely C, Dragosits K and Thurner S 2007 The prisoner's dilemma on co-evolving networks under perfect rationality *Physica D* **228** 40–8
- [58] Hanaki N, Peterhansl A, Dodds P S and Watts D J 2007 Cooperation in evolving social networks *Manage. Sci.* **53** 1036–50
- [59] Tanimoto J 2007 Dilemma solving by the coevolution of networks and strategy in a 2×2 game *Phys. Rev. E* **76** 021126
- [60] Van Segbroeck S, Santos F C, Nowé A, Pacheco J M and Lenaerts T 2008 The evolution of prompt reaction to adverse ties *BMC Evol. Biol.* **8** 287
- [61] Fu F, Hauert C, Nowak M A and Wang L 2008 Reputation-based partner choice promotes cooperation in social networks *Phys. Rev. E* **78** 026117
- [62] Van Segbroeck S, Santos F C, Lenaerts T and Pacheco J M 2009 Reacting differently to adverse ties promotes cooperation in social networks *Phys. Rev. Lett.* **102** 058105
- [63] Szolnoki A and Perc M 2009 Resolving social dilemmas on evolving random networks *Europhys. Lett.* **86** 30007
- [64] Do A-L, Rudolf L and Gross T 2010 Patterns of cooperation: fairness and coordination in networks of interacting agents *New J. Phys.* **12** 063023
- [65] Wu B, Zhou D, Fu F, Qingjun L, Wang L and Traulsen A 2010 Evolution of cooperation on stochastic dynamical networks *PLoS ONE* **5** e11187
- [66] Perc M and Szolnoki A 2009 Coevolutionary games—a mini review *Biosystems* **99** 109–25
- [67] Ohtsuki H and Nowak M A 2006 The replicator equation on graphs *J. Theor. Biol.* **243** 86–97
- [68] Nowak M A, Sasaki A, Taylor C and Fudenberg D 2004 Emergence of cooperation and evolutionary stability in finite populations *Nature* **428** 646–50
- [69] Traulsen A, Nowak M A and Pacheco J M 2006 Stochastic dynamics of invasion and fixation *Phys. Rev. E* **74** 11909
- [70] Traulsen A, Pacheco J M and Nowak M A 2007 Pairwise comparison and selection temperature in evolutionary game dynamics *J. Theor. Biol.* **246** 522–9
- [71] Traulsen A, Shores N and Nowak M A 2008 Analytical results for individual and group selection at any intensity *Bull. Math. Biol.* **70** 1410–24
- [72] Van Segbroeck S, Santos F C and Pacheco J M 2010 Adaptive contact networks change effective disease infectiousness and dynamics *PLoS Comput. Biol.* **6** e1000895