

OPEN ACCESS

Generalized integrability and volume-preserving diffeomorphisms

To cite this article: C Adam *et al* 2008 *J. Phys.: Conf. Ser.* **128** 012025

View the [article online](#) for updates and enhancements.

You may also like

- [Bulk entanglement entropy and matrices](#)
Sumit R Das, Anurag Kaushal, Gautam Mandal et al.
- [Symmetries and exact solutions of the BPS Skyrme model](#)
C Adam, C D Fosco, J M Queiruga et al.
- [On the quantum structure of space-time, gravity, and higher spin in matrix models](#)
Harold C Steinacker

ECS
The
Electrochemical
Society
Advancing solid state &
electrochemical science & technology

DISCOVER
how sustainability
intersects with
electrochemistry & solid
state science research

Generalized Integrability and Volume-preserving Diffeomorphisms

C. Adam¹, J. Sánchez-Guillén¹ and A. Wereszczyński²

¹ Departamento de Física de Partículas, Universidad de Santiago and Instituto Galego de Física de Altas Enerxías (IGFAE) E-15782 Santiago de Compostela, Spain

² Institute of Physics, Jagiellonian University, Reymonta 4, 30-059 Kraków, Poland

E-mail: adam@fpaxp1.usc.es joaquin@fpaxp1.usc.es wereszczynski@th.if.uj.edu.pl

Abstract. The concepts of generalized zero curvature conditions and integrability in higher dimensions are briefly reviewed, where integrability in this context always means the existence of infinitely many conservation laws. It turns out that the conservation laws provided by the generalized integrability are, under certain additional assumptions, generated by the volume-preserving diffeomorphisms on target space. The possible conservation laws for field theories with a three-dimensional target space are classified. Further, an explicit example is discussed in some detail.

1. Introduction

Non-linear field theories are important in many fields of physics, with applications ranging from elementary particle theory to condensed matter physics. One feature of these theories which adds to their relevance is the possibility for the existence of extended static (solitons) or stationary (Q-balls) solutions. On the other hand, non-linear field theories are notoriously difficult to analyse, where the degree of difficulty strongly depends on the dimension of the base space (space-time) on which the fields are defined. In 1+1 dimensions, an ample mathematical apparatus has been developed for the analysis of non-linear theories, among which there are the inverse scattering method, Bäcklund transformations, or the zero curvature representation for integrable systems, which generalizes the Lax pair representation of finite-dimensional integrable systems. Integrability, that is, the existence of infinitely many conserved quantities, is related to all of these methods, and seems to be crucial in the analytical treatment of nonlinear theories, like, e.g., the explicit construction of solutions.

In higher dimensions, much less is known about non-linear field theories. A general concept of integrability has not yet been developed there. One may have, however, theories which contain an integrable subsector like, e.g., in the non-linear sigma model in 2+1 dimensions, where the integrable subsector is formed by the holomorphic solitons of Belavin and Polyakov. One generalization of the zero curvature representation of Shabat and Zakharov to higher dimensions has been proposed in [1], and it was demonstrated there that this proposal leads to non-linear field theories which have either infinitely many conservation laws in the full theory, or which contain integrable subsectors, defined by some additional constraint equations on the fields, such that the solutions belonging to this subsectors have infinitely many conservation laws. This zero

curvature representation, therefore, realizes the concept of integrability in higher-dimensional non-linear field theories in a specific and well-defined manner.

These methods have later been applied to specific models and to the analytic construction of both static and time-dependent solutions. For models with infinitely many conservation laws (the so-called AFZ model and related models), static and time-dependent solutions have been constructed, e.g., in [2], [3], [4], [5], and in [6], [7], respectively. Solutions in integrable subsectors of models which are, themselves, not integrable, have been constructed, e.g., in [8], [9]–[11] (the Nicole model and versions thereof) and in [12], [13] (diverse models on base space S^3). All these models share the property that their target space has dimension two.

A wellknown nonlinear field theory with three-dimensional target space is the Skyrme model [14] with target space $SU(2)$ (or equivalently the three-sphere S^3). Further, this model contains an integrable subsector, and the simplest Skyrmion (i.e., the simplest soliton of the Skyrme model with baryon number equal to one) belongs to this integrable subsector, see [16].

In many cases, it turns out that most of the new conserved currents in models and their subsectors are Noether currents and generalizations thereof, i.e., they are related to transformations of the target space variables (see [17]). So a direct, geometric approach has been successfully undertaken to find those currents, first for models with two-dimensional target spaces, [18], [19], and later also for three-dimensional target spaces, [20].

It is the purpose of this paper to give a short overview of some of these recent results on higher-dimensional integrability mentioned above, and to present some applications. Concretely, in Section 2 we briefly review the generalized curvature condition which was proposed in Ref. [1] as a possible way to generalize integrability to higher dimensions. In Section 3 we introduce volume-preserving diffeomorphisms on target space. For the case of three-dimensional target space we then classify for a wide class of Lagrangians all possible conservation laws, where the conserved currents in all cases are Noether currents of the volume-preserving target space diffeomorphisms. This section closely follows Ref. [20], but provides a slightly more refined classification, which turns out to be useful for applications. In Section 4 we study as an explicit example the abelian projection of Yang–Mills dilaton theory, which turns out to be integrable. Here we closely follow the results of [21]. Section 5 contains a brief discussion.

2. Generalized zero curvature condition

Here we briefly review the proposal for generalized integrability of Ref. [1], to which we refer for the details. The structure needed consists of a reducible Lie algebra $\tilde{G}$ which is a direct sum of another Lie algebra G and an abelian ideal H ,

$$\tilde{G} = G \oplus H \quad (1)$$

together with a flat connection

$$A_\mu \in G, \quad \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu] = 0, \quad (2)$$

and a covariantly constant vector field

$$B_\mu \in H, \quad \partial_\mu B^\mu + [A_\mu, B^\mu] = 0 \quad (3)$$

with

$$A_\mu = A_\mu^a T^a, \quad B_\mu = B_\mu^\alpha P^\alpha, \quad [P^\alpha, P^\beta] = 0, \quad (4)$$

where T^a and P^α form a basis in G and H , respectively. Further, $\mu = 1 \dots d$ are base space indices.

To gain some intuition, let us first remark that Eq. (2) is just a generalization to higher dimensions of the zero curvature condition of Zakharov and Shabat in 1+1 dimensions. Further,

Eq. 3 in some sense just generalizes the Lax pair $\dot{L} = [L, M]$ to higher dimensions. Another important point is that Eqs. (2), (3) are not chosen arbitrarily but may, in fact be derived from a generalized curvature condition as follows.

Firstly, the zero curvature condition Eq. (2) of Zakharov and Shabat may be derived as a consequence of the path independence of the Wilson line (or parallel transport) operator

$$W = P \exp \left(\int_0^\sigma d\sigma' A_\mu \frac{dx^\mu}{d\sigma'} \right) \quad (5)$$

where P indicates the path ordering. In an analogous way, Eqs. (2), (3) may be derived from the hypersurface independence of the following hypersurface ordered operator V in d dimensions,

$$V = \tilde{P} \exp \left(\int_{\Sigma_{d-1}} d\sigma^1 \dots d\sigma^{d-1} W^{-1} \tilde{B}_{\mu_1 \dots \mu_{d-1}} W \frac{dx^{\mu_1}}{d\sigma^1} \dots \frac{dx^{\mu_{d-1}}}{d\sigma^{d-1}} \right). \quad (6)$$

Here the $d-1$ form $\tilde{B}$ is the Hodge dual of the vector (one-form) of Eq. (3) (in fact, $\tilde{B}$ is the more natural object from the point of view of generalized integrability). Further, Σ_{d-1} is a based, ordered, closed hypersurface with base point $x_0 \equiv x(\sigma_j = 0)$. $\tilde{P}$ is the hypersurface ordering, which we shall explain a bit more in a moment.

The hypersurface independence of V , in turn, may be derived from the zero curvature condition for a connection $\mathcal{A}$ in higher loop space $\Omega^n(M, x_0)$ where

$$\Omega^n(M, x_0) = \{\gamma : S^n \rightarrow M, \gamma(0, \dots, 0) = x_0\} \quad (7)$$

Explicitly, the connection $\mathcal{A}$ reads

$$\mathcal{A} = \int_{\Sigma_{d-1}} d\sigma^1 \dots d\sigma^{d-2} W^{-1} \tilde{B}_{\mu_1 \dots \mu_{d-1}} W \frac{dx^{\mu_1}}{d\sigma^1} \dots \frac{dx^{\mu_{d-2}}}{d\sigma^{d-2}} \delta x^{\mu_{d-1}} \quad (8)$$

where $\delta x^{\mu_{d-1}}$ is the differential on higher loop space which provides an arbitrary infinitesimal variation of the higher loop. A closed ordered based hypersurface Σ^{d-1} may be interpreted as a closed loop in loop space Ω^{d-2} , and this observation allows to understand the hypersurface ordering. It is just ordinary path ordering of the corresponding ordinary loop in higher loop space Ω^{d-2} .

We want to emphasize that the conditions Eqs. (2), (3) are *sufficient, local* conditions for the zero curvature condition on the connection $\mathcal{A}$ of Eq. (8), but certainly they are not the most general ones.

After this brief review of the generalized zero curvature condition, we assume that Eqs. (2), (3) hold and make the following additional simplifying assumptions that

- G is a semisimple Lie algebra (e.g. $\mathfrak{su}(2)$) with

$$[T^a, T^b] = f_c^{ab} T^c \quad (9)$$

a structure constants f_c^{ab} .

- H is a (in general, reducible) representation space of G :

$$[T^a, P^\alpha] = R^{\alpha\beta} (T^a) P^\beta \quad (10)$$

- A_μ is explicitly flat:

$$A_\mu = g^{-1} \partial_\mu g, \quad g \in \mathcal{G} \quad (11)$$

(where e.g. $\mathcal{G} = \text{SU}(2)$) such that only Eq. (3) provides a nontrivial condition ($D_\mu B^\mu = 0$).

- Under these conditions, the currents

$$J_\mu = gB_\mu g^{-1} \quad (12)$$

are automatically conserved, $\partial^\mu J_\mu = 0$, and therefore the number of the conserved currents equals the dimension of the representation space H , $\dim H$. If $\dim H = \infty$, then we say that the corresponding field theory is integrable.

In general, the conserved currents of an integrable theory may be either Noether currents or may be related to hidden symmetries. Under the assumptions Eq. (9) - (12), however, the currents J_μ turn out to be Noether currents of geometric target space transformations, where the target space is spanned by the parameters of $g \in \mathcal{G}$.

A first example for this structure is like follows.

- The Lie group $\mathcal{G}$ is $SU(2)$ where, however, its elements $g \in \mathcal{G}$ are restricted to the equator of $SU(2)$ such that the target space is two-dimensional.
- The representation space H is the space of representations of $SU(2)$ with arbitrary integer angular momentum quantum number l , but magnetic quantum number m restricted to ± 1 ,

$$H = \{\text{reps } R_{lm} \text{ of } SU(2), m = \pm 1, l = 1, \dots, \infty\} \quad (13)$$

- Then the conserved currents J_μ of Eq. (12) turn out to generate area preserving diffeomorphisms on the two-dimensional target space (which may be, e.g., the two-sphere S^2 , but this depends on the Lagrangian).

A detailed discussion of this case may be found in [17], or in [18], [19].

Another class of theories with three-dimensional target spaces is obtained when the group element g is assumed to take values in the full unrestricted group $SU(2)$. There it turns out that the resulting conservation laws are generated by some subsets of the generators of volume-preserving diffeomorphisms on that target space. This case is discussed in the next section, where also a classification of the conservation laws of these theories is given. More details may be found in Ref. [20]

3. Conservation laws for Skyrme-type models

3.1. Volume-preserving diffeomorphisms

Let us start with a three-dimensional manifold (later to be identified with target space) with local coordinates X^i and with a volume form which in local coordinates reads

$$dV = h(X^i) dX^1 \wedge dX^2 \wedge dX^3 \quad (14)$$

where h is the volume density. Further, a diffeomorphism is an infinitesimal transformation

$$X^i \rightarrow X^i + \epsilon Y^i(X^j), \quad (15)$$

where ϵ is infinitesimal, and the Y^i are arbitrary functions of the X^j . A volume-preserving diffeomorphism has to obey

$$\partial_i(hY^i) \equiv \frac{\partial}{\partial X^i}(hY^i) = 0, \quad (16)$$

in addition. The corresponding vector fields

$$v^{(Y)} = Y^i \partial_i \quad (17)$$

form a closed Lie algebra, that is

$$[v^{(Y)}, v^{(\tilde{Y})}] = v^{(\tilde{\tilde{Y}})} \quad (18)$$

such that

$$\tilde{Y}^i = (\partial_j Y^i) \tilde{Y}^j - (\partial_j \tilde{Y}^i) Y^j, \quad \partial_i (h \tilde{Y}^i) = 0 \quad (19)$$

is again a volume-preserving diffeomorphism. For later convenience we change coordinates according to

$$\begin{aligned} u &\equiv X^1 + iX^2, & \xi &\equiv X^3 \\ Y^u &\equiv Y^1 + iY^2, & Y^\xi &\equiv Y^3. \end{aligned} \quad (20)$$

These new coordinates are especially useful for a parametrization of $g \in \text{SU}(2)$,

$$g = \exp(i\xi \vec{n} \cdot \vec{\sigma}) = \cos \xi + i \sin \xi \vec{n} \cdot \vec{\sigma} \quad (21)$$

provided that we also replace the unit vector field $\vec{n}$ by a complex field u via stereographic projection

$$\vec{n} \rightarrow u = \frac{n_1 + in_2}{1 + n_3}. \quad (22)$$

Further, we assume from now on the following form of the volume density $h = h(u\bar{u}, \xi)$ for simplicity. Finally, we interpret u, ξ as target space variables of a Lagrangian field theory with general Lagrangian $\mathcal{L}(u, \bar{u}, \xi, u_\mu, \bar{u}_\mu, \xi_\mu)$, where $u_\mu \equiv \partial_\mu u$, etc., then the Noether currents corresponding to the vector fields $v^{(Y)}$ generating volume-preserving diffeomorphisms on target space are given by

$$J_\mu^{(Y)} = Y^u \Pi_\mu + Y^{\bar{u}} \bar{\Pi}_\mu + Y^\xi P_\mu \quad (23)$$

with the usual canonical four-momenta

$$\Pi_\mu \equiv \partial_{u^\mu} \mathcal{L}, \quad P_\mu \equiv \partial_{\xi^\mu} \mathcal{L}. \quad (24)$$

The charges $Q^{(Y)} = \int d^3\mathbf{r} J_0^{(Y)}$ generate a Lie algebra isomorphic to the algebra of the vector fields $v^{(Y)}$ via the Poisson bracket, as usual.

3.2. Classification of conserved currents

We now specialize to the class of Lagrangians

$$\mathcal{L}(a, b, c, \xi, d, e) \quad (25)$$

where

$$a = u\bar{u}, \quad b = u^\mu \bar{u}_\mu, \quad d = \xi^\mu \xi_\mu \quad (26)$$

$$c = (u^\mu \bar{u}_\mu)^2 - u_\mu^2 \bar{u}_\nu^2, \quad e = \xi^\mu u_\mu \xi^\nu \bar{u}_\nu. \quad (27)$$

To motivate this choice let us mention that, e.g., the Skyrme model belongs to this class. Indeed, the Skyrme model has Lagrangian $\mathcal{L}_{\text{Sk}} = \frac{m^2}{2} \mathcal{L}_2 - \lambda \mathcal{L}_4$ where

$$\begin{aligned} \mathcal{L}_2 &= \text{tr}(g^{-1} g_\mu g^{-1} g^\mu) = d + 4b \frac{\sin^2 \xi}{(1+a)^2} \\ \mathcal{L}_4 &= \text{tr}[g^{-1} g_\mu, g^{-1} g_\nu]^2 = \frac{\sin^2 \xi}{(1+a)^2} (bd - e) + \frac{\sin^4 \xi}{(1+a)^4} c \end{aligned}$$

and g is the $\text{SU}(2)$ group element of Eq. (21).

For the class of Lagrangians (25) we now want to find which subsets of the currents (23) are conserved under which conditions. The calculation is lengthy but straight forward, and we

Table 1. Case a) corresponds to the symmetry $u \rightarrow e^{i\alpha}u$.

Case b) implies that the Lagrangian can be expressed by the pullback of a certain target space metric, such that h is the Riemannian volume density of that metric. One example for this case is the Skyrme model.

One example for case d) is provided by the abelian projection of Yang–Mills dilaton theory, which is discussed in Section 4.

No integrability conditions.	
a)	no condition on $\mathcal{L}$. Generically there exists only one vector field Y : $Y^u = iu$, $Y^{\bar{u}} = -i\bar{u}$, $Y^\xi = 0$.
b)	$\mathcal{L} = \mathcal{F}(hb, h^2c, d, he)$. There exist finitely many Y generating the isometries of the target space metric.
c)	$\mathcal{L}_b = 0$ and $\mathcal{L}_e = 0$. Y form the abelian subalgebra ($\tilde{G} = \tilde{G}(a)$): $Y^u = iu\tilde{G}_a$, $Y^{\bar{u}} = -i\bar{u}\tilde{G}_a$, $Y^\xi = 0$.
d)	conditions b) and c) on $\mathcal{L}$, and factorizing $h = h_1(a)h_2(\xi)$. Y forms the non-abelian subalgebra (for $G = G(u, \bar{u})$): $Y^u = ih_1^{-1}G_{\bar{u}}$, $Y^{\bar{u}} = -ih_1^{-1}G_u$, $Y^\xi = 0$.

Table 2. The integrability condition $u^2\bar{u}_\mu^2 - \bar{u}^2u_\mu^2 = 0$ may also be expressed like $\partial^\mu(\text{mod}(u))\partial_\mu(\arg(u)) = 0$, which provides a more geometric interpretation.

Integrability condition $u^2\bar{u}_\mu^2 - \bar{u}^2u_\mu^2 = 0$	
a)	$\mathcal{L}_b = 0$ and $\mathcal{L}_e = 0$. Y forms the abelian subalgebra (for $G = G(a, \xi)$): $Y^u = ih^{-1}uG_a$, $Y^{\bar{u}} = -ih^{-1}\bar{u}G_a$, $Y^\xi = 0$.
b)	$\mathcal{L}_e = 0$. Y form the abelian subalgebra ($\tilde{G} = \tilde{G}(a)$): $Y^u = iu\tilde{G}_a$, $Y^{\bar{u}} = -i\bar{u}\tilde{G}_a$, $Y^\xi = 0$.

present the resulting classification in the following four tables. In Table 1 the fields just obey the field equations and, therefore, there is a one to one correspondence between symmetries and conservation laws. In Tables 2-4, on the other hand, the fields have to obey certain first order equations (“integrability conditions”), in addition. These integrability conditions are not of the Euler–Lagrange type and, therefore, there is no longer a one to one correspondence between symmetries and conservation laws, see Ref. [22] for a detailed discussion.

Table 3. Case a) is obeyed by many configurations of the Skyrme model. E.g., the simplest Skyrme with baryon number one as well as many ansaetze for Skyrme configurations satisfy the conditions of case a).

Case d): the “weight number” $\mathcal{W}$ is defined for monomials of first derivatives of fields as $\mathcal{W} = \text{power}(u_\mu) + \text{power}(\bar{u}_\mu) - 2\text{power}(\xi_\mu)$, which gives e.g. $\mathcal{W}(b) = 2$, $\mathcal{W}(e) = -2$.

Integrability conditions $u^\mu \xi_\mu = 0$.	
a)	no condition on $\mathcal{L}$; or $\mathcal{L} = \mathcal{F}(hb, h^2c, d, he)$. Y forms the abelian subalgebra (for $G = G(a, \xi)$): $Y^u = ih^{-1}uG_a$, $Y^{\bar{u}} = -ih^{-1}\bar{u}G_a$, $Y^\xi = 0$. <i>And the further integrability condition $u^2\bar{u}_\mu^2 - \bar{u}^2u_\mu^2$ holds.</i>
b)	$\mathcal{L}_b = 0$. Y forms the abelian subalgebra (for $G = G(a, \xi)$): $Y^u = ih^{-1}uG_a$, $Y^{\bar{u}} = -ih^{-1}\bar{u}G_a$, $Y^\xi = 0$.
c)	$\mathcal{L}_b = 0$ and $\mathcal{L} = \mathcal{F}(hb, h^2c, d, he)$. Y form the subset $Y_\xi^\xi = 0$ (is <i>not</i> a subalgebra).
d)	$\mathcal{L}_b = 0$ and $\mathcal{L} = \mathcal{F}(hb, h^2c, d, he)$ and $\mathcal{W}(\mathcal{L}) = 0$. no further condition on Y .

Table 4. Case b) is obeyed by many configurations of the Skyrme model. E.g., the simplest Skyrme with baryon number one or the rational map ansaetze for Skyrme configurations satisfy the conditions of case b).

Integrability conditions $u_\mu^2 = 0$ and $u^\mu \xi_\mu = 0$.	
a)	no condition on $\mathcal{L}$. Y forms the abelian subalgebra (for $G = G(a, \xi)$): $Y^u = ih^{-1}uG_a$, $Y^{\bar{u}} = -ih^{-1}\bar{u}G_a$, $Y^\xi = 0$.
b)	$\mathcal{L} = \mathcal{F}(hb, h^2c, d, he)$ Y form the subset $Y_\xi^\xi = 0$ (is <i>not</i> a subalgebra).
c)	$\mathcal{L} = \mathcal{F}(hb, h^2c, d, he)$ and $\mathcal{W}(\mathcal{L}) = 0$. no further condition on Y .

4. Example: Abelian projection of YM dilaton theory

Here we want to demonstrate that the abelian projection of Yang–Mills dilaton theory is integrable in our sense. It belongs, in fact, to case d) of Table 1. A more detailed discussion can be found in Ref. [21]. The Lagrangian of Yang–Mills dilaton theory is

$$\mathcal{L} = \frac{1}{4}(2\xi^\mu \xi_\mu - e^{-2\kappa\xi} F^{a\mu\nu} F_{\mu\nu}^a) \quad (28)$$

where A_μ^a is an SU(2) Yang–Mills field and $F_{\mu\nu}^a$ is the corresponding field strength. Next, we want to employ the Cho–Faddeev–Niemi decomposition of the gauge field,

$$A_\mu^a = n^a C_\mu + \epsilon^{abc} n_\mu^b n^c + W_\mu^a, \quad (29)$$

where C_μ is an abelian gauge field, n^a is a unit vector in color space, and the so-called “valence field” W_μ^a is perpendicular to n^a in color space, $n^a W_\mu^a = 0$. To be consistent, the decomposition fields have to obey the constraint

$$\partial^\mu W_\mu^a + C_\mu \epsilon^{abc} n^b W_\mu^c + n^a W_\mu^b n_\mu^b = 0. \quad (30)$$

This constraint makes that the number of degrees of freedom of the original gauge field and of the decomposition match, and further, it provides the correct behaviour under gauge transformations for the decomposition fields, which infinitesimally read

$$\begin{aligned} \delta n^a &= \epsilon^{abc} n^b \alpha^c \\ \delta W_\mu^a &= \epsilon^{abc} W_\mu^b \alpha^c \\ \delta C_\mu &= n^a \alpha_\mu^a. \end{aligned} \quad (31)$$

In a next step, we perform the abelian projection, which consists in setting the valence field equal to zero,

$$W_\mu^a = 0. \quad (32)$$

Observe that the abelian projection is gauge invariant and obeys the constraint (30). The resulting gauge field $\hat{A}_\mu^a = n^a C_\mu + \epsilon^{abc} n_\mu^b n^c$ still is a full SU(2) connection, but with abelian field strength. The resulting abelian projected Yang–Mills dilaton theory is already integrable, that is, it has infinitely many conserved currents, see Ref. [21] for details. Here we make the further simplifying assumption $C_\mu \equiv 0$ (which is no longer gauge invariant). The resulting abelian projected Lagrangian is

$$\mathcal{L}_{\text{AP}} = \frac{1}{4} (2\xi^\mu \xi_\mu - e^{-2\kappa\xi} H^{\mu\nu} H_{\mu\nu}) \quad (33)$$

with

$$H_{\mu\nu} = \epsilon^{abc} n_\mu^a n_\nu^b n_\nu^c \quad (34)$$

or, after the stereographic projection (22),

$$\begin{aligned} \mathcal{L}_{\text{AP}} &= \frac{1}{2} \xi^\mu \xi_\mu - 2e^{-2\kappa\xi} \frac{(u^\mu \bar{u}_\mu)^2 - u_\mu^2 \bar{u}_\nu^2}{(1 + u\bar{u})^4} \\ &\equiv \frac{1}{2} d - 2h^2 c \end{aligned} \quad (35)$$

where $h = h_1(a)h_2(\xi) \equiv (1+a)^{-2}e^{-\kappa\xi}$. It corresponds to case d) of Table 1 and has, therefore, infinitely many symmetries and infinitely many conservation laws.

We now want to use our explicitly integrable parametrization of the abelian projection of Yang–Mills dilaton theory to discuss the problem of static solutions. For that purpose, we should first review the known results on that issue. It is known that there exist static, sphaleron type solutions in Yang–Mills dilaton theory. For the fully nonabelian theory, solutions both for radially and cylindrically symmetric ansätze are known numerically, whereas for the abelian subsector solutions for radially and cylindrically symmetric ansätze are known analytically. In the latter case, it is further known that the energy of the analytic solutions grows linearly with a certain integer m from the ansatz (the magnetic quantum number). The latter fact points

to the existence of a Bogomolny bound in the abelian projection, but an ansatz-independent derivation of this Bogomolny bound has not yet been given in the literature.

In our integrable abelian projection of YM dilaton theory, the analytic solutions may be calculated easily by quadratures, and the Bogomolny bound may be derived explicitly. Indeed, upon introducing spherical polar coordinates (r, θ, φ) in three-dimensional base space, the ansatz $\xi = \xi(r)$, $u = v(\theta) \exp(im\varphi)$ turns out to be consistent with the static field equations because of the base space symmetries of the theory. The resulting ordinary differential equations for $\xi(r)$ and $v(\theta)$ turn out to be solvable by quadratures, such that the corresponding exact analytic solutions may be calculated easily. The solvability by quadratures of the field equations might be related to the integrability of the theory. For details we refer to [21].

Finally, the Bogomolny bound may be derived easily within our parametrization. Indeed, we find for the energy corresponding to the Lagrangian (33) for static configurations

$$\begin{aligned} E_{\text{AP}} &= \frac{1}{2} \int d^3\mathbf{r} \left((\nabla\xi)^2 + e^{-2\kappa\xi} \vec{H}^2 \right) \\ &= \frac{1}{2} \int d^3\mathbf{r} \left(\nabla\xi - e^{-\kappa\xi} \vec{H} \right)^2 + \int d^3\mathbf{r} e^{-\kappa\xi} \nabla\xi \cdot \vec{H} \\ &\geq \int d^3\mathbf{r} e^{-\kappa\xi} \nabla\xi \cdot \vec{H} \equiv E_{\text{Bog}}. \end{aligned}$$

(where $\vec{H}$ is the Hodge dual of H_{jk}) and, therefore, the Bogomolny equation

$$\nabla\xi - e^{-\kappa\xi} \vec{H} = 0. \quad (36)$$

All the analytic static solutions mentioned above satisfy this equation and are, therefore, Bogomolny solutions. Further, the Bogomolny energy E_{Bog} may be expressed by the winding number of a map $S^3 \rightarrow S^3$, see again Ref. [21].

To recapitulate, our main results for the abelian projection of YM dilaton theory are that

- there exist infinitely many symmetries and infinitely many conserved currents,
- this fact may explain the infinitely many analytic solutions (this still is a conjecture, which exploits the analogy to the lower dimensional cases),
- there exist both a Bogomolny bound and a Bogomolny equation for static configurations, and the latter is solved by all known analytic solutions.

5. Discussion and Outlook

It was the purpose of this article to briefly review some recent developments in the attempts to generalize the concept of integrability to higher-dimensional nonlinear field theories. We gave a brief introduction to the general proposal for higher-dimensional integrability of Ref. [1] and then showed how, under certain additional assumptions, this higher-dimensional integrability is related to certain geometric target space transformations (concretely, volume preserving diffeomorphisms) which provide infinitely many conservation laws. We discussed in some more detail the case of a three-dimensional target space and, as a specific example, the abelian projection of Yang–Mills dilaton theory and its static analytic solutions. Some more applications to specific theories have already been studied (see, e.g., the references quoted in the Introduction), which already demonstrates the usefulness and importance of the concept of generalized integrability for the study of higher-dimensional nonlinear field theories. There exist, however, many more applications which are still open to further investigation. One obvious application is the search for time-dependent solutions (e.g. Q-balls) in theories where till now only static solutions have been found (e.g. in the integrable submodel of Yang–Mills dilaton theory of the previous section). Another possibility for generalizations consists in the choice of

larger groups $\mathcal{G}$ instead of $SU(2)$ in the integrability construction discussed in Section 2. This leads to integrable theories with higher-dimensional target spaces. The search for an integrable submodel of Einstein Yang–Mills dilaton theory and for analytic solutions within this submodel would be an obvious candidate, especially as for this theory only numerical solutions are known so far.

The generalizations and further investigations mentioned here still deal with a connection which is trivially flat, $A_\mu = g^{-1}\partial_\mu g$, such that the zero curvature condition (2) of Section 2 is trivially fulfilled, and Eq. (3) remains the only nontrivial generalized zero curvature condition. A further possible generalization consists in treating Eq. (2) as a nontrivial condition, too, which generates nontrivial constraints on the connection. The resulting modified generalized integrability might then lead to nonlocal conserved currents and to conservation laws which are not generated by geometric transformations, as is well-known to be the case in 1+1 dimensions. This line of investigation is, however, almost completely unexplored, and the above remarks are, therefore, largely tentative at the moment.

Acknowledgments

A.W. gratefully acknowledges support from Adam Krzyżanowski Fund and Jagiellonian University (grant WRBW 41/07). C.A. and J.S.-G. thank MCyT (Spain) and FEDER (FPA2005-01963), and support from Xunta de Galicia (grant PGIDIT06PXIB296182PR and Conselleria de Educacion). Further, C.A. acknowledges support from the Austrian START award project FWF-Y-137-TEC and from the FWF project P161 05 NO 5 of N.J. Mauser.

References

- [1] Alvarez O, Ferreira LA and Sanchez-Guillen J 1998 *Nucl. Phys. B* **529**, 689
- [2] Aratyn H, Ferreira LA and Zimmerman A 1999 *Phys. Lett.* **B456** 162
- [3] Aratyn H, Ferreira LA and Zimmerman A 1999 *Phys. Rev. Lett.* **83** 1723
- [4] De Carli E and Ferreira LA 2005 *J. Math. Phys.* **46** 012703 (*Preprint hep-th/0406244*)
- [5] Wereszczynski A. 2004 *Eur. Phys. J.* **C38** 261 (*Preprint hep-th/0405155*)
- [6] Ferreira LA 2006 *J. High Energy Phys.* JHEP03(2006)075 (*Preprint hep-th/0601235*)
- [7] Riserio do Bonfim AC and Ferreira LA 2006 *J. High Energy Phys.* JHEP03(2006)097 (*Preprint hep-th/0602234*)
- [8] Nicole DA 1978 *J. Phys.* **G4** 1363
- [9] Wereszczynski A 2005 *Eur. Phys. J.* **C41** 265 (*Preprint math-ph/0504008*)
- [10] Wereszczynski A 2005 *Phys. Lett.* **B621** 201 (*Preprint hep-th/0508121*)
- [11] Adam C, Sanchez-Guillen J, Vazquez RA and Wereszczynski A 2006 *J. Math. Phys.* **47** 052302 (*Preprint hep-th/0602152*)
- [12] Ward RS 1999 *Nonlinearity* **12** 241
- [13] Adam C, Sanchez-Guillen J and Wereszczynski A 2006 *Eur. Phys. J.* **C47** 513 (*Preprint hep-th/0602008*)
- [14] Skyrme THR 1961 *Proc. R. Soc. Lond.* **A260** 127
- [15] Skyrme THR 1962 *Nucl. Phys.* **31** 556
- [16] Ferreira LA and Sanchez-Guillen J 2001 *Phys. Lett.* **B504** 195 (*Preprint hep-th/0010168*)
- [17] Babelon O and Ferreira LA 2002 *J. High Energy Phys.* JHEP11(2002)020
- [18] Adam C and Sanchez-Guillen J 2005 *Phys. Lett.* **B626** 235 (*Preprint hep-th/0508011*)
- [19] Adam C, Sanchez-Guillen J and Wereszczynski A 2006 *J. Math. Phys.* **47** 022303 (*Preprint hep-th/0511277*)
- [20] Adam C, Sanchez-Guillen J and Wereszczynski A *J. Math. Phys.* **48** 032302 (*Preprint hep-th/0610227*)
- [21] Adam C, Sanchez-Guillen J and Wereszczynski A 2007 *Preprint hep-th/0703224*
- [22] Adam C and Sanchez-Guillen J 2005 *J. High Energy Phys.* JHEP01(2005)004 (*Preprint hep-th/0412028*)