

PAPER • OPEN ACCESS

Study the bactericidal activity of Propolis in rabbit model

To cite this article: Salim Hussein Hassan and Mohammed A K Al-Saadi 2019 *J. Phys.: Conf. Ser.* **1294** 062010

View the [article online](#) for updates and enhancements.

You may also like

- [Formulation and evaluation of anti-MRSA nanoemulsion loaded with Achyrocline satureioides: a new sustainable strategy for the bovine mastitis](#)
Gabriela Tasso Pinheiro Machado, Maria Beatriz Veleirinho, Luciana Aparecida Honorato et al.
- [Comparative evaluation of antibacterial activity of caffeic acid phenethyl ester and PLGA nanoparticle formulation by different methods](#)
Tülin Arasoglu, Serap Derman and Banu Mansuroglu
- [Controllability of magnetorheological shock absorber: II. Testing and analysis](#)
Xian-Xu Bai, Xue-Cai Deng and Sheng Shen



ECS
The
Electrochemical
Society
Advancing solid state &
electrochemical science & technology

DISCOVER
how sustainability
intersects with
electrochemistry & solid
state science research

Study the bactericidal activity of Propolis in rabbit model

Salim Hussein Hassan¹ and Mohammed A K Al-Saadi ²

1 Community health Dept. Technical Institute of Karbala/ Al- Furat Al- Awsat Technical University, Iraq

2 College of Medicine/ Babylon University, Iraq

Email : inkr.salm@atu.edu.iq

Abstract. This study aims to assess the bactericidal activity of Propolis extract against methicillin resistance *Staphylococcus aureus* (MRSA) in vivo. It included (11 male white New Zealand rabbits) evenly divided into two groups (6 animals in each one). Group I was fed daily with one milliliter of 20% extract Propolis for twenty days, while group II was fed without Propolis to serve as a control group. Ten days after the last dose, all animals were injected in intraperitoneal with single dose viable MRSA antigen at 0.5×10^8 cfu/ ml. After two days, blood samples that collected from all animals by heart puncture were cultured on blood agar media to assess bactericidal power activity of extract Propolis. The results pointed to effect of this extract on survive viable MRSA bacteria as compared with survive of same bacteria in control group animals. It was concluded Propolis had bactericidal power against pathogenic bacteria that appear resistant to antibiotics like MRSA bacteria.

Key words: Propolis, MRSA, bactericidal, rabbits.

Introduction:

Propolis is a natural resinous product that honeybees collect from several plants and mix it with bees wax and salivary enzymes that is also called bee glue (Silva-Carvalho *et al.*, 2015). Bees use propolis on their hives as defense against killers and microorganisms, to repair harm, as a thermal isolator, and to build aseptic locals to avoid infections of microbes (Marcucci, 1995; De Castro and Marcucci, 2000; Bankova, 2005). Since ancient times, propolis has been used by humans to meet the wants of health and food preservation. In the last years the interest in this natural product has better due to its wide spectrum of biological and pharmacological properties (Falcon *et al.*, 2013). Propolis is a lipophilic substantial that is firm and fragile when cold however elastic, soft, and very gummy when it heated. Possesses an agreeable aromatic odor and diverse color, including brown, green, and red, among others (Fokt *et al.*, 2012; Wagh, 2013). The chemical composition of it, generally composed of 50% resin, 30% wax, 10% essential oils, 5% pollen, and 5% other elements which contain flavonoids, organic compounds like phenolic acids (cinammic and caffeic acid), minerals, fatty acids and esters of phenolic acids terpenes, and alcohols (Barlak *et al.*, 2011; Righi *et al.*, 2013). Usually, the foremost constituents of propolis are resin and volatiles, which are materials obtained from a variation of botanical processes in different fragments of plants, and beeswax, concealed by the bees (Bankova, 2005). Samples of poplar propolis (from New Zealand, Europe, North America, and Asia) are principally composed of phenolic acids, and their esters as well as flavonoids (De Castro and Marcucci, 2000; Falcao *et al.*, 2013). Thus, the characters of different kinds of propolis rendering to its plant source (Silva-carvalho *et al.*, 2015). It is stated that pathogenic bacteria have a somewhat large possible for developing antibiotic resistances (Schwaber *et al.*, 2006). These resistance bacteria are



largely due to the extensive use of antibiotics in agriculture, medicine, and in animal care, and this problem is compounded by the lack of new antibiotics to attack bacteria in different ways to circumvent the resistant genes. Therefore, finding of new antimicrobial agents (of new sources) who are effective against resistant bacteria would be of great importance, consequently propolis is widely and extensively studies all around the world (Silva, *et al.*, 2012 b). Many researchers have reported the biological activities of propolis collected from different regions of Iraq (Al-Ammar, 2001) in Najaf city; (Aziz, 2005) in Baghdad and (Al-Shamari, 2006) in Qadysia city, were reported the antimicrobial activities of different sample of Iraqi propolis. Propolis contains substances that are able to inhibit cyclooxygenase and the consequent synthesis of PGE₂ (Araujo *et al.*, 2011). Propolis extracts can act on the nonspecific immune response via inhibiting of macrophage activation, H₂O₂ production and nitric oxide production; and this inhibition has been suggested as one mechanism of anti-inflammatory effects of propolis (Orsi *et al.*, 2000). The current study was designed to detecting the bactericidal power of blood in rabbits that feeding with Propolis.

1. Materials and methods:

In present study, ethanolic extract was used and prepare by dissolving two grams of crude propolis by ten ml of dimethyl sulfoxide (DMSO) to obtain ethanolic extract propolis solution at 20% concentration, and storage at 4°C up to using for animal administration. MRSA was obtained from microbiology laboratory of Technical Institute in Karbala city that isolated from burned person who has skin lesion, and this isolate re-culture to insure the diagnosis of it. Also, the study included 11 male white New Zealand rabbits) evenly divided into two groups I and II (6 animals in each one). Group I animals were fed with extract Propolis that prepared previously for 20 days, while group II were fed without this extract as control group. Single dose of viable MRSA bacteria was injected in intraperitoneal of animals in both groups and after two days one milliliter of blood was taken from all animals by heart puncture in order to culture on blood agar media to assess the blood clearance activity.

2. Results and Discussion:

The animals that injected with viable MRSA (11 animals), were divided into two groups as described in table (1). Blood samples were cultured to assess the clearance percentage of blood to detect the effects of Propolis on survive the viable bacteria (MRSA) in blood stream. The percentage of bactericidal power in group I was higher (83%), and in group II was (60%) as illustrated in the table (1). The result indicates Propolis has defense mechanisms that involve in the clearance the blood stream from pathogenic bacteria (MRSA) were injected in the animals. The gastrointestinal tract serves as an interface between the gut and immune system, with the intestinal lining functioning as a barrier; lactobacilli colonization decreases the passage of bacteria from the gut into the bloodstream (Mandeep *et al.*, 2013).

Table (1): activity of blood bactericidal effect on MRSA in testing groups

Testing groups	No.	Number of Blood culture +ve	% of +ve blood culture	MRSA Clearance %
Group II	6	1	17	83
Group III	5	2	40	60
Total	11	3		

The inhibiting activity of Propolis may be attributed to indirect effect on DNA replication that lead to inhibition of bacteria cell divisions, so present of many products in Propolis such as phenolic and flavonoids products which may cause this inhibition of bacterial activity (Salmani and Hassan, 2011), as well as disorganizing the cytoplasmic membrane and so inhibits protein synthesis (Hegazi *et al.*,

2014). Enzyme RNA-polymerase plays important role in DNA replication is also inhibits by Propolis and this mechanism lead to inhibition of bacterial activity (Wojtyczka *et al.*, 2013). Propolis and some of its derivatives were reported to be responsible for uncoupling the energy transducing cytoplasmic membrane inhibiting bacterial motility, which might contribute to the antibacterial action (Bankova *et al.*, 2000). In a study done by (Rahman, *et al.*, 2010), were showed the Propolis inhibit a Gram's positive bacteria better than Gram's negative. The antimicrobial activity of Propolis varied depending on dosage, sample, and the extraction solvents, the DMSO extracts of all Propolis samples were more active than the acetone extracts of the same samples, against *Brucella melitensis* (Orsi, *et al.*, 2005). The usage of natural products has been one of the greatest fruitful in modern medicines. Propolis has attracted increased attention because of antimicrobial activity against a varied range of pathogenic germs (Onlen *et al.*, 2007). Also the increasing of phagocytic activity by propolis in a study by (Abd Al-Sahib, 2011; Tao *et al.*, 2014) who mentioned propolis can significantly enhance the phagocytic function. The increased phagocytic activity against *S. aureus* bacteria may be due to the presence of glucose which assists in the destroying action of macrophages (Chepulis, 2007). Zinc plays important role in immune responses, in innate immune response it act as chemo-attractants for immune cells such as PMNs chemotaxis, and so promotes monocyte adhesion to endothelial cells as well as affects the maturation of DCs. On the other hand, the adaptive immune system is also affected by zinc levels, like T-cells are especially vulnerable to zinc deficiency that cause thymic atrophy and subsequent T-cell lymphopenia (Bonaventura *et al.*, 2015). Zinc also regulates metallothionein which is a family of cysteine-rich protein, that has roles in free radical scavenging and inflammatory process (Choi *et al.*, 2015). Haase and Lother, 2009 were mentioned the zinc deficiency cause impaired innate immune cells functions by many pathways such as oxidative burst of neutrophil granules, in contrast of high levels of zinc that induce chemotaxis. In a study done by AL-Greti 2016 who stated levels of Zinc in the serum of animals were fed with Propolis as compared with control groups, and for this fact can be attributed the bactericidal power of blood against MRAS bacteria in animals were feeding with Propolis. Park *et al.* (2014) were reported that dietary zinc-methionine enhanced mononuclear phagocytic function against *Salmonella enteritidis* and influences clearance of *E.coli* from blood in young's Turkey. Zinc is also required for phagocytosis activity of macrophages (Rao *et al.*, 2014), for example, high level of zinc prompts chemotaxis of PMNs. On the other hand, zinc deficiency causes decreased phagocytosis, and oxidative burst of neutrophil and monocytes (Prasad, 2000; Haase and Rink, 2009). Nicotinamide adenine di nucleotide phosphate (NADPH) which is regulates superoxide anion production that is responsible for the destruction the pathogen after phagocytosis, is also affected by zinc deficiency (Hasegawa *et al.*, 2000).

3. References:

- [1] Abd Alsahib, Z. A. (2011). The effects of honey on phagocytic activity against *Staphylococcus aureus*. Journal of Karbala University, 9(3):335–339.
- [2] Al-Ammar, M. H. M. (2001). Effects activity of propolis components on some spp. of pathogenic bacteria.M.sc. Thesis, Kufa University, Iraq.
- [3] Al-Halbosi, M. M. and Sulaiman, G. M. (2008). The Modulatory Effect of Iraqi Propolis Extract on Mitomycin-C Induced Micronucleus Formation in Albino Male Mice. Original Article Summary: Introduction, 50(1): 77–82.
- [4] Al-Hassnawi, H.H. M. (2012). Molecular Characterization of Antibiotic Resistance and Virulence Factors of Methicillin Resistance *Staphylococcus aureus* (MRSA) Isolated from Clinical Cases in Babylon Province. Ph. D. Thesis College of Medicine Babylon University, Iraq.
- [5] Araujo, M. R.; Libério, S. ; Guerra, R. M.; Ribeiro, M. S., and Nascimento, F. R. (2011). Mechanisms of action underlying the anti-inflammatory and immunomodulatory effects of propolis: A brief review. Brazilian Journal of Pharmacology, 22(1): 208–219.

- [6] Aziz, K. E. (2005). The effect of propolis extract on some species of bacteria isolated from inflamed infections of skin. M. Sc. Thesis Collage of Science, Al-Mustansiriya University, Iraq.
- [7] Bankova, V. (2005). Recent trends and important developments in propolis research. Evidence-based Compl. and Alt. Medicine, 2 (1): 29–32.
- [8] Barlak Y., Deger O., Olak M. C., Karatayli S. C., Bozdayi A. M. and Ucesan F. Y (2011). Effect of Turkish propolis extracts on proteome of prostate cancer cell line. Proteome Science, 9 article 74.
- [9] Brooks, G.F.; Butel, J.S.; and Morse, S. A. (2013). Medical microbiology. 26th ed. Lange Medical books, London.
- [10] Brown, E. M.; Sadarangani, M.; and Finlay, B. B. (2013). The role of the immune system in governing host-microbe interactions in the intestine. Nature Immunology, 14(7): 660–667.
- [11] Chepulis, L. M. (2007). "The Effects of Honey Compared with Sucrose and a Sugar-free Diet on Neutrophil Phagocytosis and Lymphocyte Numbers after Long-term Feeding in Rats. Journal of Complementary and Integrative Medicine.4: (1), Article 8.
- [12] Choi, R., Kim, H.-T., Lim, Y., Kim, M.-J., Kwon, O., and Jeon, (2015). Serum Concentrations of Trace Elements in Patients with Tuberculosis and Its Association with Treatment Outcome. Nutrients Journal, 7(7): 5969–5981
- [13] Clinical and Laboratory Standards Institute (CLSI). (2012). Performance standards for antimicrobial susceptibility testing. Approved standard. 32(3) National Committee for Clinical Laboratory Standards, Wayne, U.S.A.
- [14] Colin Hill, Francisco Guarner, Gregor Reid, Glenn R. Gibson, Daniel J. Merenstein, and Bruno P. (2014). "Expert consensus document: The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic". Nature Reviews Gastroenterology and Hepatology, 11 (8): 506–514.
- [15] De Castro V. S., and Marcucci M. C, (2000). Propolis: recent advances in chemistry and plant origin. Apidologie, 31 (1): 3–15.
- [16] Falcao S. I., Vale, N. P. and Gomes, E. (2013). Phenolic profiling of Portuguese propolis by LC-MS spectrometry: uncommon propolis rich in flavonoid glycosides. Phytochemical Analysis Journal, 24(4):309–318.
- [17] Fokt, H.; Pereira, A.; Ferreira A.; Cunha A. and Aguiar C. (2012). How do bees prevent hive infections? The antimicrobial properties of propolis. Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology, Mendez-Vilas, Ed., Microbiology Book Series, 1(2): 481–493.
- [18] Haase, H. and Lothar, R. (2009). The immune system and the impact of Zinc during aging. Licensee Biochemical Medical Central Ltd. 6:9. Herrmann, Journal and Lagrange.
- [19] Haase, H., and Rink, L. (2009). The immune system and the impact of zinc during aging. Immunity and ageing, 6(1):6-9.
- [20] Hasegawa, H.; Suzuki, K. ; and Nakaji, S. (2000). Effects of zinc on the reactive oxygen species generating capacity of human neutrophils and on the serum opsonic activity in vitro. Luminescence 15 : 321-7.
- [21] Hegazi, A. G., Altahtawy, R. H. M., Abdou, A. M., and Allah, F. A. (2014). Egyptian Propolis 10 : It ' s Effect on Hematological Changes and Bacterial Load in Mice-Bearing Ehrlich Ascites Carcinoma and Concurrently Infected with *Staphylococcus aureus*. Academic Journal of Cancer Research 7(3): 215–223.
- [22] Hendi, N.K.K. (2011). Antimicrobial Activity and Chemical Composition of Iraqi Propolis Collected from Middle Euphrates Region Ph.D. Thesis College of Medicine Babylon University, Iraq.
- [23] Junqueira L.C. and Carneiro, J. (2005). Basic Histology. Text and atlas 11th ed. McGraw-Hill. London.

- [24] Ma, X.; Guo, Z.; Liu, Y.; Zhou, Y.; Wang, Z.; Shen, Z. and Wang, J. (2016). The Humoral and Cellular Immune Response Characteristic of Propolis Flavonoid Used as Adjuvant to Inactivated PPV Vaccine in Sows. *Biochemistry and Analytical Biochemistry Journal*, 5(1):1-5.
- [25] MacFaddin, J.F. (2000). *Biochemical tests for identification of medical bacteria*. 3rd ed. The Williams and Wilkins, Baltimore, USA.
- [26] Mandeep S., Charanjit, R. and Aruna, B. (2013). Enhancement of Immune Responses by Probiotic Properties of *Lactobacillus acidophilus* Ncdc 195. *Journal of Drug Delivery and Therapeutics*, 3(3): 56–60.
- [27] Marcucci M. (1995). Propolis: chemical composition, biological properties
- [28] Onlen, Y. ;Duran, N. ; Atik, E.; Savas, L. ; Altug, E; Yakan, S. and Aslatas, O. (2007). Antibacterial activity of propolis against MRSA and synergism with topical Mupirocin. *Journal Alternative and Complementary Medicine* 13 (7): 713-718.
- [29] Orsi RO, Funari SRC, Soares AMVC, Calvi SA, Oliveira SL, Sforcin JM, and Bankova V (2000). Immunomodulatory action of propolis on macrophage activation. *Journal Venom Animal Toxins*, 6: 205-219.
- [30] Prasad AS. (2000). Effects of zinc deficiency on Th1 and Th2 cytokine shifts. *Journal Infectious Diseases* 182(Suppl 1):S62-S68.
- [31] Rahman, M. M. ; Richardson, A. ; and Sofian-Azirun, M. (2010). Antibacterial activity of propolis and honey against *Staphylococcus aureus* and *Escherichia coli*. *African Journal of Microbiology Research*, 4(16): 1872-1878.
- [32] Rao, S. V. R., Prakash, B., and Panda, A. K. (2014). Nutritional Modulations for Optimizing Immunocompetence in Chicken. 31(4), 314–323.
- [33] Righi, A.A., Negri, G., Salatino, A. (2013). Comparative chemistry of propolis from eight Brazilian localities. *Evidence Based Complement Alternat Medicine*, 1–14.
- [34] Salmani, A. K. and Hassan, I. Q. (2011). Evaluation of the antibacterial activity of ethanol extract of Iraqi propolis in vitro on some pathogenic bacteria causing mastitis and pneumonia in cows. *Journal of Anbar Veteran Science*, 2(4): 97–104.
- [35] Schwaber, M.J; Navon-Venezia, S; Kaye, K, S; Ami, R.B; Schwartz, D. and Carmeli1, Y. (2006). Clinical and economic impact of bacteremia with extended- spectrum- β - lactamase-producing Enterobacteriaceae. *Antimicrob. Agents Chem.*, 50: 1257-1262.
- [36] Silva-carvalho, R., Baltazar, F., and Almeida-aguiar, C. (2015). Propolis : A Complex Natural Product with a Plethora of Biological Activities that can be explored for Drug Development. *Evidence-Based Complementary and Alternative Medicine*, review article.
- [37] Tao, Y., Wang, D., Hu, Y., Huang, Y., Yu, Y., and Wang, D. (2014). The immunological enhancement activity of propolis flavonoids liposome in vitro and in vivo. *Evidence-based complementary and alternative medicine*, review article 483513 pp.1-8.
- [38] Wagh, V. D. (2013). Propolis: a wonder bee's product and its pharmacological potentials. *Advances in pharmacological sciences*, 308249.
- [39] Wojtyczka, R. D., K pa, M., Idzik, D., Kubina, R., Kabała-Dzik, A., Dziedzic, A., and W sik, T. J. (2013). In Vitro Antimicrobial Activity of Ethanolic Extract of Polish Propolis against Biofilm Forming *Staphylococcus epidermidis* Strains. *Evidence-based complementary and alternative Medicine*, 590703.
- [40] Yaghoubi, S.M; Ghorbani, G.R; Soleimanian, Z. S. and Satari, R. (2007). Antimicrobial activity of Iranian propolis and its chemical composition. *DARU*, 15: 45-48.