

PAPER • OPEN ACCESS

Molecular Study and Antibiotic susceptibility patterns of some Extended Spectrum Beta-Lactamase Genes (ESBL) of *Klebsiella pneumoniae* in Urinary Tract Infections

To cite this article: Ashwak B. Al-Hashimy and Weam K. Al-Musawy 2020 *J. Phys.: Conf. Ser.* **1660** 012017

View the [article online](#) for updates and enhancements.

You may also like

- [Comparison of chitosan and SLN nano-delivery systems for antibacterial effect of cinnamon \(*Cinnamomum verum*\) oil against MDR *K. pneumoniae* and *E. coli*](#)
Masoumeh Rohani, Mehran Nemattalab, Mohammad Hedayati et al.
- [Application of nanotechnology in antimicrobial finishing of biomedical textiles](#)
Andrea Zille, Luís Almeida, Teresa Amorim et al.
- [Recent advances in anticancer and antimicrobial activity of silver nanoparticles synthesized using phytochemicals and organic polymers](#)
Irshad A Wani, Tokeer Ahmad and Ajit Khosla



ECS
The
Electrochemical
Society
Advancing solid state &
electrochemical science & technology

DISCOVER
how sustainability
intersects with
electrochemistry & solid
state science research

Molecular Study and Antibiotic susceptibility patterns of some Extended Spectrum Beta-Lactamase Genes (ESBL) of *Klebsiella pneumoniae* in Urinary Tract Infections

Ashwak B. AL-Hashimy and Weam K. Al-Musawy

Institute of Genetic Engineering and Biotechnology for Postgraduate Studies,
University of Baghdad, Baghdad, Iraq.
Ministry of health, Central Public Health Laboratory, Baghdad, Iraq.

Abstract. The present study included the collection of 121 samples from MSU for investigating the presence of *K.pneumoniae* in UTIs, the samples have been collected from Al-Shaheed Mohammed Baqir AL-Hakeem hospital and private laboratories in Baghdad province. The study was carried out through March 2019 to the beginning of June 2019. The samples were identified based on the morphological and microscopically characteristics of the colonies when they were culturing on number of culture media as well as biochemical tests, molecular identification were also used as a final diagnostic test for isolates that were positive as they belong to *K.pneumoniae* bacteria during previous tests based on the blaTEM, blaSHV and blaCTX-M genes which has specific sequences for *K.pneumoniae* bacteria as a detection gene and also consider as virulence factor so it have a synonyms mechanism to antibiotic resistance. The results of the final diagnosis showed that 38 isolates belong to target bacteria, The examination of the sensitivity of all bacterial isolates was done for selected 38 isolation towards the 16 antibiotics by a Vitek2 compact ASTN system and the isolates were resistant for a number of antibiotics used such as; Amikacin (5.26%), Imepenem (5.26%), Ertapenem (7.89%), Meropenem (10.52%), Gentamicin (21.05%), Ciprofloxacin (26.32%), Cefoxitin (39.47%), Trimethoprim/Sulfamethoxazole (50%), Ceftriaxone (52.63%), Fosfomycin (55.26%), Piperacillin/Tazobactam (57.89%), Nitrofurantoin (57.89%), Ceftazidime (65.79%), Cefuroxime (71.05%), Cefixime (73.68%) and Ampicillin (100%). The presence of Extended Spectrum Beta-Lactamase genes in 38 *K.pneumoniae* isolates were 65.8 % of the ESBL genotypes expressed blaSHV genes followed by 52.6 % blaTEM and 42.1 % for blaCTX-M.

Keywords. *Klebsiella pneumoniae*, urinary tract infection, antibiotic susceptibility, blaTEM, blaSHV, blaCTX-M.

1. Introduction

Klebsiella pneumoniae is a Gram-negative, non-motile, encapsulated, lactose-fermenting, facultative anaerobic, gas-producing and rod-shaped bacterium, *K. pneumoniae* can grow either with or without



oxygen, which make it a facultative anaerobe. *Klebsiella pneumoniae* is responsible for many infections such as pneumonias, urinary tract infections, bacteremias, and liver abscesses. *Klebsiella pneumoniae* strains have become increasingly resistant to antibiotics, creating new challenges for modern medicine. It is based on four main parts; the capsule, the lipopolysaccharide, the fimbriae, and the siderophores, have been well studied and they are important for virulence in at least one infection model [1]. Extended spectrum beta-lactamases are a group of enzymes produced by certain bacteria that are able to hydrolyze extended spectrum antibiotics belonging to the penicillin and cephalosporin groups and monobactam. ESBLs are found in Gram-negative bacteria, especially in Enterobacteriaceae and *P.aeruginosa* [2]. Extended spectrum beta-lactamases are often located on plasmids that are transferable from strain to strain and between bacterial species. Although the prevalence of ESBLs is not known, it is clearly increasing. ESBL-producing Enterobacteriaceae is the main reason behind a number of outbreaks of the contagion around the globe, which resembles a challenge to control [3]. The SHV-1 b-lactamase is accountable for up to 20% of the plasmid-mediated ampicillin resistance in *K. pneumoniae* species [4].

2. Materials and Methods

2.1. Clinical Isolates

A total of (121) clinical specimen were collected from the beginning March 2019 to the beginning of June 2019 from Al-Shaheed Mohammed Baqir Al-Hakeem Hospital in Baghdad city and private laboratories. From patients with UTIs.

Isolation and identification of *Klebsiella pneumoniae* by Traditional methods: colony morphology and growing on different medium (CHROMagar Orientation, blood and macckongy agar), catalase test, citrate utilization, Triple-sugar iron agar, Indole test, oxidase reaction, urease test, growth at 37 °C, and identify by VITEK 2 compact.

2.2. Antimicrobial Susceptibility

The drug susceptibility test was carried out for all the isolates on VITEK 2 compact system to; Gentamicin, Nitrofurantoin, Trimethoprim /Sulfamethoxazole, Piperacillin /Tazobactam, Ceftazidime, Imipenem, Meropenem, Amikacin, Ciprofloxacin, Ampicillin, Ertapenem, Ceftriaxone, Cefoxitin, Cefuroxime, Cefixime and Fosfomycin.

2.3. Molecular Detection

DNA was extracted from activated pure culture of *Klebsiella pneumoniae* bacteria using DNA Bacteria Kit (geneaid). Detection of DNA bands using Agarose gelelectrophoresis (1%). The primers used in this study are shown in Table (1). PCR amplification of for the detection of *K.pneumoniae* genus was performed on all phenotypically tested strains of *K.pneumoniae*.

Table 1. Sequences of primers used for conventional PCR to detect *K.pneumonia* ESBL genes.

Target gene	Oligonucleotide primer sequence 5' to 3'	Amplicon size (bp)	Reference
<i>bla</i> SHV-F <i>bla</i> SHV-R	TCCCATGATGAGCACCTTTAAA TCCTGCTGGCGATAGTGGAT	104	[5]
<i>bla</i> CTX-M-F <i>bla</i> CTX-M-R	TCTTCCAGAATAAGGAATCCC CCGTTTCCGCTATTACAAAC	909	[6]
<i>bla</i> TEM-F <i>bla</i> TEM-R	TCCGCTCATGAGACAATAACC TTGGTCTGACAGTTACCAATGC	931	[7]

3. PCR amplification

5x FIREPol® Master Mix is a premixed ready-to-use solution containing all reagents required for PCR (except template, primers and water). Using 20 µL of PCR reaction, 5 µl DNA template was amplified using 4 µl of 5x FIREPol® master mix (Solis BioDyne/ Europe) and 1 µl were added from each of forward and reversed primers, up to the final volume 20 µl with nucleases free water (9 µl). Negative control contained all material except DNA, that D.W. was added instead of template DNA, the PCR programs were set on Thermal-cycler (Applied BioSystem, Singapore). Amplification reactions was started according to the program described in the table (2).

Table 2. Conditions of uniplex PCR reaction for ESBL genes detection.

Target gene	PCR conditions	No. of cycles
<i>bla</i> TEM and <i>bla</i> CTX- M	Initial denaturation at 95°C for 5 min	36 cycle
	Denaturation 95 °C for 30 sec	
	Annealing 60 °C for 30 sec	
	Extension 72 °C for 1 min	
	Final extension at 72 °C for 5 min	
<i>bla</i> SHV	Initial denaturation for 5 min at 94°C	36 cycle
	Denaturation 95 °C for 30 sec	
	Annealing 50 °C for 30 sec	
	Extension 72 °C for I min	
	Final extension at 72 °C for 5 min	

4. Results and Discussion

The first or initial diagnosis that based on microscopic examination and characteristic of the bacterial colonies, a total of 121 clinical specimens of urine were cultured on selective medium CHROMagar Orientation the 38 isolates of *K.pneumoniae* appeared as mucoid metallic blue colonies as in figure (1).



Figure 1. *K.pneumoniae* on CHROMagar orientation.

The results of biochemical tests for *K.pneumoniae* were negative results for indole and oxidase tests and gave positive result for citrate utilization, catalase and urease tests. *K.pneumoniae* were non-motile and On the TSI test strains produced acids both in butt and slant along with gas production and no H₂S production.

Antimicrobial susceptibility was performed on all 38 *K. pneumoniae* isolates to 16 different antibiotics by Vitek2 compact ASTN system that showed in chart (1).




Chart 1. Susceptibility of *K. pneumoniae* to 16 different antibiotics by Vitek2 compact.

It was found that all isolates were (100 %) resistant to Ampicillin the results was matched with study of [8, 9] in Iraq and Iran respectively whom isolates gram negative bacteria from UTIs patients. Depending on the results showed in chart (1) The bacterial isolates showed resistance to second and third generation cephalosporin ranged between (50% to 73%) our results with the previous reported by Ghaima et al. [10] in Baghdad province from patients with UTIs and Mansury et al. [11] in Iran, The second cephalosporin generation (Cefoxitin, Cefuroxime) and third generation (Ceftriaxone, Cefixime and Ceftazidime) were concordant with the results of Mishra et al. [12] in India who referred that the resistance of uropathogens to cephalosporin reached to 75%. The resistant against Nitrofurantoin in current study were (57.89%) to *K.pneumoniae* back to the study of Tawgozy and Amin [13] in Erbil city which were convergent to our results, nearly results were found in United State America [14] whom found that the service members who Sustained injured and infected with MDR *K.pneumoniae* in Iraq and Afghanistan were (62 %) resistant against Nitrofurantoin. Results referred also to convergent resistant against Piperacillin /Tazobactam (57.89 %) that similar with Al-Jubori [15] study in Baghdad and Al Wutayd et al. [16] study in Kingdom Saudi Arabia, while in Duhok city was sensitive (100%) [17]. Trimethoprim/ Sulfamethoxazole and Gentamicin in this study show low resistant like Khalid and Yassin, [18] in Duhok city from the clinical specimens including (urine, wound and the middle ear) get convergent result 33% for both of them in ESBL isolates. The present research argues that *K. pneumoniae* possesses a low-level of resistance against Imipenem (10.52%), Ertapenem (7.89%), Meropenem (5.26%) and Amikacin (5.26%) that convergent many studies with UTIs infection in Iraq and worldwide include [19, 20, 21] in Iraq, Iran and India respectively.

Total DNA was extracted using genomic DNA mini Kits then detected by gel electrophoresis as show in figure (2).




Figure 2. Agarose gel electrophoresis of extracted DNA to check purity and integrity. (70 V/ 30 min.).



Figure 3. Agarose gel electrophoresis of PCR products for the resistance genes *blaSHV*. Lane L: 100bp DNA ladder; lanes 1-38: *K. pneumoniae* isolates. Positive result in 1, 2, 4, 5, 8, 9, 11, 12, 14, 16, 17, 18, 19, 20, 21, 23, 34, 26, 27, 29, 30, 32, 34, 36 and 38.



Figure 4. Agarose gel electrophoresis of PCR products for the resistance genes *blaTEM*. Lane L: 100bp DNA ladder; 1-38 *K. pneumoniae* isolate. Positive result in 1, 2, 4, 7, 9, 12, 13, 15, 16, 18, 20, 22, 23, 25, 28, 29, 31, 33, 35 and 36.




Figure 5. Agarose gel electrophoresis of PCR products for the resistance genes *blaCTX-M*. Lane L: 100bp DNA ladder; 1-38 *K. pneumoniae* isolate. Positive result in 1, 2, 5, 7, 8, 10, 12, 14, 15, 17, 20, 22, 25, 28, 31 and 34.

Table 3. Genotypes of ESBL Genes Detected in Extended Spectrum β -lactamases-producing *K.pnumoniae* Isolates.

Genotypes	<i>K.pnumoniae</i> (n=38)
One genotype	
<i>blaSHV</i>	25 (65.8 %)

<i>bla</i> TEM	20 (52.6 %)
<i>bla</i> CTX-M	16 (42.1 %)
Number of selective genes in isolates	
All Three genes	4(10.5%)
Two genes	18(47.4%)
One gene	13(34.2%)
No one	3(7.9%)

The results of PCR reaction (Figures, 2 to 5) and Table (3) revealed the results of reorganization and genotypes of ESBL genes among *K.pneumoniae* isolates have been characterized for *bla*TEM, *bla*SHV and *bla*CTX-M genes, 65.8 % of the ESBL genotypes expressed *bla*SHV genes followed by 52.6 % *bla*TEM and 42.1 % for *bla*CTX-M. The results are in agreement with a local research done by Aljanaby and Alhasnawi, [22] whom demonstrated that majority of *K.pneumoniae* isolates from patients with urinary tract infection in Al-Najaf city were expressed prevalence with *bla*SHV gene followed by *bla*TEM (55.8%) then (51.2%) for *bla*CTX-M. the majority of the isolates expressed *bla*SHV genes that agreement with Salman [23] study in Baghdad also with UTIs patients. Somewhat identical outcomes have been reported in a study conducted in Turkey among patients with ESBL-producing carbapenem-resistant *K. pneumoniae* strains the most prevalent ESBL gene were *bla*SHV (97%), followed by *bla*TEM (92%), and *bla*CTX-M-1 (62%) [24]. While the local study in Baghdad revealed to very low prevalence of *bla*CTX-M (4.5 %) [23]. The current study found the percentage of isolates that contain two genes were (47.4%) that relative with Aljanaby and Alhasnawi, [22] study who demonstrated that *K. pneumoniae* isolates from inpatients with urinary tract infection in Al-Kufa hospital in Al-Najaf province of Iraq, and gave high resistance rate toward ampicillin, 3rd generation cephalosporins and Piperacillin /Tazobactam. Our study show 3 isolates (10.5%) carry all three genes similar with local study for Aljanaby and Alhasnawi, [22] in Al-Najaf province. Also convergent with Ali and Zahab, [25] in Egypt whom found 16.7% of isolates contain (*bla*TEM, *bla*SHV and *bla*CTX-M) genes. Our research show 3 isolates (7.9%) although of absence (*bla*TEM, *bla*SHV and *bla*CTX-M) genes there are resistance to antibiotics because of other ESBL genes that the same found with [22, 25] in Iraq and Egypt respectively.

References

- [1] Paczosa, M. K. and Mecsas, J. (2016). *Klebsiella pneumoniae*: going on the offense with a strong defense. *Microbiol. Mol. Biol. Rev.*, 80(3): 629-661.
- [2] Ejaz, H.; ul-Haq, I.; Mahmood, S.; Zafar, A. and Mohsin, M. (2013). Detection of extended-spectrum β -lactamases in *Klebsiella pneumoniae*: Comparison of phenotypic characterization methods. *Pakistan Journal of Medical Science*, 29(3): 768.
- [3] Rupp, M. and Fey, P. (2003). Extended Spectrum β -Lactamase (ESBL)-producing *Enterobacteriaceae*: considerations for diagnosis, prevention and drug treatment. *Journal Drugs*, 63(4): 353-365.
- [4] Huang, Y.; Yu, X.; Xie, M.; Wang, X.; Liao, K.; Xue, W. et al. (2016). Widespread dissemination of Carbapenem-resistant *Escherichia coli* sequence type 167 strains harboring blaNDM-5 in clinical settings in China. *Antimicrobial Agents Chemotherapy*, 60(7): 4364–4368.
- [5] Roschanski, N.; Fischer, J.; Guerra, B. and Roesler, U. (2014). Development of a multiplex real-time PCR for the rapid detection of the predominant beta-lactamase genes CTX-M, SHV, TEM and CIT-type AmpCs in *Enterobacteriaceae*. *PloS one*, 9(7): 100956.
- [6] Chouchani, C.; Marrakchi, R.; Ferchichi, L.; El Salabi, A., and Walsh, T. R. (2011). VIM and IMP metallo- β -lactamases and other extended-spectrum β -lactamases in *Escherichia coli* and *Klebsiella pneumoniae* from environmental samples in a Tunisian hospital. *Apmis*, 119(10): 725-732.

- [7] Eghbalpoor, F.; Habibi, M.; Azizi, O.; Asadi Karam, M. R. and Bouzari, S. (2019). Antibiotic resistance, virulence and genetic diversity of *Klebsiella pneumoniae* in community-and hospital-acquired urinary tract infections in Iran. *Acta microbiologica immunologica Hungarica*, 1-18 pages.
- [8] Salih, M. K.; Alrabadi, N. I.; Thalij, K. M. and Hussien, A. S. (2016). Isolation of pathogenic gram-negative bacteria from urinary tract infected patients. *Open Journal of Medical Microbiology*, 6(02): 59.
- [9] Ghotaslou, R.; Sadeghi, M. R.; Akhi, M. T., Hasani, A. and Asgharzadeh, M. (2018). Prevalence and antimicrobial susceptibility patterns of ESBL, ampC and carbapenemase-producing enterobacteriaceae isolated from hospitalized patients in Azerbaijan, Iran. *Iranian journal of pharmaceutical research: IJPR*, 17(1): 79.
- [10] Ghaima, K. K., Hamid, H. H., and Hasan, S. F. (2017). Biofilm formation, antibiotic resistance and detection of mannose-resistant *Proteus*-like (MR/P) fimbriae genes in *Proteus mirabilis* isolated from UTI. *International Journal of ChemTech Research*, 10(5), 964-971.
- [11] Mansury, D.; Motamedifar, M.; Sarvari, J.; Shirazi, B. and Khaledi, A. (2016). Antibiotic susceptibility pattern and identification of extended spectrum β -lactamases (ESBLs) in clinical isolates of *Klebsiella pneumoniae* from Shiraz, Iran. *Iranian journal of microbiology*, 8(1): 55.
- [12] Mishra, M.; Debata, N. and Padhy, R. (2013). Surveillance of multi-drug resistant uropathogenic bacteria in hospitalized patient in Indian. *Asian Pacific J tropical biomedicine*, 3(4):315-324.
- [13] Tawgozy, F. H. A. and Amin, B. K. (2018). Molecular analysis of *Klebsiella Pneumoniae* isolated from UTI patients in Erbil city. *ZANCO Journal of Pure and Applied Sciences*, 30(3): 1-9.
- [14] Kiley, J.; Mende, K.; Kaiser, S. J.; Carson, L.; Lu, D. Z.; Merritt, T. et al. (2018). 1184. Resistance patterns and susceptibility analysis of *Klebsiella pneumoniae* infections in service members who sustained trauma in Iraq and Afghanistan. In *open forum infectious diseases*. 357- : (1)5 358.
- [15] Al-Jubori, S. S. (2012). Determination genetic factors controlling β -lactemase enzyme related to bla TEM and bla SHV families using the polymerase chain reaction. *Iraqi Journal of Biotechnology*, 11(2): 377-388.
- [16] Al Wutayd, O.; Al Nafeesah, A.; Adam, I. and Babikir, I. (2018). The antibiotic susceptibility patterns of uropathogens isolated in Qassim, Saudi Arabia. *The Journal of Infection in Developing Countries*, 12(11): 946-952.
- [17] Assafi, M. S.; Ibrahim, N. M.; Hussein, N. R.; Taha, A. A. and Balatay, A. A. (2015). Urinary bacterial profile and antibiotic susceptibility pattern among patients with urinary tract infection in Duhok city, Kurdistan region, Iraq. *International Journal of Pure and Applied Sciences and Technology*, 30(2): 54.
- [18] Khalid, H. M., and Yassin, N. A. (2017). Distribution of extended spectrum β -lactamase genes among *Proteus mirabilis* isolated from clinical specimens in Duhok city, Kurdistan region, Iraq. *Science Journal of University of Zakho*, 5(1), 1-6.
- [19] Hussein, N. R.; Daniel, S.; Salim, K. and Assafi, M. S. (2018). Urinary tract infections and antibiotic sensitivity patterns among women referred to Azadi teaching hospital, Duhok, Iraq. 5(2): 27-30.
- [20] Majlesi, A.; Kakhki, R. K.; Nejad, A. S. M.; Mashouf, R. Y.; Roointan, A.; Abazari, M. et al. (2018). Detection of plasmid-mediated quinolone resistance in clinical isolates of Enterobacteriaceae strains in Hamadan, West of Iran. *Saudi journal of biological sciences*, 25(3): 426-430.
- [21] Veeraraghavan, B.; Jesudason, M. R.; Prakasah, J. A. J.; Anandan, S.; Sahni, R. D.; Pragasam, A. K. et al. (2018). Antimicrobial susceptibility profiles of gram-negative bacteria causing infections collected across India during 2014–2016: Study for monitoring antimicrobial resistance trend report. *Indian journal of medical microbiology*, 36(1): 32.
- [22] Aljanaby, A. and Alhasnawi, H. (2017). Phenotypic and molecular characterization of multidrug resistant *Klebsiella pneumoniae* Isolated from Different Clinical Sources in Al-Najaf

- Province-Iraq. Pakistan Journal of Biological Sciences, 20(5): 217-232.
- [23] Salman, R. A. (2019). Study the Extended Spectrum Beta-Lactamase Genes (ESBL) in *Klebsiella pneumoniae* Isolated from Urinary Tract Infections. MSC. Thesis. University of Baghdad. Institute of Genetic Engineering and Biotechnology for Postgraduate Studies, Iraq.
- [24] Iraz, M.; Özad Düzgün, A. Sandallı, C.; Doymaz, M.; Akkoyunlu, Y.; Saral, A. et al. (2015). Distribution of β -Lactamase genes among Carbapenem-resistant *Klebsiella pneumoniae* strains isolated from patients in Turkey. *Annals of Laboratory Medicine*, 35(6):595-601.
- [25] Ali, M. S. and Zahab, A. Z. (2011). Extended Spectrum β -Lactamases in *Escherichia coli* and *Klebsiella pneumoniae*, their molecular characterization and associated risk factors. *Egyptian Journal of Medical Microbiology*, 20(3): 35-45.