## ENVIRONMENTAL RESEARCH LETTERS

#### **OPEN ACCESS**

## The role of surface energy fluxes in pan-Arctic snow cover changes

To cite this article: Xiaogang Shi et al 2011 Environ. Res. Lett. 6 035204

View the article online for updates and enhancements.

#### You may also like

- Polar amplification and elevationdependence in trends of Northern Hemisphere snow cover extent.
  <u>1971–2014</u>
  Marco A Hernández-Henríquez, Stephen J Déry and Chris Derksen
- Reconstructing spring sea ice concentration in the Chukchi Sea over recent centuries: insights into the application of the PIP<sub>as</sub> index Jung-Hyun Kim, Jong-Ku Gal, Sang-Yoon Jun et al.
- Evaluating the impact of peat soils and snow schemes on simulated active layer thickness at pan-Arctic permafrost sites Jing Tao, William J Riley and Qing Zhu



This content was downloaded from IP address 3.144.151.106 on 28/04/2024 at 23:13

# The role of surface energy fluxes in pan-Arctic snow cover changes

## Xiaogang Shi<sup>1</sup>, Pavel Ya Groisman<sup>2</sup>, Stephen J Déry<sup>3</sup> and Dennis P Lettenmaier<sup>1,4</sup>

<sup>1</sup> Department of Civil and Environmental Engineering, University of Washington, Seattle, WA, USA

<sup>2</sup> National Climatic Data Center, Asheville, NC, USA

<sup>3</sup> Environmental Science and Engineering Program, University of Northern British Columbia, Prince George, BC, Canada

E-mail: dennisl@u.washington.edu

Received 19 June 2011 Accepted for publication 26 August 2011 Published 26 September 2011 Online at stacks.iop.org/ERL/6/035204

#### Abstract

We analyze snow cover extent (SCE) trends in the National Oceanic and Atmospheric Administration's (NOAA) northern hemisphere weekly satellite SCE data using the Mann-Kendall trend test and find that North American and Eurasian snow cover in the pan-Arctic have declined significantly in spring and summer over the period of satellite record beginning in the early 1970s. These trends are reproduced, both in trend direction and statistical significance, in reconstructions using the variable infiltration capacity (VIC) hydrological model. We find that spring and summer surface radiative and turbulent fluxes generated in VIC have strong correlations with satellite observations of SCE. We identify the role of surface energy fluxes and determine which is most responsible for the observed spring and summer SCE recession. We find that positive trends in surface net radiation (SNR) accompany most of the SCE trends, whereas modeled latent heat (LH) and sensible heat (SH) trends associated with warming on SCE mostly cancel each other, except for North America in spring, and to a lesser extent for Eurasia in summer. In spring over North America and summer in Eurasia, the SH contribution to the observed snow cover trends is substantial. The results indicate that  $\Delta$ SNR is the primary energy source and  $\Delta$ SH plays a secondary role in changes of SCE. Compared with  $\Delta$ SNR and  $\Delta$ SH,  $\Delta$ LH has a minor influence on pan-Arctic snow cover changes.

Keywords: pan-Arctic, snow cover, recession, surface energy fluxes

#### 1. Introduction

The pan-Arctic domain is one of the most sensitive regions on Earth to global climate change (Manabe and Stouffer 1994, Miller and Russell 2000, Holland and Bitz 2003, Serreze *et al* 2009). As the largest single component of the cryosphere in terms of spatial extent (Armstrong and Brodzik 2001), snow cover variations in space and time have resulted in significant changes in the surface energy and water budgets over the pan-Arctic land region (Serreze *et al* 2000, Peterson *et al* 2002, Serreze *et al* 2002, Yang *et al* 2003, McClelland *et al* 2006, Rawlins *et al* 2010, Adam *et al* 2007, Shiklomanov *et al* 2007). However, the interpretation of changes in the areal extent of snow cover and its timing is complicated by the sparseness of *in situ* observations of surface radiative and turbulent fluxes, which are the most important variables affecting snow surface energy exchange processes (Cline 1997).

Previous studies have attempted to track energy balance changes associated with snow cover variations at the point scale in Finland (Kuusisto 1986, Koivusalo and Kokkonen 2002); New Zealand (Prowse and Owens 1982, Moore and Owens 1984); the southern Sierra Nevada region of California in the western United States (Marks and Dozier 1992), and at

<sup>&</sup>lt;sup>4</sup> Address for correspondence: Department of Civil and Environmental Engineering, University of Washington, Box 352700, Seattle, WA 98195-2700, USA.



Figure 1. Study domain with 100 km resolution EASE grid mesh.

the basin scale in two regions of Alaska (Robinson 1986); the Red River Valley of North Dakota and Minnesota (Dyer and Mote 2002); Trail Valley Creek of northern Canada (Marsh and Pomeroy 1996, Pohl and Marsh 2006). However, few studies have examined the large-scale factors that would provide better understanding of the relative importance of snow surface energy balance components (Male and Granger 1981, Cline 1997, Leathers et al 2004), in large-scale snow cover changes, especially for the pan-Arctic land area. On the other hand, land surface models (e.g. Liang et al 1994) have improved to the point that they may, in some cases, serve as surrogates for in situ observations. Off-line runs of these models provide sources for most or all terms in the snow surface energy balance and offer the opportunity to investigate the nature of the space-time variability of the snow surface energy budget (Betts et al 2009, Troy and Wood 2009, Shi et al 2010).

In this letter, our main objective is to identify the individual role of surface energy fluxes in the snow surface energy balance and determine which are most responsible for observed changes in snow cover extent (SCE) of the pan-Arctic land region as shown in figure 1. First, monotonic trends in satellite snow cover observations and corresponding reconstructions generated by a land surface model are analyzed using a non-parametric trend test. Subsequently, the relationships between observed SCE and modeled surface radiative and turbulent fluxes are examined. In the final part of the letter, the relative importance of each component in the snow surface energy balance is estimated.

#### 2. Data sets

#### 2.1. Snow cover extent data

Observed seasonal values of SCE were extracted from the weekly snow cover and sea ice extent version 3 product for

the northern hemisphere maintained at the National Snow and Ice Data Center (NSIDC), which combines snow cover and sea ice extent for the period from October 1966 through June 2007 (Armstrong and Brodzik 2005). The data set is based on weekly maps of continental SCE produced by the National Oceanic and Atmospheric Administration's (NOAA) National Environmental Satellite Data and Information Service (NESDIS) (Robinson et al 1993, Frei and Robinson 1999), which were derived from digitized versions of manual interpretations of Advanced Very High Resolution Radiometer (AVHRR), Geostationary Operational Environmental Satellite (GOES), and other visible band satellite data. This satellitebased data set has been regridded to the NSIDC EASE grid with a spatial resolution of 25 km. Our study is restricted to the period 1972–2006 because there are some missing charts between 1967 and 1971 (Robinson 2000). Although ending the time series in 2006 leaves out some exceptionally low Arctic spring SCE values in recent years (e.g. 2008-10), the nonparametric statistical method we used (section 3.1) is robust to modest changes in the length of the record analyzed. In addition, Greenland is not included in the analyses as its snow cover is mainly perennial in nature (Déry and Brown 2007). Brown et al (2010) have assessed this SCE record (commonly referred to as the NOAA weekly SCE record) in comparison to other available Arctic snow cover data sets. In general, their study, and others (Wiesnet et al 1987, Robinson et al 1993) have found that the NOAA weekly SCE data set is reliable for continental-scale studies of snow cover variability. It has become a widely used tool for deriving trends in climate-related studies (Groisman et al 1994, Déry and Brown 2007, Flanner et al 2009, Derksen et al 2010, Derksen and Brown 2011), notwithstanding uncertainties in some parts of the domain for certain times of the year, especially during springtime over northern Canada (Wang et al 2005). A more recent update to the data set we used (NOAA snow chart climate data record (CDR)) is now available (Brown and Robinson 2011), but the differences between the new CDR and the data set we used at the pan-Arctic scale are small.

In this study, simulated SCE was reconstructed from 1972 to 2006 using the variable infiltration capacity (VIC) model, which is a macroscale hydrologic model that solves the energy and water balance and represents ephemeral snow cover over a gridded domain (Liang et al 1994, 1996). The off-line simulations from VIC used here are at a 3 h time step in full energy balance mode (meaning that the model closes its surface energy budget), forced with daily precipitation, maximum and minimum temperatures and wind speed through 2007 at a spatial resolution (EASE grid) of 100 km, constructed using methods outlined by Adam and Lettenmaier (2008). Precipitation and temperature were from gridded observations (Willmott and Matsuura 2009) and wind speed was from the NCEP/NCAR reanalysis (Kalnay et al 1996). Precipitation was adjusted for gauge undercatch and orographic effects as described by Adam and Lettenmaier (2003) and Adam et al (2006). The snow parameterization in VIC represents snow accumulation and ablation processes using a two-layer energy and mass balance approach (Cherkauer and Lettenmaier 2003) and a canopy snow interception algorithm (Storck et al 2002,

Andreadis *et al* 2009) when an overstory is present. In the VIC model, each grid cell is partitioned into five elevation (snow) bands, which can include multiple land cover types (tiles). The snow model is then applied to each tile separately. When snow water equivalent is greater than zero, VIC assumes that snow fully covers the tile. For each grid cell, the simulated snow cover extent is calculated as the area averages of the tiles.

#### 2.2. Surface energy fluxes data

Surface energy fluxes including downward shortwave radiation (DSW) and downward longwave radiation (DLW) were calculated by using a Temperature INDex (TIND) scheme (Kimball et al 1997, Thornton and Running 1999, Shi et al 2010) wherein DSW and DLW are estimated based on relationships with the daily temperature range and daily average temperature, respectively. TIND has been commonly used in model intercomparison experiments such as the Project for Intercomparison of Land Parameterization Schemes (PILPS) (e.g. Pitman et al 1999) and land surface models, such as VIC, for long-term simulations in cases when direct observations of energy fluxes are not available. Shi et al (2010) evaluated DSW, DLW, and albedo computed in an off-line simulation of VIC embedded with TIND along with satellite data and global reanalysis products in comparison with in situ observations from the Global Energy Balance Archive (GEBA, Ohmura et al 1989) and showed that these estimates compared well with observations over the pan-Arctic land region. Compared to the in situ observations, the mean seasonal DSW from the European Centre for Medium-Range Weather Forecast (ECMWF) 40 Year Reanalysis (ERA-40), the ECMWF Interim Reanalysis (ERA-Interim), the International Satellite Cloud Climatology Project Flux Data (ISCCP-FD), and the TIND-based scheme all have small biases ( $\pm 20 \text{ W m}^{-2}$ ). ERA-40, ERA-Interim and VIC DLW deseasonalized monthly anomalies had high correlations (r =0.96, 0.97 and 0.91, respectively) with GEBA observations whereas the correlation for the satellite-based (ISCCP-FD) product was somewhat lower. VIC deseasonalized monthly albedo had similar anomaly correlations with GEBA observations as did ERA-40, ERA-Interim and ISCCP-FD estimates.

Surface net radiation (SNR) was obtained as the sum of net shortwave (SW) and longwave radiative (LW) fluxes. SW at the snow surface is a measure of the difference between DSW and upward shortwave radiation (USW). USW is the product of DSW and snow surface albedo that is assumed to decay with age based on relationships published by the US Army Corps of Engineers (1956). LW is the sum of DLW emitted by the atmosphere and the fluxes emitted upward by a melting snow surface. DLW was estimated using equation (2.42) from Bras (1990), which is based on air temperature and a function for emissivity from Tennessee Valley Authority (1972). The turbulent fluxes (sensible heat (SH) and latent heat (LH)) near the snow surface were produced using VIC's bulk aerodynamic approach, which is described in Andreadis et al (2009). The algorithm requires snow surface temperature, which is calculated by VIC's snow



**Figure 2.** Spring and summer snow cover area fraction over the pan-Arctic land areas in North America and Eurasia from VIC and satellite observations for the period of 1972–2006.

algorithm, wind speed, surface air temperature and relative humidity, the last three of which are taken from the forcing data. Bulk transfer coefficients for momentum, heat, and water vapor were calculated initially for a neutral condition (Price and Dunne 1976). Subsequently, the aerodynamic resistance in the presence of snow cover was corrected using the bulk Richardson's number for stable or unstable atmospheric conditions (Anderson 1976) as implemented in the VIC snow model (Andreadis *et al* 2009). A similar approach has been successfully applied in various settings in Arctic environments (e.g. Hinzman *et al* 1991, Woo *et al* 1999, Boike *et al* 2003). Other energy fluxes in the snow surface energy balance, such as ground heat and advective fluxes were also generated by VIC. VIC's energy flux convention is that surface energy fluxes toward the snow surface are defined as positive.

#### 3. Results

#### 3.1. Temporal analyses of SCE and surface energy fluxes

We defined April and May as spring, consistent with Groisman *et al* (1994), and a three-month window centered on July was defined as summer. Observed and simulated SCE in spring and summer expressed as a snow-covered fraction were calculated for the Eurasia and North America study domains (figure 1). Figure 2 shows North American and Eurasian snow cover fraction (SCF) during spring and summer from VIC and NOAA observations for the period of 1972–2006. The VIC simulations match the observed SCF over both North America and Eurasia quite well, with a mean absolute bias of 4.5% in spring and 0.6% for summer.

To examine long-term trends in SCE and surface energy fluxes, we used the non-parametric Mann–Kendall trend test (Mann 1945) for trend significance, and the Sen method (Sen 1968) to estimate trend slope. A 5% significance level (two-sided test) was specified. Trend tests were performed for seasonal SCE from VIC and satellite observations, and modeled radiative and turbulent fluxes averaged over the snowcovered portions of Eurasia and North America, respectively.

Table 1 summarizes trend test results for spring and summer SCE over North America and Eurasia from VIC and satellite observations. Strong negative trends were found in the NOAA satellite observations, during spring and summer in both North American and Eurasian sectors of the Arctic,

**Table 1.** Monotonic trends in spring and summer SCE over North America and Eurasia from VIC and satellite observations (OBS) in the pan-Arctic land region. The significance level (*p*-value) was calculated by two-sided Mann–Kendall test. (The unit of trend slope is in  $yr^{-1}$ .)

		,	VIC	OBS			
		<i>p</i> <	Slope	<i>p</i> <	Slope		
North America	Spring Summer	0.005 0.025	-0.0023 -0.0012	0.005 0.005	-0.0023 -0.0037		
Eurasia	Spring Summer	0.10 0.025	$-0.0006 \\ -0.0006$	0.10 0.005	$-0.0011 \\ -0.0015$		

which are statistically significant (p < 0.025) except that for Eurasia in spring the significance level is p < 0.10. VIC reproduces the same trend directions, with similar significance levels as compared with the satellite observations, for both continents in spring and summer. However, the VIC and observed trend slope magnitudes differ somewhat, especially in summer. The simulated trend slope in summer is about one third of that observed in North America, and 40% of that observed in Eurasia in summer. This discrepancy may be related to uncertainties in the NOAA SCE data set in July and August for both continents (Déry and Brown 2007) and in May and June for North America (Wang *et al* 2005). Because the trend directions and statistical significances are consistent, we chose to include July and August to maintain completeness of the summer period.

The non-parametric Mann–Kendall trend test was also applied to the surface energy inputs to the snow surface, including SNR, SH and LH. Table 2 summarizes their trends over both continents during spring and summer. Trends were computed for seasonal mean fluxes over the snow-covered portion of the regions. Strong positive trends were found in SNR during spring and summer in North America and Eurasia. Similarly, SH fluxes also had statistically significant upward trends, except for Eurasia in spring for which the direction of change was positive, but the trend was not significant at p < 0.10. LH changes were mostly negative in spring to summer, but were statistically significant at p < 0.10 only for North America in summer and Eurasia in spring.

### 3.2. Correlations between observed SCE and modeled surface energy fluxes

The Pearson's product moment correlation coefficient was used to assess relationships between satellite SCE and VICsimulated surface energy fluxes. Figure 3 shows scatterplots of the surface energy fluxes (a) SNR, (b) SH, and (c) LH from VIC against observed SCF over North America and Eurasia in spring and summer. The correlations of surface energy fluxes with SCF as shown in figure 3 are all statistically significant at p < 0.025 (two-sided test). The negative sign indicates that SNR and SH have opposite change directions with SCF. SNR has a relatively stronger relationship with SCF than SH, especially in Eurasia. For LH, the correlations are positive and are stronger in North America than in Eurasia.



**Figure 3.** Scatterplots of surface energy fluxes: (a) SNR, (b) SH, and (c) LH versus satellite SCF over North America and Eurasia during spring and summer. The sign of star means that the correlation has a significance level p < 0.025.

#### 3.3. Role of surface energy fluxes in snow cover changes

In general, the energy balance at a snow surface includes net radiative fluxes, sensible and latent heat fluxes, ground heat fluxes, and the energy transfer due to rain on snow. Over the pan-Arctic, the ground heat flux is a small component of the energy balance of a melting snowpack compared with radiative and turbulent heat fluxes. Therefore, its effects on total snowmelt can safely be ignored (Gray and Prowse 1993). Similarly, rain on snow has an important influence on the water retention characteristics of snow and water movement in the pack but is of minor importance compared with other energy fluxes (Male and Granger 1981). Surface energy fluxes toward the snow surface are defined as positive, therefore the net radiative and sensible heat fluxes usually have positive sign and supply the energy available for snowmelt. Latent heat fluxes are directed away from the snow surface and reduce the melt energy. However, it was not clear which individual component(s) of the snow surface energy budget dominates the significant spring and summer SCE recession since 1972 as shown in table 1. Therefore, we calculated the increment due to monotonic trends in SNR, SH and LH over the 35 yr period as  $\Delta$ SNR,  $\Delta$ SH and  $\Delta$ LH in order to determine the role of each term in the observed downward trends of SCE.

**Table 2.** Trend analyses for spring and summer surface energy fluxes over snow-covered North America and Eurasia generated from VIC. The significance level (*p*-value) was calculated by two-sided Mann–Kendall trend test. (The unit of trend slope is in  $W/m^2/yr$ .)

		S	NR	5	SH	LH		
		<i>p</i> <	Slope	<i>p</i> <	Slope	<i>p</i> <	Slope	
North America	Spring Summer	0.05 0.05	0.1679 0.1965	0.005 0.025	0.1328 0.0940	0.10 0.01	-0.0329 -0.0923	
Eurasia	Spring Summer	0.025 0.005	0.1362 0.2181	0.20 0.025	$0.0288 \\ 0.0700$	0.025 0.20	$-0.0436 \\ -0.0287$	

**Table 3.** Trend analyses for the terms related with SNR during spring and summer over snow-covered North America and Eurasia generated from VIC. The significance level (*p*-value) was calculated by two-sided Mann–Kendall trend test. (The unit of total change during the 35 yr period for fluxes is W m<sup>-2</sup>; the unit for  $T_{min}$  and  $T_{max}$  is °C; no unit for Albedo and RH.)

		SW		DSW		Albedo		DLW		$T_{\min}$		$T_{\rm max}$		RH	
		<i>p</i> <	Δ	<i>p</i> <	Δ	<i>p</i> <	Δ	<i>p</i> <	Δ	<i>p</i> <	Δ	<i>p</i> <	Δ	<i>p</i> <	Δ
North America	Spring Summer	0.025 0.05	7.98 9.45	0.01	-3.83 0.42	0.01 0.025	$-0.04 \\ -0.06$	0.05 0.005	5.68 2.86	0.05 0.025	1.48 0.72	0.05 0.20	1.27 0.46	0.025	-2.41 0.96
Eurasia	Spring Summer	0.05 0.005	5.50 8.31	_	0.16 0.12	0.025 0.005	$-0.03 \\ -0.04$	0.025 0.005	4.83 3.92	0.05 0.005	1.06 0.74	0.05 0.005	1.30 0.90	0.01 0.005	2.56 3.50

Figure 4 shows the relative role of surface energy fluxes averaged over the snow-covered portions of North America and Eurasia in spring and summer. It is apparent that  $\Delta$ SNR is the dominant energy source in both spring and summer, accounting for between 61.8% and 102.3% of the energy available for snow cover changes. The contribution of  $\Delta$ SH plays a secondary role (from 25.3% to 50.0%).  $\Delta$ LH is always opposite in sign with  $\Delta$ SH and  $\Delta$ SNR and almost completely cancels  $\Delta$ SH in North America during summer and Eurasia in spring. However,  $\Delta$ LH has a smaller absolute value than  $\Delta$ SH at other times, such as in North America during spring (-11.8%) and Eurasia in summer (-10.8%). Therefore, we conclude that  $\Delta$ LH has a minor influence on pan-Arctic snow cover changes compared with  $\Delta$ SNR and  $\Delta$ SH.

Figure 5 summarizes the latitudinal variations in spring and summer surface energy fluxes over North America and Eurasia. Basically,  $\Delta$ SNR has a latitudinal pattern, which in general is at a maximum in the lower latitudinal band, and then decreases with latitude poleward. Compared with  $\Delta$ SNR,  $\Delta$ LH shows a similar pattern with a negative sign in most cases. However, it sharply decreases to zero at higher latitudes. In Eurasia during summer, it even has a positive sign for the bands 70–75°N and 75–80°N.

Corresponding to the patterns in  $\Delta$ SNR and  $\Delta$ LH,  $\Delta$ SH is variable depending on the seasons and latitudinal bands. As shown in figure 5(a), the contribution of  $\Delta$ SH is much larger than  $\Delta$ SNR for the 45–50°N latitudinal band in North America, reaching 78% of the total energy attributable to SCE changes. Thereafter, this contribution decreases gradually with latitude to the 70–75°N band. At 75–80°N and 80–85°N,  $\Delta$ SH turns negative. In summer,  $\Delta$ SH over North America is effectively canceled by  $\Delta$ LH with a negative residual from 55–60°N to 70–75°N and a positive one for 75–80°N and 80– 85°N. For Eurasia,  $\Delta$ SH does not follow the same pattern as for North America. Although  $\Delta$ SH in Eurasia is about half of  $\Delta$ SNR for the latitudinal bands 45–50°N and 50–55°N during spring, its effect over the entire domain as shown in figure 4(b) is small because of the small contribution between 55–60°N and  $65-70^{\circ}$ N. In summer, the pattern was mainly dominated by the  $65-70^{\circ}$ N latitudinal band in figure 5(d) because of its area weight (up to 47.4%).

#### 4. Summary and discussion

By exploring long-term trends in satellite observations of SCE, we have shown that North American and Eurasian snow cover over the pan-Arctic declined significantly in spring and summer for the period 1972-2006. Furthermore, longterm means of seasonal SCE and their trend directions are reproduced by the VIC model, which allowed us to diagnose the causes of the observed trends. We have also shown that surface radiative and turbulent heat fluxes simulated in VIC have strong correlations with observed SCE. We find that positive trends in SNR are mostly associated with the observed and model-derived SCE trends. Modeled LH and SH trends associated with warming mostly cancel, except for North America in spring, and to a lesser extent for Eurasia in summer, when the SH contribution to the SCE trends remains substantial. Our results indicate that  $\triangle$ SNR is the primary energy source and  $\Delta$ SH plays a secondary role in changes of SCE. Compared with  $\triangle$ SNR and  $\triangle$ SH,  $\triangle$ LH only has a minor influence on pan-Arctic snow cover changes.

Changes in sensible and latent heat fluxes are mostly dominated by increases in pan-Arctic surface air temperature, which have risen at a rate almost twice as large as the global average in recent decades (Lugina *et al* 2006, Serreze and Francis 2006, Bekryaev *et al* 2010). As shown in table 3, the increases in SNR are mainly associated with increased SW and increased DLW due to warmer atmospheric temperature, whereas emitted upward longwave fluxes do not change much as the snowpack temperature is mostly isothermal during the melt period. Strong upward trends in SW mostly result from statistically significant decreasing trends in snow surface albedo, while the contribution from increased DSW trends is minor (<5%) except for North America during spring when



Figure 4. Relative role of spring and summer surface energy fluxes averaged only over snow-covered (a) North America and (b) Eurasia.



**Figure 5.** Latitudinal profiles of the changes in spring and summer surface energy fluxes over North America and Eurasia snow-covered areas. The blue line shows the weight (fraction of snow cover in each latitudinal band).

DSW decreases. VIC was forced by DSW calculated using the method of Thornton and Running (1999) based on the daily temperature range and vapor pressure. Therefore, there is a decreasing DSW trend in table 3 for North America during spring because daily minimum temperature ( $T_{min}$ ) has increased more rapidly than daily maximum temperature ( $T_{max}$ ) and the relative humidity (RH) has decreased in the mean time.

Aside from the reasons described above, the difference in surface energy fluxes between North America and Eurasia may be related to the data used to force VIC, which generally are of higher quality in North America than Eurasia because of denser observation networks (Niu and Yang 2007), especially for the snow-covered portions during summer. Therefore, further study is warranted to investigate the effects of variations in forcing variables such as precipitation, wind speed and cloud cover (not used in the Thornton and Running method), which affect both surface radiative and turbulent heat fluxes. Furthermore, our study has investigated only aggregate changes over North America and Eurasia; further study is warranted to resolve spatial variations in trends over these large subcontinental areas.

#### Acknowledgments

This work was supported by NASA grants NNX07AR18G and NNX08AU68G to the University of Washington. The authors thank Dr David A Robinson and Mr Thomas Estilow from Rutgers University and Dr Thomas L Mote from the University of Georgia for their assistance with acquisition of the data sets used in the study. The authors also thank Mr Theodore Bohn, and Drs Chunmei Zhu, Huilin Gao and Yanhong Gao of the University of Washington for technical assistance. Dr Stephen J Déry acknowledges support from the Canada Research Chairs program and the Natural Sciences and Engineering Research Council of Canada. We thank an anonymous reviewer and Dr Chris Derksen (Environment Canada) for their thorough and constructive reviews.

#### References

- Adam J C, Clark E A, Lettenmaier D P and Wood E F 2006 Correction of global precipitation products for orographic effects J. Clim. 19 15–38
- Adam J C, Haddeland I, Su F and Lettenmaier D P 2007 Simulation of reservoir influences on annual and seasonal streamflow changes for the Lena, Yenisei, and Ob' rivers *J. Geophys. Res.* 112 D24114
- Adam J C and Lettenmaier D P 2003 Adjustment of global gridded precipitation for systematic bias J. Geophys. Res. 108 D94257
- Adam J C and Lettenmaier D P 2008 Application of new precipitation and reconstructed streamflow products to streamflow trend attribution in northern Eurasia *J. Clim.* 21 1807–28
- Anderson E A 1976 A point energy and mass balance model of a snow cover *Technical Report No. 19* (Silver Spring, MD: National Oceanic and Atmospheric Administration)
- Andreadis K, Storck M P and Lettenmaier D P 2009 Modeling snow accumulation and ablation processes in forested environments *Water Resour. Res.* **45** W05429

- Armstrong R L and Brodzik M J 2001 Recent northern hemisphere snow extent: a comparison of data derived from visible and microwave sensors *Geophys. Res. Lett.* **28** 3673–6
- Armstrong R L and Brodzik M J 2005 Northern Hemisphere EASE-Grid Weekly Snow Cover and Sea Ice Extent Version 3 (Boulder, CO: National Snow and Ice Data Center, Digital Media) accessed May 2010 at http://nsidc.org/data/nsidc-0046. html
- Bekryaev R V, Polyakov I V and Alexeev V A 2010 Role of polar amplification in long-term surface air temperature variations and modern Arctic warming J. Clim. 23 3888–906
- Betts A, Köhler K M and Zhang Y 2009 Comparison of river basin hydrometeorology in ERA-interim and ERA-40 reanalyses with observations J. Geophys. Res 114 D02101
- Boike J, Roth K and Ippisch O 2003 Seasonal snow cover on frozen ground: energy balance calculations of a permafrost site near Ny-Ålesund Spitsbergen J. Geophys. Res. 108 D28163
- Bras R L 1990 *Hydrology: An Introduction to Hydrologic Science* (Boston, MA: Addison-Wesley)
- Brown R D, Derksen C and Wang L 2010 A multi-dataset analysis of variability and change in Arctic spring snow cover extent 1967–2008 J. Geophys. Res. 115 D16111
- Brown R D and Robinson D A 2011 Northern hemisphere spring snow cover variability and change over 1922–2010 including an assessment of uncertainty *Cryosphere* **5** 219–29
- Cherkauer K A and Lettenmaier D P 2003 Simulation of spatial variability in snow and frozen soil field *J. Geophys. Res.* **108** D228858
- Cline D W 1997 Snow surface energy exchanges and snowmelt at a continental midlatitude Alpine site *Water Resour. Res.* **33** 689–701
- Derksen C, Brown R D and Wang L 2010 Terrestrial snow (Arctic) in state of the climate in 2009 *Bull. Am. Meteorol. Soc.* **91** S93–4
- Derksen C and Brown R D 2011 Terrestrial snow (Arctic) in state of the climate in 2010 *Bull. Am. Meteorol. Soc.* **92** S154–5
- Déry S J and Brown R D 2007 Recent northern hemisphere snow cover extent trends and implications for the snow-albedo feedback *Geophys. Res. Lett.* 34 L22504
- Dyer J L and Mote T L 2002 Role of energy budget components on snow ablation from a mid-latitude prairie snowpack *Polar Geogr.* 26 1–29
- Flanner M G, Zender C S, Hess P G, Mahowald N M, Painter T H, Ramanathan V and Rasch P J 2009 Springtime warming and reduced snow cover from carbonaceous particles *Atmos. Chem. Phys.* 9 2481–97
- Frei A and Robinson D A 1999 Northern hemisphere snow extent: regional variability 1972–94 *Int. J. Climatol.* **19** 1535–60
- Gray D M and Prowse T D 1993 Snow and floating ice *Handbook of Hydrology* ed D R Maidment (Columbus, OH: McGraw-Hill)
- Groisman P Y, Karl T R, Knight R W and Stenchikov G L 1994 Changes of snow cover temperature and radiative heat balance over the northern hemisphere J. Clim. 7 1633–56
- Hinzman L D, Kane D L and Gieck R E 1991 Regional snow ablation in the Alaskan Arctic Northern Hydrology: Selected Perspectives ed T D Prowse and C S L Ommaney (Saskatoon, SK: National Hydrology Research Institute)
- Holland M M and Bitz C M 2003 Polar amplification of climate change in coupled models *Clim. Dyn.* 21 221–32
- Kalnay E et al 1996 The NCEP/NCAR 40-year reanalysis project Bull. Am. Meteorol. Soc. 77 437–72
- Kimball J S, Running S W and Nemani R 1997 An improved method for estimating surface humidity from daily minimum temperature Agric. For. Meteorol. 85 87–98
- Koivusalo H and Kokkonen T 2002 Snow processes in a forest clearing and in a coniferous forest J. Hydrol. **262** 145–64
- Kuusisto E 1986 The energy balance of a melting snow cover in different environment *Modeling Snowmelt-Induced Processes IAHS Publication No. 155* (Wallingford: IAHS Press)
- Leathers D J, Graybeal D, Mote T L, Grundstein A J and Robinson D A 2004 The role of airmass types and surface

energy fluxes in snow cover ablation in the central appalachians *J. Appl. Meteorol.* **43** 1887–99

- Liang X, Lettenmaier D P, Wood E F and Burges S J 1994 A simple hydrologically based model of land surface water and energy fluxes for general circulation models *J. Geophys. Res.* 99 14415–28
- Liang X, Wood E F and Lettenmaier D P 1996 Surface soil moisture parameterization of the VIC-2L model: evaluation and modifications *Glob. Planet. Change* **13** 195–206
- Lugina K M, Groisman P Y, Vinnikov K Y, Koknaeva V V and Speranskaya N A 2006 Monthly surface air temperature time series area-averaged over the 30-degree latitudinal belts of the globe, 1881–2005 *Trends Online: A Compendium of Data on Global Change* (Oak Ridge, TN: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy)
- Male D H and Granger R J 1981 Snow surface energy exchange Water Resour. Res. **173** 609–27
- Manabe S and Stouffer R J 1994 Multiple-century response of a coupled ocean–atmosphere model to an increase of atmospheric carbon dioxide *J. Clim.* **7** 5–23
- Mann H B 1945 Nonparametric tests against trend *Econometrica* 13 245–59
- Marks D and Dozier J 1992 Climate and energy exchange at the snow surface in the alpine region of the Sierra Nevada 2 snow cover energy balance *Water Resour. Res.* **28** 3043–54
- Marsh P and Pomeroy J W 1996 Meltwater fluxes at an arctic forest-Tundra site *Hydrol. Process.* **10** 1383–400
- McClelland J W *et al* 2006 A pan-Arctic evaluation of changes in river discharge during the latter half of the 20th century *Geophys. Res. Lett.* **33** L06715
- Miller J R and Russell G 2000 Projected impact of climatic change on the freshwater and salt budgets of the Arctic ocean by a GCM *Geophys. Res. Lett.* **27** 1183–6
- Moore R D and Owens I F 1984 Controls on advective snowmelt in a maritime alpine basin *J. Clim. Appl. Meteorol.* **23** 135–42
- Niu G Y and Yang Z Y 2007 An observation-based formulation of snow cover fraction and its evaluation over large North American river basins *J. Geophys. Res.* **112** D21101
- Ohmura A, Gilgen H and Wild M 1989 *Global Energy Balance Archive* (Zurich: ETH Zurich) accessed July 2010 at www.geba. ethz.ch/
- Peterson B J *et al* 2002 Increasing river discharge to the Arctic ocean Science **298** 2171–3
- Pitman A J *et al* 1999 Key results and implications from phase 1(c) of the project for intercomparison of land-surface parameterization schemes *Clim. Dyn.* **15** 673–84
- Pohl S and Marsh P 2006 Modelling the spatial-temporal variability of spring snowmelt in an arctic catchment *Hydrol. Process.* **20** 1773–92
- Price A J and Dunne T 1976 Energy balance computations of snowmelt in a subarctic area *Water Resour. Res.* **12** 686–94
- Prowse T D and Owens I F 1982 Energy balance over melting snow Craigieburn Range New Zealand J. Hydrol. (N.Z.) **21** 133–47
- Rawlins M A *et al* 2010 Analysis of the Arctic system for freshwater cycle intensification: observations and expectations *J. Clim.* 23 5715–37
- Robinson D A 1986 Initiation of spring snowmelt over Arctic lands *Cold Regions Hydrology Symposium* (Fairbanks, AK: American Water Resources Association)
- Robinson D A 2000 Weekly Northern Hemisphere Snow Maps: 1966–1999 in AMS Conf. Ser. 12 on Applied Climatology (Asheville, NC: American Meteorological Society)

- Robinson D A, Dewey K F and Heim R R 1993 Global snow cover
- monitoring: an update *Bull. Am. Meteorol. Soc.* **74** 1689–96 Sen P K 1968 Estimates of the regression coefficient based on
- Kendall's Tau J. Am. Stat. Assoc. **63** 1379–89
- Serreze M C, Barrett A P, Stroeve J C, Kindig D N and Holland M M 2009 The emergence of surface-based Arctic amplification *Cryosphere* **3** 11–9
- Serreze M C, Bromwich D H, Clark M P, Etringer A J, Zhang T and Lammers R 2002 Large-scale hydro-climatology of the terrestrial Arctic drainage system *J. Geophys. Res.* **107** 8160
- Serreze M C and Francis J A 2006 The Arctic amplification debate *Clim. Change* **76** 241–64
- Serreze M C, Walsh J E, Chapin E C, Osterkamp T, Dyugerov P, Romanovsky M V, Oechel W, Morison C J, Zhang T and Barry R G 2000 Observational evidence of recent change in the northern high-latitude environment *Clim. Change* 46 159–207
- Shi X, Wild M and Lettenmaier D P 2010 Surface radiative fluxes over the pan-Arctic land region: variability and trends J. Geophys. Res. 115 D22104
- Shiklomanov A I, Lammers R B, Rawlins M A, Smith L C and Pavelsky T M 2007 Temporal and spatial variations in maximum river discharge from a new Russian data set J. Geophys. Res. 112 G04S53
- Storck P M, Lettenmaier D P and Bolton S M 2002 Measurement of snow interception and canopy effects on snow accumulation and melt in a mountainous maritime climate Oregon United States *Water Resour. Res.* 38 11223
- Tennessee Valley Authority 1972 Heat and mass transfer between a water surface and the atmosphere *Report* 14 (Tennessee, TN: Water Resources Research Laboratory, Tennessee Valley Authority)
- Thornton P E and Running S W 1999 An improved algorithm for estimating incident daily solar radiation from measurements of temperature, humidity, and precipitation Agric. For. Meteorol. 93 211–28
- Troy T J and Wood E F 2009 Comparison and evaluation of gridded radiation products across northern Eurasia *Environ. Res. Lett.* **44** 045008
- US Army Corps of Engineers 1956 Snow Hydrology: Summary Report of the Snow Investigations (Portland, OR: North Pacific Division, US Army)
- Wang L B, Sharp M, Brown R, Derksen C and Rivard B 2005 Evaluation of spring snow covered area depletion in the Canadian Arctic from NOAA snow charts *Remote Sens*. *Environ.* 95 453–63
- Wiesnet D R, Ropelewsk C F, Kukla G J and Robinson D A 1987 A discussion of the accuracy of NOAA satellite-derived global seasonal snow cover measurements Large Scale Effects of Seasonal Snow Cover IAHS Publication No. 166 ed B E Goodison, R G Barry and J Dozier (Wallingford: IAHS Press)
- Willmott C J and Matsuura K 2009 Terrestrial Air Temperature and Precipitation: Monthly and Annual Time Series (1930–2004) (Newark, DE: University of Delaware) accessed October 2009 at http://climate.geog.udel.edu/~climate/html\_pages/archive. html
- Woo M K, Yang D and Young K L 1999 Representativeness of Arctic weather station data for the computation of snowmelt in a small area *Hydrol. Process.* **13** 1859–70
- Yang D, Robinson D, Zhao Y, Estilow T and Ye B 2003 Streamflow response to seasonal snow cover extent changes in large Siberian watersheds J. Geophys. Res. 108 4578