

LETTER • OPEN ACCESS

Public health impacts of excess NO_x emissions from Volkswagen diesel passenger vehicles in Germany

To cite this article: Guillaume P Chossière *et al* 2017 *Environ. Res. Lett.* **12** 034014

View the [article online](#) for updates and enhancements.

Related content

- [Impact of the Volkswagen emissions control defeat device on US public health](#)
Steven R H Barrett, Raymond L Speth, Sebastian D Eastham *et al.*
- [Impact of excess NO_x emissions from diesel cars on air quality, public health and eutrophication in Europe](#)
J E Jonson, J Borken-Kleefeld, D Simpson *et al.*
- [Global, regional and local health impacts of civil aviation emissions](#)
Steve H L Yim, Gideon L Lee, In Hwan Lee *et al.*

Recent citations

- [Reducing Vehicle Cold Start Emissions through Carbon Pricing: Evidence from Germany](#)
Manuel Frondel *et al*
- [Vanadium Substitution as an Effective Way to Enhance the Redox Ability of Tungstophosphoric Acid and for Application of NH₃-SCR](#)
Qi Liu *et al*
- [S. Crolius *et al*](#)

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED

27 June 2016

REVISED

12 December 2016

ACCEPTED FOR PUBLICATION

16 January 2017

PUBLISHED

3 March 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Public health impacts of excess NO_x emissions from Volkswagen diesel passenger vehicles in GermanyGuillaume P Chossière¹, Robert Malina^{1,2}, Akshay Ashok¹, Irene C Dedoussi¹, Sebastian D Eastham³, Raymond L Speth¹ and Steven R H Barrett^{1,4}¹ Laboratory for Aviation and the Environment, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge MA 02139, United States of America² Center for Environmental Sciences, Hasselt University, Martelarenlaan 42, 3500 Hasselt, Belgium³ Atmospheric Chemistry Modeling Group, Harvard University, 29 Oxford Street, Cambridge MA 02138, United States of America⁴ Author to whom any correspondence should be addressed.E-mail: sbarrett@mit.edu**Keywords:** Volkswagen, air quality, human health, dieselSupplementary material for this article is available [online](#)

Abstract

In September 2015, the Volkswagen Group (VW) admitted the use of ‘defeat devices’ designed to lower emissions measured during VW vehicle testing for regulatory purposes. Globally, 11 million cars sold between 2008 and 2015 are affected, including about 2.6 million in Germany. On-road emissions tests have yielded mean on-road NO_x emissions for these cars of 0.85 g km⁻¹, over four times the applicable European limit of 0.18 g km⁻¹. This study estimates the human health impacts and costs associated with excess emissions from VW cars driven in Germany. A distribution of on-road emissions factors is derived from existing measurements and combined with sales data and a vehicle fleet model to estimate total excess NO_x emissions. These emissions are distributed on a 25 by 28 km grid covering Europe, using the German Environmental Protection Agency’s (UBA) estimate of the spatial distribution of NO_x emissions from passenger cars in Germany. We use the GEOS-Chem chemistry-transport model to predict the corresponding increase in population exposure to fine particulate matter and ozone in the European Union, Switzerland, and Norway, and a set of concentration-response functions to estimate mortality outcomes in terms of early deaths and of life-years lost. Integrated over the sales period (2008–2015), we estimate median mortality impacts from VW excess emissions in Germany to be 1 200 premature deaths in Europe, corresponding to 13 000 life-years lost and 1.9 billion EUR in costs associated with life-years lost. Approximately 60% of mortality costs occur outside Germany. For the current fleet, we estimate that if on-road emissions for all affected VW vehicles in Germany are reduced to the applicable European emission standard by the end of 2017, this would avert 29 000 life-years lost and 4.1 billion 2015 EUR in health costs (median estimates) relative to a counterfactual case with no recall.

1. Introduction

Public health is significantly and globally affected by outdoor air pollution (WHO 2006), with the European Environment Agency (EEA) estimating that outdoor air pollution is responsible for more than 400 000 premature deaths annually in Europe (EEA 2015b). These impacts are driven primarily by population exposure to fine particulate matter with an aerodynamic diameter of 2.5 μm or less (PM_{2.5})

and, to a lesser extent, ozone, both of which have been associated with an increased risk of premature mortality by epidemiological studies (see the review study by Hoek *et al* 2013 and the multi-city analysis in WHO 2013b). Road transportation emissions account for approximately 50% of the total health impacts from ambient air pollution in Europe (OECD 2014).

Vehicle emissions standards initially introduced in 1991 (the so called Euro standards) aimed to reduce the air quality impacts of road transportation

(European Commission 1991). However, observed reductions in ambient pollution have not been as significant as expected, and the EEA noted that ‘emissions in real-life driving conditions are often higher, especially for diesel vehicles, than those measured during the approval test’ (EEA 2015b). A partial explanation for this discrepancy came to light in September 2015 when an investigation by Thompson *et al* (2014) quantified excess emissions from Volkswagen group vehicles, ultimately leading to allegations by the US Environmental Protection Agency (EPA) and admission by VW that certain vehicle models had engines equipped with software designed to reduce emissions during approval testing significantly below levels achieved under actual on-road operation (US EPA 2015).

Consequently, in October 2015 the German Federal Motor Transport Authority (KBA) ordered a mandatory recall of all 2.4 million affected passenger vehicles still in service and stated that they must be modified to comply with European regulations (KBA 2015). Following the KBA order, VW announced that the recall would begin in 2016 with completion originally scheduled for the end of the same year (VW 2015). The first vehicles were recalled at the end of January 2016 (VW 2016). For all three engine sizes affected (1.2 liter, 1.6 liter and 2.0 liter), Volkswagen will update the EA 189 engine software to ensure that on-road emissions stay below the permitted limit. In addition, for the 1.6 liter engine, VW will fit a flow rectifier to the intake duct.

In this study, we estimate excess NO_x emissions from affected cars commercialized by the brands of the Volkswagen Group that used the EA 189 1.2, 1.6 and 2.0 liter engines (Audi, Seat, Skoda, Volkswagen) within Germany, calculating health impacts and health costs throughout Europe which have already occurred between 2008 and 2015, in addition to future outcomes under different recall scenarios. We seek to estimate the number of early deaths as well as the total number of life-years lost which are at risk of resulting from excess emissions. Morbidity impacts are not evaluated or estimated.

2. Methods

Excess emissions in this paper are defined as the difference between vehicle on-road NO_x emissions and the limit value set by the applicable European Emissions Standard, Euro 5. Even though European Union regulations only require vehicles to meet this standard during the type approval test conducted in the laboratory using the New European Driving Cycle, the intent of the regulation is to limit on-road emissions and achieve real-world air quality improvements (European Commission 2007). In the absence of an official factor to account for the difference between laboratory and on-road testing, we use the regulatory limit as the point of reference for our calculations of excess emissions.

For effects occurring in the future, we first estimate impacts assuming all affected VW vehicles in Germany are modified to bring on-road emissions down to Euro 5 standards by the end of 2016, with a fleet recall occurring at a fixed rate over the course of 2016. We then calculate effects for the counterfactual case that affected vehicles remain in the fleet until their retirement without on-road emissions being reduced to Euro 5 levels. The difference between the counterfactual and recall cases yields the benefits of the recall in terms of avoided excess emissions, health impacts and health costs. The paper also quantifies the impact for a case where the recall is performed at a slower rate and completed by the end of 2017 with the same start date. It should be noted here that future health impacts under one of the ‘recall’ scenarios correspond to a reduction of on-road emissions to the Euro 5 limit value. While the absolute decrease in NO_x emissions which would result from a recall is difficult to estimate, any recall, if carried out, would reduce the average NO_x emissions rate from the current estimated value of $\sim 0.85 \text{ g km}^{-1}$ to some lower value. We take the regulatory limit (0.18 g km^{-1}) as a reasonable minimum value. It is possible to calculate the benefits of smaller reductions as a linear combination of the ‘no policy’ scenario and the ‘recall’ scenario. As detailed later in this study, the atmospheric response to excess emissions is essentially linear for the range of perturbations considered.

We augment and extend the modelling capabilities used in Barrett *et al* (2015) to estimate the impact of excess VW NO_x emissions in the US. They used an adjoint-sensitivity-based air quality modeling approach to capture the impacts of spatially disaggregate excess NO_x emissions on aggregate $\text{PM}_{2.5}$ and ozone population exposure in the US. Due to nonlinearity associated with differences in population density and distribution, background atmospheric composition and local meteorology, linear scaling of the results calculated by Barrett *et al* for the US will not necessarily yield an accurate or reliable answer for other countries. In this study we therefore use a chemistry-transport model to capture the impacts of spatially disaggregate excess NO_x emissions in Germany on spatially disaggregate $\text{PM}_{2.5}$ and ozone population exposure in Germany and other European countries. The calculated increase in population exposure due to these excess emissions is then converted into policy-relevant metrics of health impacts and costs.

The air quality and health impacts of the VW excess emissions in Germany are computed using spatially-resolved excess NO_x emissions estimates. Unless otherwise specified, NO_x emissions are reported on an NO_2 mass basis, consistent with the standards used in emissions inventories. Excess emissions are estimated from on-road measurements of the affected models, as well as the number of affected cars and the vehicle kilometers traveled. Their

Table 1. Uncertain parameters used in this study when calculating premature mortalities. Uncertainty in monetization is calculated separately.

Parameter	Central value [95% CI]	Distribution type
Emissions index (g NO _x /L)	16 [5.0, 28]	Truncated normal (bounds: 3.7, 29)
VKT growth rate (2008–2020)	0.005 [0.003, 0.007]	Triangular
VKT growth rate (2020–2040)	0.003 [0.001, 0.004]	Triangular
PM _{2.5} response bias	+11% [−15%, +91%]	Uniform
Ozone response bias	−0.050% [−18%, +35%]	Uniform
PM _{2.5} CRF (risk per 10 μg m ^{−3})	1.11 [1.05, 1.16]	Triangular
Ozone CRF (risk per 10 ppbv)	1.040 [1.013, 1.067]	Triangular

spatial distribution is inferred from the distribution of overall NO_x emissions from passenger cars in Germany, obtained from the German Environmental Protection Agency (UBA 2016). A regional chemistry-transport model is used to relate emissions to pollutant concentrations. Concentration-response functions are applied to convert the resultant population exposure to PM_{2.5} and ozone into health impacts. Given the significant uncertainties in estimating health impacts of air pollution and the associated costs, we use a probabilistic framework which propagates uncertainty through all aspects of the calculation. Public health outcomes are monetized using both the Value of Statistical Life (VSL) and Value of Life Year (VOLY) approach. In this section we describe the uncertainty estimation approach, the excess emissions calculation, the population exposure estimation, health impacts estimation and the monetary valuation of the impacts.

2.1. Uncertainty quantification

Uncertainty in input variables such as vehicle activity and emissions factors (mass of NO_x emitted per kilometer driven) is propagated through the analysis by performing a Monte Carlo simulation. For each sample, a random draw is taken for each of the parameters listed in table 1, according to the distributions shown. The total excess emissions from German light-duty VW vehicles is calculated for each sample based on the randomly drawn emissions factor and on the vehicle-kilometers traveled (VKT) growth factor. These two variables and their distributions are discussed in section 2.3. Total population exposure to PM_{2.5} and ozone is then calculated by interpolating results from the chemistry-transport model GEOS-Chem, described in section 2.4. This includes compensation for bias in the modeled response gradient for both ozone and PM_{2.5}, each drawn as a random variable.

The health impacts of variations in exposure to PM_{2.5} and ozone are calculated by applying concentration response functions (CRFs), as described in section 2.5. For each CRF, the rate of increase in relative risk per unit of exposure is again drawn as a random variable for each sample. The total mortality resulting from the excess emissions for that sample is calculated by summing the premature mortalities

calculated for PM_{2.5} and ozone, assuming independent effects. We perform one thousand independent draws of each variable to yield one thousand independent estimates of the number of premature mortalities and Years of Life Lost (YLL). The rationale for the choice of sample size is presented in the supplementary material (stacks.iop.org/ERL/12/034014/mmedia). The input variables and their distributions are discussed in the following sections and summarized in table 1. In monetizing health impacts, we perform ten additional draws of a distribution of the economic value of statistical life (VSL) and ten draws of a distribution of the Value of Life Year (VOLY), discussed in section 2.6.

Unless explicitly stated otherwise, we report the median value of the output distribution. Data sources and the uncertainties associated with the different parts of our study are detailed in the following subsections.

2.2. Affected vehicle fleet and geographic distribution

An inventory of affected vehicle models in Germany is constructed from publically available sources (Auto, Motor und Sport 2016, ADAC Online Autodatenbank 2016). It comprises cars marketed by the brands of the Volkswagen Group that used the EA 189 engine in its 1.2, 1.6, and 2.0 liter versions (Audi, Seat, Skoda, Volkswagen). We compute the number of affected cars for each relevant model that entered into service each year using annual new registration data published by the German Federal Motor Transport Authority (KBA). In order to obtain estimates for vehicle-kilometers traveled (VKT) by all affected vehicles in each year, the registration data are used as inputs to the Stochastic Transport Emissions Policy (STEP) light-duty vehicle fleet model, developed by Bastani *et al* (2012). We further calibrate the model using Germany-specific data gathered by the TRACCS project (Papadimitriou *et al* 2013). A logistic function is used to estimate the vehicle retirement rate.

The total activity of the affected vehicles is expressed in vehicles-kilometers traveled (VKT) per year. The growth rate of this quantity over time is treated as a random variable and drawn from two triangular distributions, one applicable to the years prior to 2020 and one to the years between 2020 and 2040. The total activity for each year and each sample is estimated

for each 25 km by 28 km grid cell within Germany using a spatially resolved emissions dataset of NO_x emissions from passenger cars, as reported by the German Federal Environmental Protection Agency (UBA 2016).

2.3. Emissions factors for affected vehicles

The estimation of NO_x emissions factors (mass of NO_x emitted per kilometer driven) for the affected vehicles relies on two sets of on-road measurements. The first set of measurements is from KBA (BMVI 2016) using the Real-world Driving Emissions test cycle, as announced by the European Commission's Technical Committee of Motor Vehicles (European Commission 2015). NO_x emission estimates are available for four vehicle models (VW Beetle 2.0L EA 189 Euro 5, VW Golf Plus 1.6L EA 189 Euro 5, VW Passat 2.0L EA 189 Euro 5 and VW Polo 1.2L EA 189 Euro 5). KBA published one measurement per model, yielding four on-road results.

The second set of measurements is from Thompson *et al* (2014), who tested two vehicles—a 2012 Jetta (Vehicle A) and a 2013 Passat (Vehicle B)—on several drive cycles characteristic of different types of driving (urban, rural, highway). Each cycle was driven one or two times for each vehicle, yielding a total of 17 samples.

We combined the results from both sources into a single distribution, composed of 21 samples. Each measured emissions factor expressed in grams of NO_x per kilometer is normalized by the fuel economy of the tested car (in liters of fuel per 100 kilometers). A truncated normal distribution of these factors is used to perform the draws for the Monte Carlo simulation method. The lower bound of the truncation corresponds to the Euro 5 limit of 0.18 g km⁻¹ for the average fuel economy of the fleet of affected vehicles. We obtain the emissions factor of each affected model by multiplying its fuel economy (in L/100 km) by the drawn value in 100 g L⁻¹, yielding a value in g km⁻¹, from which we compute the excess emissions factor. This approach assumes that the emissions factors scale linearly with fuel economy, and that the cars that were tested on-road constitute a representative sample of the fleet of affected vehicles. The excess emissions factor for each vehicle is multiplied by the corresponding VKT sample for this vehicle for a given year, and the sum over all vehicles yields the total amount of excess NO_x emitted each year.

Although VW have published technical details regarding the modifications to be made to each vehicle as part of the recall, at the time of publication there is no reliable data regarding the effect that these modifications have on the NO_x emissions of the affected vehicles. As such, we assume that a recalled vehicle's emissions are brought down to the Euro 5 standard of 0.18 g NO_x/km driven.

2.4. Air quality modeling

The GEOS-Chem chemistry-transport model, originally developed by Bey *et al* (2001) and since

continuously developed and updated, is used to calculate the PM_{2.5} and ozone concentrations. The model domain encompasses most of Europe, covering 15° W – 40° E and 33° N – 61° N. The resolution is 0.25° in latitude and 0.3125° in longitude (approximately 25 km × 28 km). This resolution corresponds to 759 grid cells over Germany.

The GEOS-Chem model has been extensively used at comparable resolutions to capture PM_{2.5} and ozone impacts at the ground level. Numerous studies have evaluated predicted PM_{2.5} concentrations against observations in the US (Duncan Fairlie *et al* 2007, Heald *et al* 2012, Henze *et al* 2009, Leibensperger *et al* 2012, Zhang *et al* 2012), and in Asia (Brauer *et al* 2012, 2015, Jiang *et al* 2015, Kharol *et al* 2013, van Donkelaar *et al* 2010, 2015). Barrett *et al* (2015) used the model over the US domain to predict PM_{2.5} and ozone concentrations. Protonotariou *et al* (2012) used the model over the European domain and evaluated the predicted ozone concentrations in Greece.

We use meteorological data from GEOS-FP, provided by the Global Modeling and Assimilation Office (GMAO) at NASA's Goddard Space Flight Center. Boundary conditions for this nested domain are obtained from a global GEOS-Chem run at 4° × 5° resolution, using the same meteorological source. We use the 2013 GEOS-FP data for all simulations. 2013 was a climatologically average year in Europe, with the average temperature 0.08 °C warmer than the 1995–2015 mean, and 0.16 °C below the mean of the period of interest 2008–2015 (NOAA 2016). We use the EMEP anthropogenic emissions inventory for 2012 and note that emissions reductions between 2012 and 2013 associated with other anthropogenic sources are less than 3% (EMEP 2016). Each simulation is run for a 15-month period with the first 3 months used as model spin up, during which the model is run but the output is not included in the analysis, to ensure that initial conditions do not impact the results.

A model simulation using baseline anthropogenic emissions (excluding excess emissions associated with the VW defeat device) is performed to assess the accuracy of the model in calculating PM_{2.5} and ozone concentrations. Two scenario runs are then performed where a total of 72 and 130 kilotonnes (10⁶ kg) excess NO_x emissions are added to the baseline anthropogenic emissions using the spatial distribution described in section 2.3. Excess NO_x emissions are input in the air quality model as NO. This is consistent with the fact that NO accounts for more than 80% of NO_x emissions from diesel vehicles (Carslaw and Beevers 2005, Yao *et al* 2005), and that the NO_x steady-state in the atmosphere is expected to be reached quickly with respect to the timescale of this study (3 to 30 minutes for NO concentrations of 10 to 1 ppb, following Seinfeld and Pandis 2006). The uncertainty associated with this hypothesis is further discussed in section 4. Differences in the PM_{2.5} and ozone concentrations between the baseline and scenario simulations are

attributed to the computed excess VW emissions. Given that the highest estimate for excess VW NO_x emissions is less than 0.8% of the background anthropogenic NO_x emissions in Germany, we use a linear approach in estimating the PM_{2.5} and ozone impacts of intermediate amounts of excess emissions. One additional scenario run with 520 kilotonnes of annual excess NO_x emissions was conducted in order to verify the validity of this linear approach. This is further discussed in the supplementary material.

Simulated PM_{2.5} and ozone model output concentrations over Germany are validated against observations from the European Environment Agency Air Quality e-Reporting dataset for Germany (EEA 2015a), as well as the entire European domain. Similar to Caiazzo *et al* (2013), the model normalized mean biases are used to account for the uncertainty in predicting PM_{2.5} and ozone concentrations. They are obtained from the point-to-point comparison between available measurements and the model predictions. The resulting distributions have means of 0.11 and -5.0×10^{-4} (95% CI: -0.15 to 0.91 and -0.18 to 0.35) for PM_{2.5} and ozone, respectively. This implies that the model typically overpredicts concentrations of PM_{2.5} by 11%, and underpredicts ozone by 0.050%. The reciprocals of the biases are used as multiplicative factors to correct the GEOS-Chem model predictions in the uncertainty calculations. Only the results from the comparison with the observations in Germany are used in the uncertainty calculations.

2.5. Health impacts

Population exposure to both annual PM_{2.5} and one-hour daily maximum ozone concentrations attributable to excess NO_x emissions from Volkswagen cars within Germany are calculated using the outputs of the GEOS-Chem air quality model described in section 2.4. The annual average PM_{2.5} concentrations are obtained from the hourly output of GEOS-Chem for the species that constitute PM_{2.5} (the GEOS-Chem tracers NH₄, NIT, SO₄, BPCI, BPCO, OCPI, and OCPO). One-hour daily maximum ozone concentrations during ozone season are obtained from the hourly output for the tracer O₃. Population exposure is calculated by multiplying population counts in each grid cell by the annual-average PM_{2.5} or one-hour daily maximum ozone concentration within that grid cell. The spatial distribution of population in Europe is taken from the LandScan database for 2013 (Bright *et al* 2014), which is aggregated from a 1 km resolution to the model grid cells. Country-specific population counts are obtained from the UN World Population Prospects Division for each year of the study (UN 2015) and are used to scale the population density distribution. We take into account the current (2016) member states of the European Union and, due to their geographic proximity, Switzerland and Norway. We use the term 'Europe' to refer to these states and

note that the United Kingdom is still included in future year analyses irrespective of the date the state leaves the EU.

Epidemiological concentration-response functions (CRFs) are applied to estimate premature mortality resulting from population exposure to each species. Cardiovascular premature mortality from long-term exposure to PM_{2.5} is calculated using a log-linear CRF, with a relative risk distribution taken from Hoek *et al* (2013). They find a central relative risk for cardiovascular disease mortality of 1.11 (95% CI: 1.05–1.16) per 10 $\mu\text{g m}^{-3}$ increase in PM_{2.5} exposure. Premature mortality due to exacerbation of both asthma and chronic obstructive pulmonary disease (COPD) as a result of exposure to ozone is also calculated using a log-linear CRF, this time applying the relative risk distribution from an American Cancer Society study (Jerrett *et al* 2009). This CRF has been widely used by previous studies to assess air quality impacts (Barrett *et al* 2015, Jhun *et al* 2015). Jerrett *et al* find a central relative risk of 1.04 (95% CI: 1.013–1.067) per 10 ppb increase in one-hour daily maximum ozone exposure during local ozone season. Country-specific baseline mortality rates for each disease are taken from the World Health Organization global burden of disease database (WHO 2014) and are assumed to remain constant over time. We estimate premature mortality for adults aged 30 and older given that the epidemiological studies are based on a cohort of participants aged 30 years or greater (Jerrett *et al* 2009, Krewski *et al* 2009). The age fraction for each country and each year is taken from UN population forecasts (UN 2015). Alternative PM_{2.5} CRF shapes and parameters, including the integrated exposure-response (IER) function applied in the 2010 Global Burden of Disease study (Burnett *et al* 2014, Lim *et al* 2012), a log-linear CRF from an American Cancer Society study (Krewski *et al* 2009) and an all-cause log-linear CRF reported in Hoek *et al* (2013), are evaluated to determine the sensitivity of the result to the choice of CRF. We also implemented a log-linear ozone CRF derived from the findings of Turner *et al* (2015) that link changes in exposure to the annual average of the daily maximum 8-hour average ozone concentrations to changes in mortality from respiratory and circulatory diseases. These results are presented in the supplementary material. In all cases, a cessation lag structure is applied where 30% of the mortalities due to exposure in a given year occur in the first year, 50% occur equally in years 2 through 5, and the remaining 20% occur equally over years 6 through 20, based on US EPA (2004) recommendations.

The relative risk factors for each of the CRFs are treated as independent, uncertain variables. We assume a triangular distribution with mode and 95% confidence interval taken from the corresponding epidemiological study.

We also compute the number of life-years lost for each estimated number of premature mortalities. The

number of life-years lost is the product of mortalities in each age group (for adults over 30) with the age group's corresponding standard life expectancy, obtained from UN population forecasts for the appropriate year (UN 2015).

2.6. Monetization of health impacts

Mortality effects are valued using two monetization methods that have been widely used in the literature and by government agencies. A detailed overview is given by OECD (2012). The first method relies on estimates for the monetary value of changes in mortality risk due to $\text{PM}_{2.5}$ and ozone exposure, calculated as the Value of a Statistical Life (VSL). The second uses estimates for the Value Of a Life Year lost (VOLY) due to exposure to $\text{PM}_{2.5}$ and ozone. VSL and VOLY estimates are not additive but describe two different approaches for placing a monetary value on the health impacts of air pollution.

A distribution of VSL estimates for Europe is only available in year 2010 USD (OECD 2012), therefore this distribution is converted to 2015 EUR by accounting for the USD/EUR purchasing power parity, changes in economic growth and inflation. The supplementary material contains detailed information on the conversion. We estimate a lower-bound VSL for 2015 of 1.82 million EUR and a higher bound of 5.48 million EUR, to which we fit a triangular distribution with an adjusted OECD base value of 3.65 million EUR as mode. For other years, we adjust VSL distributions for forecast changes in GDP per capita compared to 2015. The supplementary material shows the VSL distributions for all years considered in the analysis. Annual health costs from changes in premature mortality are calculated by multiplying the annual incidences of premature mortality due to excess NO_x emissions with a year-specific VSL estimate, assuming the mortality lag structure recommended by the EPA for air-quality impacts (US EPA 2011).

For VOLY, we use a year-2015 mean value of 133 000 EUR per year of life lost, with a standard deviation of 16 000 EUR derived from recommendations by European agencies (EEA 2014, EC4MACS 2012). For other years, we adjust VOLY distributions for forecast changes in GDP per capita compared to 2015 in the same fashion as for the VSL. Annual health costs from life-years lost due to exposure to $\text{PM}_{2.5}$ and ozone in each year and each sample are calculated by multiplying the number of life-years lost by a year-specific estimate for the VOLY drawn from a year-specific normal distribution. Monetization with the VOLY method is therefore sensitive to age, with early deaths in younger age groups resulting in greater estimated costs than if they occur in older age groups.

Total health costs from historical excess emissions during the years 2008 to 2015 are expressed in year-2015 EUR using a social rate of time preference of 3 percent (discount rate), as recommended by the EU ExternE methodology (Bickel and Friedrich 2005) and

by the US EPA (US EPA 2014). Total health costs because of additional incidences of premature mortality or life-years lost occurring in future years are expressed in year-2015 EUR with the same social rate of time preference as for historical costs. We follow Bickel and Friedrich (2005) and also calculate results for lower and upper bounds of discount rates as sensitivity analyses (0% and 6%, respectively). These alternative discount scenarios are presented in the supplementary material.

3. Results

We report results for excess emissions, health impacts and health costs for historical excess emissions from the affected fleet as well as for future emissions from this fleet.

3.1. Excess emissions

Based on the drive cycle tests by KBA (BMVI 2016) and Thompson *et al* (2014), we estimate the overall NO_x emissions indices for the affected vehicles to be 16 grams per liter fuel consumed. The associated 95% confidence interval (95% CI) is 5.0 to 28 g L^{-1} . The central estimate is equivalent to 0.85 g km^{-1} for the given fleet distribution, compared to the Euro 5 standard of 0.18 g km^{-1} . The total excess emissions for the affected fleet over time are shown in figure 1. The emissions factor is multiplied by the fuel economy of each model affected in order to obtain the mass of NO_x emitted per kilometer driven.

The total vehicle-kilometers traveled (VKT) by the affected vehicles between 2008 and 2015 is 354 billion km (95% CI: 349 to 360), corresponding to emissions of 240 kilotonnes of NO_x (95% CI: 28 to 440) above the total emissions that would have occurred if on-road emissions of all affected vehicles were equivalent to the Euro 5 NO_x limit of 0.18 g km^{-1} . In the absence of a recall that brings on-road emissions down to Euro 5 limit values, future emissions from 2016 onward would be 560 kilotonnes of excess NO_x emissions (95% CI: 67 to 1100), as a result of total expected future VKT for the affected vehicles from 2016 onward of 838 billion km (95% CI: 823 to 854). If, instead, on-road emissions from all affected vehicles are reduced to Euro 5 standards by the end of 2016, future excess NO_x emissions would be reduced to 29 kilotonnes (95% CI: 3.4 to 54). If the recall takes longer and is not complete until the end of 2017, future excess NO_x emissions increase to 59 kilotonnes (95% CI: 7.0 to 110).

Figure 2 represents the spatial distribution of the aggregated excess NO_x emissions between 2008 and 2015. It is assumed to follow the same spatial pattern as passenger cars NO_x emissions, as reported and gridded by UBA for the year 2010 (UBA 2016).




Figure 1. Annual excess NO_x emissions for the affected cars in kilotonnes. The results up to 2015 are estimates of the historical excess NO_x emissions and the results from 2016 onward assume no sales of new vehicles from September 2015 on and no return to recall of affected vehicles. The blue curve represents the median value, and the shaded area the 95% confidence interval.





Figure 2. Spatial distribution of the aggregated excess NO_x emissions over 8 years (2008 to 2015) in tonnes per km^2 based on data provided by UBA (UBA 2016). This figure is at the resolution of the original UBA data, $0.1^\circ \times 0.1^\circ$. It shows the median value of the estimated aggregate excess emissions, namely 240 kilotonnes. Emission density peaks at 11.4 tonnes/ km^2 . For comparison, the UBA inventory for the year 2010 only reports a total of 230 kilotonnes of NO_x emitted from passenger cars, with a peak emission density of 11 tonnes/ km^2 .

3.2. Exposure and health impacts

For each year within the scope of our analysis and each draw of the Monte Carlo simulation, the linearity assumption allows us to compute the increase in the concentrations of $\text{PM}_{2.5}$ and ozone over Europe, as well as the corresponding increase in population exposure. Figures 3(a) and (b) below present an example of the variation of the concen-

tration of $\text{PM}_{2.5}$ and ozone, respectively, corresponding to annual excess emissions of 48 kilotonnes of NO_x , the approximate median value for 2013. Figures 3(c) and (d) show the corresponding population exposure to $\text{PM}_{2.5}$ and ozone respectively, and are obtained by multiplying the concentration of $\text{PM}_{2.5}$ or ozone in each grid cell by the population count in the cell.



The excess NO_x emissions result in reduced ozone concentrations in almost all countries impacted by the excess emissions (with the exception of the region of the Alps between Switzerland and Austria). This is attributed to the high concentration of NO_x relative to volatile organic compound (VOC) concentrations at these locations. Under these conditions, an increase in NO_x emissions inhibits the production of ozone due to the removal of oxidants from the atmosphere via chemical reaction with NO_x (Seinfeld and Pandis 2006). These atmospheric conditions are denoted as VOC-limited chemical regimes and have been established for Europe by previous studies (e.g. Beekman and Vautard 2010, Martin *et al* 2004). In other locations with higher VOC concentrations, it appears that long-range transported NO_x from Germany dominates ozone production.

After applying the concentration response functions to the computed population exposures to PM_{2.5} and ozone, we are able to evaluate the number of premature mortalities that resulted from the estimated excess NO_x emissions.

Table 2 summarizes the health impacts associated with excess emissions in Germany and in Europe. It

features a retrospective analysis for the period 2008 to 2015 and three prospective scenarios, the first one without a recall of the affected cars, the second one with a recall completed by the end of 2016 and the third one with a recall completed by the end of 2017.

The benefits of a recall completed at a constant rate over the course of 2016 are obtained by comparing the scenario assuming that all affected cars are driven during their whole lifecycle without any modification to the scenario where all cars are brought down to the Euro 5 standard by the end of 2016. We also computed the benefit of a recall completed by the end of 2017. The mortality reductions from reduced ozone exposure amount to approximately 10% of the aggregate estimates shown in this table.


Figure 4 presents detailed estimates for the number of early deaths occurring in the ten countries that were most affected by excess emissions released between 2008 and 2015. The results for other European countries, as well as the distribution of the number of life-years lost per country, are presented in the supplementary material.

Table 2. Estimated historical and forecast impacts of excess NO_x emissions. The median estimates are reported with 95% CI in parentheses. Results after 2016 assume no new sales of affected vehicles. Note that the sums of medians may not equal the median of sums. All values are rounded consistently with the order of magnitude of the standard deviation of the distribution considered.

		Germany	Europe (excluding Germany)	Europe
2008 to 2015 estimated impacts	Exposure to PM _{2.5} ^a	14 (1.8; 27)	17 (0.23; 35)	31 (0.41; 62)
	Exposure to ozone ^a	−11 (−13; −2.1)	−4.2 (8.1; −0.50)	−16 (−30; −1.8)
	Early deaths	500 (54; 1200)	660 (71; 1600)	1200 (130; 2800)
	Number of life-years lost	5600 (610; 14 000)	7700 (830; 19 000)	13 000 (1400; 32 000)
	Mortality costs (VSL) ^b	1.5 (0.17; 3.7)	2.0 (0.23; 4.8)	3.5 (0.40; 8.5)
	Mortality costs (VOLY) ^b	0.80 (0.088; 1.9)	1.1 (0.12; 2.6)	1.9 (0.21; 4.9)
Forecast impacts from 2016 assuming no recall	Exposure to PM _{2.5} ^a	34 (4.4; 67)	44 (0.57; 87)	77 (10; 150)
	Exposure to ozone ^a	−27 (−52; 3.2)	−11 (−21; −1.3)	−38 (−73; −4.5)
	Early deaths	1200 (130; 3000)	1600 (180; 4000)	2900 (310; 7000)
	Number of life-years lost	14 000 (1500; 34 000)	19 000 (2100; 46 000)	33 000 (3600; 80 000)
	Mortality costs (VSL) ^b	3.8 (0.44; 9.4)	5.1 (0.58; 12)	8.9 (1.0; 22)
	Mortality costs (VOLY) ^b	2.0 (0.22; 4.8)	2.7 (0.30; 6.6)	4.6 (0.52; 11)
Benefit of a recall completed by the end of 2016	Exposure to PM _{2.5} ^a	32 (4.2; 63)	41 (0.54; 82)	74 (9.6; 150)
	Exposure to ozone ^a	−26 (−49; −3.1)	−10 (−20; −1.2)	−36 (−69; −4.3)
	Early deaths	1200 (130; 2800)	1600 (170; 3800)	2700 (300; 6600)
	Number of life-years lost	13 000 (1400; 32 000)	18 000 (2000; 44 000)	31 000 (3400; 76 000)
	Mortality costs (VSL) ^b	3.6 (0.42; 8.8)	4.8 (0.55; 12)	8.4 (0.97; 21)
	Mortality costs (VOLY) ^b	1.9 (0.20; 4.5)	2.5 (0.28; 6.2)	4.4 (0.49; 11)
Benefit of a recall completed by the end of 2017	Exposure to PM _{2.5} ^a	30 (4.0; 60)	39 (0.51; 78)	69 (9.1; 140)
	Exposure to ozone ^a	−25 (−47; −2.9)	−9.7 (−18; −1.2)	−34 (−65; −4.1)
	Early deaths	1100 (120; 2700)	1500 (160; 3600)	2600 (280; 6200)
	Number of life-years lost	12 000 (1400; 30 000)	17 000 (1900; 42 000)	29 000 (3200; 72 000)
	Mortality costs (VSL) ^b	3.4 (0.39; 8.3)	4.5 (0.52; 11)	7.9 (0.91; 19)
	Mortality costs (VOLY) ^b	1.7 (0.19; 4.2)	2.4 (0.27; 5.8)	4.1 (0.46; 10)

^a Exposure is in million of people over 30 years old times $\mu\text{g m}^{-3}$ (or times ppbv for ozone).

^b Mortality costs are expressed in billion 2015 EUR.



3.3. Health costs

Monetized health impacts of the excess NO_x emissions are shown in table 2. Costs calculated using the VOLY method are approximately 50%

lower on average than those calculated using the VSL, with an average estimate of 11 life-years lost per premature mortality. This difference is due to the prevalence of cardiovascular and respiratory disease

in the older population, resulting in the majority of air-quality-related mortalities occurring in higher age brackets with lower life expectancies (in terms of life-years remaining).

Using the VOLY method, we find that past excess emissions have already caused median health costs of 1.9 billion EUR, and that median additional costs of 4.6 billion EUR are expected under a 'no recall' scenario. 95% of these costs are avoided if the recall is completed by the end of 2016; this is reduced to 90% if the recall is not completed until the end of 2017. The costs presented here do not take into account non-mortality impacts such as new cases of chronic bronchitis or increased hospital admissions, but these are expected to be small relative to costs associated with premature mortality (Barrett *et al* 2015).

Regardless of the valuation method used, we find that the majority of monetary damages are incurred outside of Germany, with approximately 60% of all mortality costs exported to other countries within the area of interest. This implies that there might also be significant health costs within Germany due to diesel cars operating outside of Germany which are not accounted for in this study.

4. Discussion and conclusion

We compare our results to an existing study by Oldenkamp *et al* (2016) which estimates Europe-wide health impacts of VW excess emissions using a simplified modeling approach. The authors calculate point estimates for health impacts from spatially unresolved, constant relationships between excess emissions and mortality impacts. Using this approach, the authors link estimated historical excess NO_x emissions of 491.7 kilotonnes to 5000 early deaths, and estimated future emissions in the absence of vehicle modifications of 802.2 kilotonnes to an additional 8200 early deaths. We use a chemistry-transport model to estimate spatially resolved population exposure in each European country as a result of the additional NO_x emissions in Germany. Using this approach yields an estimate for premature mortality per unit of excess emissions that is 50% lower than the value used in Oldenkamp *et al* (2016), i.e. 5.1 vs. 10.2 early deaths per kilotonne of excess emissions.

In the case of the US, Barrett *et al* (2015) report median excess NO_x emissions of 36.7 kilotonnes for 2008 from 2015 from 482 000 affected cars sold, leading to 59 additional early deaths at the median. Similarly, Oldenkamp *et al* (2016) found 33.8 kilotonnes of excess NO_x over the same period, yielding 59 premature mortalities. Holland *et al* (2015), found 45.1 kilotonnes of NO_x and 46.1 mortalities. In our study, we find that while the number of VW cars sold in Germany is 440% higher than the number sold in the US, excess emissions in Germany are 540% higher

than in the US, which is a result of annual kilometer driven per vehicle in Germany being 19 percent higher than in the US. We note however that our estimate is for the average diesel vehicle in Germany, whereas Barrett *et al* (2015) compute the annual mileage of an average American vehicle.

Further comparing the results of these studies to our results, we find that each unit of NO_x emitted in Europe results in 5 times as many premature mortalities (per capita) as in the US. This difference can be explained by the combination of Europe's greater population density and its more NO_x-sensitive background conditions. A study by Koo *et al* (2013) found that in Europe, a combination of high ammonia emissions and relatively low sulfur emissions results in a higher sensitivity of population-weighted PM_{2.5} to NO_x emissions than any other region in the world apart from Eastern China, where population densities are comparable to or greater than Europe. This aspect is discussed further in the supplementary material.

4.1. Limitations

Some sources of uncertainty in this study have not been quantified. The fleet model does not account for possible differences in distance traveled by vehicle model considered. Vehicle kilometers traveled for a given year are assumed to depend only on the year and the age of the vehicle. However, we note that the TRACCS report (Papadimitriou *et al* 2013) distinguished between four categories of diesel passenger cars (small, lower-medium, upper-medium, executive) and found no significant differences among these categories in terms of vehicle kilometers traveled, when controlling for year and vehicle age (less than 5% for the average vehicle). We also assume that the spatial distribution of the excess NO_x emissions and population remains constant over time.

The excess NO_x emissions are input into the air quality model as pure NO. While it is usually considered that NO accounts for more than 80% of NO_x emissions from diesel vehicles (Carslaw and Beevers 2005, Yao *et al* 2005), it appears that the NO₂ to NO_x ratio in primary emissions from diesel vehicles is increasing over time (Alvarez *et al* 2008). However, the NO_x steady-state is reached in about 3 to 30 minutes for ambient NO concentrations of 10 ppb to 1 ppb respectively (Seinfeld and Pandis 2006), which means that the initial NO₂ to NO_x ratio is not expected to significantly influence PM_{2.5} production. Ashok *et al* (2013) have shown in the case of aviation that the maximum sensitivity of population PM_{2.5} exposure to the NO₂ to NO_x ratio in emissions was 2.75%. Since PM_{2.5} is the main driver of the health impacts estimated in this study, we do not expect our results would vary significantly if we modified the NO₂ to NO_x ratio in the primary emissions. As for ozone, the fact that excess emissions are input as pure NO results in increased nighttime titration of ozone. However, we find that this does not occur during daylight hours due

to the high background ozone concentrations. Since the ozone health impact metrics are based on exposure to 1-hour daily maximum and 8-hour daily maximum ozone concentrations which occur during the daylight hours, the calculated ozone impacts are insensitive to the primary NO_2 to NO_x ratio.

We assume the toxicity of different PM species to be equal, consistent with standard practice. However, it is worth noting that the particle composition of $\text{PM}_{2.5}$ is suspected to influence the related health impacts, as pointed out by Hoek *et al* (2013) and WHO (2013b). Since NO_x emissions impact ammonium nitrate most strongly, any differential toxicity of ammonium nitrate relative to the basket of urban PM for which CRFs are derived is not captured. The standard GEOS-Chem chemical mechanism does not include a detailed secondary organic aerosol model, and these compounds are therefore not included in our health impact assessment.

Although some epidemiological studies have been published which find a link between increased exposure to NO_2 and premature mortality independent of exposure to other pollutants, there remains substantial disagreement with regard to the specific health outcomes affected and their magnitude. If we include the health impacts of changes in exposure to NO_2 due to VW excess emissions following the recommendations of the HRAPIE group (WHO 2013a), we find that excess NO_x emissions in Germany released between 2008 and 2015 caused 1 300 (95% CI: 160 to 2800) additional premature mortalities due to exposure to NO_2 , including 1 200 in Germany. The associated health costs are estimated to be 2 billion EUR (median VOLY estimate). Future emissions in the absence of modification of affected cars are expected to cause 3200 additional premature mortalities from exposure to NO_2 in Europe (including 2800 in Germany) and 5.2 billion EUR in associated health costs. Including NO_2 impacts would approximately double the total impacts estimated earlier, with Germany bearing the great majority of this additional burden. It should also be noted that NO_2 is a precursor to ozone and secondary particulate matter (these effects are accounted for in our study).

When using a CRF derived from the findings of Turner *et al* (2015), we observed greater reductions in mortality due to reduced ozone exposure than our central estimate (derived from Jerrett *et al* 2009). Integrated excess emissions over the period 2008 to 2015 are estimated to have averted 28 premature mortalities (95% CI: 1.7 to 85) with the Jerrett *et al* 2009 CRF, while the Turner CRF predicts 78 avoided mortalities (95% CI: 0.28 to 640). The total impacts in Europe of past excess emissions (from exposure to both $\text{PM}_{2.5}$ and O_3) using the Turner CRF as the central estimate would then be 1000 premature mortalities (95% CI: 93 to 2700). This result is to be compared to the 1,200 mortalities obtained with our central CRF (95 % CI: 130 to 2800). Using the Turner

CRF as the central ozone estimate, we find that future emissions in the absence of any modification to the affected cars are predicted to cause 2600 (95% CI: 230 to 6 700) additional premature mortalities in Europe. This is to be compared to 2900 (95% CI: 310 to 7 000) additional premature mortalities estimated with the Jerrett CRF as our central ozone estimate. In conclusion, changing our central estimate for ozone-related health impacts from Jerrett *et al* (2009) to Turner *et al* (2015) yields 10% lower mortality estimates. The supplementary material provides results obtained with different $\text{PM}_{2.5}$ CRFs.

In addition, our assessment does not include morbidity impacts, which would increase the health costs of the estimated excess emissions but typically account for less than 10% of total health costs associated with air quality impacts (US EPA 2011). It should also be mentioned here that although the epidemiological studies used here and in the supplementary material are widely used in the literature (Barrett *et al* 2015 for instance), they represent only a small portion of the health literature on $\text{PM}_{2.5}$ and ozone and mortality.

We also note that no potential environmental benefits of excess NO_x emissions are included in the analysis. An example could be the reduced use of diesel exhaust fluid in the selective catalytic reduction, which could lead to a lower risk of ammonia leakage, a phenomenon which has been known to contribute to the $\text{PM}_{2.5}$ health impacts of the road transportation sector at approximately equal magnitude as NO_x (Dedoussi and Barrett 2014).

Finally, we note that the assessment of model performance and the corresponding correction by reciprocal biases use the baseline scenario and the officially reported emissions data. These might not fully account for real on-road emissions, and this issue might introduce a second-order effect that is not accounted for. However, given the magnitude of the perturbation (less than 0.8% of total NO_x emissions), this effect is not expected to alter our results significantly.

4.2. Wider implications

In January 2016, and following orders by the KBA (2015), VW started to recall affected vehicles with the aim of reducing their NO_x emissions to an as-yet unspecified level. The modification costs per car have been reported to amount to 60 EUR, excluding the value added tax (FAZ 2016). We assume that the vehicle modification can be completed within 1 hour, including time required to travel to and from the modification site. Applying an average value-of-time of 40 EUR per hour (Korzhenevych *et al* 2014), modification of one vehicle incurs a total cost of approximately 100 EUR. The overall cost of the recall would then be 240 million EUR. This estimate does not include other costs such as increased fuel expenses or social costs of additional carbon emissions if the vehicle modification results in a reduction in fuel

economy. Assuming that the modification reduces on-road emissions to the Euro 5 standard, the total cost of the recall amounts to 5% of the median 4.4 billion EUR health costs (using the VOLY approach) which we estimate will be avoided if the recall is completed by the end of 2016. Our analysis suggests that the recall will remain net beneficial as long as the per-vehicle cost is below 1800 EUR.

Although this analysis was centered on Volkswagen-manufactured cars operated in Germany, NO_x emissions from diesel vehicles in excess of the existing standards constitute a broader and ongoing issue in the European Union. On-road measurements have shown that vehicles from several other manufacturers emit more NO_x than the applicable limits (Carslaw *et al* 2011, Franco *et al* 2014, BMVI 2016), despite the absence of any identified measures intended to circumvent emissions tests. These discrepancies have motivated the development of on-road emissions testing procedures (Weiss *et al* 2013, European Commission 2015) which were approved by the European Council in February 2016. The VW vehicles modeled in this study comprise 2.6 million of the 13 million diesel passenger vehicles operating in Germany in 2012 (Eurostat 2016). Consequently, there will be additional health impacts due to excess emissions from vehicles produced by other manufacturers that are not accounted for in our study. In addition, there were a total of 86 million diesel passenger vehicles operating in the EU28 in 2010 (Papadimitriou *et al* 2013), making excess diesel NO_x emissions a public health concern for all of Europe. On the modeling side, current road transport NO_x inventories are also likely to be affected, if they are based on measurements from laboratory test cycles (e.g. EEA 2013), given that emissions from real-world driving conditions could be significantly different than those measured during a laboratory test cycle (Franco *et al* 2014).

Acknowledgments

We would like to thank the German Umweltbundesamt, in particular Mr Stefan Feigenspan and Mr Michael Kotzulla, for their valuable help in obtaining data on the spatial distribution of NO_x emissions in Germany.

References

- Allgemeiner Deutscher Automobil Club (ADAC, German automobile club) 2016 Autodatenbank (www.adac.de/infotestrat/autodatenbank/Autodaten/default.aspx) (in German)
- Alvarez R, Weilenmann M and Favez J-Y 2008 Evidence of increased mass fraction of NO₂ within real-world NO_x emissions of modern light vehicles—derived from a reliable online measuring method *Atmos. Environ.* **42** 4699–4707
- Ashok A, Lee I H, Arunachalam S, Waitz I A, Yim S H L and Barrett S R H, 2013. Development of a response surface model of aviation's air quality impacts in the United States *Atmos. Environ.* **77** 445–52
- Auto, Motor und Sport 2016 All about the VW scandal, 28 January 2016 (www.auto-motor-und-sport.de/news/ea189-motor-mit-betrugssoftware-diese-autos-sind-betroffen-9982008.html) (in German)
- Barrett S R H, Speth R L, Eastham S D, Dedoussi I C, Ashok A, Malina R and Keith D W 2015 Impact of the Volkswagen emissions control defeat device on US public health *Environ. Res. Lett.* **10** 114005
- Bastani P, Heywood J B and Hope C 2012 The effect of uncertainty on US transport-related GHG emissions and fuel consumption out to 2050 *Transport. Res. A-Pol* **46** 517–48
- Beekmann M and Vautard R 2010 A modelling study of photochemical regimes over Europe: robustness and variability *Atmos. Chem. Phys.* **10** 10067–84
- Bey I, Jacob D J, Yantosca R M, Logan J A, Field B D, Fiore A M, Li Q, Liu H Y, Mickley L J and Schultz M G 2001 Global modeling of tropospheric chemistry with assimilated meteorology: model description and evaluation *J. Geophys. Res.* **106** 23073–95
- Bickel P and Friedrich R (ed) 2005 Impact pathway approach: monetary valuation *ExternE—Externalities of Energy: Methodology 2005 Update* (Luxembourg: European Commission) ch 7
- BMVI Bundesministerium für Verkehr und digitale Infrastruktur 2016 Bericht der Untersuchungskommission 'Volkswagen'
- Brauer M *et al* 2012 Exposure assessment for estimation of the global burden of disease attributable to outdoor air pollution *Environ. Sci. Technol.* **46** 652–60
- Brauer M *et al* 2015 Ambient air pollution exposure estimation for the Global Burden of disease 2013 *Environ. Sci. Technol.* **50** 79–88
- Bright E A, Coleman P R, Rose A N and Urban M L 2014 *LandScan 2013™ High Resolution global Population Data Set* (www.ornl.gov/landscan/)
- Burnett R T *et al* 2014 An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure *Environ. Health Perspect.* **122** 397–403
- Caiazzo F, Ashok A, Waitz I A, Yim S H L and Barrett S R H 2013 Air pollution and early deaths in the United States. Part I: Quantifying the impact of major sectors in 2005 *Atmos. Environ.* **79** 198–208
- Carslaw D C and Beevers S D 2005 Estimations of road vehicle primary NO₂ Exhaust emission fractions using monitoring data in London *Atmos. Environ.* **39** 167–77
- Carslaw D C, Beevers S D, Tate J E, Westmoreland E J and Williams M L 2011 Recent evidence concerning higher NO_x emissions from passenger cars and light duty vehicles *Atmos. Environ.* **2011** 45 7053–63
- Dedoussi I C and Barrett S R H 2014 Air pollution and early deaths in the United States. Part II: Attribution of PM_{2.5} exposure to emissions species, time, location and sector *Atmos. Environ.* **99** 610–7
- Duncan Fairlie T, Jacob D J and Park R J 2007 The impact of transpacific transport mineral dust in the United States *Atmos. Environ.* **41** 1251–66
- EC4MACS European Consortium for Modelling of Air Pollution and Climate Strategies 2012 eds M Holland, A Wagner and A Hunt, The ALPHA Benefit Assessment Model, March 2012
- EEA European Environment Agency 2013 *EMEP Technical Guidance to Prepare National Emission Inventories* (Luxembourg: European Environment Agency) Publications Office (<http://publications.europa.eu/10.2800/92722>)
- EEA 2014 Costs of air pollution from European industrial facilities 2008–2012—an updated assessment, *EEA Technical Report* No 20/2014, Publications Office of the European Union
- EEA 2015a Air Quality e-Reporting, Data for Germany, 2013, published on May 29, 2015, archived on May 16, 2016 (www.eea.europa.eu/data-and-maps/data/aqereporting#tab-data-by-country) (Accessed: 12 May 2016)

- EEA 2015b *Air Quality in Europe—2015 Report*, EEA Report No 5/2015, Publications Office of the European Union
- EMEP European Monitoring and Evaluation Programme 2016 Centre on Emission Inventories and Projections, WebDab Emissions Database, last updated May 30, 2016 (www.ceip.at/ms/ceip_home1/ceip_home/webdab_emepdatabase/reported_emissiondata/) (Accessed: 1 June 2016)
- European Commission 1991 Council Directive 91/441/EEC of 26 June 1991 amending Directive 70/220/EEC on the approximation of the laws of the Member States relating to measures to be taken against air pollution by emissions from motor vehicles (<http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:31991L0441&from=EN>) (Accessed: 19 June 2016)
- European Commission 2007 Regulation (EC) No 715/2007 of the European Parliament and of the Council of 20 June 2007 on type approval of motor vehicles with respect to emissions from light passenger and commercial vehicles (Euro 5 and Euro 6) and on access to vehicle repair and maintenance information (<http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2007:171:0001:0016:EN:PDF>) (Accessed: 30 September 2016)
- European Commission 2015 Press release: ‘Commission welcomes Member States’ agreement on robust testing of air pollution emissions by cars’, October 28, 2015 (http://europa.eu/rapid/press-release_IP-15-5945_en.htm) (Accessed: 23 May 2016)
- Eurostat 2016 Custom data obtained from the road transportation database (<http://ec.europa.eu/eurostat/web/transport/data/database>)
- Franco V, Sánchez F P, German J and Mock P 2014 Real-world exhaust emissions from modern diesel cars (www.automotive-online.nl/upload/files/feitencijfers/1443626509ICCT_PEMS-study_diesel-cars_20141013.pdf)
- Frankfurter Allgemeine Zeitung (FAZ) 2016 VW zahlt Werkstätten 60 Euro pro Software-Update (www.faz.net/aktuell/wirtschaft/vw-abgasskandal/nach-vw-abgasskandal-startet-rueckrufaktion-manipulierter-autos-14039092.html) (Accessed: 2 June 2016) (in German)
- Heald C L *et al* 2012 Atmospheric ammonia and particulate inorganic nitrogen over the United States *Atmos. Chem. Phys. Discuss.* **12** 10295–312
- Henze D K, Seinfeld J H and Shindell D T 2009 Inverse modeling and mapping US air quality influences of inorganic PM 2.5 precursor emissions using the adjoint of GEOS-chem. *Atmos. Chem. Phys.* **9** 5877–903
- Hoek G, Krishnan R M, Beelen R, Peters A, Ostro B, Brunekreef B and Kaufman J D 2013 Long-term air pollution exposure and cardio- respiratory mortality: a review *Environ. Health* **12** 43
- Holland S P, Mansur E, Muller N and Yates A 2015 Damages and expected deaths due to excess NO_x emissions from 2009–2015 Volkswagen diesel vehicles *Environ. Sci. Technol.*
- Jerrett M, Burnett R T, Pope C A, Ito K, Thurston G, Krewski D, Shi Y, Calle E and Thun M 2009 Long-term ozone exposure and mortality *New Engl. J. Med.* **360** 1085–95
- Jhun I, Coull BA, Schwartz J, Hubbell B and Koutrakis P 2015 The impact of weather changes on air quality and health in the United States in 1994–2012 *Environ. Res. Lett.* **10** 084009
- Jiang X *et al* 2015 Revealing the hidden health costs embodied in Chinese exports *Environ. Sci. Technol.* **49** 4381–8
- KBA Kraftfahrtbundesamt 2015 Kraftfahrt-Bundesamt ordnet den Rückruf von 2, 4 Millionen Volkswagen an, Press release October 16, 2015, Flensburg
- Kharol S K, Martin R V, Philip S, Vogel S, Henze D K, Chen D, Wang Y, Zhang Q and Heald C L 2013 Persistent sensitivity of Asian aerosol to emissions of nitrogen oxides *Geophys. Res. Lett.* **40** 1021–26
- Koo J, Wang Q, Henze D K, Waitz I and Barrett S R H 2013 Spatial sensitivities of human health risk to intercontinental and high-altitude pollution *Atmos. Environ.* **71** 140–7
- Korzhenyevych A, Dehnen N, Bröcker J, Holtkamp M, Meier H, Gibson G, Varma A and Cox V 2014 Update of the handbook on external costs of transport. European Commission DG MOVE
- Krewski D *et al* 2009 Extended follow-up and spatial analysis of the American Cancer Society study linking particulate air pollution and mortality *HEI Research Report 140* (Boston, MA: Health Effects Institute)
- Leibensperger E *et al* 2012 Climatic effects of 1950–2050 changes in US anthropogenic aerosols-part 1: aerosol trends and radiative forcing *Atmos. Chem. Phys.* **12** 3333–48
- Lim S S *et al* 2012 A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010 *Lancet* **380** 2224–60
- Martin R V, Fiore A M and Van Donkelaar A 2004 Space-based diagnosis of surface ozone sensitivity to anthropogenic emissions *Geophys. Res. Lett.* **31** L06120
- NOAA 2016 Climate at a Glance National Centers for Environmental Information (NCEI) (www.ncdc.noaa.gov/cag/time-series/global/europe/land/ytd/12/1995-2015)
- OECD 2012 Recommended value of a statistical life: numbers for policy analysis *Mortality Risk Valuation in Environment, Health and Transport Policies* (Paris: OECD Publishing) ch 6 (<https://doi.org/10.1787/9789264130807-en>)
- OECD 2014 Reviewing the evidence of and calculating the cost of the health impacts of air pollution *The Cost of Air Pollution: Health Impacts of Road Transport* (Paris: OECD Publishing) ch2 (<https://doi.org/10.1787/9789264210448-en>)
- Oldenkamp R, Van Zelm R and Huijbregts M A J 2016 Valuing the human health damage caused by the fraud of Volkswagen *Environ. Pollut.* **212** 121–7
- Papadimitriou G, Ntziachristos L, Wüthrich P, Notter B, Keller M, Fridell E, Winnes H, Styhre L and Sjödin A 2013 *Transport data collection supporting the quantitative analysis of measures relating to transport and climate change (TRACCS)* final report prepared for the Directorate-General for Climate Action, European Commission
- Protonotariou A P, Bossioli E, Tombrou M, Mihalopoulos N, Biskos G, Kalogiros J, Kouvarakis G and Amiridis V 2012 Air pollution in eastern mediterranean: Nested-Grid GEOS-CHEM model results and airborne observations ed G Costas Helmis and T Panagiotis *Nastos Advances in Meteorology, Climatology and Atmospheric Physics* (Berlin Heidelberg: Springer) pp 1203–9 (http://doi.org/10.1007/978-3-642-29172-2_168)
- Seinfeld J H and Pandis S N 2006 *Chemistry of the troposphere Atmospheric Chemistry and Physics: From air Pollution to Climate Change* (Hoboken, NJ: John Wiley and Sons) ch 6
- Thompson G J, Carder D K, Besch M C, Thiruvengadam A and Kappanna H K 2014 In-use emissions testing of light-duty diesel vehicles in the United States (www.theicct.org/sites/default/files/publications/WVU_LDDV_in-use_ICCT_Report_Final_may2014.pdf)
- Turner M C *et al* 2015 Long-term ozone exposure and mortality in a Large prospective study *Am. J. Respir. Crit. Care Med.* **193** 1134–42
- UBA Umweltbundesamt 2016 NO_x emissions from passenger cars for year 2010, gridded with the tool GRETA, custom data obtained from UBA in 2016
- UN 2015 *World Population Prospects: The 2015 Revision, custom data acquired via website* United Nations, Department of Economic and Social Affairs, Population Division 2015 (<http://esa.un.org/unpd/wpp/DataQuery/>)
- US EPA 2004 Advisory Council on Clean Air Compliance Analysis Response to Agency Request on Cessation Lag (Washington, DC)
- US EPA 2011 The Benefits and Costs of the Clean Air Act: 1990 to 2020 Final report of US Environmental Protection Agency Office of Air and Radiation

- US EPA 2014 Guidelines for Preparing Economic Analysis National Center for Environmental Economics Office of Policy (<http://yosemite.epa.gov/ee/epa/eed.nsf/webpages/Guidelines.html>)
- US EPA 2015 Notice of Violation (18 September 2015) sent by EPA to Volkswagen Group of America, Inc. (www.epa.gov/sites/production/files/2015-10/documents/vw-nov-cao-09-18-15.pdf)
- van Donkelaar A, Martin R V, Brauer M, Kahn R, Levy R, Verduzco C and Villeneuve P J 2010 Global estimates of ambient fine particulate matter concentrations from satellite-based aerosol optical depth: development and application *Environ. Health Perspec.* **118** 847–55
- van Donkelaar A, Martin R V, Brauer M and Boys B L 2015 Use of satellite observations for long-term exposure assessment of global concentrations of fine particulate matter *Environ. Health Perspec.* **123** 135–43
- VW Volkswagen 2015 Technical measures for the EA 189 diesel engines affected presented to the German Federal Motor Transport Authority, Press Release, Nov 25 2016, Wolfsburg
- VW Volkswagen 2016 Volkswagen making progress as planned with first wave of technical measures for diesel engines, press release, Feb 18 2016, Wolfsburg
- Weiss M, Bonnel P, Hummel R and Steininger N 2013 A complementary emissions test for light-duty vehicles: assessing the technical feasibility of candidate procedures *Report EUR 25572 EN. European Commission Joint Research Centre*
- WHO 2006 Health Risks of Particulate Matter from Long-range Transboundary Air Pollution (www.euro.who.int/__data/assets/pdf_file/0006/78657/E88189.pdf)
- WHO 2013a Health risks of air pollution in Europe – HRAPIE project (http://www.euro.who.int/__data/assets/pdf_file/0006/238956/Health-risks-of-air-pollution-in-Europe-HRAPIE-project,-Recommendations-for-concentration-response-functions-for-cost-benefit-analysis-of-particulate-matter,-ozone-and-nitrogen-dioxide.pdf)
- WHO 2013b Review of evidence on health aspects of air pollution – REVIHAAP project: technical report. Copenhagen, WHO Regional Office for Europe (www.euro.who.int/__data/assets/pdf_file/0004/193108/REVIHAAP-Final-technical-report.pdf)
- WHO 2014 Disease and Injury Country Mortality Estimates for 2012 (World Health Organization)
- Yao X, Lau N T, Chan C K and Fang M 2005 The use of tunnel concentration profile data to determine the ratio of NO₂/NO_x directly emitted from vehicles *Atmos. Chem. Phys. Discuss.* **5** 12723–40
- Zhang L, Jacob D J, Knipping E M, Kumar N, Munger JW, Carouge C C, van Donkelaar A, Wang Y X and Chen D 2012 Nitrogen deposition to the United States: distribution, sources, and processes *Atmos. Chem. Phys.* **12** 4539–54