

LETTER • OPEN ACCESS

## Drinking water quality from rural handpump-boreholes in Africa

To cite this article: D J Lapworth *et al* 2020 *Environ. Res. Lett.* **15** 064020

View the [article online](#) for updates and enhancements.

You may also like

- [How does gendered vulnerability shape the adoption and impact of sustainable livelihood interventions in an era of global climate change?](#)  
Maia Call and Samuel Sellers
- [Assessment of land use change in the Thuma forest reserve region of Malawi, Africa](#)  
Mihla Phiri and Harrington Nyirenda
- [Evidence, ideology, and the policy of community management in Africa](#)  
Luke Whaley, Donald John MacAllister, Helen Bonsor et al.



The Breath Biopsy® Guide  
Fourth edition

FREE

DOWNLOAD THE FREE E-BOOK

BREATH BIOPSY

OWLSTONE MEDICAL

## Environmental Research Letters



## LETTER

## Drinking water quality from rural handpump-boreholes in Africa

## OPEN ACCESS

## RECEIVED

27 November 2019

## REVISED

10 March 2020

## ACCEPTED FOR PUBLICATION

16 March 2020

## PUBLISHED

27 May 2020

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



D J Lapworth<sup>1,12</sup> , A M MacDonald<sup>2</sup> , S Kebede<sup>3</sup>, M Owor<sup>4</sup>, G Chavula<sup>5</sup>, H Fallas<sup>2</sup>, P Wilson<sup>6</sup>, J S T Ward<sup>1</sup> , M Lark<sup>7</sup>, J Okullo<sup>4</sup>, E Mwathunga<sup>8</sup>, S Banda<sup>8</sup>, G Gwengweya<sup>8</sup>, D Nedaw<sup>3</sup>, S Jumbo<sup>9</sup>, E Banks<sup>10</sup> , P Cook<sup>10</sup> and V Casey<sup>11</sup>

<sup>1</sup> British Geological Survey, Maclean Building, Wallingford OX10 8BB, United Kingdom

<sup>2</sup> British Geological Survey, Lyell Centre, Edinburgh EH14 4AP, United Kingdom

<sup>3</sup> Department of Earth Sciences, Addis Ababa University, Addis Ababa, Ethiopia

<sup>4</sup> Department of Geology and Petroleum Studies, Kampala, Makerere University, Uganda

<sup>5</sup> Department of Civil Engineering, University of Malawi, Blantyre, Malawi

<sup>6</sup> Geological Survey of Northern Ireland, British Geological Survey, Department of Enterprise, Trade & Investment, Dundonald House, Upper Newtownards Road, Belfast BT4 3SB, United Kingdom

<sup>7</sup> School of Biosciences, University of Nottingham, Sutton Bonington Campus, Leicestershire LE12 5RD, United Kingdom

<sup>8</sup> Chancellor College, University of Malawi, Zomba, Malawi

<sup>9</sup> WaterAid Malawi, Lilongwe, Malawi

<sup>10</sup> National Centre for Groundwater Research and Training, College of Science & Engineering, Flinders University, Adelaide, SA 5001, Australia

<sup>11</sup> WaterAid UK, Vauxhall, London SE11 5JD, United Kingdom

<sup>12</sup> Author to whom any correspondence should be addressed

E-mail: [djla@bgs.ac.uk](mailto:djla@bgs.ac.uk)

**Keywords:** groundwater, water quality, boreholes, handpumps, SDG 6, Africa, climate

Supplementary material for this article is available [online](#)

## Abstract

Groundwater provides a vital source of drinking water for rural communities in many parts of Africa, particularly in the dry season when there are few safe alternative sources. This paper summarises results from a study ( $n = 428$ ) assessing dry season water quality, both microbiological and inorganic chemistry, in handpump equipped boreholes (HPBs) across the Ethiopia Highlands ( $n = 142$ ), Malawi ( $n = 162$ ) and Uganda ( $n = 124$ ) using a stratified, randomised sampling design. This study seeks to examine general water quality by randomly sampling rural groundwater supplies across larger areas with different geology and climate. The majority, 72%, of HPBs surveyed provide good quality dry season drinking water as defined by WHO drinking water quality criteria. Within this overall picture, the most notable constraints were from thermotolerant coliforms (TTCs), which exceeded the WHO drinking water guideline of zero colony forming units (cfu/100 ml) in 21% of sites (range 0–626 cfu/100 ml). TTC contamination was found to have a significant and positive correlation with annual average rainfall ( $\rho = 0.2$ ,  $p = 0.00003$ ). Across all three countries, WHO health based chemical drinking water quality values were exceeded at 9% of sites and were found for manganese (4%), fluoride (2.6%) and nitrate (2.5%); arsenic concentrations were below the guideline value of  $10 \mu\text{g l}^{-1}$  (range  $< 0.5$ – $7 \mu\text{g l}^{-1}$ ). The high percentage of Mn exceedances ( $14\% \pm 5.2\% > 400 \mu\text{g l}^{-1}$ ) found in drinking water sources in Uganda challenges the decision by WHO not to formalise a health-based guideline for Mn. While the overall level of microbiological contamination from HPBs is low, results from this study strongly suggest that at a national and regional level, microbiological contamination rather than chemical contamination will provide a greater barrier to achieving targets set for improved drinking water quality under the UN-SDG 6. Efforts should be made to ensure that boreholes are properly sited and constructed effectively to reduce pathogen contamination.

## 1. Introduction

Groundwater is the major source of drinking water in Africa (Foster *et al* 2008, Pavelic *et al* 2012, Gaye and Tindimugaya 2019) and use of groundwater for drinking water is increasing due to population and economic growth (Vörösmarty *et al* 2005, United Nations Department of Economic and Social Affairs Population Division 2019) and the search for climate resilient water supplies (Howard *et al* 2016). Boreholes equipped with handpumps (HPBs) are the main method by which people access groundwater in rural areas (JMP 2019). Household supplies of water come from a wide range of different sources including HPBs, shallow hand-dug wells, springs, ephemeral rivers, harvested rainfall and piped water supplies, depending on local/seasonal availability, hydrogeological conditions and the level of investment in water infrastructure (UNICEF and WHO 2015). Compared to surface water sources or shallow hand-dug wells, HPBs usually tap deeper aquifers that are more resilient to inter-annual climate variability (Chilton and Foster 1995, Macdonald *et al* 2009, Lapworth *et al* 2013, Taylor *et al* 2013). For many rural communities in Africa, HPBs are an integral part of household drinking water supply and are often the only source in the dry season, or longer periods of drought (Calow *et al* 2010, Macdonald *et al* 2019).

Understanding the quantity and accessibility of groundwater available for rural community supplies in Africa has been the primary focus of regional research to date (e.g. Macdonald *et al* 2012; Bonsor *et al* 2018, Cuthbert *et al* 2019). In contrast, the quality (both chemical and microbiological) of groundwater resources in Africa has tended to focus on areas with known natural water quality problems such as arsenic and fluoride (Reimann *et al* 2003, Edmunds and Smedley 2005, 2013, Rango *et al* 2013), contamination from mining (Smedley 1996, Von Der Heyden and New 2004) or urbanisation (Lapworth *et al* 2017a). Regional groundwater quality information is nevertheless essential for assessing the availability of safe drinking water across Africa (Bartram and Cairncross 2010, Hunter *et al* 2010, Parker *et al* 2010). HPBs are often reported to have better water quality compared to other groundwater sources (Parker *et al* 2010, Sorensen *et al* 2015a). However, the baseline water quality of HPBs sources in Africa remains understudied compared to other continents globally. Africa wide meta-analyses of groundwater quality have been undertaken for selected water quality parameters, including nitrate (Ouedraogo and Vanclooster 2016), fluoride (Kut *et al* 2016), arsenic (Ahoulé *et al* 2015) and faecal contamination (Bain *et al* 2014). These meta-studies have highlighted that the majority of published studies on groundwater quality in Africa have (i) limited geographical and geological scope, (ii) studies rarely consider paired observations of microbiological and chemical water

quality, and (iii) results are often reported from a mixture of different groundwater source types (e.g. Smedley 1996, Reimann *et al* 2003, Parker *et al* 2010, Sorensen *et al* 2015b, Bretzler *et al* 2017).

Groundwater sources in Africa are commonly used for drinking water and cooking with no (or very limited) treatment and as such understanding the raw chemical and microbiological quality of these sources remains a key priority from a human health perspective (Bain *et al* 2014). HPBs are a critical component of 'improved' drinking water sources (JMP 2018). Therefore, characterising HPB drinking water quality is required to ensure the provision of safe drinking water in this region and contribute to assessing progress towards the UN Sustainable Development Goal (SDG) 6 (UN 2015). This type of information underpins future investment in HPB, and other improved infrastructure on the Joint Monitoring Programme (JMP) drinking water service ladder (JMP 2018), as well as aesthetic considerations, water treatment options and management of this finite freshwater resource.

The objective of this paper is to investigate the drinking water quality from HPBs in sub-Saharan Africa, specifically in eastern and southern Africa, using paired microbiological and chemical water quality data. A large subsample of rural HPBs from three countries (Ethiopia, Malawi and Uganda) were included in this study, which represent many of the major hydrogeological/climate settings found in populated Africa. As far as the authors are aware, this is the largest multi-country water quality survey of HPBs undertaken in Africa. Drinking water quality is compared against World Health Organisation (WHO) criteria to assess current performance against SDG6 (UN 2015) drinking water targets. This study aims to (i) quantify the key health-based water quality exceedances that may affect HPB users; (ii) assess the relative proportion of water quality exceedances from microbiological and inorganic chemical parameters; and (iii) investigate the relationship between HPB water quality and aquifer type, the length of the dry season and annual average rainfall.

## 2. Methods

### 2.1. Experimental design and study area


The three countries, Ethiopia, Malawi and Uganda, were chosen because they cover a range of the geological and climatic conditions found in Africa and also rely on HPBs for much of their rural water supply (JMP 2018). Both climate and geology may have an impact on the water quality of HPBs, i.e. the geochemistry of groundwater and likelihood of microbiological contamination. Using a stratified randomised approach, we anticipate that the results from this study are not constrained or unduly influenced by localised conditions or anomalies, for example natural variations in geology or localised climate

anomalies. The planning for this study was undertaken through detailed consultation with national ministries and district level water officers to obtain the most comprehensive record of HPBs at the district and community level. Sampling was undertaken by a two-stage stratified random design. The primary sample units, which were stratified, were Woredas in Ethiopia and Districts in Malawi and Uganda. These primary units were determined largely by accessibility, but some primary units that were known to have very few HPBs were excluded. In Ethiopia the stratification was based on a combination of aquifer class (fractured igneous or porous igneous). The Woredas in each of these categories were then divided into poverty classes ('better off' or 'poorer'). In Malawi and Uganda the stratification was by poverty class (again, 'better off' or 'poorer'). Random selection of primary units within the strata was done without replacement using the Rao–Hartley–Cochran (RHC) sampling procedure (Cochran 1977). Within the selected districts or woredas, communities were then selected by simple random sampling and from each one a HPB was selected for examination.

Figure 1 shows the location of the HPB survey sites as well as the geology and climate of the study area. HPB sample sites were collected from four key aquifer types found in Africa (Macdonald *et al* 2012): crystalline basement, consolidated sediments, unconsolidated sediments, and volcanic rocks (both fractured and porous igneous rocks). Note that the known high fluoride areas of the rift valley in Ethiopia were not sampled, partly due to the low coverage of rural water supply handpumps in this area compared to other areas. Sites also encompassed a range of climates, spanning regions with zero dry months/years through to dry seasons of up to 6 months/years. Sampling was undertaken during dry periods to facilitate access to remote field areas. As a one-off survey, this also minimises seasonal impacts on water quality and characterises water quality under conditions when there are few alternative sources of drinking water. Annual average rainfall data (1901–2012) for each location was obtained from (Jones and Harris 2013).

## 2.2. HPB sampling and groundwater analysis

HPBs were purged, by a minimum of three borehole volumes, prior to sampling to obtain a representative groundwater sample from the aquifer. Field measurements of pH and specific electrical conductivity (SEC) were made and stable readings obtained prior to sampling. *In-situ* field measurements of turbidity and alkalinity were also undertaken. Two samples for dissolved inorganic chemistry, one for anions and one for cations, were filtered in the field (<0.45 micron) and stored in air-tight Nalgene bottles. Anion sample bottles were filled to the top to exclude air. Major anions were analysed by ion chromatography, samples for major cation and trace



element analysis by ICP-OES and ICP-MS were preserved using analysis grade nitric acid (1%v/v) (e.g. Lapworth *et al* 2013). Inorganic chemical analysis was undertaken by CSIRO Land and Water Analytical Services, Adelaide, South Australia using accredited analytical methods. Total dissolved solids (TDS) was calculated by summing the major anion and cations. Field blank samples were collected using the same procedure. Samples were stored in a cool box



in the field and then transferred to a field refrigerator the same day for storage during fieldwork prior to analysis.

We assess microbiological water quality using thermotolerant coliforms (TTCs), an indicator of faecal contamination (see Sorensen *et al* 2015a). HPB samples were collected in the field using sterilised 250 ml sample bottles and processed using a Delagua® incubator in the field within 7 h of sampling. Samples were stored in a cool box in the field prior to processing. TTCs were isolated and enumerated using the membrane filtration method and Membrane Lauryl Sulphate Broth (MLSB, Oxoid Ltd) as the selective medium. Processed samples were incubated at 44°C for between 18–24 h prior to counting of colony forming units (cfu). Typically, 100 ml of sample was filtered through 0.45 micron nitrocellulose membrane giving a detection limit of 1 cfu/100 ml. We examined incubation plates within 15 min after being removed from the incubator. All cream to yellow colonies with a diameter greater than 1 mm were considered to be TTCs and were counted. We carried out daily blanks and repeat samples for TTCs to check for cross-contamination and quantify precision. A few over-range TTC samples that were ‘too numerous to count’ were diluted and incubated within 24 h. A comprehensive water quality screening for TTCs and 13 potentially hazardous inorganic parameters (F, NO<sub>3</sub>, B, Se, Cr, Mn, Ni, Cu, As, Mo, Cd, Sb, and Pb) was undertaken in all HPBs.

### 2.3. Statistical analysis

The estimation of national mean values for the variables of interest, or for indicator variables which exceeded WHO health-based thresholds, was based on the two-stage cluster sample design described above. Stratum means and variances were combined to give an estimate of the mean at a national scale, and an associated error, based on stratum relative areas (see Cochran 1977 for a full account of how this is done for RHC stratified random two-stage sampling). In the first instance the stratum relative areas were computed from the number of water points in each stratum. This is the most straight-forward estimate, but it relates only to the original sample domain, i.e. to the set of Woredas or districts in each country deemed to be available for sampling. An alternative estimate was computed in each country (up-scaled country estimate), based on the estimated relative areas of the strata at a national scale (igneous aquifer types only in Ethiopia). This requires the assumption that the stratum mean for the sample domain is an unbiased estimate of the equivalent stratum mean at a national scale. The relative areas of the strata at the national scale were computed from information on shallow well numbers in Ethiopia, numbers of villages in Uganda, and numbers of rural enumeration areas from the Third Integrated Household Survey of Malawi (NSO 2012).


Box-plots used in the paper were generated using the ‘cenboxplot’ function from the NADA package in the open source software R v. 3.6.1 (R Core Team 2019). Percentiles in the grey area are estimated using regression on order statistics (ROS), box whiskers extend to 1.5 × interquartile range, and outliers are shown as small circles. Where insufficient data were available to generate a box-plot, dot-plots were used instead. Spearman’s rank correlation coefficient and p values where stated were calculated using R.

## 3. Results

### 3.1. Health based water quality status of HPBs

The proportion of sites that exceed WHO health based chemical and microbiological guideline values (WHO 2017), as well as TDS (which has an aesthetic threshold) for both country sampling domains and up-scaled country estimates are summarised in figure 2. A summary of the results and exceedances (and standard errors) of WHO criteria for all water quality parameters investigated are presented in the supplementary information (table S1 is available online at “(stacks.iop.org/ERL/15/064020/mmedia)”). Overall, WHO health based chemical drinking water quality values were exceeded at 9% of sites and were only found for manganese (4%), fluoride (2.6%) and nitrate (2.5%). Rates of TTC detections above the WHO drinking water guideline value of zero cfu vary considerably between countries (range 13%–24%). However, they are higher than the rates for chemical exceedances of WHO drinking water guidelines (range 0%–14%) in all three countries (figure 2).

Health based chemical drinking water exceedances were only found for manganese, fluoride and nitrate (figure 2); arsenic was not detected above the WHO guideline value of 10 µg l<sup>-1</sup>. For sampling domain estimates, fluoride accounted for the largest proportion of chemical drinking water exceedances (>1.5 mg l<sup>-1</sup>) in Malawi (6.6% ± 2.8%). Manganese (>400 µg l<sup>-1</sup>) accounted for the largest proportion in Uganda (13.8% ± 5.2%) and nitrate (>50 mg l<sup>-1</sup> NO<sub>3</sub>) the largest proportion also in Uganda (4.6% ± 3.2%). TDS was found to be >1000 mg l<sup>-1</sup> at 8.1% ± 2.6% of sites in Malawi and 1% ± 0.6% and 1% ± 1% in Highland Ethiopia and Uganda, respectively. Overall, sampling domain estimates were comparable with up-scaled country estimates, within error, for each survey country. Sampling domain based exceedance estimates for Ethiopia were the highest (23.6% ± 6.5%) of any country, and were significantly higher than those from Malawi (13% ± 3%). However, they were within the standard error for Uganda. Malawi was found to have significantly higher fluoride and TDS exceedances compared to the other countries for both sampling domain and up-scaled country estimates. However, the national estimate for Ethiopia was



**Figure 2.** Summary exceedance of WHO drinking water criteria from HPB. (a) Comparison of exceedances based on sampling domain, (b) comparison of exceedances based on up-scaled country estimates. Health based thresholds for TTCs, F, Mn, NO<sub>3</sub>, as well as TDS based on a taste acceptability thresholds of 1000 mg l<sup>-1</sup> (WHO 2017).

based on data from Highland regions and specifically excluded the rift aquifer system. Uganda had significantly higher exceedance rates for manganese compared to Ethiopia and Malawi. Exceedances for nitrate were not significantly different for the three countries used in this study, for both sampling domain and up-scaled country estimates. In addition, zinc exceedances (on aesthetic grounds, i.e. >3000 µg l<sup>-1</sup>) were significantly lower (zero) for Malawi compared to the other countries (table S1), high Zn concentrations (>1000 µg l<sup>-1</sup>) were only observed in sites that use India Mark II pumps, this is due to a combination of the corrosive nature (pH < 6.5) of some groundwaters in this study and galvanised pump components used in this pump model (figures S1 and S3).


Figure 3 shows mean and standard error plots for each country, estimated using sampling domain, for water quality parameters which exceed the WHO health based drinking water criteria (TTC, F, NO<sub>3</sub> and Mn) as well as TDS where aesthetic criteria are exceeded. With the exception of TTCs and Mn, mean estimates were all below WHO health and aesthetic guideline values. The differences in water quality issues, which dominate in each of the three countries are apparent in figure 3. Mean TTC values are higher for Highland Ethiopia (15 ± 3 cfu/100 ml) compared to Uganda (6.3 ± 6.5 cfu/100 ml) and Malawi (0.2 ± 0.4 cfu/100 ml). However, mean values are only significantly different between Ethiopia Highlands and Malawi due to the high standard error in the Uganda estimates (figure 3(a)). Mean fluoride concentrations are below 0.5 mg l<sup>-1</sup> (500 µg l<sup>-1</sup>) in all three countries and were comparable for Ethiopia

(402 ± 89 µg l<sup>-1</sup>) and Malawi (336 ± 87 µg l<sup>-1</sup>), and significantly lower for Uganda (116 ± 16 µg l<sup>-1</sup>). Mean nitrate concentrations were comparable for all three countries, and were below 15 mg l<sup>-1</sup> NO<sub>3</sub>. Mean manganese concentrations in Uganda were significantly higher compared to Highland Ethiopia and Malawi (figure 3(b)). In Uganda, mean concentrations (150 ± 39 µg l<sup>-1</sup>) were below the current health-based WHO value of 400 µg l<sup>-1</sup>, but significantly above the aesthetic threshold of 100 µg l<sup>-1</sup> (WHO 2017). Mean TDS concentrations were <500 mg l<sup>-1</sup> in all three countries and were found to be significantly different in each country. Mean TDS concentrations were highest for Malawi (428 ± 45 mg l<sup>-1</sup>) and almost twice the mean concentrations for Uganda (266 ± 23 mg l<sup>-1</sup>); Malawi also had the highest outlier values and exceeded taste-aesthetic criteria of 1000 mg l<sup>-1</sup>.

### 3.2. Variations in water quality with the number of dry months, annual average rainfall and aquifer type

Figure 4 summarises, as box-and-whisker plots and cross-plots, the variation in health based water quality parameters with the number of dry months, average annual rainfall and aquifer type. There are no obvious visible associations between the number of dry months and either microbiological or inorganic water quality or the number of sites that exceed WHO drinking water thresholds.

Low numbers of high (outlier) values for TTC and other inorganic health based water quality parameters are distributed across the range of climate settings



**Figure 3.** Distributions for key water quality parameters estimated for sampling domains. Mean and standard error for (a) TTC (cfu/100 ml) and nitrate ( $\text{mg l}^{-1}$ ), (b) fluoride ( $\mu\text{g l}^{-1}$ ), manganese ( $\mu\text{g l}^{-1}$ ), and TDS ( $\text{mg l}^{-1}$ ).



used in this study (figures 4(a)–(e)). There are, however, significantly higher TTCs (figure 4(f)) at locations with higher annual average rainfall totals ( $p = 0.2$ ,  $p = 0.00003$ , Spearman's rank correlation), the relationship is particularly clear for sites where annual rainfall exceeds  $1000 \text{ mm a}^{-1}$ , but no significant correlation for the other health based water quality indicators in relation to average rainfall totals were found (figures 4(g)–(j)). TTC exceedances are observed in all aquifer types, with overall lower counts found in unconsolidated sediments compared to the other aquifer types (figure 4(k)). Median F concentrations are comparable and below  $0.5 \text{ mg l}^{-1}$  for all aquifer types, whilst most of the outliers, which exceed the WHO threshold of  $1.5 \text{ mg l}^{-1}$  are from basement and unconsolidated sediments (figure 4(l)). Highest median nitrate concentrations are found in fractured igneous aquifers and are higher than those from porous igneous and unconsolidated sediments. However, outliers  $>50 \text{ mg l}^{-1}$  are

found in all aquifer types except unconsolidated sediments. Unconsolidated sediments have the highest TDS concentrations, with median values  $>500 \text{ mg l}^{-1}$  (figure 4(o)).

Figure 5 shows cross-plots of TTCs vs TDS for all three countries. It is apparent that in all three countries the highest TTC detections are found in samples with typically low TDS, and low values for TTCs were found in samples with higher TDS.

#### 4. Discussion

This study shows that the water quality in the majority of HPBs included in this survey is of acceptable drinking water quality based on WHO criteria. Health based exceedances were only found for TTCs, F, Mn and NO<sub>3</sub> and rates for exceedances for individual inorganic parameters were  $<10\%$  across all three countries. This contrasts with the more widely reported issue of high groundwater nitrate concentrations



elsewhere globally due to leaching of anthropogenic N sources (USGS 1999, Shand and Edmunds 2008, Gu *et al* 2013, Ascott *et al* 2017, Padilla *et al* 2018). The majority of groundwaters with F concentrations  $>1.5 \text{ mg l}^{-1}$  (9 out of 11 sites in this study) were found in Malawi, the other two sites were from the

Ethiopian Highlands where the concentrations were below  $4 \text{ mg l}^{-1}$  for all but one sample. High F groundwaters are well documented in both countries and have been linked to hydrothermal sources (Bath 1980, Reimann *et al* 2003, Rivett *et al* 2019). They are also found in high concentrations in other parts of the East



African Rift system (Malago *et al* 2017). Arsenic was not found above the WHO limit of  $10 \mu\text{g l}^{-1}$  in any HPB in this study, even though As has been reported to be an issue in some other parts of Africa (e.g. Smedley 1996, Smedley *et al* 2007, Ahoulé *et al* 2015), including in some of the countries that were surveyed as part of this study (e.g. Bamuwanye *et al* 2017). This perhaps reflects the more purposeful nature of sampling designs in many published studies (i.e. targeting likely As hot-spots), which contrasts with the randomised design that was used in this study. However, overall the evidence from this study shows that As contamination is not likely to be a widespread issue across this region and is more likely associated with particular geological settings, for example F and As hot-spots in the Rift Valley (e.g. Rango *et al* 2017).

For a small but significant proportion of sites, manganese was found to exceed the WHO health based drinking water criterion of  $400 \mu\text{g l}^{-1}$  in all three countries (figure 2). Highest manganese concentrations were found in Uganda, up to  $1550 \mu\text{g l}^{-1}$  (95th percentile of  $566 \text{ mg l}^{-1}$ , table S1) and these high concentrations are consistent with previous studies in Uganda and Malawi (e.g. Bath 1980, Taylor and Howard 1994). In this study, mean up-scaled country estimates were  $34 \pm 12 \mu\text{g l}^{-1}$ ,  $48 \pm 7 \mu\text{g l}^{-1}$  and  $150 \pm 39 \mu\text{g l}^{-1}$  for Ethiopia, Malawi and Uganda, respectively; all considerably lower than the WHO health based guideline value of  $400 \mu\text{g l}^{-1}$ . Manganese exceedance was significantly higher for Uganda ( $13.8\% \pm 5.2\%$ ) compared to  $2.1\% \pm 1.1\%$  and  $1.5\% \pm 1.4\%$  for Malawi and the Ethiopian Highlands, respectively. Elevated manganese has often been reported as a co-contaminant with high arsenic (Mitchell *et al* 2011), and is often linked to reducing and low pH conditions (e.g. Homoncik *et al* 2010). However, no association between high Mn and high As or Fe or low pH was found in this study (figure S1), in contrast to other studies globally (Edmunds and Smedley 1996, Buschmann *et al* 2007). For communities reliant on groundwater sources with values approaching or exceeding  $400 \mu\text{g l}^{-1}$ , the findings from this study challenge the decision by WHO (2017) that a formal drinking water guideline value for Mn ( $400 \mu\text{g l}^{-1}$ ) is not required. Since water is aesthetically poor at  $100 \mu\text{g l}^{-1}$ , and the mean concentrations in Uganda were found to be significantly higher than this aesthetic threshold ( $150 \pm 39 \mu\text{g l}^{-1}$ ) and were being used as drinking water sources, some review of the guideline is required. Other studies in Ghana (Rossiter *et al* 2010) and elsewhere globally (e.g. Homoncik *et al* 2010) have also shown a high proportion (11% and 9%, respectively) of rural drinking water supplies above  $400 \mu\text{g l}^{-1}$ .

Occurrences of dry season microbiological contamination were found to be low for HPBs. Overall, 79% of HPBs were free from TTCs, and <10% of sites are classified as medium (10 to <100 cfu/100 ml)

or high (100 to <1000 cfu/100 ml) risk, based on WHO classifications. Although the exceedances were low overall, HPBs were more than twice as likely to fail based on microbiological contamination (i.e. TTCs > 0 cfu) compared to inorganic (WHO) water quality criteria (figure 2). The occurrence of TTCs in HPBs varied considerably between country, with detection rates (>0 cfu/100 mL) in the Ethiopian Highlands ( $23.6\% \pm 6.5\%$ ) almost twice as high as Malawi ( $13\% \pm 3\%$ ). These results are in stark contrast to a recent survey by the Central Statistical Agency in Ethiopia that showed 85% of boreholes surveyed had TTCs > 0 cfu/100 ml (CSAE 2017), which may be due to differences in sampling methodology. (Parker *et al* 2010) found higher, but comparable, TTC detection rates (30%, compared to  $18.4\% \pm 4.6\%$  for this study) for HPBs as part of a large study in Uganda. The higher rates may be due to the fact that sampling by Parker *et al* (2010) was undertaken in May–July, which included the end of the long rainfall season in Uganda, rather than between June–August (the driest part of the year), which was the case in this study. However, it is also possible that the differences in sampling design account for the differences in rates of faecal indicators detected in HPBs in Uganda.

No significant relationship between TTC and/or inorganic health based water quality in HPBs and the length of dry season (range 0–6 months) was found in this study (figure 4). However, the highest TTCs were found for locations with highest total annual rainfall (figure 4(f)). The HPBs in this study typically draw water from <20 mbgl, and the relatively high (i.e. >30 years) mean residence times of HPBs (Lapworth *et al* 2013) in Africa and the degree of protection provided by HPBs to surface contamination are such that microbiological contamination levels are expected to be low compared to alternative groundwater sources. However, given that highest TTCs were observed in locations with highest total annual rainfall suggests that there may be a higher risk of contamination from TTCs under wetter conditions at selected sites, combined with the presence of a proximal source of contamination such as a pit latrine, or a faulty borehole sanitary seal (e.g. Back *et al* 2018). The reasons for this are uncertain, but could be linked to more intense rainfall and/or flooding under wetter conditions (Bridgman *et al* 1995, Wu *et al* 2016) and the activation of rapid recharge pathways and/or shallower groundwater tables under wetter climate conditions (Gotkowitz *et al* 2016, Cuthbert *et al* 2019). It has been previously suggested that under wetter climatic conditions, microbiological contamination may be higher (Gotkowitz *et al* 2016), and with higher contamination during the wet season (e.g. Sorensen *et al* 2015a, 2015b). However, the effects are highly uncertain (Macdonald *et al* 2009). While this study provides evidence to support the proposition that there may be a link between wetter

climate and microbiological contamination in HPBs, further studies are required to better understand this relationship.

Sites with low TDS have the most TTC contamination (figure 5). This supports the hypothesis that higher TTC contamination is related to sites with a higher proportion of rapid modern recharge, and hence lower TDS due to more limited water-rock interaction. This finding was consistent across all three countries, which adds weight to the notion that this TTC contamination relates to rapid recharge pathways. In more arid regions it is also possible that evaporative enrichment during recharge, contributes to higher dissolved constituents and TDS in the unsaturated zone (Scanlon *et al* 2006, Green *et al* 2011, Gurdak *et al* 2012, Kløve *et al* 2014). The locations included in this study were not particularly arid, so this association between climate and TDS was not observed (figure 4). Equally, land use in the rural study areas was rain fed agriculture, where groundwater recharge is likely to be dominated by diffuse processes (Lapworth *et al* 2013). Therefore, the effects of evaporative enrichment from irrigation return flows would be limited (Foster *et al* 2018). Improved construction and maintenance of water points is critical but can be very challenging to monitor and undertake in this region, despite the more obvious benefits to HPB functionality to help reduce TTC contamination within HPBs.

No exceedances in health-based water quality were observed to be strongly associated with a particular aquifer type. Low level exceedances for TTCs, F,  $\text{NO}_3$ , Mn and TDS were found for all the major aquifer types including those classed as basement, igneous and sedimentary (figure 4). HPBs in consolidated sediments and fractured igneous aquifers had no F concentrations exceeding the WHO of  $1.5 \text{ mg l}^{-1}$ . However, a small number of exceedances were found in basement, porous igneous and unconsolidated sedimentary aquifers. In this study the higher median TDS values observed in unconsolidated sediments were largely accounted for by sites in the Lower Shire Valley in southern Malawi (see figure S2), which are likely linked to low hydraulic gradients and therefore higher residence times allowing mineral dissolution as well as evaporative enrichment in groundwater due to the combination of shallow groundwater tables and high surface temperatures (Bath 1980, Monjerezi *et al* 2011, Rivett *et al* 2019). Similarly, elevated TDS groundwaters have been reported in Namibia due to mineral dissolution processes (Li *et al* 2018).

A limitation of this study is the fact that uranium, a potential contaminant of concern (Brugge and Oldmixon 2005) was not quantified. Naturally high uranium concentrations have been found in both basement and sedimentary settings globally (e.g. Hess *et al* 1985, Smedley *et al* 2006, Lapworth *et al* 2017b, Coyte *et al* 2018). Several studies in Africa

have found naturally occurring high uranium concentrations (Vogel *et al* 1999, Silliman *et al* 2007, van Wyk and Coetzee 2008), and further work on the distribution of U in groundwater in Africa is required.

## 5. Conclusions and future perspectives

This study provides an assessment of the baseline drinking water quality from rural HPBs in Sub-Saharan Africa across a range of different climates and aquifer geology. The results from this study across three countries in Africa show that the majority of drinking water from rural HPBs is found to be of good quality, based on health-based criteria, and certainly better quality than most alternative sources available in rural Africa which confirms earlier smaller scale assessments (Parker *et al* 2010, Pritchard *et al* 2016, Macdonald *et al* 2019). There are no strong links between either dry season length, aquifer geology and water quality based on an assessment of results using 14 parameters with health-based drinking water criteria. Significantly, faecal contamination, assessed using TTC indicators, was found to be the greater barrier to achieving good quality drinking water status under SDG 6 compared to inorganic chemical criteria, and affected 21% of HPBs surveyed overall, with considerable variability between countries (13%–24%). The highest TTC counts were found at sites with the highest total annual rainfall, supporting the idea that there may be a link between climate and increased contamination from faecal sources even in improved sources such as HPBs. High TTCs in low TDS waters also suggest that more attention should be given to the quality of HPB construction, and local land-use, to reduce contamination through rapid transit routes to the aquifer and the production zone of the HPB. The only inorganic health based water quality parameters found to exceed WHO guideline values were manganese, fluoride, and nitrate. The significantly higher number of exceedances in Uganda ( $13.8\% \pm 5.2\%$ ), all used for drinking water supply, with  $\text{Mn} > 400 \mu\text{g l}^{-1}$ , and the mean Mn concentration of  $150 \mu\text{g l}^{-1}$ , challenges the current proposition that water  $>100 \mu\text{g l}^{-1}$  is not used for drinking water due to aesthetic constraints, and highlights the need for a health-based formal guideline value for Mn. This study, carried out in 3 countries in Sub-Saharan Africa, shows that water quality challenges for meeting SDG 6 drinking water targets using rural HPBs do occur, but are constrained to a minority of sites and could be further reduced by better HPB construction and maintenance. In most cases HPBs are a source of drinking water of good status during the dry season, when there are limited options for alternative sources. Further work is needed to extend this type of assessment in terms of geographical coverage as well as assessing other water quality parameters not included in this

study (e.g. uranium). Although considerable effort is required to plan randomised studies effectively, given the significant cost of undertaking an assessment of equivalent scale, using a randomised approach will generate a more robust evidence base with which to assess drinking water quality status in HPBs in Africa.

## Acknowledgments

This work is part of the Hidden Crisis project within the UPGro research programme, grant number NE/M008606/1, and is co-funded by Natural Environment Research Council (NERC), UK Department for International Development (DFID), and the Economic and Social Research Council (ESRC). DJL was also funded in part by HyCRISTAL (NE/M020452/1). BGS authors published with the permission of the Executive Director of the British Geological Survey (BGS-UKRI). Significant thanks go to the WaterAid Country programme staff within each of the three countries who were instrumental to facilitating the survey and ensuring sensitive mobilisation of communities: Gossa Wolde and Gashaye Chekol (WaterAid Ethiopia); Gloria Berochan and Jacinta Nekesa (WaterAid Uganda); and Sella Jumbo and Lughano Mogha (WaterAid Malawi). Thanks also go to the social science field researchers within each country who assisted with the survey work: Yehualaseshet Tadesse (Ethiopia), Katusiime Felece (Uganda).

The data that support the findings of this study will be openly available at DOI 10.5285/bca0d930-a10d-4ae8-9afc-ad1bf73de35a (UPGro 2022, Hidden Crisis Project, Survey 1 dataset: detailed functionality assessments of hand pump boreholes in Ethiopia, Uganda and Malawi) following a delay of 24 months from the date of publication.

## ORCID iDs

D J Lapworth  <https://orcid.org/0000-0001-7838-7960>

A M MacDonald  <https://orcid.org/0000-0001-6636-1499>

J S T Ward  <https://orcid.org/0000-0003-2868-8604>

E Banks  <https://orcid.org/0000-0002-2431-7649>

## References

- Ahoulé D G, Lalanne F, Mendret J, Brosillon S and Maïga A H 2015 Arsenic in African waters: a review *Water Air Soil Pollut.* **226** 302
- Ascott M J, Gooddy D C, Wang L, Stuart M E, Lewis M A, Ward R S and Binley A M 2017 Global patterns of nitrate storage in the vadose zone *Nat. Commun.* **8** 1–7
- Back J O, Rivett M O, Hinz L B, Mackay N, Wanangwa G J, Phiri O L and Miller A V 2018 Risk assessment to groundwater of pit latrine rural sanitation policy in developing country settings *Sci. Total Environ.* **613** 592–610
- Bain R *et al* 2014 Global assessment of exposure to faecal contamination through drinking water based on a systematic review *Tropical Med. Int. Health* **19** 917–27
- Bamuwamye M, Ogwok P, Tumuhairwe V, Eragu R, Nakisozi H and Ogwang P E 2017 Human health risk assessment of heavy metals in Kampala (Uganda) drinking water *J. Food Res.* **6** 6–16
- Bartram J and Cairncross S 2010 Hygiene, sanitation, and water: forgotten foundations of health *PLoS Med.* **7** e1000367
- Bath A H 1980 Hydrochemistry in groundwater development: report on an advisory visit to Malawi. British Geological Survey Report, WD/OS/80/20.
- Bonsor H, MacDonald A, Casey V, Carter R and Wilson P 2018 The need for a standard approach to assessing the functionality of rural community water supplies *Hydrogeology J.* **26** 367–370
- Bretzler A, Lalanne F, Nikiema J, Podgorski J, Pfenninger N, Berg M and Schirmer M 2017 Groundwater arsenic contamination in Burkina Faso, West Africa: predicting and verifying regions at risk *Sci. Total Environ.* **584** 958–70
- Bridgman S, Robertson R M P, Syed Q, Speed N, Andrews N and Hunter P R 1995 Outbreak of Cryptosporidiosis associated with a disinfected groundwater supply *Epidemiol. Infect.* **115** 555–66
- Brugge D and Oldmixon B 2005 Exposure pathways and health effects associated with chemical and radiological toxicity of natural uranium: a review *Rev. Environ. Health* **20** 177–94
- Buschmann J, Berg M, Stengel C and Sampson M L 2007 Arsenic and manganese contamination of drinking water resources in Cambodia: coincidence of risk areas with low relief topography *Environ. Sci. Technol.* **41** 2146–52
- Calow R C, Macdonald A M, Nicol A L and Robins N S 2010 Ground water security and drought in Africa: linking availability, access, and demand *Groundwater* **48** 246–56
- Central Statistical Agency of Ethiopia 2017 Drinking water quality in Ethiopia – results from the 2016 Ethiopia socioeconomic survey
- Chilton P J and Foster S S D 1995 Hydrogeological characterisation and water-supply potential of basement aquifers in tropical Africa *Hydrogeol. J.* **3** 36–49
- Cochran W G 1977 *Sampling Techniques* 3rd edn (New York: Wiley)
- Coyte R M, Jain R C, Srivastava S K, Sharma K C, Khalil A, Ma L and Vengosh A 2018 Large-scale uranium contamination of groundwater resources in India *Environ. Sci. Technol. Lett.* **5** 341–7
- Cuthbert M O *et al* 2019 Observed controls on resilience of groundwater to climate variability in sub-Saharan Africa *Nature* **572** 230–4
- Edmunds W M and Smedley P L 1996 Groundwater geochemistry and health: an overview *Geol. Soc. Lond. Spec. Publ.* **113** 91–105
- Edmunds W M and Smedley P L 2005 Fluoride in natural waters *Essentials of Medical Geology: Impacts of the Natural Environment on Public Health* ed O Selenis (London: Academic)
- Edmunds W M and Smedley P L 2013 Fluoride in natural waters *Essentials of Medical Geology* ed O Selinus (Berlin: Springer) pp 311–36
- Foster S, Pulido-Bosch A, Vallejos Á, Molina L, Llop A and Macdonald A M 2018 Impact of irrigated agriculture on groundwater-recharge salinity: a major sustainability concern in semi-arid regions *Hydrogeol. J.* **26** 2781–91
- Foster S S, Tuinhof A and Garduño H 2008 Groundwater in Sub-Saharan Africa: a strategic overview of developmental issues *Applied Groundwater Studies in Africa* eds S Adelana, A M Macdonald (Boca Raton, FL: CRC Press) pp 19–34
- Gaye C B and Tindimugaya C 2019 Challenges and opportunities for sustainable groundwater management in Africa *Hydrogeol. J.* **27** 1099–110
- Gotkowitz M B, Bradbury K R, Borchardt M A, Zhu J and Spencer S K 2016 Effects of climate and sewer condition on

- virus transport to groundwater *Environ. Sci. Technol.* **50** 8497–504
- Green T R, Taniguchi M, Kooi H, Gurdak J J, Allen D M, Hiscock K M, Treidel H Aureli A *et al* 2011 Beneath the surface of global change: impacts of climate change on groundwater *J. Hydrol.* **405** 532–60
- Gu B, Ge Y, Chang S X, Luo W and Chang J 2013 Nitrate in groundwater of China: sources and driving forces *Global Environ. Change* **23** 1112–21
- Gurdak J J, McMahon P B and Bruce B W 2012 Vulnerability of groundwater quality to human activity and climate change and variability, High Plains aquifer, USA *Climate Change Effects on Groundwater Resources: A Global Synthesis of Findings and Recommendations* ed H Treidel, J Martin-Bordes, J Gurdak (London: CRC Press) vol 27, p 145
- Hess C T, Michel J, Horton T R, Prichard H M and Coniglio W A 1985 The occurrence of radioactivity in public water supplies in the United States *Health Phys.* **48** 553–86
- Homoncik S C, Macdonald A M, Heal K V, Dochartaigh B É Ó and Ngwenya B T 2010 Manganese concentrations in Scottish groundwater *Sci. Total Environ.* **408** 2467–73
- Howard G, Calow R, Macdonald A and Bartram J 2016 Climate change and water and sanitation: likely impacts and emerging trends for action *Annu. Rev. Environ. Resour.* **41** 253–76
- Hunter P R, Macdonald A M and Carter R C 2010 Water supply and health *PLoS Med.* **7** e1000361
- Joint Monitoring Programme 2018 Drinking water services Joint Monitoring Programme 2019 JMP Database detailing drinking water by country and source type
- Jones P D and Harris I C 2013 University of East Anglia Climatic Research Unit: CRU TS3.21: climatic Research Unit (CRU) Time-Series (TS) Version 3.21 of high resolution gridded data of month-by-month variation in climate (Jan. 1901–Dec. 2012) NCAS British Atmospheric Data Centre
- Kløve B, Ala-Aho P, Bertrand G, Gurdak J J, Kupfersberger H, Kværner J and Uvo C B 2014 Climate change impacts on groundwater and dependent ecosystems *J. Hydrol.* **518** 250–66
- Kut K M K, Sarswat A, Srivastava A, Pittman J C U and Mohan D 2016 A review of fluoride in african groundwater and local remediation methods *Groundwater Sustain. Dev.* **2** 190–212
- Lapworth D J, Krishan G, Macdonald A M and Rao M S 2017b Groundwater quality in the alluvial aquifer system of northwest India: new evidence of the extent of anthropogenic and geogenic contamination *Sci. Total Environ.* **599** 1433–44
- Lapworth D J, Macdonald A M, Tijani M N, Darling W G, Goody D C, Bonsor H C and Araguás-Araguás L J 2013 Residence times of shallow groundwater in West Africa: implications for hydrogeology and resilience to future changes in climate *Hydrogeol. J.* **21** 673–86
- Lapworth D J, Nkhuwa D C W, Okotto-Okotto J, Pedley S, Stuart M E, Tijani M N and Wright J 2017a Urban groundwater quality in sub-Saharan Africa: current status and implications for water security and public health *Hydrogeol. J.* **25** 1093–116
- Li Z, Wang G, Wang X, Wan L, Shi Z, Wanke H, Uugulu S and Uahengo C I 2018 Groundwater quality and associated hydrogeochemical processes in Northwest Namibia *J. Geochem. Explor.* **186** 202–14
- Macdonald A M *et al* 2019 Groundwater and resilience to drought in the Ethiopian Highlands *Environ. Res. Lett.* **14** 095003
- Macdonald A M, Bonsor H C, Dochartaigh B É Ó and Taylor R G 2012 Quantitative maps of groundwater resources in Africa *Environ. Res. Lett.* **7** 024009
- Macdonald A M, Calow R C, Macdonald D M, Darling W G and Dochartaigh B E 2009 What impact will climate change have on rural groundwater supplies in Africa? *Hydrol. Sci. J.* **54** 690–703
- Malago J, Makoba E and Muzuka A N 2017 Fluoride levels in surface and groundwater in Africa: a review *Am. J. Water Sci. Eng.* **3** 1–17
- Mitchell E, Frisbie S and Sarkar B 2011 Exposure to multiple metals from groundwater—a global crisis: geology, climate change, health effects, testing, and mitigation *Metallomics* **3** 874–908
- Monjerezi M, Vogt R D, Aagaard P and Saka J D 2011 Hydro-geochemical processes in an area with saline groundwater in lower Shire River valley, Malawi: an integrated application of hierarchical cluster and principal component analyses *Appl. Geochem.* **26** 1399–413
- National Statistical Office (NSO) and Ministry of Economic Planning and Development (MoEPD) 2012 Third integrated household survey 2010–2011 (Malawi National Statistical Office, Zomba, Malawi, and The World Bank, Washington, DC) (<http://microdata.worldbank.org/>)
- Ouedraogo I and Vanclooster M 2016 A meta-analysis and statistical modelling of nitrates in groundwater at the African scale *Hydrol. Earth Syst. Sci.* **20** 2353–81
- Padilla F M, Gallardo M and Manzano-Agugliaro F 2018 Global trends in nitrate leaching research in the 1960–2017 period *Sci. Total Environ.* **643** 400–13
- Parker A H, Youten R, Dillon M, Nussbaumer T, Carter R C, Tyrrel S F and Webster J 2010 An assessment of microbiological water quality of six water source categories in north-east Uganda *J. Water Health* **8** 550–60
- Pavelic P, Giordano M, Keraita B, Ramesh V and Rao T (eds) 2012 *Groundwater Availability and Use in Sub-Saharan Africa: A Review of 15 Countries* (Colombo: International Water Management Institute (IWMI)) p 274
- Pritchard M, Edmondson A, Craven T and Mkandawire T 2016 Development of sustainable drinking water quality solutions for rural communities in the developing world *Sustainable Ecological Engineering Design* eds M Dastbaz, C Gorse (Berlin: Springer) pp 259–77
- R Core Team 2019 R: A Language and Environment for Statistical Computing (Vienna: R Foundation for Statistical Computing) (<https://www.r-project.org/>)
- Rango T, Vengosh A, Dwyer G and Bianchini G 2013 Mobilization of arsenic and other naturally occurring contaminants in groundwater of the Main Ethiopian Rift aquifers *Water Res.* **47** 5801–18
- Rango T, Vengosh A, Jeuland M, Whitford G M and Tekle-Haimanot R 2017 Biomarkers of chronic fluoride exposure in groundwater in a highly exposed population *Science of the Total Environment* **596** 1–11
- Reimann C, Bjorvatn K, Frengstad B, Melaku Z, Tekle-Haimanot R and Siewers U 2003 Drinking water quality in the Ethiopian section of the East African Rift Valley I—data and health aspects *Sci. Total Environ.* **311** 65–80
- Rivett M O *et al* 2019 Responding to salinity in a rural African alluvial valley aquifer system: to boldly go beyond the world of hand-pumped groundwater supply? *Sci. Total Environ.* **653** 1005–24
- Rossiter H M, Owusu P A, Awuah E, Macdonald A M and Schäfer A I 2010 Chemical drinking water quality in Ghana: water costs and scope for advanced treatment *Sci. Total Environ.* **408** 2378–86
- Scanlon B R, Keese K E, Flint A L, Flint L E, Gaye C B, Edmunds W M Simmers I *et al* 2006 Global synthesis of groundwater recharge in semiarid and arid regions *Hydrol. Process.* **20** 3335–70
- Shand P and Edmunds W M 2008 The baseline inorganic chemistry of European groundwaters *Natural Groundwater Quality*, ed W M Edmunds and P Shand (Oxford: Blackwell) pp 22–58
- Silliman S E, Boukari M, Crane P, Azonsi F and Neal C R 2007 Observations on elemental concentrations of groundwater in central Benin *J. Hydrol.* **335** 374–88
- Smedley P L 1996 Arsenic in rural groundwater in Ghana: part special issue: hydrogeochemical studies in sub-Saharan Africa *J. Afr. Earth Sci.* **22** 459–70



- Smedley P L, Knudsen J and Maiga D 2007 Arsenic in groundwater from mineralised Proterozoic basement rocks of Burkina Faso *Appl. Geochem.* **22** 1074–92
- Smedley P L, Smith B, Abesser C and Lapworth D J 2006 Uranium occurrence and behaviour in British groundwater *British Geological Survey Commissioned Report CR/06/050N* (<http://nora.nerc.ac.uk/id/eprint/7432/>)
- Sorensen J P R et al 2015a In-situ tryptophan-like fluorescence: a real-time indicator of faecal contamination in drinking water supplies *Water Res.* **81** 38–46
- Sorensen J P R, Lapworth D J, Read D S, Nkhuwa D C W, Bell R A, Chibesa M, Chirwa M, Kabika J, Liemisa M and Pedley S 2015b Tracing enteric pathogen contamination in sub-Saharan African groundwater *Sci. Total Environ.* **538** 888–95
- Taylor R G and Howard K W F 1994 Groundwater quality in rural Uganda: hydrochemical considerations for the development of aquifers within the basement complex of Africa *In Groundwater Quality* ed H Nash and G J H McCall (London: Chapman & Hall) pp 31–44
- Taylor R G et al 2013 Ground water and climate change *Nat. Clim. Change* **3** 322
- UNICEF and WHO 2015 *Progress on sanitation and drinking water: 2015 update and MDG assessment* (New York: UNICEF) p 80
- United Nations 2015 Resolution adopted by the General Assembly: transforming our world: the 2030 Agenda for sustainable Development *UN Resolution A/RES/70/1*
- United Nations Department of Economic and Social Affairs Population Division 2019 *World Population Prospects 2019* (United Nations), Online Edition
- US Geological Survey—USGS 1999 The quality of our nation's waters; nutrient and pesticides *USGS Report Circular 1225*, p 82
- van Wyk N and Coetzee H 2008 The distribution of uranium in groundwater in the Bushmanland and Namaqualand areas, Northern Cape Province, South Africa *Uranium, Mining and Hydrogeology* eds B J Murkel, A Hasche-Berger (Berlin: Springer) pp 639–44
- Vogel J C, Talma A S, Heaton T H E and Kronfeld J 1999 Evaluating the rate of migration of an uranium deposition front within the Uitenhage Aquifer *J. Geochem. Explor.* **66** 269–76
- Von Der Heyden C J and New M G 2004 Groundwater pollution on the Zambian Copperbelt: deciphering the source and the risk *Sci. Total Environ.* **327** 17–30
- Vörösmarty C J, Douglas E M, Green P A and Revenga C 2005 Geospatial indicators of emerging water stress: an application to Africa *AMBIO* **34** 230–7
- WHO 2017 *Guidelines for Drinking-Water Quality* 4th edn (Geneva: WHO) pp 631
- Wu J, Yunus M, Islam M S and Emch M 2016 Influence of climate extremes and land use on fecal contamination of shallow tubewells in Bangladesh *Environ. Sci. Technol.* **50** 2669–76