ENVIRONMENTAL RESEARCH

LETTER • OPEN ACCESS

Climate change-induced peatland drying in Southeast Asia

To cite this article: Nathan C Dadap et al 2022 Environ. Res. Lett. 17 074026

View the article online for updates and enhancements.

You may also like

- <u>Water level and soil moisture monitoring</u> for peatland fire risk indicator B Kartiwa, S H Adi, H Sosiawan et al.
- <u>The role of moisture regime and hydro-</u> topography on tropical wetland ecosystem management
- L M Nugraha, P B Putra, B A Marpaung et al.
- Determination of soil moisture reduction rate on peatlands in South Sumatera due to the 2019 extreme dry season
 M Irfan, E Koriyanti, Awaluddin et al.



This content was downloaded from IP address 18.226.159.76 on 21/05/2024 at 12:38

ENVIRONMENTAL RESEARCH LETTERS

LETTER

OPEN ACCESS

CrossMark

RECEIVED 22 April 2022

REVISED 13 June 2022

ACCEPTED FOR PUBLICATION 16 June 2022

PUBLISHED 30 June 2022

Original content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence.

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Climate change-induced peatland drying in Southeast Asia

Nathan C Dadap^{1,*}[®], Alexander R Cobb²[®], Alison M Hoyt¹[®], Charles F Harvey³[®], Andrew F Feldman^{4,5}[®], Eun-Soon Im⁶[®] and Alexandra G Konings¹[®]

- ¹ Department of Earth System Science, Stanford University, Stanford, CA, United States of America
- ² Center for Environmental Sensing and Modeling, Singapore-MIT Alliance for Research and Technology, Singapore, Singapore
 - ³ Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, United States of America
 - ⁴ Biospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, United States of America
 - NASA Postdoctoral Program, NASA Goddard Space Flight Center, Greenbelt, MD, United States of America
 - ⁵ Department of Civil and Environmental Engineering, Hong Kong University of Science and Technology, Hong Kong, People's Republic of China
 - * Author to whom any correspondence should be addressed.

E-mail: ndadap@gmail.com

Keywords: tropical peatland, soil moisture, climate change, SMAP, neural network Supplementary material for this article is available online

Abstract

When organic peat soils are sufficiently dry, they become flammable. In Southeast Asian peatlands, widespread deforestation and associated drainage create dry conditions that, when coupled with El Niño-driven drought, result in catastrophic fire events that release large amounts of carbon and deadly smoke to the atmosphere. While the effects of anthropogenic degradation on peat moisture and fire risk have been extensively demonstrated, climate change impacts to peat flammability are poorly understood. These impacts are likely to be mediated primarily through changes in soil moisture. Here, we used neural networks (trained on data from the NASA Soil Moisture Active Passive satellite) to model soil moisture as a function of climate, degradation, and location. The neural networks were forced with regional climate model projections for 1985-2005 and 2040–2060 climate under RCP8.5 forcing to predict changes in soil moisture. We find that reduced precipitation and increased evaporative demand will lead to median soil moisture decreases about half as strong as those observed during recent El Niño droughts in 2015 and 2019. Based on previous studies, such reductions may be expected to accelerate peat carbon emissions. Our results also suggest that soil moisture in degraded areas with less tree cover may be more sensitive to climate change than in other land use types, motivating urgent peatland restoration. Climate change may play an important role in future soil moisture regimes and by extension, future peat fire in Southeast Asian peatlands.

1. Introduction

Peatlands in Insular Southeast Asia contain globally significant carbon stores, estimated at 67 GtC (Page *et al* 2011, Warren *et al* 2017). This carbon is maintained through high water tables that prevent peat oxidation or ignition (Hirano *et al* 2009, Dommain *et al* 2010). However, in the last half a century, degradation has threatened these carbon stores, as only ~6% of peat forests remain in pristine condition (Miettinen *et al* 2016) and widespread drainage has occurred (Dadap *et al* 2021). The resulting drier peat is vulnerable to oxidation (Hooijer *et al* 2012, Jauhiainen *et al* 2012), leading to emissions as large as $155 \pm 30 \text{ Mt C yr}^{-1}$ in 2015 (Hoyt *et al* 2020) or about 70% of combined fossil fuel emissions in Malaysia (63 Mt C yr}^{-1}) and Indonesia (149 Mt C yr}^{-1}) that year (Miettinen *et al* 2017, Andrew and Peters 2021).

Climate also affects peatland carbon loss. During drought years, large-scale burning of peatlands (Van Der Werf *et al* 2008, Field *et al* 2016, Taufik *et al* 2017) also leads to globally significant carbon emissions because dry peat is more flammable. For example, fires associated with the 1997 El Niño Southern Oscillation led to an estimated 0.81–2.56 GtC emitted, 13%–40% of global mean annual fossil

fuel emissions at the time (Page et al 2002). Although fire has been a phenomenon in Southeast Asian peatlands for at least 30 000 years (Goldammer et al 1989, Anshari et al 2001), the frequency and scale of these fires has increased dramatically in recent decades (Page and Hooijer 2016). In the second half of the 20th century, periodic droughts only led to large increases in fire during periods when degradation rates were high (Field et al 2009). This evidence suggests that the combined effects of degradation and climate on the soil moisture and groundwater levels in peatlands mediate peat fire (Taufik et al 2017, Dadap et al 2019). Specifically, degradation can worsen the sensitivity of tropical peatland emissions to meteorological drought (Siegert et al 2001), further motivating restoration and conservation efforts (Jaenicke et al 2010, Leifeld and Menichetti 2018, Goldstein et al 2020).

Given that fire emissions in Southeast Asian peatlands have historically been largest during drought conditions attributable to El Niño Southern Oscillation and the Indian Ocean Dipole (Van Der Werf et al 2008), future emissions may also be influenced by long-term trends associated with climate change (Li et al 2007). Regional climate simulations have shown that average rainfall will likely decrease in Southeast Asia in future decades (Li et al 2007, Tangang et al 2020), especially during the dry season (Kang et al 2019). Additionally, changes in solar radiation, atmospheric humidity, and temperature may also affect the peat water balance. Understanding how future climate will affect peat vulnerability is necessary to inform management, restoration, and conservations efforts. However, the sensitivity of peatland moisture to climate change is likely highly variable across the region. Several factors influence how different hydroclimatological conditions affect peat moisture including the initial distribution of water table depth, water uptake differences between vegetation types (Hirano et al 2015, Manoli et al 2018), canal properties including their depth, width, and spatial pattern (Page et al 2009, Cobb et al 2020, Dadap et al 2021), microtopography, hydraulic properties of the peat and its macropores (Mezbahuddin et al 2015, Baird et al 2017, Cobb et al 2017), and bulk density (Sinclair et al 2020). Because the distribution of these factors across the region is poorly understood and highly uncertain, it is not feasible to parameterize physical hydrologic models (or using land surface simulations from existing regional climate models (RCMs)) to understand how climate change affects peat moisture across this region.

Here, we instead used observations and a statistical modeling approach to estimate how climate change will influence peat hydrological conditions in the coming decades. In particular, we considered surface soil moisture, which has previously been shown to be closely related to peat fire risk (Dadap *et al* 2019) and for which observations are widely available across Southeast Asian peatlands using data from the Soil Moisture Active Passive (SMAP) satellite (Entekhabi et al 2010, McColl et al 2017). Here we use the term 'soil moisture' to describe peat moisture content in the upper ground surface, but note that due to the high organic matter in peatlands, this moisture is not contained in the soil matrix in the same way it is in mineral soils. In tropical peatlands, surface soil moisture is closely connected to water table depth (Hirano et al 2014, Dadap et al 2019), the most commonly used metric of peat moisture levels for fire risk studies (e.g. Wösten et al 2008, Hooijer et al 2012). However, future soil moisture regimes are unknown. To our knowledge, only one study has attempted to model future soil moisture (Li et al 2007), but the coarse resolution (>2 degree) of the general circulation models used in that study cannot account for the complex topography and land-ocean-atmosphere interactions associated with this region, nor for the effects of variations in land use and peat properties. In this study, we instead used machine learning to build a statistical model that predicts soil moisture variations across the region as a function of several climate factors. The statistical model was then used to analyze the impact of climate change on soil moisture across the region, including its spatial distribution and variation with land use type.

2. Methods

2.1. Approach

Our general approach in this study was to train statistical models (neural networks) to learn relationships between climate, degradation, location, and soil moisture in Southeast Asian peatlands under present climate. Neural networks have been shown to be a viable and in some cases superior alternative to stateof-the-art models when forecasting hydrologic variables in data scarce regions (e.g. Hsu *et al* 1995, Kratzert *et al* 2019). The trained neural networks were then used with projections of future climate to predict future soil moisture. This approach is illustrated in figure 1. Such a climate sensitivity approach has been used previously to understand features of hydrologic projections (Short Gianotti *et al* 2020).

Here, we directly predict simplified soil moisture statistics to avoid the need for explicit simulation of soil moisture timeseries in the future. These variables were: (a) mean dry season soil moisture ($\rm sm_{dry\,season}$) and (b) percent low soil moisture ($\rm pct_{low\,sm}$), defined here as the percent of time in a given year that the soil moisture is below 0.2 cm³ cm⁻³. For mean soil moisture, we focus on the dry season only because that is more closely tied to fire risk. Here, we assumed that dry season timing will remain the same in the future period. Previous work using both laboratory measurements (Frandsen 1997, Huang *et al* 2015) and SMAP soil moisture (Dadap *et al*



network predictions. The model is first trained on ERA5 climate and SMAP soil moisture data. Predictions are then calculated for reference (1985–2005) and future (2040–2060) time periods using climate data from a RCM forced by three global circulation models. Input climate data are bias-corrected to ERA5 reanalysis data using quantile mapping.

2019, figure 3) showed that peat ignition probability (at laboratory scale) and burned area (at remote sensing scales) sharply increase when soil moisture is below a threshold value of about $0.2 \text{ cm}^3 \text{ cm}^{-3}$. Thus, the pct_{low sm} statistic represents the fraction of a given year when the peat is at high fire risk and captures the non-linear response of fire to soil moisture.

2.2. Study area

This study focused on peatlands in Insular Southeast Asia, an area spanning \sim 157 000 km² on Sumatra, Borneo, and Peninsular Malaysia. All analyses were limited to pixels covered by at least 50% peatlands, as determined from 30 m land cover maps (Miettinen *et al* 2016), and were performed on the 9 km EASE-Grid resolution of the SMAP data (Brodzik *et al* 2012).

2.3. Data sources

2.3.1. Soil moisture data

Soil moisture data from SMAP are available every 2–3 d at 9 km resolution from 2015 to present. Each pixel represents a distinct soil moisture observation. An example SMAP soil moisture timeseries from one pixel is shown in Supplementary figure 1. We used

soil moisture retrieved from the Multi-Temporal Dual Channel Algorithm (MT-DCA) (Konings et al 2016, 2017, Feldman et al 2021), which is a separate dataset from the SMAP baseline science data products. Because the MT-DCA retrievals rely on a dielectric mixing model that was developed for mineral soils (Mironov et al 2004), an empirical correction was applied to account for the high organic matter content of the peat (Bircher et al 2016). Measurements with potentially high error associated with radio frequency interference, urban areas, and precipitation were excluded from the dataset. Microtopography and the presence of organic material on the peat may add error to the soil moisture retrievals, as the presence of litter can affect L-band soil moisture retrievals even in less densely vegetation sites (Kurum et al 2012). Thick vegetation can also block remote sensing measurement of soil moisture where present. Furthermore, no in situ validation of SMAP data has been performed in this region, which remains a limitation of using SMAP data in this context. However, there is evidence that soil moisture retrievals have sufficient accuracy in this region, since triple collocation-based (statistical) error analysis of SMAP soil moisture in the region previously showed that retrieval precision

is likely on par with the SMAP mission target error of $0.04 \text{ cm}^3 \text{ cm}^{-3}$ (Dadap *et al* 2019).

2.3.2. Input features

Input features were chosen to capture the possible effects of climate, degradation, and location on soil moisture (Supplementary table 1). Climate variables included precipitation and potential evapotranspiration (PET) to represent water supply and evaporative demand; PET was calculated from radiation and temperature using the Priestly-Taylor method. These were represented in the neural networks with mean dry season PET, mean dry season precipitation, mean annual precipitation and precipitation entropy. Precipitation entropy (calculated as the Shannon entropy of monthly precipitation) was included because it is a descriptor of rainfall seasonality (Feng et al 2013), or the degree to which rainfall is distributed between the wet and dry seasons. A smaller entropy value indicates larger seasonal differences in precipitation. Although PET might deviate from actual evapotranspiration (ET), only PET was included here since the RCM and reanalysis data may not capture the differences in water use strategies (and thus, the actual/potential ET ratio) in different land use types.

Because the study area is dominated by coastal areas and topographic complexity, a high resolution simulation is necessary for more accurate prediction of climate variables (Im and Eltahir 2018). Here, we used 25 km regional climate data from the Coordinated Regional Climate Downscaling Experiment-Common Regional Experiment as inputs to the neural networks for the reference (1985-2005) and future periods (2040–2060) (Giorgi et al 2021, Im et al 2021). These data are driven by three global circulation models under Representative Concentration Pathway 8.5 forcing (Meinshausen et al 2011), then downscaled using the Regional Climate Model version 4.7.0 (RegCM4.7.0) developed at the Abdus Salam International Centre for Theoretical Physics. This results in three different RCM realizations corresponding to the three GCMs. See Supplementary text 1 for more information on the climate data.

Peatland degradation features used in the neural network model included the percent of different land use types, tree cover fraction, drainage canal density, fire area, and fire count. These factors are likely to change significantly in the future, but it is difficult to predict how they will change due to shifting economic incentives and regulations (Suwarno et al 2018, Schoneveld et al 2019, Humpenöder et al 2020). We therefore only considered changes in climate variables in this study, but incorporated these additional land use and fire inputs to account for their effect on the soil moisture-climate relationship. Location descriptors including latitude, longitude, region, and distance from the edge of the peat dome were also used as predictors to account for possible spatial autocorrelated factors affecting soil moisture,

such as land use history, peat physical properties, and land management practices. See Supplementary Text 1 and Supplementary table 1 for more information on the input features and neural network structure.

2.4. Neural network prediction of soil moisture

The neural networks were trained using remotely sensed soil moisture from SMAP over the 2015–2020 period. To determine how soil moisture statistics were affected by climate change, the neural networks were then run with a set of regional climate predictions dynamically downscaled from three global climate predictions for a reference (1985–2005) and future time period (2040–2060). To reduce the effect of biases in the global circulation models downscaled by a RCM, all climate inputs were bias-corrected to match the statistics of an observation-driven dataset, here the European Centre for Medium-Range Weather Forecasts ERA5 reanalysis product (Hersbach *et al* 2019).

We compared predictions of smdry season and pctlow sm between the reference (1985-2005) and future periods (2040-2060). In each case, degradation and location input features were held constant while climate features changed based on bias-corrected RCM predictions. Bias correction of the climate data was necessary because there are biases between the RCM simulations and the pseudo-observational ERA5 data. These differences in distributions would otherwise result in projections of soil moisture incorrectly attributed to changing climate that are instead due to differences between ERA5 and the RCM. We used quantile mapping to correct these biases (Reichle and Koster 2004, Miao et al 2016). Specifically, we matched reference period RCM data to ERA5 data from the same time period, and then applied the same correction to future period RCM data. A separate quantile mapping was applied to each of the three RCM realizations (corresponding to each global circulation model). Both RCM and ERA5 data used for bias-correction were downscaled to 9 km resolution from their original 25 and 30 km grids, respectively, using nearest neighbor resampling.

2.5. Neural network models assessment

The neural networks' performances were evaluated in two different ways using cross-validation (CV). First, to assess overall model performance on unseen data, 5-fold 'random' cross validation was performed. This means that a model was trained on a random selection of 80% of the data, then predictions on the unseen 20% of the data were compared to observations. The training data and testing data were cycled through till all data had been tested in this manner. Alternatively, to assess the models' abilities in predicting interannual variability, 6-fold 'temporal' cross validation was performed, meaning that the model was trained on 5 years of data, then tested on the remaining 6th year of data. Prediction accuracy was then assessed using the coefficient of determination (R^2) (which varies from 0 to 1 with 1 indicating higher agreement between the model prediction and observation), bias, and root-mean-squared error.

3. Results and discussion

3.1. Soil moisture models assessment

Cross validation for both soil moisture variables, sm_{dry season} and pct_{low sm}, demonstrated that the neural network models could predict out-ofsample data accurately (table 1, Supplementary figure 2). The sm_{dry season} model achieved a CV mean $R^2 = 0.83$, RMSE = 0.08 cm³ cm⁻³, and a bias of 0.001 cm³ cm⁻³ on randomly sampled test data. Similarly, the pct_{low sm} model achieved a CV mean $R^2 = 0.73$, RMSE = 16%, and a bias of 0.8% on random test data. When the two networks were crossvalidated using a full year's worth of held-out data, R^2 decreased only a slight amount ($\Delta R^2 \approx 0.1$ in both cases), suggesting the networks were able to predict soil moisture behavior on unseen years of data, including simulated future years.

3.2. RCM predicts drier future atmospheric conditions

RCM projections show overall drying in the study region, as dry season precipitation is projected to decrease across 89% of the area (figure 2(a)), while PET is projected to increase across 98% (figure 2(b)). The median change in dry season precipitation is -0.79 mm d^{-1} and the median PET change is $+0.38 \text{ mm d}^{-1}$ between the reference (1985–2005) and future (2040–2060) periods (Supplementary figure 3(a)). Geographically, there are larger decreases in dry season precipitation in southern Sumatra and larger increases in dry season PET in the southern parts of the study region (figure 2). Because ET is the dominant water flux out of peatlands (e.g. Hirano *et al* 2015, Cobb and Harvey 2019), increased PET is expected to lead to decreases in soil moisture.

Annual precipitation is projected to decrease by \sim 0.5–2 mm d⁻¹ in the study region (figure 2(c), Supplementary figure 3(b)). Precipitation seasonality, as captured by precipitation entropy, exhibited a mixed change in signal by latitude in Sumatra: generally decreasing south of the equator and increasing north of it (figure 2(d), Supplementary figure 3(b)). Decreasing entropy suggests higher seasonality, which may cause drier sm_{dry season}, as precipitation may be less evenly distributed between the dry and wet seasons. These results are consistent with those of Kang et al (2019), who found that Aug–Oct precipitation (corresponding to the dry season across most of the study area) generally decreased while Nov-Jan precipitation generally increased. While our model did not account for possible changes in the timing of the dry season, only relatively minor changes are

projected in the timing of the monsoon in this region (Ashfaq *et al* 2020). Overall distributions of climate features shifted under future climate (Supplementary figure 3), but these shifts generally did not extend far beyond the ranges observed under future climate. This builds confidence that the neural networks trained using present climate-soil moisture relationships can accurately assess the impact of future climate scenarios.

3.3. Climate changes cause substantially drier soils Both soil moisture variables exhibited drier conditions under 2040–2060 climate projections compared to 1985-2005 climate, consistent with the changes in climate forcing. Median smdry season was projected to decrease during the future period by 0.023 cm³ cm⁻³ (figures 3(a) and (c)). For context, this decrease is nearly half the magnitude of the 0.056 cm^3 cm^{-3} decrease in median dry season soil moisture observed by SMAP during the 2015 and 2019 El Niño years relative to non-El Niño years between 2015 and 2020. Recent El Niño years have been associated with a nonlinear increase in fire activity (Yin et al 2016), suggesting that the magnitude of climate-change induced soil moisture drying, absent other changes, could significantly increase fire risk in the region. However, the impacts of climate change relative to recent El Niño years differ geographically. Here, we found that the predicted soil drying due to climate change is generally greater than impacts observed during recent El Niño droughts north of the equator, while the opposite is true south of the equator in the study region (figures 4(a) and (b)).

The pct_{lowsm} variable, a more direct measure of fire risk than sm_{dry season}, increases over almost the entire region. Our neural network projected a median increase in pctlow sm of 3% (from 12.5% to 15.5%) (figures 3(b) and (d)), suggesting that extremely dry conditions associated with high fire risk will be more prevalent in the future. To estimate how large the pct_{low sm} associated impact on burned area might be, we consider a single average burned area associated with dry soil moisture (below 0.2 cm³ cm⁻³) and another average burned area for wet soil moisture conditions (as calculated from the curve in figure 3(a)of Dadap et al 2019). The increase of the 3% in pctlow sm would then correspond to a 10% increase in burned area due to future climate change. This calculation, though highly simplified, illustrates the outsized increase in fire risk associated with even small increases in pct_{low sm} driven by climate change.

Drought conditions during recent El Niño years have been attributed primarily to precipitation drought (e.g. Field *et al* 2016), but our model suggests that future changes in sm_{dry season} are also affected by increased evaporative demand (i.e. increasing PET). This is evident from the higher feature importance of PET compared to precipitation inputs for both neural networks (Supplementary figure 4).

Table 1. Cross-validation ('CV') results \pm standard deviation across folds. Temporal CV was performed by holding out one year of data at a time for the test set, and training on the other years. For example, the data would be trained on 2015–2019 data and evaluated on unseen 2020 data. This was then repeated for all six years of data. Random CV involved random selection of data from all years (across all pixel-times) when performing five-fold cross validation.

Model	Random CV train R^2	Random CV test R ²	Temporal CV train R^2	Temporal CV test R^2
sm _{dry season} pct _{low sm}	$\begin{array}{c} 0.95 \pm 0.01 \\ 0.92 \pm 0.02 \end{array}$	$\begin{array}{c} 0.83 \pm 0.02 \\ 0.73 \pm 0.03 \end{array}$	$\begin{array}{c} 0.90 \pm 0.08 \\ 0.91 \pm 0.03 \end{array}$	$\begin{array}{c} 0.73 \pm 0.12 \\ 0.64 \pm 0.13 \end{array}$



Figure 2. Mean change in climate variables between reference (1985–2005) and future (2040–2060) periods for (a) dry season precipitation, (b) dry season PET, (c) annual precipitation and (d) precipitation entropy. Red indicates drier dry season conditions; note the colorbar is reversed in (b). Non-peat areas are shown in gray. These four variables make up the input climate features in the neural networks.

Consistent with this finding, running the model with future (2040–2060) PET but with reference (1985–2005) precipitation resulted in a decrease in median $sm_{dry season}$ that was 0.008 cm³ cm⁻³, or 36% of the change when precipitation drivers were included. Thus, our results suggest that increased evaporative demand will play a significant role in driving soil moisture changes under climate changes. Land-atmosphere feedbacks may further exacerbate soil drought and atmospheric aridity under future climate (Zhou *et al* 2019).

3.4. Degraded areas are more sensitive to climate change

To better understand where soil moisture changes will occur, we separated model predictions by land use (here determined by the majority land use type in each pixel). During the reference period (1985–2005), pristine forest was predicted to have the wettest median $\text{sm}_{dry \text{ season}}$, while open undeveloped was the driest (figure 4(a)). Nevertheless, reference period distributions of $\text{sm}_{dry \text{ season}}$ were generally found to have little variation across land uses (figure 4(a)). This was somewhat surprising, as land use is often used as a proxy for hydrologic disturbance (e.g. Miettinen *et al* 2017, Taufik *et al* 2020). However, our model predictions were mostly consistent with a meta-analysis of *in situ* soil moisture measurements, which show similar soil moisture magnitudes across land use types and large variation within land uses (Supplementary figure 5, Supplementary table 2). Such high variability of soil moisture within land use types is likely due to differences in precipitation regimes, peat physical properties, drainage density, and more Kurnianto *et al* 2019, Aldrian and Dwi Susanto 2003, Dadap *et al* 2021.

Degraded land use types (including degraded forest, open undeveloped, smallholder plantation, and industrial plantation) exhibit larger magnitudes of drying than pristine forest (figures 5(c) and (d)). In particular, open undeveloped areas are predicted to experience the largest changes, while pristine forests are predicted to experience the smallest changes. Open undeveloped areas generally have the lowest starting soil moistures, suggesting that the driest areas will dry further than wetter areas. The differences in soil moisture changes by land use type could be caused by (a) climate changing more in certain land



Figure 3. Changes in soil moisture variables between reference (1985–2005) and future (2040–2060) time periods. (a) Probability distributions for $sm_{dry \, season}$ smoothed by a kernel density estimator. (c) Cumulative distributions for $pct_{low \, sm}$. For (a) and (b), thin lines denote individual GCM climate projections while the thick line denotes mean distribution across GCMs. (c) and (d) Histograms showing per-pixel change in $sm_{dry \, season}$ and $pct_{low \, sm}$ due to climate change.



Figure 4. Comparison of future climate impacts with present day El Niño. (a) Difference in predicted $\Delta sm_{dry \, season}$ due to climate change vs $\Delta sm_{dry \, season}$ observed during recent El Niño years (2015 & 2019). (b) Same as in (a) but for $\Delta pct_{low \, sm}$. Non-peat areas are shown in gray.

use types and/or (b) certain land use types are inherently more sensitive to changes in climate. However, the former does not appear to be a major factor, because the soil moisture changes ($\Delta sm_{dry\,season}$ and $\Delta pct_{low\,sm}$) vary independently of the changes in climate variables ($\Delta precip$ and ΔPET) when grouped by land use type (figure 6), except for increases in PET with decreases in $sm_{dry\,season}$. This suggests that land use could affect the sensitivity of soil moisture response to climate change. Our results further suggest that tree cover affects soil moisture sensitivity to climate change. We regressed $\Delta \text{sm}_{\text{dry season}}$ and $\Delta \text{pct}_{\text{low sm}}$ with the input metrics that capture peatland degradation (tree cover, canal density, and fire), and found significant relationships for both variables only with tree cover (Supplementary figure 6). These relationships suggest that areas with less tree cover are more sensitive to climate changes (i.e. will experience more drying) than areas with more tree cover. This increased sensitivity with



Figure 5. Soil moisture distributions grouped by land use type for (a) $sm_{dry season}$ and (b) $pct_{low sm}$ during reference (1985–2005) and future (2040–2060) periods. Box denotes inter-quartile range and median. Change in median (c) $sm_{dry season}$ and (d) $pct_{low sm}$ from reference to future periods.



climate variables (dry season PET and dry season precipitation). Changes in soil moisture do not appear to vary with changes in climate. Note the signs for sm_{dry season} and for dry season PET denote negative change.

less tree cover can be explained by a number of possible mechanisms. First, tree cover reduces the solar radiation reaching the ground surface. In areas with less or shorter vegetation, this effect is minimized, and atmospheric conditions are more likely to determine changes in soil evaporation (Fan *et al* 2019, Ohkubo *et al* 2021). Deforested areas are also more likely to contain degraded soils with increased hydrophobicity (Bechtold *et al* 2018, Perdana *et al* 2018). This in turn could decrease rainfall infiltration, increase soil evaporation, and decrease the capillary connection with the water table and the surface soil, making degraded areas more sensitive to climate changes. Furthermore, reduced hydraulic diversity (Anderegg *et al* 2018), shallower roots, or less stomatal regulation (Manoli *et al* 2018) are characteristic of agricultural areas that have lower tree cover fraction.

It should also be noted that SMAP soil moisture measurement could be affected by differences in peat microtopography by land use type, complicating comparisons of soil moisture between land use types. For example, the duff and litter layers that form the hummock and hollow topography endemic to pristine peatlands are often replaced by a denser, flatter surface when graded or converted to agricultural use (Lim *et al* 2012). These differences could in turn affect the profile of soil moisture measurement relative to the groundwater table. For example, Sakabe

et al (2018) found high variability in surface soil moisture within pristine forests based on the location of measurement: hummocks averaged 0.06 cm³ cm⁻³ while hollows averaged 0.54 cm³ cm⁻³, but the drier value would not necessarily imply higher fire risk. Such small-scale spatial variability would be averaged to a single measurement by SMAP, which integrates measurements over 9 km pixels. However, this variability would not exist in land use types where the ground surface is generally flatter. Thus, in situ validation studies are needed to better understand how to interpret differences in SMAP retrievals between land use types and their implications for fire risk and carbon emissions. Nonetheless, comparisons within land use types would not be affected by this potential issue, and the predicted drying trends observed in all land use types underscores the consistent prediction of drying due to climate change.

4. Conclusions

Our model projections suggest that future drier climatic conditions across Southeast Asia will lead to lower mean soil moisture and more frequent periods with dangerously dry peat conditions that would lead to increased fire risk. The median predicted decreases in soil moisture are nearly half the magnitude of those experienced during high-fire drought years associated with El Niño under current climate, portending more prevalent fire risk due to climate change. More research is needed to understand the impact of changes in El Niño severity or changes in dry season length, two factors that were not considered in this study. In contrast to recent droughts, future drier soil conditions also appear to be driven by increased evaporative demand in addition to reduced precipitation. Further work is needed to assess the combined and interacting impacts of changing land use-which will mediate how evaporative demand changes will affect future ET and thus ultimate soil conditions-and changing climate, thus requiring the development of detailed land use change scenarios. Our findings suggest that more degraded peatlands with lower tree cover may be especially sensitive to climate change, motivating the importance of restoration in not only reducing current carbon emissions and fire risk, but also towards lessening the impacts from future climate change. Degradation is understood to be a critical determinant of peatland hydrology, but our results suggest that climate change will also play an important role in determining future soil moisture regimes.

Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: https://github.com/ndadap/future-sm-peatlands, doi: 10.5281/zenodo.6740137.

Acknowledgments

N C Dadap was supported by NASA Headquarters under the NASA Earth and Space Science Fellowship Program-Grant 80NSSC18K1341. This work is supported by NSF under Award EAR-1923478 to A G Konings and C F Harvey. This research was also supported by the National Research Foundation (NRF), Prime Minister's Office, Singapore under its Campus for Research Excellence and Technological Enterprise (CREATE) program and Grant No. NRF2016-ITCOO1-021. The Center for Environmental Sensing and Modeling (CENSAM) is an interdisciplinary research group (IRG) of the Singapore MIT Alliance for Research and Technology (SMART). A F Feldman was supported by an appointment to the NASA Postdoctoral Program at the NASA Goddard Space Flight Center, administered by Oak Ridge Associated Universities under contract with NASA.

ORCID iDs

Nathan C Dadap lo https://orcid.org/0000-0001-7586-6210

Alexander R Cobb () https://orcid.org/0000-0002-3128-3002

Alison M Hoyt in https://orcid.org/0000-0003-0813-5084

Charles F Harvey b https://orcid.org/0000-0002-7759-4447

Andrew F Feldman b https://orcid.org/0000-0003-1547-6995

Eun-Soon Im [©] https://orcid.org/0000-0002-8953-7538

Alexandra G Konings https://orcid.org/0000-0002-2810-1722

References

- Aldrian E and Dwi Susanto R 2003 Identification of three dominant rainfall regions within Indonesia and their relationship to sea surface temperature *Int. J. Climatol.* 23 1435–52
- Anderegg W R L *et al* 2018 Hydraulic diversity of forests regulates ecosystem resilience during drought *Nature* **561** 538–41
- Andrew R and Peters G 2021 The global carbon project's fossil CO₂ emissions dataset Figshare Preprint (https://doi.org/ 10.6084/m9.figshare.16729084.v1)
- Anshari G, Peter Kershaw A and Van Der Kaars S 2001 A late pleistocene and holocene pollen and charcoal record from peat swamp forest, Lake Sentarum wildlife reserve, West Kalimantan, Indonesia *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **171** 213–28
- Ashfaq M *et al* 2020 Robust late twenty-first century shift in the regional monsoons in RegCM-CORDEX simulations *Clim. Dyn.* **57** 1463–88
- Baird A J, Low R, Young D, Swindles G T, Lopez O R and Page S 2017 High permeability explains the vulnerability of the carbon store in drained tropical peatlands *Geophys. Res. Lett.* 44 1333–9
- Bechtold M, Dettmann U, Wöhl L, Durner W, Piayda A and Tiemeyer B 2018 Comparing methods for measuring water retention of peat near permanent wilting point *Soil Sci. Soc. Am. J.* 82 601–5

- Bircher S, Demontoux F, Razafindratsima S, Zakharova E, Drusch M, Wigneron J and Kerr Y H 2016 L-band relative permittivity of organic soil surface layers—a new dataset of resonant cavity measurements and model evaluation *Remote Sens.* 8 1024
- Brodzik M J, Billingsley B, Haran T, Raup B and Savoie M H 2012 EASE-grid 2.0: incremental but significant improvements for earth-gridded data sets *ISPRS Int. J. Geo-Inf.* **1** 32–45

Cobb A R, Dommain R R, Tan F, Heng N H E H and Harvey C F 2020 Carbon storage capacity of tropical peatlands in natural and artificial drainage networks *Environ. Res. Lett.* 15 23–25

Cobb A R and Harvey C F 2019 Scalar simulation and parameterization of water table dynamics in tropical peatlands *Water Resour. Res.* **55** 9351–77

Cobb A R, Hoyt A M, Gandois L, Eri J, Dommain R, Abu Salim K, Kai F M, Haji Su'ut N S and Harvey C F 2017 How temporal patterns in rainfall determine the geomorphology and carbon fluxes of tropical peatlands *Proc. Natl Acad. Sci.* E5187–96

- Dadap N C, Cobb A R, Hoyt A M, Harvey C F and Konings A G 2019 Satellite soil moisture observations predict burned area in Southeast Asian peatlands *Environ. Res. Lett.* 14 094014
- Dadap N C, Hoyt A M, Cobb A R, Oner D, Kozinski M, Fua P V, Rao K, Harvey C F and Konings A G 2021 Drainage canals in Southeast Asian peatlands increase carbon emissions *AGU Adv.* **2** 1–14
- Dommain R, Couwenberg J and Joosten H 2010 Hydrological self-regulation of domed peatlands in south-east Asia and consequences for conservation and restoration *Mires Peat* **6** 1–17

Entekhabi D et al 2010 The Soil Moisture Active Passive (SMAP) mission Proc. IEEE 98 704–16

- Fan Y, Meijide A, Lawrence D M, Roupsard O, Carlson K M, Chen H-Y Y, Röll A, Niu F and Knohl A 2019 Reconciling canopy interception parameterization and rainfall forcing frequency in the community land model for simulating evapotranspiration of rainforests and oil palm plantations in Indonesia J. Adv. Model. Earth Syst. 11 732–51
- Feldman A F, Konings A G, Piles M and Entekhabi D 2021 The multi-temporal dual channel algorithm (MT-DCA) (version 4) (https://doi.org/10.1109/jstars.2021.3124857)
- Feng X, Porporato A and Rodriguez-Iturbe I 2013 Changes in rainfall seasonality in the tropics Nat. Clim. Change 3 811–5
- Field R D et al 2016 Indonesian fire activity and smoke pollution in 2015 show persistent nonlinear sensitivity to El Niño-induced drought Proc. Natl Acad. Sci. 113 9204–9

Field R D, Van Der Werf G R and Shen S S P P 2009 Human amplification of drought-induced biomass burning in Indonesia since 1960 *Nat. Geosci.* 2 185–8

Frandsen W H 1997 Ignition probability of organic soils *Can. J. For. Res.* **27** 1471–7

Giorgi F *et al* 2021 The CORDEX-CORE EXP-I initiative: description and highlight results from the initial analysis *Bull. Am. Meteorol. Soc.* **103** E293–E310

Goldammer J G, Seibert B and Kehutanan F 1989 Natural rain forest fires in Eastern Borneo during the pleistocene and holocene *Naturwissenschaften* **76** 518–20

Goldstein A *et al* 2020 Protecting irrecoverable carbon in Earth's ecosystems *Nat. Clim. Change* **10** 287–95

Hersbach H *et al* 2019 ERA5 monthly averaged data on single levels from 1979 to present

Hirano T, Jauhiainen J, Inoue T and Takahashi H 2009 Controls on the carbon balance of tropical peatlands *Ecosystems* 12 873–87

Hirano T, Kusin K, Limin S and Osaki M 2014 Carbon dioxide emissions through oxidative peat decomposition on a burnt tropical peatland *Glob. Change Biol.* **20** 555–65

Hirano T, Kusin K, Limin S and Osaki M 2015 Evapotranspiration of tropical peat swamp forests *Glob. Change Biol.* **21** 1914–27

Hooijer A, Page S, Jauhiainen J, Lee W A, Lu X X, Idris A and Anshari G 2012 Subsidence and carbon loss in drained tropical peatlands *Biogeosciences* **9** 1053–71

- Hoyt A M, Chaussard E, Seppalainen S and Harvey C 2020 Widespread subsidence and carbon emissions across Southeast Asia peatlands *Nat. Geosci.* **13** 435–40
- Hsu K, Gupta H V and Sorooshian S 1995 Artificial neural network modeling of the rainfall-runoff process *Water Resour. Res.* **31** 2517–30

Huang X, Rein G and Chen H 2015 Computational smoldering combustion: predicting the roles of moisture and inert contents in peat wildfires *Proc. Combust Inst.* 35 2673–81

Humpenöder F, Karstens K, Lotze-Campen H, Leifeld J, Menichetti L, Barthelmes A and Popp A 2020 Peatland protection and restoration are key for climate change mitigation *Environ. Res. Lett.* **15** 104093

Im E S *et al* 2021 Emergence of robust anthropogenic increase of heat stress-related variables projected from CORDEX-CORE climate simulations *Clim. Dyn.* **57** 1629–44

Im E S and Eltahir E A B 2018 Simulation of the diurnal variation of rainfall over the western maritime continent using a regional climate model *Clim. Dyn.* **51** 73–88

Jaenicke J, Wösten H, Budiman A and Siegert F 2010 Planning hydrological restoration of peatlands in Indonesia to mitigate carbon dioxide emissions *Mitig. Adapt. Strateg. Glob. Change* 15 223–39

Jauhiainen J, Hooijer A and Page S E 2012 Carbon dioxide emissions from an acacia plantation on peatland in Sumatra, Indonesia *Biogeosciences* **9** 617–30

Kang S, Im E S and Eltahir E A B 2019 Future climate change enhances rainfall seasonality in a regional model of western maritime continent *Clim. Dyn.* **52** 747–64

Konings A G, Piles M, Rötzer K, McColl K A, Chan S K and Entekhabi D 2016 Vegetation optical depth and scattering albedo retrieval using time series of dual-polarized L-band radiometer observations *Remote Sens. Environ.* 172 178–89

Konings A G, Williams A P and Gentine P 2017 Sensitivity of grassland productivity to aridity controlled by stomatal and xylem regulation *Nat. Geosci.* **10** 284–8

Kratzert F, Klotz D, Herrnegger M, Sampson A K, Hochreiter S and Nearing G S 2019 Toward improved predictions in ungauged basins: exploiting the power of machine learning *Water Resour. Res.* 55 11344–54

Kurnianto S, Selker J, Boone Kauffman J, Murdiyarso D and Peterson J T 2019 The influence of land-cover changes on the variability of saturated hydraulic conductivity in tropical peatlands *Mitig. Adapt. Strateg. Glob. Change* 24 535–55

Kurum M, O'Neill P E, Lang R H, Cosh M H, Joseph A T and Jackson T J 2012 Impact of conifer forest litter on microwave emission at L-band *IEEE Trans. Geosci. Remote Sens.* 50 1071–84

Leifeld J and Menichetti L 2018 The underappreciated potential of peatlands in global climate change mitigation strategies *Nat. Commun.* **9** 1071

Li W, Dickinson R E, Fu R, Niu G-Y Y, Yang Z-L L and Canadell J G 2007 Future precipitation changes and their implications for tropical peatlands *Geophys. Res. Lett.* 34 L01403

Lim K H, Lim S S, Parish F and Suharto R 2012 RSPO Manual on Best Management Practices (Bmps) for Existing Oil Palm Cultivation on Peat (Kuala Lumpur: RSPO)

Manoli G, Meijide A, Huth N, Knohl A, Kosugi Y, Burlando P, Ghazoul J and Fatichi S 2018 Ecohydrological changes after tropical forest conversion to oil palm *Environ. Res. Lett.* 13 064035

McColl K A, Alemohammad S H, Akbar R, Konings A G, Yueh S and Entekhabi D 2017 The global distribution and dynamics of surface soil moisture *Nat. Geosci.* **10** 100–4

Meinshausen M *et al* 2011 The RCP greenhouse gas concentrations and their extensions from 1765 to 2300 *Clim. Change* **109** 213–41

Mezbahuddin M, Grant R F and Hirano T 2015 How hydrology determines seasonal and interannual variations in water table depth, surface energy exchange, and water stress in a tropical peatland: modeling versus measurements J. Geophys. Res. Biogeosci. 120 2132–57

- Miao C, Su L, Sun Q and Duan Q 2016 A nonstationary bias-correction technique to remove bias in GCM simulations J. Geophys. Res. Atmos. 121 5718–35
- Miettinen J, Hooijer A, Vernimmen R, Liew S C and Page S E 2017 From carbon sink to carbon source: extensive peat oxidation in insular Southeast Asia since 1990 *Environ. Res. Lett.* 12 024014
- Miettinen J, Shi C and Liew S C 2016 Land cover distribution in the peatlands of Peninsular Malaysia, Sumatra and Borneo in 2015 with changes since 1990 *Glob. Ecol. Conserv.* 6 67–78
- Mironov V L, Dobson M C, Kaupp V H, Komarov S A and Kleshchenko V N 2004 Generalized refractive mixing dielectric model for moist soils *IEEE Trans. Geosci. Remote Sens.* 42 773–85
- Ohkubo S, Hirano T and Kusin K 2021 Influence of fire and drainage on evapotranspiration in a degraded peat swamp forest in central Kalimantan, Indonesia *J. Hydrol.* **603** 126906
- Page S E and Hooijer A 2016 In the line of fire: the peatlands of Southeast Asia *Phil. Trans. R. Soc.* B **371** 20150176
- Page S E, Rieley J O and Banks C J 2011 Global and regional importance of the tropical peatland carbon pool *Glob. Change Biol.* **17** 798–818
- Page S E, Siegert F, Rieley J O, Boehm H-D V, Jaya A and Limin S 2002 The amount of carbon released from peat and forest fires in Indonesia during 1997 *Nature* 420 61–65
- Page S *et al* 2009 Restoration ecology of lowland tropical peatlands in Southeast Asia: current knowledge and future research directions *Ecosystems* **12** 888–905
- Perdana L R et al 2018 Hydrophilic and hydrophobic characteristics of dry peat IOP Conf. Ser.: Earth Environ. Sci. 105 012083
- Reichle R H and Koster R D 2004 Bias reduction in short records of satellite soil moisture *Geophys. Res. Lett.* **31** 2–5
- Sakabe A, Itoh M, Hirano T and Kusin K 2018 Ecosystem-scale methane flux in tropical peat swamp forest in Indonesia *Glob. Change Biol.* **24** 5123–36
- Schoneveld G C, Ekowati D, Andrianto A and Van Der Haar S 2019 Modeling peat- and forestland conversion by oil palm smallholders in Indonesian Borneo *Environ. Res. Lett.* 014006
- Short Gianotti D J, Akbar R, Feldman A F, Salvucci G D and Entekhabi D 2020 Terrestrial Evaporation and moisture

drainage in a warmer climate *Geophys. Res. Lett.* **47** e2019GL086498

- Siegert F, Ruecker G, Hinrichs A and Hoffmann A A 2001 Increased damage from fires in logged forests during droughts caused by El Niño *Nature* **414** 437–40
- Sinclair A L, Graham L L B, Putra E I, Saharjo B H, Applegate G, Grover S P and Cochrane M A 2020 Effects of distance from canal and degradation history on peat bulk density in a degraded tropical peatland *Sci. Total Environ.* 699 134199
- Suwarno A, van Noordwijk M, Weikard H P and Suyamto D 2018 Indonesia's forest conversion moratorium assessed with an agent-based model of land-use change and ecosystem services (LUCES) *Mitig. Adapt. Strateg. Glob. Change* 23 211–29
- Tangang F *et al* 2020 Projected future changes in rainfall in Southeast Asia based on CORDEX–SEA multi-model simulations *Clim. Dyn.* **55** 1247–67
- Taufik M, Minasny B, McBratney A B, Van Dam J, Jones P and Van Lanen H 2020 Human-induced changes in Indonesia peatlands increase drought severity *Environ. Res. Lett.* 15 084013
- Taufik M, Torfs P J J F, Uijlenhoet R, Jones P D, Murdiyarso D and Van Lanen H A J 2017 Amplification of wildfire area burnt by hydrological drought in the humid tropics *Nat. Clim. Change* 7 428–31
- Van Der Werf G R, Randerson J T, Giglio L, Gobron N and Dolman A J 2008 Climate controls on the variability of fires in the tropics and subtropics *Glob. Biogeochem. Cycles* 22 GB3028
- Warren M, Hergoualc'h K, Kauffman J B, Murdiyarso D and Kolka R 2017 An appraisal of Indonesia's immense peat carbon stock using national peatland maps: uncertainties and potential losses from conversion *Carbon Balance and Management* 12 12
- Wösten J H M, Clymans E, Page S E, Rieley J O and Limin S H 2008 Peat-water interrelationships in a tropical peatland ecosystem in Southeast Asia *Catena* 73 212–24
- Yin Y *et al* 2016 Variability of fire carbon emissions in equatorial Asia and its nonlinear sensitivity to El Niño *Geophys. Res. Lett.* **43** 10472–9
- Zhou S, Park Williams A, Berg A M, Cook B I, Zhang Y, Hagemann S, Lorenz R, Seneviratne S I and Gentine P 2019 Land–atmosphere feedbacks exacerbate concurrent soil drought and atmospheric aridity *Proc. Natl Acad. Sci. USA* 116 18848–53