

TOPICAL REVIEW • OPEN ACCESS


## Taking stock of and advancing knowledge on interaction archetypes at the nexus between land, biodiversity, food and climate

To cite this article: Diana Sietz and Regina Neudert 2022 *Environ. Res. Lett.* **17** 113004

View the [article online](#) for updates and enhancements.

You may also like

- [Multi-tier archetypes to characterise British landscapes, farmland and farming practices](#)  
Cecily E D Goodwin, Luca Bütikofer, Jack H Hatfield et al.
- [Archetypes of community-based pond aquaculture in Indonesia: applying the social-ecological systems framework to examine sustainability tradeoffs](#)  
Ben Nagel, Nurliah Buhari and Stefan Partelow
- [The representative structure of graphene oxide nanoflakes from machine learning](#)  
Benyamin Motevalli, Amanda J Parker, Baichuan Sun et al.



**The Breath Biopsy® Guide**  
Fourth edition

FREE

DOWNLOAD THE FREE E-BOOK

BREATH BIOPSY

OWLSTONE MEDICAL

ENVIRONMENTAL RESEARCH  
LETTERS

## TOPICAL REVIEW

## OPEN ACCESS

RECEIVED  
31 May 2022REVISED  
9 October 2022ACCEPTED FOR PUBLICATION  
14 October 2022PUBLISHED  
14 November 2022

Original content from  
this work may be used  
under the terms of the  
[Creative Commons  
Attribution 4.0 licence](#).

Any further distribution  
of this work must  
maintain attribution to  
the author(s) and the title  
of the work, journal  
citation and DOI.

Taking stock of and advancing knowledge on interaction  
archetypes at the nexus between land, biodiversity, food  
and climateDiana Sietz<sup>1,\*</sup> and Regina Neudert<sup>2</sup> <sup>1</sup> Thünen Institute of Biodiversity, Federal Research Institute for Rural Areas, Forestry and Fisheries, Braunschweig, Germany<sup>2</sup> Greifswald University, Germany

\* Author to whom any correspondence should be addressed.

E-mail: [diana.sietz@thuenen.de](mailto:diana.sietz@thuenen.de)**Keywords:** nexus archetype, pattern, social-ecological system, land system, knowledge transfer, scaling

## Abstract

Global challenges related to land, biodiversity, food and climate interact in diverse ways depending on local conditions and the broader context in which they are embedded. This diversity challenges learning and integrated decision-making to sustainably transform the nexus, that is to say the interactions between these land-based challenges. Providing aggregated insights, archetype analysis has revealed recurrent patterns within the multitude of interactions, i.e. interaction archetypes that are essential to enhance the understanding of nexus relations. This paper synthesises the state of knowledge on interaction or nexus archetypes related to land, biodiversity, food and climate based on a systematic literature review. It focusses on the coverage of thematic aspects, regional distribution, social dimensions and methodologies. The results show that consideration of comprehensive land–biodiversity–food–climate interactions is rare. Furthermore, there are pronounced regional knowledge gaps, social dimensions are inadequately captured, and methodological shortcomings are evident. To enhance the investigation of interaction archetypes, we have framed a future research agenda providing directions to fully capture interactions across space and time, better use the potential of scenario archetypes and up-scale transformative actions. These advances will constructively contribute insights that help to achieve the ambitious objective to sustainably transform the nexus between land, biodiversity, food and climate.

## 1. Introduction

The post-2020 sustainability framework urgently demands that the challenges of land degradation, biodiversity decline, food insecurity and climate change must be addressed (United Nations, 2014, European Commission 2020, 2021, UNCBD 2021). Clearly, it is impossible to solve these land-based challenges separately through siloed, sectoral management approaches. Instead, the scope of action needs to be radically extended to specifically focus on the nexus, i.e. interactions between all relevant thematic aspects (Biggs *et al* 2015, Lal 2016, O'Connor *et al* 2021). For example, enhancing biodiversity and associated ecosystem services, such as water purification, pollination and biological pest control, are fundamental for re-configuring agricultural systems so that production can be sustained while ensuring

climate adaptation and mitigation (Chappell and Lavalle 2011, IPBES 2019). However, despite empirical evidence, integrated actions at large scale remain the exception and policies are misaligned (Schmidt-Traub *et al* 2019). Therefore, a central question is: what can we learn from specific local ways in which land use, biodiversity change, food production and climate dynamics co-evolve to overcome fragmentation and scale up integrative actions?

In response to this question, archetype analysis (Oberlack *et al* 2019, Sietz *et al* 2019) has revealed recurrent, i.e. archetypal patterns of factors and processes that repeatedly determine land-based interactions within the multitude of local conditions (Brzezina *et al* 2017, Hartel *et al* 2018, Dasgupta *et al* 2019, Bahri *et al*, 2020). We define interaction or nexus archetypes as recurrent patterns

of co-evolving land, biodiversity, food and climate dynamics. Interaction archetypes provide insights into functionally similar interactions focusing on an intermediate level of abstraction (Oberlack *et al* 2019) in between the particularities of single cases and panacea perspectives (Schellnhuber *et al* 2002, Basurto and Ostrom 2009). Through cumulative learning, they can enhance understanding and inform action to sustainably transform, i.e. reconfigure land–biodiversity–food–climate nexus relations. Yet, archetype research has so far often focussed on particular phenomena separately (Eisenack *et al* 2021), thereby leaving valuable potential to realise the full benefits the approach offers for tailoring policies and actions to particular contexts untapped. Highlighting this potential, archetype analysis has been proposed to advance future research on biodiversity, ecosystem services and related feedbacks within land systems, particularly in the context of scenario development (IPBES 2016). Future studies seeking to realise the benefits of archetype analysis should pay in-depth attention to validating their findings, a currently under-utilised opportunity to demonstrate their credibility and hence usefulness for informing decision-making (Piemontese *et al* 2022). Clearly defined application requirements, including intended users, spatial scale and temporal aspects of application, are prerequisites for determining the degree to which archetypes suitably respond to knowledge needs and sustainability challenges.

Sustainably transforming land-based interactions requires a sound understanding of social-ecological system dynamics (Ostrom 2009, McGinnis and Ostrom 2014, Díaz *et al* 2019). Social-ecological systems have been framed as complex adaptive systems highlighting that human land use is intrinsically tied to ecological dynamics (Allen and Garmestani 2015, Folke *et al* 2016). This reflects the close interdependence of all human and non-human beings and their environment that lies at the core of most Indigenous Peoples' notion of a good life (Acosta Espinosa 2011, Gregor Barié 2014). Hence, the capacity of all knowledge systems needs to be embraced to inform the transformation of complex adaptive systems. In complex systems, critical thresholds, cross-scale linkages and feedback loops imply non-linear behaviour and associated patterns of system behaviour (Holling and Gunderson 2002, Scheffer and Carpenter 2003, Walker *et al* 2006, Sietz *et al* 2017a, Van Den Elsen *et al* 2020). Especially the social dimensions, including governance, human behaviour and human decision-making that shape complexity in land use (Ostrom 2009) deserve greater attention. Understanding their current influence and mechanisms is crucial for sustainability transformations. Lack of integration of complexity and the role of social dimensions in knowledge and action frameworks increases uncertainty in scenarios about future

interaction trajectories (O'Connor *et al* 2021) undermining chances to reverse the persistent failure to achieve global sustainability goals (UN 2021).

This paper aims to take stock of and enhance knowledge about interaction archetypes to support learning and decision-making on the sustainable (re)configuration of land-based nexus relations. Based on a systematic literature review, we synthesise existing knowledge on interaction archetypes related to land, biodiversity, food and climate interlinkages in order to provide a state-of-the-art knowledge source. In particular, we capture the thematic aspects addressed, regional coverage, social dimensions and methodologies used to identify and apply interaction archetypes. Based on this synthesis, we frame an agenda for future research on interaction archetypes structured around three major frontiers. These frontiers provide directions to comprehensively investigate archetypical land–biodiversity–food–climate interactions, advance analysis of nexus scenario archetypes and support up-scaling and transfer of successful transformative action.

## 2. Methodology

As the basis for our synthesis, we conducted a systematic literature review (Petticrew and Roberts 2006, Booth *et al* 2012). We searched peer-reviewed studies listed in the Web of Science and Scopus databases (last access 1 April 2021) using the following combined terms applied to titles, abstracts and keywords: *archetyp\** AND ((*land* OR *farm\** OR *agric\**) AND (*biodiv\** OR '*ecosyst\* service\**' OR *food* OR *climat\**)). These searches yielded 280 unique references. We limited this set to only those studies that met the following inclusion criteria: focus on at least one type of interaction between land and biodiversity, food and climate, identification or application of archetypes and related interactions, sound methodological description, interdisciplinary approach and comprehensive interpretation of results. We excluded the subject areas of medicine, biotechnology, atmospheric science, sanitation and architecture as these subject areas are beyond our focus on land-based interactions. Since archetype analysis is an emerging field of scientific research (Oberlack *et al* 2019, Sietz *et al* 2019, Eisenack *et al* 2021), we focused our analysis on scientific literature. Using this selection procedure, we identified 50 thematic studies and two synthesis studies on archetype analysis. Although the selected thematic studies included research on syndromes, the predecessor of archetypes, capturing unsustainable patterns of human-nature interactions (Lüdeke *et al* 2004, Sietz *et al* 2006, Manuel-Navarrete *et al* 2007), we did not comprehensively cover syndrome studies. We were specifically interested in revealing the state of knowledge on archetype analysis as a rapidly

advancing field of research capturing both sustainable and unsustainable patterns (Oberlack *et al* 2019) that supports decision-making on sustainability with evidence-based and contextualised generalisations.

We coded the 50 thematic studies according to the assessed types of interactions, their regional coverage, the main land systems addressed, social dimensions, archetype definitions, methods used to investigate archetypical interactions, data sources used, units of analysis and publication year. These categories emerged inductively during coding and subsequent analysis and provide important insights for advancing the knowledge on interaction archetypes. Investigating the type of interactions and regional coverage (section 3.1) yields insights into thematic and regional knowledge gaps. The social dimensions (section 3.2) critically shape current land use and provide insights into the potential for sustainability transformations. Archetype definitions (section 3.3) reveal variants and gaps in the conceptual foundations of archetype analysis. Investigating methods (section 3.4) provides insights into different methodologies yielding diverse types and qualities of insights. Lastly, we give a short overview of the temporal development of the research field (section 3.5). The two synthesis studies were dealt with separately providing the basis for evaluating archetype definitions and methodological portfolios. These studies were not a part of the quantitative synthesis. We drew lessons from a structured discussion of the synthesised findings to elaborate the future research agenda as a basis to enable better understanding and the sustainable management of land interactions with biodiversity, food and climate.

### 3. Results and discussion

#### 3.1. Interaction types and their regional distribution

The synthesis of interaction types and regional coverage revealed insights into thematic and spatial knowledge and gaps in this knowledge. The reviewed studies primarily investigated archetypical interactions of land in combination with food (82%) and less so in combination with biodiversity (50%) and climate (48%; figure 1). These foci are a good reflection of the global debate on sustainable land and food systems (UN 2014, European Commission 2020, 2021, UNCBD 2021). Knowledge on archetypical interactions is offered for specific subregions defined by IPBES (2020). These subregions serve as a reference in our synthesis to directly feed insights into global assessments of the importance and trends of biodiversity and nature's contributions to people (e.g. Nexus Assessment and Transformative Change Assessment of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services IPBES). Some studies presented global insights either at least partly covering all IPBES (2020) subregions

or providing a generic perspective with general applicability.



Few of the studies (20%) considered the full set of land–biodiversity–food–climate interactions (figure 1). These studies addressed agricultural land systems mainly in combination with various other land uses and sectors such as forestry, nature conservation, urban areas, tourism and energy (e.g. Jäger *et al* 2015, Dasgupta *et al* 2019, Harrison *et al* 2019). These interactions were mainly investigated in central and western Europe and in Asia, including multi-regional coverage.

Focussing on less comprehensive interrelations, 16% of the studies exclusively considered land–biodiversity–food interactions (figure 1), explored in agricultural land systems, often in conjunction with nature conservation, recreation and forestry (e.g. Levers *et al* 2018, Li *et al* 2018). These studies covered primarily Europe and the global scale. A comparable share of 18% of all studies assessed only land–food–climate interactions, again focussing on agriculture (e.g. Nyam *et al* 2020, Egerer *et al* 2021) sometimes in combination with food chains, forestry and land conversion (e.g. Weissteiner *et al* 2011, Vaclavik *et al* 2013, Loiseau *et al* 2020). These studies covered mostly Europe, Africa and Asia.

Alongside the earlier interactions, specific foci on biodiversity and climate interactions are increasingly receiving attention, in various combinations (18% of the studies; figure 1). Most of these studies were published between 2018–2021 (14% of the studies; see also section 3.5). They covered either only a few regions or the global scale. For example, archetypical land–biodiversity–climate interactions (6% of the studies) were examined only at global scale (figure 1, e.g. Haberman and Bennett 2019, Higuera *et al* 2019). In contrast, land–biodiversity interactions (8% of the studies) were investigated in western Africa, southeastern Asia and at global scale (e.g. Magliocca *et al* 2019, Valette *et al* 2020). Moreover, knowledge is available on land–climate interrelations (4% of the studies) in equal parts for western Africa, the Americas and northeastern Asia (Keys and Wang-Erlandsson 2018, Evers *et al* 2019).

The largest share of the studies (28%) exclusively assessed land–food archetypes disregarding other thematic aspects (figure 1), with insights mostly published since 2015 (22% of the studies; see also section 3.5). Insights into archetypical land–food interactions are available for all subregions but the Caribbean, addressing agriculture (e.g. Zare *et al* 2017) often in combination with water management and land acquisition (e.g. Gohari *et al* 2013, Oberlack *et al* 2016, Turner *et al* 2016).

In a regional perspective, most insights are available for central and western Europe, the global scale and western Africa (figure 2). A global scale extent enables regional comparisons encompassing diverse social-ecological contexts in which




land–biodiversity–food–climate interactions evolve. However, it inevitably only captures regional specificities to some extent. This means that regional aspects of underlying processes, such as land use intensity, biodiversity management and food production, are only captured in a generalised way. In particular, spatially and temporally well-resolved quantitative data and qualitative information on underlying processes are often not available at global

scale, thereby largely constraining efforts to gain a comprehensive understanding of archetypical interactions across the globe.

Beyond the insights derived at global scale, it is particularly disadvantageous that there is low coverage of specific regions such as central and eastern Africa, Mesoamerica, South America and southern Asia (e.g. Locatelli *et al* 2017, Vidal Merino *et al* 2019). These regions include areas that bear high





risk of conflicts between biodiversity, food security and weather extremes. Risks arise particularly in those regions that include hotspots of biodiversity threats, food insecurity and future agricultural expansion (Delzeit *et al* 2017, Molotoks *et al* 2017). For example, the Peruvian Andes have not only been highlighted as a megadiverse biodiversity hotspot (Rodríguez and Young 2000) but also as an area with high risk of agricultural expansion due to large uncultivated areas suited for agriculture (Molotoks *et al* 2017). The Caribbean is least covered, revealing the most pronounced regional knowledge gaps. Questions remain, for example, about the social-ecological consequences of colonialism and their persistence until today, e.g. related to the depletion of native flora and fauna including wild food species caused by large-scale sugar cane plantations and the introduction of exotic taxa (Paravisini-Gebert 2016, Wallman 2018). These findings indicate the urgent requirement to advance our regional knowledge on nexus problems and potential solutions to address land degradation, biodiversity loss, food insecurity and aggravating impacts of climate change in synergistic ways.

### 3.2. Social dimensions of archetypal interlinkages

Insights into the social components (Ostrom 2009) of interaction archetypes help to recognise leverage points suited to trigger transformative change. Considering three social dimensions, firstly, we captured governance referring to organisations, institutions and processes that shape policies and policy-making. Secondly, we took into account the human behaviour related to the characteristics and decision-making logic of stakeholders who manage land and benefit from the products of natural resource use. Thirdly, we recorded land use decisions as the central connector between natural resources and stakeholders.


Overall, only about half of the reviewed studies ( $n = 26$ ) investigated how governance shaped archetypal interactions (figure 3). In addition, human behaviour of stakeholders was an important

topic of analysis in just 19 studies, while only 16 studies analysed land use decisions. Regarding combinations of social dimensions, merely 11 studies analysed all three social dimensions together (e.g. Banson *et al* 2016, Brzezina *et al* 2017, Almekinders *et al* 2019). Moreover, two studies (Levers *et al* 2018, Li *et al* 2018) focussed on human behaviour in combination with land use decisions, while one study (Oberlack *et al* 2016) investigated governance and human behaviour at the same time. In contrast, 14 studies did not investigate any of the social dimensions considered in our review.

The limited consideration of social dimensions, and above all combinations of these, greatly restricts our knowledge and capacity to (re)organise land use in a way that fosters synergies and minimises trade-offs between land, biodiversity, food and climate concerns. For example, this concerns the allocation of costs and benefits among stakeholders and also impedes balancing stakeholders' interests. This resembles the inadequate consideration of the ways in which governance affects other land-based interactions such as framed in the water-energy-food nexus (Urbinnati *et al* 2020). A better understanding of the ways in which governance, human behaviour and land use decisions typically co-evolve and influence land-biodiversity-food-climate interactions is clearly needed to develop response options that adequately address social-ecological dynamics. Consideration of sudden changes, feedbacks, historical trajectories and dynamics of the broader social context (Ostrom 2009, Cumming *et al* 2020) is important to effectively manage leverage points for transformative change.

### 3.3. Archetype definitions

Variants of and gaps in archetype definitions yield information on the conceptual foundations of the studies reviewed. Archetypes were defined from the perspectives of diverse meanings, features and reference literature. The meanings used to describe and



evaluate archetypes included case typologies, building blocks and scenario archetypes (Oberlack *et al* 2019). These were further differentiated according to specific features underlying the archetype analysis encompassing recurrent patterns, system archetypes, syndromes and scenarios.

In terms of specific meanings, the majority of studies (58%, figure 4) used a case typology approach capturing similarities across a set of cases. In contrast, around a quarter (28%) applied a building blocks approach in which individual cases can be characterised by one or a combination of several archetypes. Taking yet another perspective, a small proportion of studies (14%) investigated scenario archetypes depicting common features of possible future interaction trajectories.

Studies that used a case typology approach most frequently identified recurrent patterns (figure 4) in factors and processes that shape the interactions between land, biodiversity, food and climate (e.g. Adenle and Speranza 2021). Reference was mainly made to archetype definitions used in sustainability research (Oberlack *et al* 2019, Sietz *et al* 2019) and land system science (Vaclavik *et al* 2013). Some studies related to other schools of thought, e.g. regularities in economies (Sadoulet and Dejanvry 1992) and prototypes (Theodosiou *et al* 2012; based on Eugster and Leisch 2009).


In contrast, the majority of studies applying a building blocks approach used system archetypes as diagnostic tools to establish or synthesise insights into characteristic structures and dynamics of social-ecological systems (figure 4). These studies either diagnosed system dynamics or tested a hypothesised system change (e.g. Nyam *et al* 2020). Reference was specifically made to the field of system archetypes

(e.g. Senge 2006) investigating typical causal interlinkages reappearing across many cases such as the 'limits to growth' system archetype (Gohari *et al* 2013). System archetypes offer particular potential for analysing interactions between land, biodiversity, food and climate due to their focus on reappearing systemic interlinkages.

Among the studies approaching archetypes as case typologies and building blocks, only two studies used syndromes, i.e. recurrent non-sustainable human-nature interactions, as specific features (figure 4). They described change processes as archetypal dynamic and co-evolutionary patterns of human-nature interactions (e.g. Weissteiner *et al* 2011), referring to Downing and Lüdeke (2002) and Müller *et al* (2014). These studies take integrative perspectives capturing complex sets of factors at multiple levels and in multiple sectors, aiming to overcome single-faceted sectoral approaches.

Most studies that investigated scenario archetypes expectedly referred to scenarios as specific features (figure 4). These referenced literature on 'scenario archetypes' or 'scenario families' (e.g. Pedde *et al* 2019) or the Global Environmental Outlook-4 (UNEP 2007, Dasgupta *et al* 2019). There is clearly potential to extend and deepen the use of scenario archetypes in the future as they hold great potential for visioning about desired future development (Kubiszewski *et al* 2016, Pereira *et al* 2018) and factors that foster or impede specific interaction pathways.

Overall, about one quarter of the reviewed studies (28%) neither provided an archetype definition nor reference literature. The proportion of studies without reference literature was found to be particularly high in those using a case typology approach (figure 4). This may impede the drawing



of comprehensive lessons aimed at informing transformative action, hence necessitating improvement in future research.

### 3.4. Methodologies

The reviewed studies employed methodologies resting on various methods, data sources and units of analysis. Influencing the type of insights gained from a study, these methodological aspects serve to reflect on epistemological and normative underpinnings, data requirements and implications in view of specific purposes and research questions (Sietz *et al* 2019). The methods used to identify and apply interaction archetypes stem from a broad methodological portfolio which has grown and diversified since the synthesis of the archetype methods applied in sustainability research (Sietz *et al* 2019). Overall, the reviewed studies used mainly qualitative methods (44%) and to a lesser extent quantitative (30%) and mixed methods, i.e. combining qualitative and quantitative approaches (26%). 36% of the studies used more than one method.

Among the purely qualitative studies (figure 5), the most frequently used method was qualitative system modelling (e.g. Turner *et al* 2016, Bahri 2020), often used to identify system archetypes (see section 3.3). Moreover, expert and stakeholder assessments, theoretical conceptualisations (e.g. Felgenhauer 2015, Hartel *et al* 2018) and qualitative classifications (e.g. Bay-Larsen *et al* 2018) also played a prominent role. Expert and stakeholder assessments were always used in combination with other methods, mainly with qualitative system modelling (e.g. Queenan *et al* 2020).


In the solely quantitative studies, cluster analysis was the most relevant method (e.g. Lim-Camacho *et al* 2017, Sietz *et al* 2017b, Tessier *et al* 2021). Less frequently used methods included meta-analysis (Gerst *et al* 2015, Lee and Lautenbach 2016), machine-learning algorithms (e.g. Adenle and Speranza 2021) and measures of statistical distance or similarity (e.g. Vaclavik *et al* 2016).

The mixed method studies most frequently combined qualitative system modelling with quantitative integrative modelling and meta-analysis with qualitative classification. Further studies combined machine-learning algorithms with an expert/stakeholder assessment and qualitative classification with quantitative integrative modelling. Spatial statistics and rule-based classifications used to identify social-ecological system archetypes (Sietz *et al* 2019) were not found in this review.


The reviewed studies used one or several types of data sources (figure 6). Firstly, existing data were used whereby available statistical, census and remote sensing data were drawn on. Secondly, literature information was extracted from scientific publications. Thirdly, participatory data collection took place. This involved processes in which stakeholders took part in the data collection. Fourthly, self-collected data were gathered through interviews and surveys of experts and stakeholders.

The most frequently used data source was scientific literature, followed by existing data, participatory data collection and self-collected data (figure 6). Most studies (64%) used only one data source. In contrast, 24% of the studies involved two sources, e.g. drawing on existing data in combination with





**Figure 6.** Data sources and combinations of these sources used in the reviewed studies. ( $n$  = number of studies; Note: Dotted lines indicate combinations of data sources in non-adjacent cells.)



**Figure 7.** Units of analysis considered in the reviewed studies.

literature (e.g. Rocha *et al* 2020), combining literature and participatory data collection (e.g. Moraine *et al* 2017) or linking participatory and self-collected data (e.g. Karrasch *et al* 2019). Moreover, 6% of the studies used three data sources, including a combination of literature, participatory and self-collected data (e.g. Armatas *et al* 2018). Taking an even more comprehensive approach, 6% of the studies combined all four data sources (e.g. Bahri 2020).


Regarding units of analysis (figure 7), functional relationships most frequently underlay the archetype analyses (44%), often used to establish causal loop diagrams and investigate system archetypes (e.g. Banson *et al* 2016, Turner *et al* 2017, Bahri 2020). Since we reviewed studies of archetypical interactions, the units of analysis in nearly all studies had a functional dimension, but the studies focussing explicitly on functional relationships put this aspect in the foreground. Spatially explicit units were also prominently used (30%) such as grid cells and landscapes (e.g. Fischer *et al* 2017). Other units of analysis encompassed scenarios (12%; e.g. Thompson 2018, Leventon *et al* 2019), individual stakeholders

(10%; e.g. Bay-Larsen *et al* 2018) and generic units such as conservation areas (4%, e.g. Whitelaw *et al* 2014).

Taken together, qualitative methods, functional relationships and information provided in scientific literature mostly informed the investigations on archetypical interactions, in various combinations. The knowledge derived provides valuable insights into principal relationships between factors and processes that recurrently shape the interactions. However, limited use of quantitative methods as well as participatory and self-collected data on stakeholders and scenarios leaves considerable unexplored potential to better understand the complexity determining the interactions and adaptive management options.

### 3.5. Publication year

The publication year provides information on the temporal development of research on archetypical interactions. The reviewed studies have been published since 1992, yet the majority (88%) have appeared since 2015 (figure 8). Until 2013, studies captured only archetypical land–food–climate



and land–food interactions. Full sets of land–biodiversity–food–climate interactions were published between 2015–2019, mainly coinciding with less comprehensive land–biodiversity–food and land–food interactions. Specific foci on land–biodiversity interactions have been published since 2019 and land–climate interactions have regained attention since 2018.

#### 4. Future research agenda

Our synthesis demonstrates that knowledge on archetypal interactions is limited in terms of inter-linked thematic aspects, regional coverage, social dimensions and methodologies. The future agenda highlights research directions to capture archetypal interactions more comprehensively, close regional knowledge gaps, make better use of the potential of scenario archetypes and inform the up-scaling of actions.


##### 4.1. Comprehensively capturing archetypal interactions across regions and scales

The next generation of interaction archetypes needs to pay attention to the full set of interactions between land, biodiversity, food and climate (see sections 3.1 and 3.2; figure 9(a)) to enable a solid understanding of the ways in which all involved thematic aspects typically co-evolve. Moving from fragmented perspectives to interaction archetypes helps to better understand recurrent challenges and opportunities (figure 9(a)). Investigation of system archetypes (see section 3.3) provides a particularly suitable way to synthesise feedback mechanisms and diagnose or test hypotheses about archetypal dynamics. However, greater efforts are needed to trace patterns over time. Particular consideration needs to be dedicated to integrated analytical approaches that capture complexity, i.e. interdependency and feedback mechanisms (Scheffer and Carpenter 2003, Suding 2011, Cumming *et al* 2020). Important characteristics to be addressed encompass non-linear dynamics including thresholds, resilience, time lags and reversibility (figure 9(b); Holling 1973, Walker and Meyers 2004,

Walker *et al* 2006, Van Den Elsen *et al* 2020). Stability landscapes (figure 9(b)) are useful depictions for exploring non-linear interaction dynamics as a basis for analysing future scenario archetypes and adaptive management options. In the absence of alternative stable states, an existing, less sustainable stable state (see stability landscape I in figure 9(b)) requires adaptive management that substantially reconfigures the current social-ecological conditions so that another, more sustainable stable state emerges and a shift can occur. Yet, if another, more sustainable stable state exists, then there is potential for transformative change. In this case, interactions can be tipped out of the basin of attraction of their current state to shift to the alternative stable state (see stability landscape II in figure 9(b)). In contrast, more sustainable archetypal interactions require efforts to maintain or further improve their current state (see stability landscape III in figure 9(b)).

Systematic collection of time series data on interactions is key to be able to explore and effectively manage nexus relations. Participatory and self-collected data (see section 3.4) offer particular opportunities to capture relevant processes in necessary detail. Established long-term observation systems can provide important lessons for monitoring land–biodiversity–food–climate interactions. They highlight the importance of coordinated, comprehensive and sustained data collection to inform decision making and actions on interconnected challenges and opportunities (Bestelmeyer *et al* 2011, GEO 2015). Integration of diverse data sources, such as derived from remote sensing, *in-situ* experiments and citizen observatories, offers a powerful tool for filling functional and regional knowledge gaps (see sections 3.1 and 3.2). Predictable long-term funding is a prerequisite for ensuring that data collection, synthesis and monitoring can be continued at time scales that are sufficiently long.

In future research, attention needs to be paid to under-researched regions, above all the Caribbean, Mesoamerica, central Africa and Oceania (see section 3.1). This encompasses current hot-spots of change at the interface between biodiversity



conservation, agriculture and weather impacts (Molotoks *et al* 2017, Costello *et al* 2021). But in-depth research is also needed in regions with newly emerging risks and opportunities associated with future food system change and climate change (Sietz and Feola 2016, Leach *et al* 2020, O'Neill *et al* 2021). Interaction archetypes will ideally be investigated in spatially explicit ways. Spatial statistics (Kehoe *et al* 2015, Delzeit *et al* 2017), currently missing in the methodological portfolio (see section 3.4), provide suitable avenues to capture spatial interactions.

Trends observed at one spatial scale may evolve differently at higher or lower scales (Sietz 2014, Chase *et al* 2019). Hence, systematic understanding of the interlinking and nesting of feedbacks (Sietz *et al* 2017b) across spatial, temporal and decision-making scales (O'Connor *et al* 2021) is essential. Future research demands methodological innovation (e.g. spatially coordinated observations, multi-scale models considering pattern-process relationships; IPBES 2016) that support cross-scale synthesis. Varying the extent of the spatial scale and the resolution (Sietz 2014) are important considerations to explore non-linearity in the strength and direction of interactions. As demonstrated by ecological research, synergies and trade-offs may occur repeatedly at specific spatial or temporal scales, e.g. driven by recurrent weather extremes (Gonzalez *et al* 2020).

Advancing validation in future studies of interaction archetypes is important to strengthen

the credibility of findings. Framed as a multi-dimensional procedure, validation should address conceptual validity (i.e. salience of research framing), construct validity (i.e. attribute selection) and application validity (i.e. usefulness of results for application by users), among other dimensions (Piemontese *et al* 2022). This requires the systematic planning of validation and the collection of validation data, alongside data on drivers of change, responses and action levers, as well as multi-stakeholder engagement.

#### 4.2. Tapping the potential of scenario archetypes

Projecting possible future interaction pathways is key to identifying leverage points for adaptive management. Yet, the inherent complexity in interactions including uncertainty and surprise challenges stakeholders to make and adapt management decisions (Holling 1973, 1978). The basis of adaptive management, including governance and human behaviour (section 3.2), needs to be urgently addressed to provide insights into the conditions under which transformative change can most effectively be initiated or reinforced (Allen and Garmestani 2015). For example, analysis of scenario archetypes (see section 3.3) can help to explore the innovative potential of agroecological re-organisation of agriculture (Sietz *et al* 2022), a decoupling of resource use from habitat degradation and circular economic approaches. In particular, consideration of transformative future pathways tailored to the

characteristics of agricultural land systems (Sietz *et al* 2022) can advance the analytical power of scenario archetype models. The baseline period needs to be carefully chosen (e.g. climate baseline, Liersch *et al* 2020) as a reference for analysing interaction scenarios. Analysis of archetypical system dynamics has substantial potential to inform scenario analysis and adaptive management, above all when advancing its use in quantitative approaches (see sections 3.3 and 3.4), as it focuses on non-linear behaviour of complex systems and systemic feedbacks (figure 9(b)). It supports adaptive management in its objective to use emerging windows of opportunities (Sietz *et al* 2017a), create synergies and balance trade-offs providing a vision and direction for action.

Future scenario research is urged to integrate indigenous and local knowledge and cultural practices that have been the basis for the implementation of adaptive governance and successful navigation of transformative change. Important questions relate to the role that unity between humans and nature (IPBES 2019) can play in sustainably governing social-ecological interactions. Moreover, what implications may the Rights of Nature as part of Sumak Kawsay (Buen Vivir, 'good living') codified in the Constitution of Ecuador (Acosta Espinosa 2011, Kauffman and Martin 2017) have? What insights does nature's entitlement with rights as a collective subject of interest as framed in the Bolivian Law of Mother Earth (Pachamama; Gregor Barié 2014) offer for adaptively governing interactions in the future? Addressing these questions is essential to (re)design inclusive socio-ecological policies.

### 4.3. Scaling up successful actions

Interaction archetypes support learning from the multitude of observations and can inform decision-makers on the up-scaling potential of the insights derived. Up-scaling refers to the expansion and transfer of successful actions (Linn 2012) as an integral part of systematically fostering desired trends and synergies while balancing trade-offs. Up-scaling involves testing, transferring and refining actions in regions with similar interactions based on the assumption that similar drivers and outcomes could be addressed by comparable actions (Sietz *et al* 2011). In particular, future research needs to expand the systematic investigation of potential transfer regions (Vaclavik *et al* 2016, Piemontese *et al* 2020) and discussion of transferability of successful actions (Sietz *et al* 2017b) using spatially explicit interaction archetypes in order to inform scaling considerations.

Different stakeholders' roles in scaling, e.g. their decision making and behaviour (see section 3.2) including expectations and evaluation of interactions and outcomes, need to be better understood to ensure successful expansion and refinement of actions. Scenario archetypes (see section 4.2) need to be appropriately inclusive to support stakeholders in exploring

processes that may be triggered by up-scaling activities but cannot be fully anticipated.

Although strategies may diffuse spontaneously, systematic use of leverage points is needed to support the scaling of actions and sustain their success. This requires the identification of archetypical changes in highly influential social-ecological system properties. There are clear opportunities for investigating archetypical levers emerging at the interface between global sustainability goals. For example, climate adaptation priorities outlined in National Adaptation Plans established under the United Nations Framework Convention on Climate Change can support the achievement of Sustainable Development Goals. The future research agenda framed in this paper supports advanced analysis of archetypical leverage points as an indispensable means to design and implement the large scale action needed to transform the land–biodiversity–food–climate nexus.

### Data availability statement

All data that support the findings of this study are included within the article.

### Acknowledgments

We thank Christoph Oberlack for his feedback on the graphical representation of nexus interactions. This paper contributes to the Global Land Programme (GLP, <https://glp.earth>).

### ORCID iDs

Diana Sietz  <https://orcid.org/0000-0002-2309-2134>

Regina Neudert  <https://orcid.org/0000-0003-0910-794X>

### References

Archetype studies reviewed in this paper are listed in part B of the references.

#### A) General references

- Acosta Espinosa A 2011 Riesgos y amenazas para el Buen Vivir *Ecuador Debate* **84** 51–55
- Allen C R and Garmestani A S 2015 *Adaptive Management of Social-Ecological Systems* (Dordrecht, Netherlands: Springer) ([https://doi.org/10.1007/978-94-017-9682-8\\_14](https://doi.org/10.1007/978-94-017-9682-8_14))
- Basurto X and Ostrom E 2009 Beyond the tragedy of the commons *Economia delle Fonti di Energia e dell'Ambiente* **52** 55–60
- Bestelmeyer B T *et al* 2011 Analysis of abrupt transitions in ecological systems *Ecosphere* **2** art129
- Biggs E M *et al* 2015 Sustainable development and the water–energy–food nexus: a perspective on livelihoods *Environ. Sci. Policy* **54** 389–97
- Booth A, Papaioannou D and Sutton A 2012 *Systematic Approaches to a Successful Literature Review* (London: Sage)
- Chappell M J and Lavalle L A 2011 Food security and biodiversity: can we have both? An agroecological analysis *Agric. Hum. Values* **28** 3–26



- Chase J M *et al* 2019 Species richness change across spatial scales *Oikos* **128** 1079–91
- Costello M J, Vale M M, Kiessling W, Maharaj S, Price J and Hirak G 2021 *IPCC WGII Sixth Assessment Report. Cross-Chapter Paper 1: Biodiversity Hotspots. Final Draft 1 October 2021*
- Cumming G S *et al* 2020 Advancing understanding of natural resource governance: a post-Ostrom research agenda *Curr. Opin. Environ. Sustain.* **44** 26–34
- Delzeit R, Zabel F, Meyer C and Václavík T 2017 Addressing future trade-offs between biodiversity and cropland expansion to improve food security *Reg. Environ. Change* **17** 1429–41
- Díaz S *et al* 2019 Pervasive human-driven decline of life on Earth points to the need for transformative change *Science* **366** eaax3100
- Downing T E and Ludeke M 2002 International desertification, social geographies of vulnerability and adaption *Global Desertification: Do Humans Cause Deserts?* ed J F Reynolds and D M Stafford Smith (Berlin: Dahlem University Press) pp 233–52
- Eisenack K, Oberlack C and Sietz D 2021 Avenues of archetype analysis: roots, achievements, and next steps in sustainability research *Ecol. Soc.* **26** 31
- Eugster M J A and Leisch F 2009 From spider-man to hero—archetypal analysis in R *J. Stat. Softw.* **30** 1–23
- European Commission 2020 Farm to fork strategy for a fair, healthy and environmentally-friendly food system *European Commission* (available at: [https://ec.europa.eu/food/horizontal-topics/farm-fork-strategy\\_de](https://ec.europa.eu/food/horizontal-topics/farm-fork-strategy_de)) (Accessed 25 February)
- European Commission 2021 EU biodiversity strategy for 2030: bringing nature back into our lives *European Commission, Directorate-General for Environment, Publications Office* (available at: <https://data.europa.eu/doi/10.2779/048>) (Accessed 25 February)
- Folke C, Biggs R, Norström A V, Reyers B and Rockström J 2016 Social-ecological resilience and biosphere-based sustainability science *Ecol. Soc.* **21** 41
- GEO 2015 Strategic plan 2016–2025 implementing GEOSS: reference document (available at: [www.earthobservations.org/documents/GEO\\_Strategic\\_Plan\\_2016\\_2025\\_Implementing\\_GEOSS\\_Reference\\_Document.pdf](http://www.earthobservations.org/documents/GEO_Strategic_Plan_2016_2025_Implementing_GEOSS_Reference_Document.pdf))
- Gonzalez A *et al* 2020 Scaling-up biodiversity-ecosystem functioning research *Ecol. Lett.* **23** 757–76
- Gregor Barié C 2014 Nuevas narrativas constitucionales en Bolivia y Ecuador: el buen vivir y los derechos de la naturaleza *Latinoam. Rev. Estud. Latinoam.* **59** 9–40
- Holling C S 1973 Resilience and stability of ecological systems *Annu. Rev. Ecol. Syst.* **4** 1–23
- Holling C S 1978 *Adaptive Environmental Assessment and Management* (New York: Wiley)
- Holling C S and Gunderson L H 2002 *Panarchy: Understanding Transformations in Human and Natural Systems* (Washington, DC: Island Press)
- IPBES 2020 IPBES regions and sub-regions (1.1) IPBES technical support unit on knowledge and data (Zenodo) (<https://doi.org/10.5281/zenodo.3928281>)
- IPBES 2019 *Global Assessment Report of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services* ed E S Brondízio, J Settele, S Diaz and H T Ngo (Bonn: IPBES secretariat) p 1144
- IPBES 2016 The methodological assessment report on scenarios and models of biodiversity and ecosystem services *Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services* ed S Ferrier *et al* (Bonn: IPBES secretariat) p 348
- Kauffman C M and Martin P L 2017 Can rights of nature make development more sustainable? Why some Ecuadorian lawsuits succeed and others fail *World Dev.* **92** 130–42
- Kehoe L, Kuemmerle T, Meyer C, Levers C, Václavík T and Kreft H 2015 Global patterns of agricultural land-use intensity and vertebrate diversity *Divers. Distrib.* **21** 1308–18
- Kubiszewski I, Anderson S J, Costanza R and Sutton P C 2016 The future of ecosystem services in Asia and the Pacific *Asia Pacific Policy Stud.* **3** 389–404
- Lal R 2016 Global food security and nexus thinking *J. Soil Water Conserv.* **71** 85A–90A
- Leach M, Nisbett N, Cabral L, Harris J, Hossain N and Thompson J 2020 Food politics and development *World Dev.* **134** 105024
- Liersch S *et al* 2020 One simulation, different conclusions—the baseline period makes the difference! *Environ. Res. Lett.* **15** 104014
- Linn J F 2012 *Scaling up in Agriculture, Rural Development, and Nutrition: Lesson on Scaling-Up: Opportunities and Challenges for the Future. 2020 vision briefs* vol 19 (International Food Policy Research Institute (IFPRI))
- Ludeke M K B, Petschel-Held G and Schellnhuber H-J 2004 Syndromes of global change: the first panoramic view *GAIA—Ecol. Perspect. Sci. Soc.* **13** 42–49
- Manuel-Navarrete D, Gómez J J and Gallopín G 2007 Syndromes of sustainability of development for assessing the vulnerability of coupled human–environmental systems. The case of hydrometeorological disasters in Central America and the Caribbean *Glob. Environ. Change* **17** 207–17
- Mcginnis M D and Ostrom E 2014 Social-ecological system framework: initial changes and continuing challenges *Ecol. Soc.* **19** 30
- Molotoks A, Kuhnert M, Dawson T P and Smith P 2017 Global hotspots of conflict risk between food security and biodiversity conservation *Land* **6** 67
- Müller D, Sun Z, Vongvisouk T, Pflugmacher D, Xu J and Mertz O 2014 Regime shifts limit the predictability of land-system change *Glob. Environ. Change* **28** 75–83
- O'Connor M I *et al* 2021 Grand challenges in biodiversity-ecosystem functioning research in the era of science-policy platforms require explicit consideration of feedbacks *Proc. R. Soc. B* **288** 20210783
- O'Neill B *et al* 2021 *IPCC WGII Sixth Assessment Report. Chapter 16: Key Risks Across Sectors and Regions. Final Draft 1 October 2021*
- Ostrom E 2009 A general framework for analyzing sustainability of social-ecological systems *Science* **325** 419–22
- Paravisini-Gebert L 2016 Food, biodiversity, extinctions: Caribbean fauna and the struggle for food security during the conquest of the new world *J. West Indian Literature* **24** 11–26 (available at: [www.jstor.org/stable/90003602](http://www.jstor.org/stable/90003602))
- Pereira L M, Hichert T, Hamann M, Preiser R and Biggs R 2018 Using futures methods to create transformative spaces: visions of a good Anthropocene in southern Africa *Ecol. Soc.* **23** 19
- Petticrew M and Roberts H 2006 *Systematic Reviews in the Social Sciences: A Practical Guide* (Oxford: Blackwell Publishing)
- Piemontese L *et al* 2022 Validity and validation in archetype analysis: practical assessment framework and guidelines *Environ. Res. Lett.* **17** 025010
- Rodríguez L O and Young K R 2000 Biological diversity of peru: determining priority areas for conservation *J. Hum. Environ.* **29** 329–37
- Scheffer M and Carpenter S R 2003 Catastrophic regime shifts in ecosystems: linking theory to observation *Trends Ecol. Evol.* **18** 648–56
- Schellnhuber H J, Ludeke M K B and Petschel-Held G 2002 The syndromes approach to scaling describing global change on an intermediate functional scale *Integr. Assess.* **3** 201–19
- Schmidt-Traub G, Obersteiner M and Mosnier A 2019 Fix the broken food system in three steps *Nature* **569** 181–3
- Senge P M 2006 *The Fifth Discipline: The Art and Practice of the Learning Organization* (New York: Crown Business)
- Sietz D 2014 Regionalisation of global insights into dryland vulnerability: better reflecting smallholders' vulnerability in Northeast Brazil *Glob. Environ. Change* **25** 173–85



- Sietz D and Feola G 2016 Resilience in the rural Andes: critical dynamics, constraints and emerging opportunities *Reg. Environ. Change* **16** 2163–9
- Sietz D, Fleskens L and Stringer L C 2017a Learning from non-linear ecosystem dynamics is vital for achieving land degradation neutrality *Land Degrad. Dev.* **28** 2308–14
- Sietz D, Klimek S and Dauber J 2022 Tailored pathways toward revived farmland biodiversity can inspire agroecological action and policy to transform agriculture *Commun. Earth Environ.* **3** 211
- Sietz D, Lüdeke M K B and Walther C 2011 Categorisation of typical vulnerability patterns in global drylands *Glob. Environ. Change* **21** 431–40
- Sietz D, Untied B, Walkenhorst O, Lüdeke M K B, Mertins G, Petschel-Held G and Schellnhuber H J 2006 Smallholder agriculture in Northeast Brazil: assessing heterogeneous human-environmental dynamics *Reg. Environ. Change* **6** 132–46
- Suding K N 2011 Toward an era of restoration in ecology: successes, failures, and opportunities ahead *Annu. Rev. Ecol. Evol. Syst.* **42** 465–87
- UN 2014 Report of the open working group of the general assembly on sustainable development goals, A/68/970 (United Nations)
- UN 2021 The sustainable development goals report 2021 (New York: United Nations Statistics Division Development Data and Outreach Branch)
- UNCBD 2021 First draft of the post-2020 global biodiversity framework *CBD/WG2020/3/3* (available at: [www.cbd.int/doc/c/abb5/591f/2e46096d3f0330b08ce87a45/wg2020-03-03-en.pdf](http://www.cbd.int/doc/c/abb5/591f/2e46096d3f0330b08ce87a45/wg2020-03-03-en.pdf))
- UNEP 2007 *Global environmental outlook 4: environment for development* (Valletta)
- Urbinnati A M, Benites-Lazaro L L, Carvalho C M D and Giatti L L 2020 The conceptual basis of water-energy-food nexus governance: systematic literature review using network and discourse analysis *J. Integr. Environ. Sci.* **17** 21–43
- van Den Elsen E *et al* 2020 Advances in understanding and managing catastrophic ecosystem shifts in Mediterranean ecosystems *Front. Ecol. Evol.* **8** 561101
- Walker B H, Holling C S, Carpenter S C and Kinzig A P 2004 Resilience, adaptability and transformability *Ecol. Soc.* **9** 5
- Walker B, Gunderson L, Kinzig A, Folke C, Carpenter S and Schultz L 2006 A handful of heuristics and some propositions for understanding resilience in social-ecological systems *Ecol. Soc.* **11** 13
- Walker B and Meyers J A 2004 Thresholds in ecological and social-ecological systems: a developing database *Ecol. Soc.* **9** 3
- Wallman D 2018 Histories and trajectories of socio-ecological landscapes in the lesser Antilles: implications of colonial period zooarchaeological research *Environ. Archaeol.* **23** 13–22
- B) Archetype studies reviewed in this paper**
- Adenle A A and Speranza C I 2021 Social-ecological archetypes of land degradation in the Nigerian Guinea Savannah: insights for sustainable land management *Remote Sens.* **13** 32
- Almekinders C J M, Stone G D, Baranski M, Carney J A, Hanspach J, Krishna V V, Ramirez-Villegas J, Van Etten J and Zimmerer K S 2019 Socioecological Interactions amid Global Change *Agrobiodiversity: Integrating Knowledge for a Sustainable Future (Strüngmann Forum Reports vol 24)* ed K S Zimmerer, S De Haan and J R Lupp (Cambridge, MA: MIT Press) pp 117–43
- Armatas C A, Campbell R M, Watson A E, Borrie W T, Christensen N and Venn T J 2018 An integrated approach to valuation and tradeoff analysis of ecosystem services for national forest decision-making *Ecosyst. Serv.* **33** 1–18
- Bahri M 2020 Analysis of the water, energy, food and land nexus using the system archetypes: a case study in the Jatiluhur reservoir, West Java, Indonesia *Sci. Total Environ.* **716** 137025
- Banson K E, Nguyen N C and Bosch O J H 2016 Using system archetypes to identify drivers and barriers for sustainable agriculture in Africa: a case study in Ghana *Syst. Res. Behav. Sci.* **33** 79–99
- Bay-Larsen I, Risvoll C, Vestrum I and Bjorkhaug H 2018 Local protein sources in animal feed—perceptions among arctic sheep farmers *J. Rural Stud.* **59** 98–110
- Brzezina N, Biely K, Helfgott A, Kopainsky B, Vervoort J and Mathijs E 2017 Development of organic farming in Europe at the crossroads: looking for the way forward through system archetypes lenses *Sustainability* **9** 821
- Dasgupta R, Hashimoto S and Gundimeda H 2019 Biodiversity/ecosystem services scenario exercises from the Asia-Pacific: typology, archetypes and implications for sustainable development goals (SDGs) *Sustain. Sci.* **14** 241–57
- Egerer S, Cotera R V, Celliers L and Costa M M 2021 A leverage points analysis of a qualitative system dynamics model for climate change adaptation in agriculture *Agric. Syst.* **189** 103052
- Evers C R, Ager A A, Nielsen-Pincus M, Palaiologou P and Bunzel K 2019 Archetypes of community wildfire exposure from national forests of the western US *Landsc. Urban Plan.* **182** 55–66
- Felgenhauer T 2015 Addressing the limits to adaptation across four damage-response systems *Environ. Sci. Policy* **50** 214–24
- Fischer J, Abson D J, Bergsten A, Collier N F, Dorresteijn I, Hanspach J, Hylander K, Schultner J and Senbeta F 2017 Reframing the food-biodiversity challenge *Trends Ecol. Evol.* **32** 335–45
- Gerst M D, Cox M E, Locke K A, Laser M and Kapuscinski A R 2015 A taxonomic framework for assessing governance challenges and environmental effects of integrated food-energy systems *Environ. Sci. Technol.* **49** 734–41
- Gohari A, Eslamian S, Mirchi A, Abedi-Koupaei J, Bavani A M and Madani K 2013 Water transfer as a solution to water shortage: a fix that can backfire *J. Hydrol.* **491** 23–39
- Haberman D and Bennett E M 2019 Ecosystem service bundles in global hinterlands *Environ. Res. Lett.* **14** 084005
- Harrison P A *et al* 2019 Synthesizing plausible futures for biodiversity and ecosystem services in Europe and Central Asia using scenario archetypes *Ecol. Soc.* **24** 27
- Hartel T, Fagerholm N, Torralba M, Balazsi A and Plieninger T 2018 Forum: social-ecological system archetypes for European Rangelands *Rangeland Ecol. Manage.* **71** 536–44
- Higuera P E *et al* 2019 Integrating subjective and objective dimensions of resilience in fire-prone landscapes *Bioscience* **69** 379–88
- Jäger J, Rounsevell M, Harrison P, Omann I, Dunford R, Kammerlander M and Pataki G 2015 Assessing policy robustness of climate change adaptation measures across sectors and scenarios *Clim. Change* **128** 395–407
- Karrasch L, Klenke T and Kleyer M 2019 Land-use elements and attributed ecosystem services: an archetype approach to land-use evaluation at the German North Sea coast *Ecol. Soc.* **24** 13
- Keys P W and Wang-Erlandsson L 2018 On the social dynamics of moisture recycling *Earth Syst. Dyn.* **9** 829–47
- Lee H and Lautenbach S 2016 A quantitative review of relationships between ecosystem services *Ecol. Indic.* **66** 340–51
- Leventon J, Schaal T, Velten S, Loos J, Fischer J and Newig J 2019 Landscape-scale biodiversity governance: scenarios for reshaping spaces of governance *Environ. Policy Governance* **29** 170–84
- Livers C *et al* 2018 Archetypical patterns and trajectories of land systems in Europe *Reg. Environ. Change* **18** 715–32
- Li Q R, Babu T S A, Sieber S and Zander P 2018 Assessing divergent consequences of payments for ecosystem services on rural livelihoods: a case-study in China's Loess Hills *Land Degrad. Dev.* **29** 3549–70

- Lim-Camacho L, Ariyawardana A, Lewis G K, Crimp S J, Somogyi S, Ridoutt B and Howden S M 2017 Climate adaptation of food value chains: the implications of varying consumer acceptance *Reg. Environ. Change* **17** 93–103
- Locatelli B, Lavarello S, Sloan S, Tappeiner U and Geneletti D 2017 Characteristic trajectories of ecosystem services in mountains *Front. Ecol. Environ.* **15** 150–9
- Loiseau E, Colin M, Alaphilippe A, Coste G and Roux P 2020 To what extent are short food supply chains (SFSCs) environmentally friendly? Application to French apple distribution using life cycle assessment *J. Clean. Prod.* **276** 124166
- Magliocca N R, Khuc Q V, Ellicott E A and De Bremond A 2019 Archetypal pathways of direct and indirect land-use change caused by Cambodia's economic land concessions *Ecol. Soc.* **24** 25
- Moraine M, Duru M and Therond O 2017 A social-ecological framework for analyzing and designing integrated crop-livestock systems from farm to territory levels *Renew. Agric. Food Syst.* **32** 43–56
- Nyam Y S, Kotir J H, Jordaan A and Ogundeyi A A 2020 Identifying behavioural patterns of coupled water-agriculture systems using system archetypes *Syst. Res. Behav. Sci.* **39** 305–23
- Oberlack C *et al* 2019 Archetype analysis in sustainability research: meanings, motivations, and evidence-based policy making *Ecol. Soc.* **24** 26
- Oberlack C, Tejada L, Messerli P, Rist S and Giger M 2016 Sustainable livelihoods in the global land rush? Archetypes of livelihood vulnerability and sustainability potentials *Glob. Environ. Change-Hum. Policy Dimens.* **41** 153–71
- Pedde S, Kok K, Holscher K, Oberlack C, Harrison P A and Leemans R 2019 Archotyping shared socioeconomic pathways across scales: an application to central Asia and European case studies *Ecol. Soc.* **24** 30
- Piemontese L, Castelli G, Fetzner I, Barron J, Liniger H, Harari N, Bresci E and Jaramillo F 2020 Estimating the global potential of water harvesting from successful case studies *Glob. Environ. Change-Hum. Policy Dimens.* **63** 102121
- Queenan K, Sobratee N, Davids R, Mabhaudhi T, Chimonyo M, Slotow R, Shankar B and Hasler B 2020 A systems analysis and conceptual system dynamics model of the livestock-derived food system in South Africa: a tool for policy guidance *J. Agric. Food Syst. Commun. Dev.* **9** 275–98
- Rocha J, Malmberg K, Gordon L, Brauman K and Declerck F 2020 Mapping social-ecological systems archetypes *Environ. Res. Lett.* **15** 034017
- Sadoulet E and Dejanvry A 1992 Agricultural trade liberalization and low-income countries—a general equilibrium-multimarket approach *Am. J. Agric. Econ.* **74** 268–80
- Sietz D, Frey U, Roggero M, Gong Y Q, Magliocca N, Tan R, Janssen P and Vaclavik T 2019 Archetype analysis in sustainability research: methodological portfolio and analytical frontiers *Ecol. Soc.* **24** 34
- Sietz D, Ordóñez J C, Kok M T J, Janssen P, Hilderink H B M, Tittmonell P and van Dijk H 2017b Nested archetypes of vulnerability in African drylands: where lies potential for sustainable agricultural intensification? *Environ. Res. Lett.* **12** 095006
- Tessier L, Bijttebier J, Marchand F and Baret P V 2021 Identifying the farming models underlying Flemish beef farmers' practices from an agroecological perspective with archetypal analysis *Agric. Syst.* **187** 103013
- Theodosiou T, Valsamidis S, Hatziliadis G and Nikolaidis M 2012 Measuring, archotyping and mining production data *J. Syst. Inform. Technol.* **14** 318–35
- Thompson P B 2018 Four sociotechnical imaginaries for future food systems *Professionals in Food Chains* ed S Springer and H Grimm (Wageningen Academic Publishers) pp 187–91
- Turner B L, Menendez H M, Gates R, Tedeschi L O and Atzori A S 2016 System dynamics modeling for agricultural and natural resource management issues: review of some past cases and forecasting future roles *Resources* **5** 40
- Turner B L, Wuellner M, Nichols T, GATES R, Tedeschi L O and Dunn B H 2017 A systems approach to forecast agricultural land transformation and soil environmental risk from economic, policy, and cultural scenarios in the north central United States (2012–2062) *Int. J. Agric. Sustain.* **15** 102–23
- Vaclavik T, Langerwisch F, Cotter M, Fick J, Hauser I, Hotes S, Kamp J, Settele J, Spangenberg J H and Seppelt R 2016 Investigating potential transferability of place-based research in land system science *Environ. Res. Lett.* **11** 095002
- Vaclavik T, Lautenbach S, Kuemmerle T and Seppelt R 2013 Mapping global land system archetypes *Glob. Environ. Change-Hum. Policy Dimens.* **23** 1637–47
- Valette M, Vinceti B, Gregorio N, Bailey A, Thomas E and Jalonen R 2020 Beyond fixes that fail: identifying sustainable improvements to tree seed supply and farmer participation in forest and landscape restoration *Ecol. Soc.* **25** 30
- Vidal Merino M, Sietz D, Jost F and Berger U 2019 Archetypes of climate vulnerability: a mixed-method approach applied in the Peruvian Andes *Clim. Dev.* **11** 418–34
- Weissteiner C J, Boschetti M, Bottcher K, Carrara P, Bordogna G and Brivio P A 2011 Spatial explicit assessment of rural land abandonment in the Mediterranean area *Glob. Planet. Change* **79** 20–36
- Whitelaw P A, King B E M and Tolkach D 2014 Protected areas, conservation and tourism—financing the sustainable dream *J. Sustain. Tour.* **22** 584–603
- Zare F, Bagheri A and Elsayah S 2017 Using system archetypes for problem framing and a qualitative analysis: a case study in Iranian water resource management *22nd Int. Congress on Modelling and Simulation (Hobart, Tasmania, Australia)* pp 1433–9