PAPER • OPEN ACCESS

Artificial Neural Network Model to Predict Crude Protein and Crude Fiber from Physical Properties of Feedstuffs

To cite this article: MM Sholikin et al 2019 IOP Conf. Ser.: Earth Environ. Sci. 372 012049

View the article online for updates and enhancements.

You may also like

 Low optical dosage heating-reduced viscosity for fast and large-scale cleanup of spilled crude oil by reduced graphene oxide melamine nanocomposite adsorbents

Guangqiao Xu, Li Zhang, Wei Yu et al.

- <u>The effect of barnacles powder as a</u> <u>source of protein feed substitution against</u> the digestibility of crude protein and crude fiber on male rabbits rex B Agustono, F Maulana, S H Warsito et al.
- The Effect of Different Levels of Dietary Crud Protein During Different Age Stages on the Productive Performance of ROSS 308 A A Ismael and Z J M Saed





DISCOVER how sustainability intersects with electrochemistry & solid state science research



This content was downloaded from IP address 3.145.156.250 on 06/05/2024 at 08:46

Artificial Neural Network Model to Predict Crude Protein and Crude Fiber from Physical Properties of Feedstuffs

MM Sholikin¹, MD Alifian¹, FM Purba¹, A Jayanegara² and Nahrowi²

¹Graduate School of Nutrition and Feed Science, Faculty of Animal Science, Bogor Agricultural University, Bogor, Indonesia

²Department of Nutrition and Feed Technology, Faculty of Animal Science, Bogor Agricultural University, Bogor, Indonesia

E-mail: sholikinshino@gmail.com

Abstract. The aim of this research was to build artificial neural networks model to predict crude protein and crude fiber content from physical properties of feedstuffs. The 91 data were obtained from https://repository.ipb.ac.id using keywords, e.g., sifat fisik and pakan. To reduce the dimensional of the data had been transformed. The independent variables consist of specific gravity (SG), bulk density (BD), compacted bulk density (CBD) and angle of repose (AoR). The dependent variable was crude protein (CP) and crude fiber (CF). Artificial neural networks (ANN) model built by R programing language 3.6.0 using library R-base and neuralnet. The correlation and accuracy used to compare predicted and actual. ANN model of crude fiber has an accuracy of 75.08% and Pearson's signification correlation (0.7529; P <0.01). ANN model of crude fiber has an accuracy of 75.08% and Pearson's signification correlation (0.7529; P <0.01). The artificial neural networks model generally can perform better to predict crude protein and crude fiber from physical properties of feedstuffs.

Keywords: artificial neural networks model, crude fiber, crude protein, physical properties, feedstuffs.

1. Introduction

Fluctuate the database of several feedstuffs due to extream environmental change, increasing of a large number of feedstuffs database (especially, in physical and chemical properties), and non-linear interaction on each parameter. They will become crowded and wasted if not be defined the knowledge in it. Using computation technique may proper for this due to these problems. Computation techniques, such response surface modelling and artificial neural networks, are used increasingly in feed science because they are the robust, accurate, and flexible technique for prediction analysis.[1] These techniques have been used for optimizing of antibacterial activity extraction process from Hermetia illucens.[2] Artificial neural networks have been used for the prediction of nutritional requirements of digestible lysine, methionine, and threonine in broiler chicks.[3] ANN has been developed as a model that mimics the nervous system. ANN is mainly used in solving difficult problems.[4] ANN properties for learning from database patterns make it a powerful program based

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1 on incomplete domain compilation or compilation of partial inaccuracies or conflicting data.[4] The objective of this research was to build artificial neural networks model to predict crude protein and crude fiber content from physical properties of feedstuffs and validate it using correlation and accuracy value.

2. Materials and Methods

Collecting and Transforming Data

A total of 91 data were obtained from https://repository.ipb.ac.id database using keywords, e.g., *sifat fisik* and *pakan*. The data included specific gravity (SG), bulk density (BD), compacted bulk density (CBD), angle of repose (AoR), crude protein (CP), and crude fiber (CF). CP and CF converted in 100% dry matter (DM). The summary database is presented in **Table 1**.

Table 1. Summary database of physical properties, crude protein, and crude fiber of feedstuffs (n = 91).

Parameter	Feedstuffs			
Specific gravity (kg/m ³)	1.2427	±	0.2184	
Bulk density (kg/m ³)	0.3950	±	0.1425	
Compacted bulk density (kg/m ³)	0.5350	±	0.1336	
Angle of repose (°)	36.7012	<u>+</u>	6.4289	
Crude protein (%DM)	18.1645	<u>+</u>	10.0826	
Crude fiber (%DM)	15.2704	<u>+</u>	9.1858	
DM, dry matter.				

Then, data was split into data train (75%, n = 69) and data test (25%, n = 22). After that, the data train and data test were normalize using minimum and maximum transformation. The data transformation formula is as follows.

$$\mathbf{x}' = (x - \min(x)) \div (\max(x) + \min(x)),$$

where x = initial value; x' = transformed value; max(x) = maximum value; min(x) = minimum value.

Build, Fitting, and Validate Model

The artificial neural networks model was built on R-base 3.6.0 programming language and used "library (neuralnet)".[5, 6] Four input parameter consists of SG, BD, CBD, and AoR perform the ANN model. Four hidden layers (hly), which are hly1 (n = 40), hly2 (n = 15), hly3 (n = 9), and hly4 (n = 3) to build ANN model of predicted CP content of feedstuffs. While ANN model of predicted CF content of feedstuffs used three hidden layers, there are hly 1 (n = 3), hly2 (n = 15), and hly3 (n = 6). The four and three hidden layers of ANN model follows this formula,

$$CP(x) = w_0 + \sum_{l=0}^{L} w_l \cdot f\left(w_{0l} + \sum_{k=0}^{K} w_{kl} \cdot f\left(w_{0k} + \sum_{j=0}^{J} w_{jkl} \cdot f\left(w_{0j} + \sum_{i=0}^{I} w_{ijkl} \cdot x_i\right)\right)\right),$$

$$CF(x) = w_0 + \sum_{k=0}^{K} w_k \cdot f\left(w_{0k} + \sum_{j=0}^{J} w_{jk} \cdot f\left(w_{0j} + \sum_{i=0}^{I} w_{ijk} \cdot x_i\right)\right),$$

where w_0 denotes the intercept of the output neuron and w_{0l} the intercept of the lth hidden neuron. Additionally, w_l denotes the synaptic weight corresponding to the synapse starting at the lth hidden neuron and leading to the output neuron, $w_l = (w_{1l}, ..., w_{nl})$ the vector of all synaptic weights corresponding to the synapses leading to the lth hidden neuron, and $x = (x_1, ..., x_n)$ the vector of all covariates [6].

The 1st Animal Science and Food Technology Conference (AnSTC) 2019	IOP Publishing
IOP Conf. Series: Earth and Environmental Science 372 (2019) 012049	doi:10.1088/1755-1315/372/1/012049

The fitting plot used to compare predicted and actual value. Validation of the ANN model has used Pearson correlation and accuracy value. Pearson correlation significance marked as ns, *, and ** for P>0.05, P<0.05, and P<0.01, respectively.

3. Results and Discussion

Physical properties of feedstuffs, e.g, specific gravity, bulk density, compacted bulk density, and angle of repose can affect chemical properties, e.g, dry matter, organic matter, crude protein, crude fibre, and ether extract. Increase of specific gravity may increase quality of feedstuffs, especially crude protein and starch. Beside that lower specific gravity cause decrease feedstuffs and increase crude fibre content.

Validation	Crude protein	Crude fiber	
Accuracy	0.8315	0.7508	
Correlation	0.7112**	0.7529**	
$n_{0} D 0.05 * D 0.05 * D 0.01$			

Table 2. Validation of artificial neural networks model of feedstuffs (n = 22).

ns, P>0.05; *, P<0.05; **, P<0.01.

ANN model validation of crude protein and crude fiber using accuracy and Pearson's correlation values are present in **Table 3**. ANN model of crude protein has an accuracy of 83.15% in predicting the correct output. Pearson's significance correlation (0.7112; P < 0.01) of the crude protein ANN model, this states that there is a strong relationship between predictive value and actual value. ANN model of crude fiber has an accuracy of 75.08% in predicting the correct output. Pearson's signification correlation (0.7529; P < 0.01), this states that there is a strong relationship between predictive value and actual value.

ANN model of crude protein with four hidden layers (40, 15, 9, 3) has a positive correlation (Figure 1). Pearson correlation value, between predictions and actual 0.7112, this means there is a strong relationship between the two variables. This model can be used to estimate crude protein content of feedstuffs.



Predicted CP (%DM)



IOP Conf. Series: Earth and Environmental Science **372** (2019) 012049 doi:10.1088/1755-1315/372/1/012049



Predicted CF (%DM)

Figure 2. Plot between predicted (∇) and actual (\blacklozenge) crude fiber (CF) content of feedstuffs.

ANN model of crude fiber with three hidden layers (3, 15, 6) has a positive correlation (**Figure 2**). Pearson correlation value, between predictions and actual 0.7529, this means there is a strong relationship between the two variables. This model can be used to estimate crude fiber content of feedstuffs.

4. Conclusions

The artificial neural networks model generally can perform better to predict crude protein and crude fiber from physical properties of feedstuffs.

References

- [1] S. Shahinfar, H Mehrabani-Yeganeh, C Lucas, A Kalhor, M Kazemian, and KA Weigel. Prediction of breeding values for dairy cattle using artificial neural networks and neuro-fuzzy systems. *Computational and Mathematical Methods in Medicine*. ID:127130.
- [2] M. M. Sholikin, M. D Alifian, A Jayanegara, and Nahrowi. 2019. Optimization of the *Hermetia illucens* Larvae Extraction Process with Response Surface Modelling and Its Amino Acid Profile and Antibacterial Activity. *IOP Conf. Ser.: Mater. Sci. Eng.* 546 062030.
- [3] M. Mehri. 2012. Development of artificial neural network models based on experimental data of response surface methodology to establish the nutritional requirements of digestible lysine, methionine, and threonine in broiler chicks. *Poultry Sci.* 91:3280-3285.
- [4] Y. Huang. 2009. Advances in artificial neural networks-methodological development and application. *Algorithms* 2:973-1007.
- [5] R-Team. 2019. R: A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing*.
- [6] F. Günther and S. Fritsch. 2010. Neuralnet: Training of neural networks. *The R Journal* 2:30-38.