



PAPER • OPEN ACCESS

# Transverse momentum spectra of hadrons in high energy pp and heavy ion collisions

To cite this article: Kapil Saraswat *et al* 2018 *J. Phys. Commun.* **2** 035003

View the [article online](#) for updates and enhancements.

## You may also like

- [Transverse momentum distributions in proton–proton collisions at LHC energies and Tsallis thermodynamics](#)  
M D Azmi and J Cleymans
- [Entropy per Rapidity in Pb-Pb Central Collisions using Thermal and Artificial Neural Network \(ANN\) Models at LHC Energies](#)  
D. M. Habashy, Mahmoud Y. El-Bakry, Werner Scheinast et al.
- [SECTORS AND LARGE-SCALE MAGNETIC FIELD STRENGTH FLUCTUATIONS IN THE HELIOSHEATH NEAR 110 AU: VOYAGER 1, 2009](#)  
L. F. Burlaga and N. F. Ness



## PAPER

## OPEN ACCESS

## RECEIVED

1 December 2017

## REVISED

31 January 2018

## ACCEPTED FOR PUBLICATION

16 February 2018

## PUBLISHED

1 March 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



# Transverse momentum spectra of hadrons in high energy pp and heavy ion collisions

Kapil Saraswat<sup>1,4</sup> , Prashant Shukla<sup>2,3</sup> and Venktesh Singh<sup>1</sup>

<sup>1</sup> Department of Physics, Institute of Science, Banaras Hindu University, Varanasi 221005, India

<sup>2</sup> Nuclear Physics Division, Bhabha Atomic Research Center, Mumbai 400085, India

<sup>3</sup> Homi Bhabha National Institute, Anushakti Nagar, Mumbai 400094, India

<sup>4</sup> The author to whom any correspondence should be addressed.

E-mail: [kapilsaraswatbhu@gmail.com](mailto:kapilsaraswatbhu@gmail.com), [pshuklabarc@gmail.com](mailto:pshuklabarc@gmail.com) and [venkaz@yahoo.com](mailto:venkaz@yahoo.com)

**Keywords:** QGP, hadron spectra, tsallis distribution, energy loss

## Abstract

We present a study of transverse momentum ( $p_T$ ) spectra of unidentified charged particles in pp collisions at RHIC and LHC energies from  $\sqrt{s} = 62.4$  GeV to 13 TeV using Tsallis/Hagedorn function. The power law of Tsallis/Hagedorn form gives very good description of the hadron spectra in  $p_T$  range from 0.2 to 300 GeV/c. The power index  $n$  of the  $p_T$  distributions is found to follow a function of the type  $a + b/\sqrt{s}$  with asymptotic value  $a = 6.8$ . The parameter  $T$  governing the soft bulk contribution to the spectra remains almost same over wide range of collision energies. We also provide a Tsallis/Hagedorn fit to the  $p_T$  spectra of hadrons in pPb and different centralities of PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. The data/fit shows deviations from the Tsallis distribution which become more pronounced as the system size increases. We suggest simple modifications in the Tsallis/Hagedorn power law function and show that the above deviations can be attributed to the transverse flow in low  $p_T$  region and to the in-medium energy loss in high  $p_T$  region.

## 1. Introduction

The light hadrons are the most abundant particles produced in the pp and heavy ion collisions. The transverse momentum ( $p_T$ ) spectra of hadrons can be used to infer the particle production mechanism in pp collisions. In heavy ion collisions, additional final state effects such as collective flow [1, 2], recombination [2, 3] and jet-quenching [4] in different  $p_T$  ranges are superimposed over the hadron spectra. The hadron  $p_T$  spectra in pp collisions are successfully described by Tsallis distribution [5, 6] in terms of only two parameters, the Tsallis parameter  $T$  and the parameter  $q$  which governs the degree of non-thermalization. It is well known [7–9] that the functional form of the Tsallis distribution, which describes near-thermal systems is essentially the same as the power law function by Hagedorn which is applicable to the QCD hard scatterings [10, 11]. There are numerous studies which show that the Tsallis/Hagedorn distribution gives an excellent description of  $p_T$  spectra of all identified hadrons measured in pp collisions at SPS, RHIC and LHC energies [7, 12–14]. The work in [15] makes a comparative study of various forms of Tsallis distributions available in the literature by fitting the hadron  $p_T$  spectra measured at RHIC and LHC. The recent papers [16–18] present a study of the Tsallis parameters for  $p_T$  distributions of pions produced in pp collisions as a function of  $\sqrt{s}$  ranging between 6.3 GeV and 7 TeV. The CMS papers [19, 20] present studies on measured identified particle spectra using Tsallis distribution at  $\sqrt{s} = 0.9, 2.76, 7$  and 13 TeV. There are many attempts to use the Tsallis distribution in heavy ion collisions after taking into account the transverse collective flow, which is a final state effect [21–23]. The work in [24] studies the  $p_T$  spectra of the strange hadrons production in pp collision at  $\sqrt{s} = 7$  TeV, pPb collision at  $\sqrt{s_{NN}} = 5.02$  TeV and PbPb collision at  $\sqrt{s_{NN}} = 2.76$  TeV using the Tsallis distribution, which includes the transverse flow.

The Tsallis distribution is applied to unidentified light charged hadron spectra measured in pp collisions at LHC over a wide  $p_T$  range upto 200 GeV/c [25]. The work in [26] studies the  $p_T$  spectra of both jets and hadrons

in pp collisions. They find that the power index  $n$  is 4–5 for jet production and 6–10 for hadron production. The work in [27, 28] uses Tsallis power law to fit the hadron spectra in wide  $p_T$  range measured in pp collisions. Looking more closely at the data/fit they suggest that deviations of the data from power law fit follows a log-periodic oscillation, which could imply a complex exponent of the power law. It was further suggested [29, 30] that the oscillations in data/fits in PbPb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV are similar to those in pp data in the same range of transverse momenta.

In this work, we study transverse momentum ( $p_T$ ) spectra of unidentified charged particles in pp collisions at  $\sqrt{s} = 0.0624, 0.2, 0.9, 2.36, 2.76, 5.02, 7$  and 13 TeV using Tsallis/Hagedorn function. The statistical and systematic errors are added in quadrature and are used in the fits. The parameters of such fits are studied as a function of beam energies. We also study the spectra of identified charged particles in pp collisions albeit at smaller  $p_T$ . Since the aim of the work is to obtain a function describing the hadron spectra in wide  $p_T$  range, we choose the unidentified particles for which the measurements are available at very high  $p_T$  upto 200 GeV/c. We also make a Tsallis/Hagedorn fit to the  $p_T$  spectra of hadrons in pPb and different centralities of PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. We suggest simple modifications in the Tsallis/Hagedorn power law function to include transverse flow and in-medium energy loss in the hadronic spectra.

## 2. Tsallis/Hagedorn distribution function and the modification

The transverse mass ( $m_T = \sqrt{p_T^2 + m^2}$ ) distribution of particles produced in hadronic collisions can be described by the Hagedorn function which is a QCD-inspired summed power law [10] given as

$$E \frac{d^3N}{dp^3} = A \left( 1 + \frac{m_T}{p_0} \right)^{-n}. \quad (1)$$

This function describes both the bulk spectra in the low  $m_T$  region and the particles produced in QCD hard scatterings reflected in the high  $p_T$  region. Let us compare this function with the Tsallis distribution [5, 6] of thermodynamic origin given by

$$E \frac{d^3N}{dp^3} = C_n m_T \left( 1 + (q - 1) \frac{m_T}{T} \right)^{-1/(q-1)}. \quad (2)$$

The Tsallis distribution describes near-thermal systems in terms of Tsallis parameter  $T$  and the parameter  $q$ , which measures degree of non-thermalization [31]. The functions in equation (1) and in equation (2) have similar mathematical forms with  $n = 1/(q - 1)$  and  $p_0 = nT$ . Larger values of  $n$  correspond to smaller values of  $q$ . Both  $n$  and  $q$  have been interchangeably used in Tsallis distribution [6, 12, 32–34]. Phenomenological studies suggest that, for quark-quark point scattering,  $n \sim 4$  [35, 36], which grows larger if multiple scattering centers are involved. The study in [15] suggests that both the forms given in equation (1) and in equation (2) give equally good fit to the hadron spectra in pp collisions. We use equation (2) in case of pp collisions.

Tsallis/Hagedorn function is able to describe  $p_T$  spectra in pp collisions practically at all generations of proton colliders. There have been many attempts to use the Tsallis distribution in heavy ion collisions as well by including the transverse collective flow [21–23]. In addition, in heavy ion collisions, particle spectra at high  $p_T$  are known to be modified due to in-medium energy loss. The Tsallis/Hagedorn distribution can be modified by including these final state effects in different  $p_T$  regions as follows:

$$E \frac{d^3N}{dp^3} = A_1 \left[ \exp \left( -\frac{\beta p_T}{p_1} \right) + \frac{m_T}{p_1} \right]^{-n_1} : p_T < p_{T_{th}}. \quad (3a)$$


$$E \frac{d^3N}{dp^3} = A_2 \left[ \frac{B}{p_2} \left( \frac{p_T}{q_0} \right)^\alpha + \frac{m_T}{p_2} \right]^{-n_2} : p_T > p_{T_{th}}. \quad (3b)$$

The first function (equation (3a)) is shown to govern the thermal and collective part of the hadron spectrum with the temperature  $T = p_1/n_1$  and the average transverse flow velocity  $\beta$  [22].

The second function (equation (3b)) is obtained after shifting the distribution in equation (1) by energy loss  $\Delta m_T$  in the medium as

$$E \frac{d^3N}{dp^3} = A_2 \left[ 1 + \frac{m_T + \Delta m_T}{p_2} \right]^{-n_2}. \quad (4)$$

The energy loss  $\Delta m_T$  is proportional to  $p_T$  at low  $p_T$  and in general can be parameterized similar to the work in [37] as



$$\Delta m_T = B \left( \frac{p_T}{q_0} \right)^\alpha. \quad (5)$$

Here, the parameter  $\alpha$  quantifies different energy loss regimes for light quarks in the medium [38, 39]. The parameter  $B$  is proportional to the medium size and  $q_0$  is an arbitrary scale set as 1 GeV. Using equation (5) in equation (4) and ignoring 1 we get equation (3b) applicable for high  $p_T$ . In our study, we find that this function describes the particle spectra at  $p_{Tth}$  above 7 GeV/c. Fits to the data would constrain the value of  $B/p_2$  and thus  $p_2$  is not an independent parameter. The empirical parton energy loss in nuclear collisions at RHIC energies is found to be proportional to  $p_T$  [40].

### 3. Results and discussions

Figure 1 shows the invariant yields of the charged particles as a function of  $p_T$  for pp collisions at  $\sqrt{s} = 62.4$  and 200 GeV measured by the PHENIX experiment [41, 42]. The solid curves are the Tsallis distributions fitted to the spectra. The Tsallis distribution function gives good description of the data for both the collision energies which can be inferred from the values of  $\chi^2/\text{NDF}$  given in the table 1.


Figure 2 shows the invariant yields of the charged particles as a function of  $p_T$  for pp collisions at  $\sqrt{s} = 0.9, 2.76, 7$  and 13 TeV measured by the ALICE experiment [43, 44]. The solid curves are the Tsallis distributions. The Tsallis distribution function gives good description of the data for collision energies which can be inferred from the values of  $\chi^2/\text{NDF}$  given in the table 1.

Figure 3 shows the invariant yields of the charged particles as a function of  $p_T$  for pp collisions at  $\sqrt{s} = 0.9, 2.36, 2.76, 5.02$  and 7 TeV measured by the CMS experiment [45–48]. The solid curves are the Tsallis distributions fitted to the spectra. The Tsallis distribution function gives good description of the data for all collision energies which can be inferred from the values of  $\chi^2/\text{NDF}$  gives in the table 1.

Figure 4 shows the Tsallis parameter  $n$  for the charged particles as a function of the pp collision energy  $\sqrt{s}$ . The value of  $n$  decreases as we move from RHIC to LHC energies. The decreasing value of  $n$  shows that number of quarks participating for a produced particle are reduced for higher energy collisions. The parameter  $n$  can be parametrized by a function of the type

$$n(\sqrt{s}) = a + \frac{b}{\sqrt{s}}. \quad (6)$$

Here  $a = 6.81 \pm 0.06$  and  $b = 59.24 \pm 3.53$  GeV with  $\chi^2/\text{NDF} = 0.78$ . The QCD point scattering for pions production gives  $n = 4$ . Earlier studies have suggested that the value of  $n$  is larger for baryons as compared to



**Table 1.** The parameters of the Tsallis function obtained by fitting the charged particle spectra in pp collision at RHIC and LHC energies.

| $\sqrt{s}$ | Experiment | $n$              | $q$             | $T$<br>(MeV)      | $\frac{\chi^2}{\text{NDF}}$ |
|------------|------------|------------------|-----------------|-------------------|-----------------------------|
| 62.4 GeV   | PHENIX     | $14.28 \pm 0.69$ | $1.07 \pm 0.05$ | $102.87 \pm 5.98$ | 0.49                        |
| 200 GeV    | PHENIX     | $11.13 \pm 0.45$ | $1.09 \pm 0.04$ | $98.27 \pm 6.50$  | 0.49                        |
| 900 GeV    | ALICE      | $8.69 \pm 0.13$  | $1.12 \pm 0.02$ | $84.71 \pm 2.58$  | 0.16                        |
| 2.76 TeV   | ALICE      | $7.95 \pm 0.07$  | $1.13 \pm 0.01$ | $84.57 \pm 2.21$  | 0.17                        |
| 7 TeV      | ALICE      | $7.48 \pm 0.05$  | $1.13 \pm 0.01$ | $84.81 \pm 2.08$  | 0.26                        |
| 13 TeV     | ALICE      | $7.03 \pm 0.14$  | $1.14 \pm 0.02$ | $81.40 \pm 3.20$  | 0.38                        |
| 900 GeV    | CMS        | $8.72 \pm 0.22$  | $1.11 \pm 0.03$ | $83.24 \pm 4.19$  | 0.12                        |
| 2.36 TeV   | CMS        | $7.79 \pm 0.36$  | $1.13 \pm 0.05$ | $78.61 \pm 6.82$  | 0.21                        |
| 2.76 TeV   | CMS        | $7.98 \pm 0.21$  | $1.13 \pm 0.03$ | $89.66 \pm 13.64$ | 0.02                        |
| 5.02 TeV   | CMS        | $7.67 \pm 0.02$  | $1.13 \pm 0.00$ | $87.58 \pm 2.26$  | 0.79                        |
| 7 TeV      | CMS        | $7.55 \pm 0.16$  | $1.13 \pm 0.02$ | $86.46 \pm 12.60$ | 0.02                        |



that for mesons. Since the bulk of the particles produced in the pp collisions are predominantly pions, we can consider the unidentified particle spectra as that of pions.

Figure 5 shows the Tsallis temperature parameter  $T$  for the charged particles as a function of  $\sqrt{s}$  in pp collision. The parameter  $T$  can be parametrized by a function of the type

$$T(\sqrt{s}) = c + \frac{d}{\sqrt{s}}. \quad (7)$$

Here  $c = 0.082 \pm 0.002$  GeV and  $d = 0.151 \pm 0.048$  (GeV)<sup>2</sup> with  $\chi^2/\text{NDF} = 0.71$ . The parameter  $T$  slowly decreases with the collision energy.

We also present such analysis for each identified particle separately although in a much smaller range of  $p_T$  limited by measured data. The identified particle  $p_T$  spectra measured at RHIC and LHC energies [33, 49–55] are fitted with the Tsallis distribution and its parameters  $n$  and  $T$  are obtained. Figure 6 shows the Tsallis parameter  $n$  for the identified charged particles as a function of  $\sqrt{s}$ . The panel (a) is for charged pions, (b) is for neutral pions, (c) is for charged kaons and (d) is for protons. The solid curve is the fit given by function in equation (6). The asymptotic values of parameter  $n$  are  $6.41 \pm 0.10$  for charged pions,  $7.23 \pm 0.48$  for neutral pions,  $6.72 \pm 0.18$  for charged kaons and  $8.76 \pm 0.36$  for protons. The value of  $\chi^2/\text{NDF}$  is not good for charged pions and kaons because the value of  $n$  at  $\sqrt{s} = 900$  GeV is away from the fit line. This study shows that the asymptotic value of  $n$  for unidentified charged particles is closer to what is obtained for pions and kaons. Figure 7 shows the Tsallis



temperature parameter  $T$  for the identified particles as a function of  $\sqrt{s}$  in pp collision. The panel (a) is for charged pions, (b) is for neutral pions, (c) is for charged kaons and (d) is for protons. The behavior of the parameter  $T$  is similar to what is shown in figure 5.

Figure 8 shows the invariant yields of the charged particles as a function of  $p_T$  for pPb collisions and for many centralities of PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV measured by the CMS experiment [47]. The solid curves are the fitted Tsallis distributions (equation (2)). Figure 9 shows the ratio of the data and the fitted Tsallis distribution as a function of  $p_T$  for pp, pPb and PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. As we move from




Figure 5. The Tsallis temperature parameter  $T$  for the charged particle as a function of the centre of mass energy  $\sqrt{s}$  of pp collision.






Figure 6. The Tsallis parameter  $n$  as a function of the centre of mass energy  $\sqrt{s}$  of pp collision for (a) charged pions (b) neutral pions (c) charged kaons and (d) protons. The solid curve represents the function  $(a + b/\sqrt{s})$ .


peripheral to central PbPb collisions, the data show increasing deviations from the Tsallis fit. The  $\chi^2/\text{NDF}$  values of the Tsallis fit are given in the table 2. The pp and pPb data also show some deviations from the fit. Interestingly, the deviation pattern in pp and pPb looks similar to that in PbPb collisions with deviation magnitude increasing with system size. It was suggested in [28, 29] that the ratio of the data to the Tsallis fit shows a log oscillation function and which can be parametrized by a function of the form



$$f(p_T) = a + b \cos[c \log(p_T + d) + e]. \quad (8)$$

Here  $a, b, c, d$  and  $e$  are the fit parameters. We observe from figure 9 that the log oscillation function although loaded with five parameters does not describe the deviation pattern specially for more central PbPb collisions.






**Table 2.** The  $\chi^2/\text{NDF}$  of the Tsallis function obtained by fitting the charged particle spectra in pp, pPb and PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV.


| System                      | PbPb | PbPb | PbPb  | PbPb  | PbPb  | PbPb  | pPb  | pp   |
|-----------------------------|------|------|-------|-------|-------|-------|------|------|
| Centrality (%)              | 0–5  | 5–10 | 10–30 | 30–50 | 50–70 | 70–90 | —    | —    |
| $\frac{\chi^2}{\text{NDF}}$ | 5.93 | 6.11 | 3.89  | 3.11  | 1.84  | 1.20  | 1.22 | 0.79 |

Figure 10 shows the invariant yields of the charged particles as a function of  $p_T$  for pPb and PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV measured by the CMS experiment [47, 56]. The solid curves are the modified Tsallis distributions given by equation (3). Figure 11 shows the ratio of the data and the fit function by the modified Tsallis distribution as a function of  $p_T$  for pp, pPb and PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. The ratio of the data and the fit function shows that the modified Tsallis distribution function gives excellent description of the measured data in full  $p_T$  range for all the systems. The parameters of the modified Tsallis distribution are given in the table 3. The values of the first set of parameters ( $n_1, p_1, \beta$ ) increase with increasing system size for PbPb collisions. It shows that degree of thermalization (governed by  $n_1$ ) and the transverse flow (governed by  $\beta$ ) increase with system size. While fitting the second function, we fix the parameter  $n_2 = 7.7$  guided by pp value. The exponent  $\alpha$  which decides the variation of the energy loss of partons as a function of their energy remains within 0.4 to 0.7. The parameter  $B$  is proportional to system size and increases as we move from pp to the most central PbPb collisions. To summarize, the functions given in equations (3) give excellent description of the hadron spectra over wide range of  $p_T$  with its parameters indicating different physics effects in the collisions.

The modified function (equation (3)) is intended for PbPb systems to cover medium effects. We have applied it for small systems as well. The charged particle spectrum in pp collision when fitted with the Tsallis (equation (2)) gives  $\chi^2/\text{NDF} = 0.79$ . The  $\chi^2/\text{NDF}$  value improves to 0.31 when fitting is performed with the modified Tsallis (equation (3)). The pp system shows a small transverse flow and energy loss effect. It is not surprising since recent experiments have measured effects of collectivity in pp collisions when analysis on high multiplicity events was performed [57].



**Figure 10.** The invariant yields of the charged particles as a function of the transverse momentum  $p_T$  for pPb and PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV measured by the CMS [47, 56]. The solid curves are the modified Tsallis distributions (equation (3)).



**Figure 11.** The ratio of the charged particle yield data and the fit function (Modified Tsallis distribution equation (3)) as a function of the transverse momentum  $p_T$  for pp, pPb and PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV.

#### 4. Conclusion

We carried out an analysis of transverse momentum spectra of the unidentified charged particles in pp collisions at RHIC and LHC energies from  $\sqrt{s} = 62.4$  GeV to 13 TeV using Tsallis distribution function. The power law

**Table 3.** The parameters of the modified Tsallis function equation (3) obtained by fitting the charged particle spectra in pp, pPb and PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV.

| System             | $n_1$           | $p_1$<br>(GeV/c) | $\beta$         | $\alpha$        | $B$<br>(GeV/c)  | $\frac{\chi^2}{\text{NDF}}$ |
|--------------------|-----------------|------------------|-----------------|-----------------|-----------------|-----------------------------|
| PbPb<br>(0%–5%)    | $8.28 \pm 0.92$ | $1.38 \pm 0.17$  | $0.62 \pm 0.06$ | $0.47 \pm 0.03$ | $5.72 \pm 0.23$ | 0.29                        |
| PbPb<br>(5%–10%)   | $8.13 \pm 0.93$ | $1.35 \pm 0.18$  | $0.60 \pm 0.06$ | $0.42 \pm 0.04$ | $5.45 \pm 0.33$ | 0.68                        |
| PbPb<br>(10%–30%)  | $7.73 \pm 0.81$ | $1.28 \pm 0.14$  | $0.62 \pm 0.08$ | $0.52 \pm 0.03$ | $5.01 \pm 0.20$ | 0.14                        |
| PbPb:<br>(30%–50%) | $7.03 \pm 0.65$ | $1.11 \pm 0.11$  | $0.61 \pm 0.11$ | $0.45 \pm 0.04$ | $4.25 \pm 0.22$ | 0.36                        |
| PbPb<br>(50%–70%)  | $6.64 \pm 0.72$ | $0.96 \pm 0.12$  | $0.50 \pm 0.08$ | $0.58 \pm 0.03$ | $3.73 \pm 0.17$ | 0.06                        |
| PbPb<br>(70%–90%)  | $6.66 \pm 1.20$ | $0.90 \pm 0.26$  | $0.34 \pm 0.06$ | $0.59 \pm 0.04$ | $2.95 \pm 0.16$ | 0.63                        |
| pPb                | $7.78 \pm 1.70$ | $1.34 \pm 0.95$  | $0.14 \pm 0.05$ | $0.66 \pm 0.08$ | $2.95 \pm 0.73$ | 0.32                        |
| pp                 | $7.78 \pm 1.70$ | $1.09 \pm 0.41$  | $0.14 \pm 0.07$ | $0.60 \pm 0.03$ | $2.87 \pm 0.10$ | 0.31                        |

of Tsallis/Hagedorn form gives very good description of the hadron spectra in the  $p_T$  range 0.2 to 300 GeV/c. The power index  $n$  of the  $p_T$  distributions is found to follow a function of the type  $a + b/\sqrt{s}$  with asymptotic value  $a = 6.81$ . The Tsallis parameter  $T$  governing the soft bulk contribution to the spectra also decreases slowly with the collision energy. We also provide a Tsallis fit to the  $p_T$  spectra of hadrons in different centralities of PbPb collisions at  $\sqrt{s_{NN}} = 5.02$  TeV. The measured charged particle  $p_T$  spectra in PbPb collisions show deviations from Tsallis form which become more pronounced as the system size increases. We suggest simple modifications of the Tsallis function incorporating transverse flow in the low  $p_T$  region and in-medium energy loss in the high  $p_T$  region. This function gives excellent description of charged particle spectra in pp, pPb and PbPb collisions with its parameters having potentials to quantify various in-medium effects in all systems.

## ORCID iDs

Kapil Saraswat  <https://orcid.org/0000-0002-1211-5652>

Prashant Shukla  <https://orcid.org/0000-0001-8118-5331>

Venkatesh Singh  <https://orcid.org/0000-0003-4535-6230>

## References

- [1] Hirano T and Nara Y 2004 *Phys. Rev. C* **69** 034908
- [2] Fries R J, Greco V and Sorensen P 2008 *Ann. Rev. Nucl. Part. Sci.* **58** 177
- [3] Fries R J 2004 *J. Phys. G* **30** S853
- [4] Wang X N 2004 *Phys. Lett. B* **579** 299
- [5] Tsallis C 1988 *J. Statist. Phys.* **52** 479
- [6] Biro T S, Purcsel G and Urmossy K 2009 *Eur. Phys. J. A* **40** 325
- [7] Khandai P K, Sett P, Shukla P and Singh V 2013 *Int. J. Mod. Phys. A* **28** 1350066
- [8] Wong C Y and Wilk G 2012 *Acta Phys. Polon. B* **43** 2047
- [9] Wong C Y and Wilk G 2013 *Phys. Rev. D* **87** 114007
- [10] Hagedorn R 1983 *Riv. Nuovo Cim.* **6N10** 1
- [11] Blankenbecler R and Brodsky S J 1974 *Phys. Rev. D* **10** 2973
- [12] Adare A et al (PHENIX Collaboration) 2011 *Phys. Rev. D* **83** 052004
- [13] Sett P and Shukla P 2014 *Adv. High Energy Phys.* **2014** 896037
- [14] Zheng H, Zhu L and Bonasera A 2015 *Phys. Rev. D* **92** 074009
- [15] Zheng H and Zhu L 2016 *Adv. High Energy Phys.* **2016** 9632126
- [16] Parvan A S, Teryaev O V and Cleymans J 2017 *Eur. Phys. J. A* **53** 102
- [17] Parvan A S 2016 *Eur. Phys. J. A* **52** 355
- [18] Azmi M D and Cleymans J 2014 *J. Phys. G* **41** 065001
- [19] Chatrchyan S et al (CMS Collaboration) 2012 *Eur. Phys. J. C* **72** 2164
- [20] Sirunyan A M and (CMS Collaboration) 2017 *Phys. Rev. D* **96** 112003
- [21] Tang Z, Xu Y, Ruan L, van Buren G, Wang F and Xu Z 2009 *Phys. Rev. C* **79** 051901
- [22] Khandai P K, Sett P, Shukla P and Singh V 2014 *J. Phys. G* **41** 025105
- [23] Sett P and Shukla P 2015 *Int. J. Mod. Phys. E* **24** 1550046
- [24] Saraswat K, Shukla P, Kumar V and Singh V 2017 *Eur. Phys. J. A* **53** 84
- [25] Cleymans J and Azmi M D 2016 *J. Phys. Conf. Ser.* **668** 012050

- [26] Wong C Y, Wilk G, Cirto L J L and Tsallis C 2015 *Phys. Rev. D* **91** 114027
- [27] Wilk G and Wlodarczyk Z 2014 *Physica A* **413** 53
- [28] Wilk G and Wlodarczyk Z arXiv:1403.3508 [hep-ph]
- [29] Rybczynski M, Wilk G and Wlodarczyk Z 2015 *EPJ Web Conf.* **90** 01002
- [30] Wilk G and Wlodarczyk Z 2015 *Entropy* **17** 384
- [31] Wilk G and Wlodarczyk Z 2000 *Phys. Rev. Lett.* **84** 2770
- [32] Cleymans J and Worku D 2012 *Eur. Phys. J. A* **48** 160
- [33] Adare A *et al* (PHENIX Collaboration) 2011 *Phys. Rev. C* **83** 064903
- [34] Abelev B I *et al* (STAR Collaboration) 2007 *Phys. Rev. C* **75** 064901
- [35] Blankenbecler R, Brodsky S J and Gunion J F 1975 *Phys. Rev. D* **12** 3469
- [36] Brodsky S J, Pirner H J and Raufeisen J 2006 *Phys. Lett. B* **637** 58
- [37] Spousta M 2017 *Phys. Lett. B* **767** 10
- [38] Baier R, Schiff D and Zakharov B G 2000 *Ann. Rev. Nucl. Part. Sci.* **50** 37
- [39] De S and Srivastava D K 2012 *J. Phys. G* **39** 015001  
De S and Srivastava D K 2013 *J. Phys. G* **40** 049502 arXiv:1107.5659 [nucl-th]
- [40] Wang G and Huang H Z 2009 *Phys. Lett. B* **672** 30
- [41] Adare A *et al* (PHENIX Collaboration) 2012 *Phys. Rev. D* **86** 092006
- [42] Adler S S *et al* (PHENIX Collaboration) 2005 *Phys. Rev. Lett.* **95** 202001
- [43] Abelev B B *et al* (ALICE Collaboration) 2013 *Eur. Phys. J. C* **73** 2662
- [44] Adam J *et al* (ALICE Collaboration) 2016 *Phys. Lett. B* **753** 319
- [45] Chatrchyan S *et al* (CMS Collaboration) 2012 *Eur. Phys. J. C* **72** 1945
- [46] Khachatryan V *et al* (CMS Collaboration) 2010 *J. High Energy Phys.* JHEP02(2010)041
- [47] Khachatryan V *et al* (CMS Collaboration) 2017 *J. High Energy Phys.* JHEP04(2017)039
- [48] Chatrchyan S *et al* (CMS Collaboration) 2011 *J. High Energy Phys.* JHEP08(2011)086
- [49] Agakishiev G *et al* (STAR Collaboration) 2012 *Phys. Rev. Lett.* **108** 072302
- [50] Aamodt K *et al* (ALICE Collaboration) 2011 *Eur. Phys. J. C* **71** 1655
- [51] Abelev B B *et al* (ALICE Collaboration) 2014 *Phys. Lett. B* **736** 196
- [52] Adam J *et al* (ALICE Collaboration) 2015 *Eur. Phys. J. C* **75** 226
- [53] Adler S S *et al* (PHENIX Collaboration) 2003 *Phys. Rev. Lett.* **91** 241803
- [54] Abelev B B *et al* (ALICE Collaboration) 2012 *Phys. Lett. B* **717** 162
- [55] Abelev B B *et al* (ALICE Collaboration) 2014 *Eur. Phys. J. C* **74** 3108
- [56] Khachatryan V *et al* (CMS Collaboration) 2015 *Eur. Phys. J. C* **75** 237
- [57] Khachatryan V *et al* (CMS Collaboration) 2017 *Phys. Lett. B* **765** 193