



LEGWORK: A Python Package for Computing the Evolution and Detectability of Stellar-origin Gravitational-wave Sources with Space-based Detectors

T. Wagg^{1,2,3} , K. Breivik⁴ , and S. E. de Mink^{2,3,5} ¹ Department of Astronomy, University of Washington, Seattle, WA 98195, USA; tomjwagg@gmail.com² Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA³ Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Straße 1, D-85741 Garching, Germany⁴ Center for Computational Astrophysics, Flatiron Institute, 162 Fifth Avenue, New York, NY, 10010, USA⁵ Anton Pannekoek Institute for Astronomy and GRAPPA, University of Amsterdam, NL-1090 GE Amsterdam, The Netherlands

Received 2021 November 16; revised 2022 March 3; accepted 2022 March 7; published 2022 June 28

Abstract

We present LEGWORK (LISA Evolution and Gravitational Wave Orbit Kit), an open-source Python package for making predictions about stellar-origin gravitational-wave sources and their detectability in LISA or other space-based gravitational-wave detectors. LEGWORK can be used to evolve the orbits of sources due to gravitational-wave emission, calculate gravitational-wave strains (using post-Newtonian approximations), compute signal-to-noise ratios, and visualize the results. It can be applied to a variety of potential sources, including binaries consisting of white dwarfs, neutron stars, and black holes. Although we focus on double compact objects, in principle LEGWORK can be used for any system with a user-specified orbital evolution, such as those affected by a third object or gas drag. We optimized the package to make it efficient for use in population studies, which can contain tens of millions of sources. This paper describes the package and presents several potential use cases. We explain in detail the derivations of the expressions behind the package as well as identify and clarify some discrepancies currently present in the literature. We hope that LEGWORK will enable and accelerate future studies triggered by the rapidly growing interest in gravitational-wave sources.

Unified Astronomy Thesaurus concepts: [Open source software \(1866\)](#); [Gravitational waves \(678\)](#); [Gravitational wave detectors \(676\)](#); [Compact objects \(288\)](#); [Orbital evolution \(1178\)](#); [White dwarf stars \(1799\)](#); [Neutron stars \(1108\)](#); [Stellar mass black holes \(1611\)](#)

Software reviewed by the [Journal of Open Source Software JOSS](#)

1. Introduction

The planned space-based gravitational-wave detector LISA (Laser Interferometer Space Antenna) will present an entirely new view of gravitational waves by focusing on lower frequencies ($10^{-5} < f/\text{Hz} < 10^{-1}$) than ground-based detectors. This will enable the study of many new source classes including mergers of supermassive black holes (e.g., Begelman et al. 1980; Klein et al. 2016; Bellovary et al. 2019), extreme mass ratio inspirals (e.g., Berti et al. 2006; Barack & Cutler 2007; Babak et al. 2017; Moore et al. 2017), and cosmological GW backgrounds (e.g., Bartolo et al. 2016; Caprini et al. 2016; Caldwell et al. 2019). However, this frequency regime is also of interest for the detection of local stellar-mass binaries during their inspiral phase. LISA is expected to detect Galactic stellar-origin binaries containing combinations of white dwarfs, neutron stars, and black holes, ranging from the numerous double white dwarf population, to the rare but loud double black hole population.

The potential to detect stellar-origin sources with LISA has been studied in the past by various groups (e.g., Nelemans et al. 2001; Liu 2009; Liu & Zhang 2014; Ruiter et al. 2010; Belczynski et al. 2010; Yu & Jeffery 2010; Nisanke et al. 2012). The recent direct detection of gravitational waves with ground-based detectors has led to renewed interest in this topic (e.g., Korol et al. 2017, 2018, 2019, 2020; Christian & Loeb 2017; Kremer et al. 2017, 2018; Lamberts et al. 2018, 2019;

Fang et al. 2019; Andrews et al. 2020; Lau et al. 2020; Breivik et al. 2020a, 2020b; Roebber et al. 2020; Chen et al. 2020; Sesana et al. 2020; Shao & Li 2021).

Each of these studies requires making estimates of the signal-to-noise ratio (S/N) of individual binary systems and possibly the slow gravitational-wave inspiral that led to the present-day parameters. So far, most studies have made use of custom-made codes that have not been made publicly available.

We believe that the large renewed interest in LISA and the stellar-origin sources it may detect will lead to many more studies in the near future that would need similar computations. This leads to a significant amount of redundancy, which at best results in extra work for each individual—and at worst leads to an increased chance of introducing mistakes and inconsistencies when translating the necessary expressions to software.

LEGWORK is an open-source Python package designed to streamline the process of making predictions of LISA detection rates for stellar-origin binaries such that it is as fast, reliable, and simple as possible. With LEGWORK, one can evolve the orbits of a binary or a collection of binaries and calculate their strain amplitudes for any range of frequency harmonics. One can compute the sensitivity curve for LISA or other future gravitational-wave detectors (e.g., TianQin’s curve, or that of a custom instrument) and use it to compute the S/N of a collection of sources. Furthermore, LEGWORK provides tools to visualize all of the results with easy-to-use plotting functions. Finally, LEGWORK is fully tested to check for consistency in the derivations described below.

Specifically, we implement the post-Newtonian expressions by Peters & Mathews (1963) and Peters (1964) for the evolution of binary orbits due to the emission of gravitational



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

waves, equations for the strain amplitudes, and S/Ns of binaries from various papers (e.g., Flanagan & Hughes 1998; Finn & Thorne 2000; Cornish & Larson 2003; Barack & Cutler 2004; Moore et al. 2015) and approximations for the LISA and TianQin sensitivity curves given in Robson et al. (2019) and Huang et al. (2020), respectively. The post-Newtonian expressions are approximately of order 0.5, as they account for the orbital evolution from energy loss due to the emission of GWs, but do not include other effects such as spin-orbit coupling. We find this is an excellent approximation for stellar-origin sources in the LISA band, as any higher-order terms are either zero or negligible.

The open-source nature of the project means that new users as well as seasoned experts in the field can work together in a collaborative setting to consider new features and enhancements to the package as well as check the implementation. At the same time, with our thorough online documentation, derivations, and tutorials, we hope LEGWORK can make this functionality more accessible to the broader scientific community.

We note that LEGWORK is not the only initiative of this kind. We highlight the “Gravitational Wave Universe Toolbox” presented by Yi et al. (2021), which was developed to simulate observations on the GW universe with different detectors covering the full gravitational-wave spectrum and source classes. We also highlight “GWPlotter” by Moore et al. (2015), which provides an interactive plotting tool to compare the sensitivity of different gravitational-wave detectors. LEGWORK differs from the tools listed here, as we have tried to provide tools that are optimized to rapidly make predictions for large populations of stellar-origin sources and we have focused on space-based detectors.

LEGWORK has been developed with stellar-origin binary population studies in mind. We highlight two recent papers that use our package. Wagg et al. (2021) investigates several populations of potential LISA sources (double black holes, black hole neutron stars, and double neutron stars). They use LEGWORK to evolve each synthesized source over the age of the Milky Way, make predictions about the LISA detectable population, and explore how it varies with different binary physics assumptions. Additionally, Thiele et al. (2021) examine the implications of assuming a metallicity-dependent binary fraction for the formation of close double white dwarfs (WDWDs) in the LISA frequency band. They use LEGWORK to calculate the S/N of WDWDs, and in particular, apply a custom noise model that combines the LISA sensitivity curve from Robson et al. (2019) with a new fit for the Galactic confusion noise based on their WDWD population.

LEGWORK can be installed with pip or obtained from the GitHub repository.⁶ All examples shown in this paper and code to reproduce the figures are available in the repository. Instructions for installation and basic usage are provided in the online documentation,⁷ which contains the most up-to-date instructions. LEGWORK is jointly published in the *Journal of Open Source Software* (Wagg et al. 2022).

The paper is organized as follows. In Section 2, we give an overview of the capabilities of the LEGWORK package and its various modules. We detail the derivations of the equations relevant for LEGWORK in Section 3. In Section 4, we outline




Figure 1. Package structure of LEGWORK. Each box represents a module and describes its function. The arrows indicate the interdependencies of the modules. </>

some example use cases of LEGWORK, to demonstrate its use. Finally, in Section 5 we conclude and summarize our work.

2. Package Overview

The LEGWORK package is composed of seven modules that each focus on a particular aspect of calculations useful for gravitational-wave sources that are detectable by space-based detectors. In Figure 1, we illustrate the general structure of the package with each of its modules. The `source` module is the central module of the package and provides a simple interface to the functions in the rest of the modules. For more complex analyses, users may want to interact directly with individual modules, particularly those in the top row of Figure 1, as they comprise the core functionality of LEGWORK. Below, we explain the capabilities of each module in detail.

Evol handles the orbital evolution of a binary due to the emission of gravitational waves. It includes functions for computing the merger times of both circular and eccentric binaries. In addition, you can use this module to evolve binary orbit parameters forward in time with any number or arrangement of time steps. We discuss the relevant equations in Section 3.2.

Strain contains two functions that compute a binary’s gravitational-wave strain and characteristic strain amplitude, respectively. Each of these functions is capable of computing the strain for an array of binaries at any number of time steps and evaluated at any number of frequency harmonics. We discuss the relevant equations in Section 3.3.

PSD is used for evaluating the effective noise power spectral density of a detector at different frequencies. The module currently contains the LISA and TianQin sensitivity curves that can be tweaked by adjusting parameters such as the observation time, response function, and even the arm length. For the Galactic confusion noise, a user can choose one of three models (Robson et al. 2019; Huang et al. 2020; Thiele et al. 2021), use their own custom model, or turn it off entirely. Additionally, this module allows the user to specify a custom detector sensitivity curve. We discuss the relevant equations in Section 3.4.

SNR uses the functions in `evol`, `strain`, and `psd` to compute the S/N of sources. It contains four functions that cover the permutations of whether a source is circular or eccentric and stationary in frequency space on the timescale of the mission or evolving. We discuss the relevant equations in Section 3.5.

⁶ <https://github.com/TeamLEGWORK/LEGWORK>

⁷ <https://legwork.readthedocs.io/en/latest/>

Visualization contains several wrappers for plotting one- and two-dimensional distributions with histograms, scatter plots, and kernel density estimator (KDE) plots in order to quickly analyze a collection of sources. In addition, it provides functions for plotting sources directly onto a sensitivity curve.

Source provides a direct and simple interface to the functions in other modules through the `Source` Class. You can instantiate this Class with an array of sources and use it to compute their strains or S/Ns directly. Moreover, depending on the user's choice of allowed gravitational-wave luminosity error, the Class dynamically decides on the number of frequency harmonics needed to capture the full signal of each binary and at what eccentricity to no longer consider a binary circular. This Class also provides a quick means of evolving the sources, visualizing the parameters of each source, and allowing you to plot the binaries on the sensitivity curve.

Utils is a collection of miscellaneous utility functions mainly consisting of conversions between variables as well as constants and expressions from Peters (1964). We discuss the relevant equations in Section 3.1.

2.1. Units and Automated Testing

To ensure stability with physical units, all quantities included in LEGWORK use the `astropy.units` module (Astropy Collaboration et al. 2013, 2018). This means that all inputs to LEGWORK can be given in the units of the user's choice and will be automatically converted.

Furthermore, all of the source code in LEGWORK is fully tested with continuous integration in the LEGWORK GitHub repository. We employ several unit tests to ensure consistency between each of the use cases described below. For example, we require that the S/N calculation for circular and stationary binaries produces consistent output whether LEGWORK uses the stationary and circular approximations or not. Similarly, we verify that the antenna patterns described below produce the expected values when averaged over source positions, inclinations, and polarizations.

2.2. Optimizations

We developed LEGWORK with an emphasis on increasing the efficiency of these computations in order to make simulations of large populations of systems tractable. We ensured that the entirety of LEGWORK is vectorized and thus scales well with larger populations. In addition, we made a several specific optimizations to further increase the speed of calculations.

First, we find that the runtime of calculating strains and S/Ns for large populations of sources is mainly limited by the computation of (1) the relative gravitational-wave power in each harmonic for eccentric systems (see Equation (5)) as well as (2) the sensitivity curve of the given detector. Therefore, in order to significantly reduce the runtime of strain and S/N computations, by default LEGWORK automatically interpolates these functions upon instantiation of any `Source` class with a large number of sources. Thereafter, all functions use the tabulated values instead of calculating them exactly.

In certain cases, one can apply approximations in place of the general S/N calculations. Although it is possible to use each of these approximations directly through the `snr` module, the `Source` module will automatically apply the most appropriate function for each individual source. LEGWORK

dynamically classifies each source as one of four types, which cover the permutations of whether a binary is effectively circular or eccentric and whether or not it is stationary in frequency space on the timescale of the LISA mission. Then, when computing the S/N, it applies a different function to each type of the source. This avoids computing unnecessary, time-intensive integrals.

Moreover, for increasingly eccentric sources, gravitational waves are emitted in an increasing number of higher-frequency harmonics. Although the total S/N is formally calculated as the sum of the S/N over an infinite number of harmonics, in practice it is sufficient to only consider a subset. In the `Source` module, the user can provide `gw_lum_tol`, a maximum allowable tolerance for the accuracy of the gravitational-wave luminosity. Given this tolerance, LEGWORK automatically calculates the required number of harmonics to satisfy this tolerance and thus minimize the computation time.

3. Derivations

In this section, we present a derivation of the equations used in LEGWORK. We emphasize that these are not new derivations—on the contrary, they are in fact given frequently in the literature. However, they are often incomplete or unclear, and in some cases they contain spurious constant factors that arise from invalid combinations of previous work. Here, we aim to present a clear, clean, and concise explanation of the expressions we use in LEGWORK.

A `</>` symbol present before an equation directly links to the relevant online LEGWORK documentation for the implementation of that equation, which additionally contains a link to exact code used to reproduce the equation. These derivations are also given in more detail in the LEGWORK documentation, where we show each of the intervening steps in more detail.

3.1. Conversions and Definitions (`utils`)

We start these derivations by defining some useful conversions and definitions. The chirp mass of a binary is the mass quantity measured by LISA and is given by `</>`

$$\mathcal{M}_c \equiv \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}}, \quad (1)$$

where m_1 and m_2 are the primary and secondary masses of the binary.

It is often convenient to convert between orbital frequency, f_{orb} , and the semimajor axis, a , of a binary, and this can be accomplished with Kepler's third law, `</>`

$$a = \left(\frac{G(m_1 + m_2)}{(2\pi f_{\text{orb}})^2} \right)^{1/3}, \quad (2)$$

where G is the gravitational constant. Inversely, `</>`

$$f_{\text{orb}} = \frac{1}{2\pi} \sqrt{\frac{G(m_1 + m_2)}{a^3}}. \quad (3)$$

For circular binaries, gravitational-wave emission occurs at twice the orbital frequency ($f_{\text{GW}} = 2f_{\text{orb}}$). However, for eccentric binaries, we need to consider all frequency harmonics of gravitational-wave emission. These are defined such that the

n th harmonic frequency is

$$f_n \equiv n \cdot f_{\text{orb}}. \quad (4)$$

It will be important to know the relative gravitational-wave power radiated into the n th harmonic for a binary with eccentricity e for the strain and S/N calculations. This is given by (Peters & Mathews 1963, Equation (20))

$$g(n, e) = \frac{n^4}{32} \left\{ \left[J_{n-2}(ne) - 2eJ_{n-1}(ne) + \frac{2}{n}J_n(ne) + 2eJ_{n+1}(ne) - J_{n+2}(ne) \right]^2 + (1 - e^2)[J_{n-2}(ne) - 2J_n(ne) + J_{n+2}(ne)]^2 + \frac{4}{3n^2}[J_n(ne)]^2 \right\}, \quad (5)$$

where $J_n(v)$ is the Bessel function of the first kind. Thus, the *sum* of $g(n, e)$ over all harmonics gives the factor by which the gravitational-wave emission is stronger for a binary of eccentricity e over an equivalent circular binary. This enhancement factor is (Peters & Mathews 1963, Equation (17))

$$F(e) \equiv \sum_{n=1}^{\infty} g(n, e) = \frac{1 + \frac{73}{24}e^2 + \frac{37}{96}e^4}{(1 - e^2)^{7/2}}. \quad (6)$$

A useful rule of thumb is that $F(0.5) \approx 5.0$, or in words, a binary with eccentricity 0.5 loses energy to gravitational waves at 5 times the rate of an equivalent circular binary.

3.2. Orbital Evolution (*evol*)

3.2.1. Circular Binaries

For a circular binary, the orbital evolution due to gravitational-wave emission can be calculated analytically, as the rate at which the separation of the binary shrinks is simply a function of its mass and the current separation (Peters 1964, Equation (5.6))

$$\frac{da}{dt}_{e=0} = -\frac{\beta}{a^3}, \quad (7)$$

where the constant β is defined as

$$\beta(m_1, m_2) \equiv \frac{64}{5} \frac{G^3}{c^5} m_1 m_2 (m_1 + m_2), \quad (8)$$

where c is the speed of light, m_1 is the primary mass, and m_2 is the secondary mass. This gives the semimajor axis of a circular binary as a function of time, t , as (Peters 1964, Equation (5.9))

$$a(t, m_1, m_2) = [a_0^4 - 4t\beta(m_1, m_2)]^{1/4}, \quad (9)$$

where a_0 is the initial semimajor axis. Moreover, we can solve for the merger time, the time until the binary will merge, by setting the final semimajor axis in Equation (9) to zero:

$$t_{\text{merge, circ}} = \frac{a_0^4}{4\beta}. \quad (10)$$

3.2.2. Eccentric Binaries

The orbital evolution is more complex for eccentric binaries, since the semimajor axis and eccentricity both evolve

simultaneously and depend on one another. The final expression cannot be solved analytically and requires numerical integration. The semimajor axis, a , and eccentricity, e , are related as (Peters 1964, Equation (5.11))

$$a(e) = c_0 \frac{e^{12/19}}{(1 - e^2)} \left(1 + \frac{121}{304} e^2 \right)^{870/2299}, \quad (11)$$

where c_0 satisfies the initial conditions such that $a(e_0) = a_0$. The time derivative of the eccentricity, e , is

$$\frac{de}{dt} = -\frac{19}{12} \frac{\beta}{c_0^4} \frac{e^{-29/19} (1 - e^2)^{3/2}}{[1 + (121/304)e^2]^{1181/2299}}, \quad (12)$$

which we can integrate to find $e(t)$ and convert to $a(t)$ using Equation (11).⁸

Inverting this function and applying the fact that we know that $e \rightarrow 0$ when the binary merges gives the merger time (Peters 1964, Equation (5.14))

$$t_{\text{merge}} = \frac{12}{19} \frac{c_0^4}{\beta} \int_0^{e_0} \frac{[1 + (121/304)e^2]^{1181/2299}}{e^{-29/19} (1 - e^2)^{3/2}} de. \quad (13)$$

For very small or very large eccentricities, we approximate this integral using the following expressions (given in unlabeled equations after Peters 1964, Equation (5.14))

$$t_{\text{merge}, e^2 \ll 1} = \frac{c_0^4}{4\beta} \cdot e_0^{48/19}, \quad (14)$$

$$t_{\text{merge}, (1-e^2) \ll 1} = \frac{768}{425} \frac{a_0^4}{4\beta} (1 - e_0^2)^{7/2}. \quad (15)$$

The standard threshold employed by LEGWORK for small eccentricities is $e = 0.15$, and that for large eccentricities is $e = 0.9999$ (as this approximates t_{merge} with an error below roughly 2%), though we note that this can be customized by the user if desired.

In addition, we implement the fit to Equation (13) from Mandel (2021) that approximates the merger time as

$$t_{\text{merge}} \approx t_{\text{merge, circ}} (1 - e_0^2)^{7/2} \times (1 + 0.27e_0^{10} + 0.33e_0^{20} + 0.2e_0^{1000}), \quad (16)$$

which gives t_{merge} with an error below 3% for eccentricities below 0.9999. We additionally add a rudimentary polynomial fit to further reduce this error to below 0.5%. The user may specify whether to use this fit or perform the full integral when calculating merger times in LEGWORK.

3.3. Strains (*strain*)

3.3.1. Characteristic Strain

The strength of a gravitational wave in a detector at any one moment is determined by the strain amplitude, h_0 . However, for stellar-origin sources at millihertz frequencies, the signal can be present in the detector for many years. This means that the n th harmonic of the binary will spend approximately f_n/\dot{f}_n seconds (or f_n^2/\dot{f}_n cycles) in the vicinity of a frequency f_n (Finn & Thorne 2000). This leads to the signal “accumulating” at the frequency f_n .

⁸ https://legwork.readthedocs.io/en/latest/api/legwork.evol.evol_ecc.html

Therefore, to account for the integration of the signal over the mission, we instead use the “characteristic” strain amplitude of the n th harmonic, $h_{c,n}$, which is the term present in the general S/N equation. This can be related to the strain amplitude in the n th harmonic, $h_{0,n}$, as (e.g., Finn & Thorne 2000; Moore et al. 2015)⁹

$$h_{c,n}^2 = \left(\frac{f_n^2}{\dot{f}_n} \right) h_{0,n}^2, \quad (17)$$

The characteristic strain represents the strain measured by the detector over the duration of the mission (approximated as a single broadband burst), while the strain amplitude is the strength of the GW emission at each instantaneous moment. For a stellar-mass binary, the characteristic strain in the n th harmonic is given by (e.g., Barack & Cutler 2004, Equation (56); Flanagan & Hughes 1998, Equation (5.1))

$$h_{c,n}^2 = \frac{1}{(\pi D_L)^2} \left(\frac{2G}{c^3} \frac{\dot{E}_n}{\dot{f}_n} \right), \quad (18)$$

where D_L is the luminosity distance to the source (note that, for Milky Way sources or any sources with redshift ~ 0 , this is simply the distance to the source), \dot{E}_n is the power radiated in the n th harmonic, and \dot{f}_n is the rate of change of the n th harmonic frequency.

The power radiated in the n th harmonic can be expressed as (Peters & Mathews 1963, Equation (19))

$$\dot{E}_n = \frac{32}{5} \frac{G^4}{c^5} \frac{m_1^2 m_2^2 (m_1 + m_2)}{a^5} g(n, e), \quad (19)$$

where $g(n, e)$ is given in Equation (5). By substituting a for f_{orb} (using Equation (3)) and applying the definition of the chirp mass (Equation (1)), we obtain a more useful form for making gravitational-wave predictions:

$$\dot{E}_n(\mathcal{M}_c, f_{\text{orb}}, e) = \frac{32}{5} \frac{G^{7/3}}{c^5} (2\pi f_{\text{orb}} \mathcal{M}_c)^{10/3} g(n, e). \quad (20)$$

The last term needed to define the characteristic strain in Equation (18) is the rate of change of the n th harmonic frequency as a result of gravitational-wave inspiral, which we can write as

$$\dot{f}_n = \frac{df_n}{da} \frac{da}{dt}. \quad (21)$$

We can find an expression for df_n/da by substituting Equation (3) into Equation (4) and differentiating

$$\frac{df_n}{da} = -\frac{3n}{4\pi} \frac{\sqrt{G(m_1 + m_2)}}{a^{5/2}}. \quad (22)$$

The rate at which the semimajor axis decreases is (Peters 1964, Equation (5.6))

$$\frac{da}{dt} = -\frac{64}{5} \frac{G^3 m_1 m_2 (m_1 + m_2)}{c^5 a^3} F(e). \quad (23)$$

Substituting Equations (22) and (23) into Equation (21) gives an expression for \dot{f}_n

$$\dot{f}_n = \frac{48n}{5\pi} \frac{G^{7/2}}{c^5} (m_1 m_2 (m_1 + m_2)^{3/2}) \frac{F(e)}{a^{11/2}}, \quad (24)$$

which, as above with \dot{E}_n , we can recast using Kepler’s third law and the definition of the chirp mass: \langle / \rangle

$$\dot{f}_n = \frac{48n}{5\pi} \frac{(G\mathcal{M}_c)^{5/3}}{c^5} (2\pi f_{\text{orb}})^{11/3} F(e). \quad (25)$$

With definitions of both \dot{E}_n and \dot{f}_n , we are now in a position to find an expression for the characteristic strain by plugging Equation (20) and Equation (25) into Equation (18): \langle / \rangle

$$h_{c,n}^2 = \frac{2^{5/3}}{3\pi^{4/3}} \frac{(G\mathcal{M}_c)^{5/3}}{c^3 D_L^2} \frac{1}{f_{\text{orb}}^{1/3}} \frac{g(n, e)}{nF(e)}. \quad (26)$$

3.3.2. Strain

In order to obtain an expression for the strain amplitude of gravitational waves in the n th harmonic, we can use Equation (17) and plug in Equations (25) and (26) \langle / \rangle

$$h_n^2 = \frac{2^{28/3}}{5} \frac{(G\mathcal{M}_c)^{10/3}}{c^8 D_L^2} \frac{g(n, e)}{n^2} (\pi f_{\text{orb}})^{4/3}. \quad (27)$$

3.3.3. Amplitude Modulation for Orbit-averaged Sources

Because the LISA detectors are not stationary and instead follow an Earth-trailing orbit, the antenna pattern of LISA is not isotropically distributed or stationary. For sources that have unknown positions, inclinations, and polarizations, we use an average for the detector. However, for sources where these quantities are known, we can consider the amplitude modulation of the strain due to the average motion of LISA’s orbit.

We write that the position of the source on the sky is given by the ecliptic coordinates (θ, ϕ) , the inclination of a source is ι , and the polarization of a source (determined by its orientation relative to the detector) is given by ψ . We follow the results of Cornish & Larson (2003) to define the amplitude modulation. However, we adapt their expression to remain in the frequency domain and follow the conventions of more recent papers (e.g. Babak et al. 2021, Equation (67)) to write the amplitude modulation as \langle / \rangle

$$A_{\text{mod}}^2 = \frac{1}{4} (1 + \cos^2 \iota)^2 \langle F_+^2 \rangle_{\text{orb}} + \cos^2 \iota \langle F_\times^2 \rangle_{\text{orb}}, \quad (28)$$

where $\langle F_+^2 \rangle_{\text{orb}}$ and $\langle F_\times^2 \rangle_{\text{orb}}$, the orbit-averaged detector responses, are defined as \langle / \rangle , \langle / \rangle

$$\begin{aligned} \langle F_+^2 \rangle_{\text{orb}} &= \frac{1}{4} (\cos^2 2\psi \langle D_+^2 \rangle_{\text{orb}} \\ &\quad - \sin 4\psi \langle D_+ D_\times \rangle_{\text{orb}} \\ &\quad + \sin^2 2\psi \langle D_\times^2 \rangle_{\text{orb}}), \end{aligned} \quad (29)$$

$$\begin{aligned} \langle F_\times^2 \rangle_{\text{orb}} &= \frac{1}{4} (\cos^2 2\psi \langle D_\times^2 \rangle_{\text{orb}} \\ &\quad + \sin 4\psi \langle D_+ D_\times \rangle_{\text{orb}} \\ &\quad + \sin^2 2\psi \langle D_+^2 \rangle_{\text{orb}}), \end{aligned} \quad (30)$$

⁹ Note that this is factor of 2 different from Finn & Thorne (2000). This is because the factor of 2 is already included in the Robson et al. (2019) sensitivity curve and so is removed here.

and \langle / \rangle , \langle / \rangle , \langle / \rangle

$$\langle D_+ D_\times \rangle_{\text{orb}} = \frac{243}{512} \cos \theta \sin 2\phi (2 \cos^2 \phi - 1) \times (1 + \cos^2 \theta), \quad (31)$$

$$\langle D_\times^2 \rangle_{\text{orb}} = \frac{3}{512} (120 \sin^2 \theta + \cos^2 \theta + 162 \sin^2 2\phi \cos^2 \theta), \quad (32)$$

$$\langle D_+^2 \rangle_{\text{orb}} = \frac{3}{2048} [487 + 158 \cos^2 \theta + 7 \cos^4 \theta - 162 \sin^2 2\phi (1 + \cos^2 \theta)^2]. \quad (33)$$

The orbital motion of LISA smears the source frequency by roughly 10^{-4} mHz due to the antenna pattern changing as the detector orbits, the Doppler shift from the motion, and the phase modulation from the $+$ and \times polarizations in the antenna pattern. Generally, the modulation reduces the strain amplitude because the smearing in frequency reduces the amount of signal buildup at the true source frequency.

We note that the amplitude modulation is only implemented in LEGWORK for quasi-circular binaries, to remain consistent with the calculation in Cornish & Larson (2003). Since the expected use case of LEGWORK is estimation of the detectability of large populations of millihertz stellar-remnant binaries, for which predictions are uncertain by orders of magnitude in some cases, an extension of Cornish & Larson (2003) that includes eccentric binaries is out of the current scope of LEGWORK.

3.4. Sensitivity Curves (*psd*)

3.4.1. LISA

For the LISA sensitivity curve, we use the equations from Robson et al. (2019). The *effective* strain spectral density of the noise is defined as \langle / \rangle

$$S_n(f) \equiv \frac{P_n(f)}{\mathcal{R}(f)} + S_c(f), \quad (34)$$

where $P_n(f)$ is the power spectral density of the detector noise and $\mathcal{R}(f)$ is the sky- and polarization-averaged signal response function of the instrument. Alternatively, if we expand out $P_n(f)$, approximate $\mathcal{R}(f)$, and simplify, we find (Robson et al. 2019, Equation (1))

$$S_n(f) = \frac{10}{3L^2} \left(P_{\text{OMS}}(f) + \frac{4P_{\text{acc}}(f)}{(2\pi f)^4} \right) \times \left(1 + \frac{6}{10} \left(\frac{f^2}{f_*^2} \right) \right) + S_c(f), \quad (35)$$

where $L = 2.5$ Gm is the detector arm length, $f_* = c/2\pi L = 19.09$ mHz is the transfer frequency,

$$P_{\text{OMS}}(f) = (1.5 \times 10^{-11} \text{ m})^2 \times \left(1 + \left(\frac{2 \text{ mHz}}{f} \right)^4 \right) \text{ Hz}^{-1} \quad (36)$$

is the single-link optical metrology noise (Robson et al. 2019, Equation (10)),

$$P_{\text{acc}}(f) = (3 \times 10^{-15} \text{ ms}^{-2})^2 \left(1 + \left[\frac{0.4 \text{ mHz}}{f} \right]^2 \right) \times \left(1 + \left[\frac{f}{8 \text{ mHz}} \right]^4 \right) \text{ Hz}^{-1} \quad (37)$$

is the single-test mass acceleration noise (Robson et al. 2019, Equation (11)), and

$$S_c(f) = A f^{-7/3} e^{-f^\alpha + \beta f \sin(\kappa f)} \times [1 + \tanh(\gamma(f_k - f))] \text{ Hz}^{-1} \quad (38)$$

is the galactic confusion noise (Robson et al. 2019, Equation (14)), where the amplitude A is fixed as 9×10^{-45} and the various parameters change over time and are listed in Table 1 of Robson et al. (2019). LEGWORK allows the user to opt to use the Robson et al. (2019) confusion noise or a custom function for the confusion noise, or even to remove the confusion noise entirely.

3.4.2. TianQin

For the TianQin sensitivity curve, we use the power spectral density given in Equation (13) of Huang et al. (2020) \langle / \rangle

$$S_N(f) = \frac{10}{3L^2} \left[\frac{4S_a}{(2\pi f)^4} \left(1 + \frac{10^{-4} \text{ Hz}}{f} \right) + S_x \right] \times \left[1 + 0.6 \left(\frac{f}{f_*} \right)^2 \right], \quad (39)$$

where $L = \sqrt{3} \times 10^5$ km is the arm length, $S_a = 1 \times 10^{-30} \text{ m}^2 \text{ s}^{-4} \text{ Hz}^{-1}$ is the acceleration noise, $S_x = 1 \times 10^{-24} \text{ m}^2 \text{ Hz}^{-1}$ is the displacement measurement noise, and $f_* = c/2\pi L$ is the transfer frequency. Note that Equation (39) includes an extra factor of 10/3 compared to Equation (13) of Huang et al. (2020). Huang et al. (2020) absorb this factor into the waveform rather than include it in the power spectral density. We include it to match the convention used by Robson et al. (2019) for the LISA sensitivity curve (see the factor of 10/3 in Equation (35)) so that the sensitivity curves can be compared fairly.

3.5. S/N for LISA (*snr*)

We note that this section draws heavily from Flanagan & Hughes (1998) Section 2.3.

3.5.1. Defining General S/N

In order to calculate the S/N for a given source of gravitational waves (GWs) in a six-link LISA detector, we need to consider the following parameters:

1. position of the source on the sky: (θ, ϕ)
2. direction from the source to the detector: (ι, β)
3. orientation of the source, which fixes the polarization of the GW: ψ
4. the distance from the source to the detector: D_L

Then, assuming a matched filter analysis of the GW signal $s(t) + n(t)$ (where $s(t)$ is the signal and $n(t)$ is the noise), which

relies on knowing the shape of the signal, the S/N, ρ , is given in the frequency domain as

$$\rho^2(D_L, \theta, \phi, \psi, \iota, \beta) = \frac{\langle s(t)^* s(t) \rangle}{\langle n(t)^* n(t) \rangle}, \quad (40)$$

$$= 2 \int_{-\infty}^{+\infty} \frac{|\tilde{s}(f)|^2}{P_n(f)} df, \quad (41)$$

$$= 4 \int_0^\infty \frac{|\tilde{s}(f)|^2}{P_n(f)} df, \quad (42)$$

where $\tilde{s}(f)$ is the Fourier transform of the signal, $s(t)$, and $P_n(f)$ is the one-sided power spectral density of the noise defined as $\langle n(t)^* n(t) \rangle = \int_0^\infty \frac{1}{2} P_n(f) df$ (see Robson et al. 2019, Equation (2)). Here, $\tilde{s}(f)$ is implicitly also dependent on $D_L, \theta, \phi, \psi, \iota$, and β as

$$|\tilde{s}(f)|^2 = |F_+(\theta, \phi, \psi) \tilde{h}_+(f, D_L, \iota, \beta) + F_\times(\theta, \phi, \psi) \tilde{h}_\times(f, D_L, \iota, \beta)|^2, \quad (43)$$

where $F_{+, \times}$ are the “plus” and “cross” antenna patterns of the LISA detector to the “plus” and “cross” strains, $h_{+, \times}$. Note throughout that any parameters discussed with the subscript $x_{+, \times}$ refer to both x_+ and x_\times .

3.5.2. Average over Position and Polarization

Now, we can consider averaging over different quantities. In LISA’s case, when averaged over all angles and polarizations, the antenna patterns are orthogonal and thus $\langle F_+ F_\times \rangle = 0$. This means we can rewrite Equation (43) as

$$|\tilde{s}(f)|^2 = |F_+(\theta, \phi, \psi) \tilde{h}_+(f, D_L, \iota, \beta)|^2 + |F_\times(\theta, \phi, \psi) \tilde{h}_\times(f, D_L, \iota, \beta)|^2, \quad (44)$$

which can then be applied to Equation (40) to give

$$\langle \rho \rangle_{\theta, \phi, \psi}^2 = 4 \int_0^\infty df \int \frac{d\Omega_{\theta, \phi}}{4\pi} \int \frac{d\psi}{\pi} \frac{|F_+ \tilde{h}_+|^2 + |F_\times \tilde{h}_\times|^2}{P_n(f)}. \quad (45)$$

From Robson et al. (2019), we can write the position and polarization average of the signal response function of the instrument, \mathcal{R} , as

$$\mathcal{R} = \langle F_+ F_+^* \rangle = \langle F_\times F_\times^* \rangle,$$

$$\text{where } \langle F_{+, \times} F_{+, \times}^* \rangle = \int \frac{d\Omega_{\theta, \phi}}{4\pi} \int \frac{d\psi}{\pi} |F_{+, \times}|^2. \quad (46)$$

Then, combining Equations (45) and (46), we find¹⁰

$$\langle \rho \rangle_{\theta, \phi, \psi}^2 = 4 \int_0^\infty df \mathcal{R}(f) \left(\frac{|\tilde{h}_+|^2 + |\tilde{h}_\times|^2}{P_n(f)} \right). \quad (47)$$

3.5.3. Average over Orientation

Now, we can average over the orientation of the source: (ι, β), noting that the averaging is independent of the distance D_L . With this in mind, we can rewrite $|\tilde{h}_+|^2 + |\tilde{h}_\times|^2$ in terms of two functions $|\tilde{H}_+|^2$ and $|\tilde{H}_\times|^2$, where $\tilde{h}_{+, \times} = \tilde{H}_{+, \times}/D_L$. Given this,

averaging over the source direction gives

$$\langle \rho \rangle_{(\theta, \phi, \psi), (\iota, \beta)}^2 = \frac{4}{D_L^2} \int_0^\infty df \mathcal{R}(f) \int \frac{d\Omega_{\iota, \beta}}{4\pi} \frac{|\tilde{H}_+|^2 + |\tilde{H}_\times|^2}{P_n(f)}, \quad (48)$$

where we would like to express $\tilde{H}_{+, \times}(f)^2$ in terms of the energy spectrum of the GW. To do this, we note that the local energy flux of GWs at the detector is given by (e.g., Press & Thorne 1972, Equation (6))

$$\frac{dE}{dAdt} = \frac{1}{16\pi} \frac{c^3}{G} \left[\left(\frac{dh_+}{dt} \right)^2 + \left(\frac{dh_\times}{dt} \right)^2 \right], \quad (49)$$

where the bar indicates an average over several cycles of the wave, which is appropriate for LISA sources. We can transform Equation (49) using Parseval’s theorem, where we can write

$$\int_{-\infty}^{+\infty} dt \int dA \frac{dE}{dAdt} = \int_0^\infty df \times \int dA \frac{c^3}{G} \frac{\pi f^2}{2} (|\tilde{h}_+|^2 + |\tilde{h}_\times|^2). \quad (50)$$

Note that the factor of frequency squared comes from the Fourier transform of the square of the time derivative in Equation (50). Now, since $A = D_L^2 \Omega$ and $|\tilde{h}_{+, \times}|^2 = |\tilde{H}_{+, \times}|^2/D_L^2$, we find

$$|\tilde{h}_{+, \times}|^2 dA = |\tilde{H}_{+, \times}|^2 d\Omega_{\iota, \beta}. \quad (51)$$

Then we can write Equation (50) in terms of $|\tilde{H}_{+, \times}|^2$ as

$$\int_{-\infty}^{+\infty} dt \int dA \frac{dE}{dAdt} = \int_0^\infty df \frac{\pi f^2 c^3}{2G} \times \int d\Omega (|\tilde{H}_+|^2 + |\tilde{H}_\times|^2). \quad (52)$$

Alternatively, by using Equation (51) and performing a Fourier transform, we can also write that

$$\int_{-\infty}^{+\infty} dt \int dA \frac{dE}{dAdt} = \int_0^\infty df \int d\Omega \frac{dE}{d\Omega df}. \quad (53)$$

From inspection of Equation (52) and Equation (53), we can write the spectral energy flux as

$$\int d\Omega \frac{dE}{d\Omega df} = \frac{\pi f^2 c^3}{2G} \int d\Omega (|\tilde{H}_+|^2 + |\tilde{H}_\times|^2). \quad (54)$$

3.5.4. Fully Averaged S/N

We are now in a position to write an expression for the fully averaged S/N. Note that, for brevity, we write $\langle \rho \rangle^2$ when referring to $\langle \rho \rangle_{(\theta, \phi, \psi), (\iota, \beta)}^2$. The application of Equation (54) to Equation (48) yields

$$\langle \rho \rangle^2 = \frac{4G}{c^3 D_L^2} \int_0^\infty df \frac{1}{P_n(f)/\mathcal{R}(f)} \int \frac{d\Omega}{4\pi} \frac{dE}{d\Omega df} \frac{2}{\pi f^2}. \quad (55)$$

This simplifies nicely to

$$\langle \rho \rangle^2 = \frac{2G}{\pi^2 c^3 D_L^2} \int_0^\infty df \frac{dE}{df} \frac{1}{f^2 P_n(f)/\mathcal{R}(f)}. \quad (56)$$

Finally, noting that $dE/df = dE/dt \times dt/df = \dot{E}/\dot{f}$, we can use the definition of the characteristic strain from Equation (18)

¹⁰ Note that this is written in Flanagan & Hughes (1998) for the LIGO response function, which is $\mathcal{R} = \langle F_{+, \times} \rangle^2 = 1/5$.

to finish up our position, direction, and orientation/polarization-averaged S/N as

$$\langle \rho \rangle^2 = \int_0^\infty df \frac{h_c^2}{f^2 P_n(f)/\mathcal{R}(f)} = \int_0^\infty df \frac{h_c^2}{f^2 S_n(f)}, \quad (57)$$

where we have defined the effective power spectral density of the noise as $S_n(f) = P_n(f)/\mathcal{R}(f)$.

It is also important to note that this is only the S/N for a circular binary for which we need only consider the $n=2$ harmonic. In the general case, a binary could be eccentric and requires a sum over *all* harmonics. Thus, we can generalize Equation (57) to eccentric binaries with $\langle \rho \rangle$

$$\langle \rho \rangle^2 = \sum_{n=1}^\infty \langle \rho_n \rangle^2 = \sum_{n=1}^\infty \int_0^\infty df_n \frac{h_{c,n}^2}{f_n^2 S_n(f_n)}, \quad (58)$$

where $h_{c,n}$ is defined in Equation (26) and S_n in Equation (35).

3.5.5. S/N Approximations

Although Equation (58) can be used for every binary, it can be useful to consider different cases in which we can avoid unnecessary sums and integrals. There are four possible cases for binaries in which we can use increasingly simple expressions for the S/N. Binaries can be circular and stationary in frequency space.

Circular binaries emit only in the $n=2$ harmonic, and so the sum over harmonics can be removed. Stationary binaries have $f_{n,i} \approx f_{n,f}$, and so the small interval allows one to approximate the integral. Note that we refer to nonstationary binaries as “evolving” here, though many papers also use the term “chirping.”

For an evolving and eccentric binary, no approximation can be made and the S/N is found using Equation (58).

For a evolving and circular binary, the sum can be removed and so the S/N is found as $\langle \rho \rangle$

$$\langle \rho \rangle_{e,e}^2 = \frac{1}{4} \int_{f_{2,i}}^{f_{2,f}} \frac{h_{c,2}^2}{f_2^2 S_n(f_2)} df_2. \quad (59)$$

For a stationary and eccentric binary, we can approximate the integral:

$$\langle \rho \rangle_{e,s}^2 = \frac{1}{4} \sum_{n=1}^\infty \lim_{\Delta f \rightarrow 0} \int_{f_n}^{f_n + \Delta f_n} \frac{h_{c,n}^2}{f_n^2 S_n(f_n)} df_n, \quad (60)$$

$$= \frac{1}{4} \sum_{n=1}^\infty \frac{\Delta f_n \cdot h_{c,n}^2}{f_n^2 S_n(f_n)}, \quad (61)$$

$$= \frac{1}{4} \sum_{n=1}^\infty \frac{\dot{f}_n \Delta T \cdot h_{c,n}^2}{f_n^2 S_n(f_n)}, \quad (62)$$

$$= \frac{1}{4} \sum_{n=1}^\infty \left(\frac{\dot{f}_n}{f_n^2} h_{c,n}^2 \right) \frac{T_{\text{obs}}}{S_n(f_n)}, \quad (63)$$

where we have applied (17) to convert between strains. This gives the following expression: $\langle \rho \rangle$

$$\langle \rho \rangle_{e,s}^2 = \frac{1}{4} \sum_{n=1}^\infty \frac{h_n^2 T_{\text{obs}}}{S_n(f_n)}. \quad (64)$$

Finally, for a stationary and circular binary, the S/N is $\langle \rho \rangle$

$$\langle \rho \rangle_{c,s}^2 = \frac{1}{4} \frac{h_2^2 T_{\text{obs}}}{S_n(f_2)}. \quad (65)$$

For S/N calculations that take into account the amplitude modulation due to LISA’s orbital motion (Section 3.3.3), we apply the calculations as described above but include the modulation in either the strain or characteristic strain as is appropriate.

4. Use Cases

In this section, we demonstrate LEGWORK’s range of capabilities through a series of example use cases. The plots and results in each subsection are reproduced directly in online demos in the LEGWORK documentation, which are each based on individual Jupyter notebooks. These tutorials are linked at the start of each subsection with $\langle \rho \rangle$ symbols.

4.1. Computing the S/N of a Binary System $\langle \rho \rangle$

The most fundamental use case of LEGWORK is to compute the S/N of an individual binary system. This can be accomplished in LEGWORK with only two lines of code—one line to set up the source and another to compute the S/N. As an example, one could consider a binary with the parameters

$$m_1 = m_2 = 10 M_\odot, d = 8 \text{ kpc}, \\ f_{\text{orb}} = 10^{-4} \text{ Hz}, e = 0.2,$$

where m_1, m_2 are the primary and secondary masses, d is the distance to the source, f_{orb} is the orbital frequency, and e is the eccentricity. As we do not specify a position, polarization, or inclination, LEGWORK will calculate the S/N averaged over these quantities. If a user specifies the position, then if either of the polarization or inclination are not specified, they will be randomly generated. LEGWORK does not allow users to specify polarizations or inclinations without positions, as we presume that the user must know the position if they know the polarization or inclination of the source. One can now instantiate a source in LEGWORK using these parameters (leaving the mission parameters as the default values, such that we compute the S/N for a 4 yr LISA mission). As shown in the linked demo, LEGWORK quickly computes that the S/N of this binary is 4.49. In the background, LEGWORK decides which S/N approximation is most applicable given the eccentricity of the binary and whether it is stationary in frequency space.

This can be generalized to a population of many binary systems with ease. Instead of inputting single values for each parameter, one can input arrays of values where each entry corresponds to a different binary. As an example, we can take the same parameters as above for three different binaries but vary the primary as $m_1 = [5, 10, 15] M_\odot$. Using LEGWORK, we find that the S/N for each of these cases is $\rho = [2.47, 4.49, 7.85]$. This also need not be limited to a 4 yr LISA mission. With LEGWORK, we can additionally specify various parameters for the detector. For example, using LEGWORK, we can find that, for a 5 yr TianQin mission with no confusion noise, the S/N for each of these cases is $\rho = [1.07, 1.95, 3.41]$.




Figure 2. The horizon distance for circular, stellar-mass binaries in a 4 yr LISA mission. The filled contours indicate the horizon distance for different orbital frequencies and chirp masses. We add white dotted contours at 8 kpc, 50 kpc, 800 kpc, and 40 Mpc to highlight the distances to the center of the Milky Way, the Magellanic Clouds, the Andromeda galaxy, and the nearest ground-based gravitational-wave detection (GW170817; Abbott et al. 2017), respectively. The diagonal black line shows the frequencies and chirp masses at which the merger time is equal to the observation time. This line emphasizes the sharp contrast for the horizon distance for binaries that merge before the LISA mission finishes observing (to the right of the line). The horizontal black lines indicate the approximate location of some common double compact-object types on this plot, with the assumed masses labeled below in solar masses. [⟨⟩](#)

4.2. Horizon Distance [⟨⟩](#)

A common question to consider with stellar-mass sources in LISA is how far away a certain source could be detected. In other words, what is the horizon distance beyond which a source no longer has an S/N greater than some chosen threshold. We can explore this question using LEGWORK.

Let us compute the horizon distance for a grid of chirp masses and orbital frequencies. First, we can recall that the S/N of a source is inversely proportional to its distance from LISA and so we can find the horizon distance, D_{hor} , as

$$D_{\text{hor}} = \frac{\rho(D)}{\rho_{\text{detect}}} \cdot D, \quad (66)$$

where $\rho(D)$ is the S/N at some distance D and ρ_{detect} is the threshold above which we consider a source to be detectable. For the purpose of this example, we will set $\rho_{\text{detect}} = 7$. We can then, as a function of chirp mass and frequency, determine the maximum distance for which the source of interest is detected based on this S/N threshold.

This can be most efficiently accomplished using LEGWORK’s [Source](#) class, since LEGWORK can then compute the merger times and S/N of each source with only two lines of code. We convert the S/Ns to horizon distances using Equation (66) and plot the result in Figure 2. The shape of the LISA sensitivity curve is clearly reflected in the horizon

distance. This is because circular sources only emit at a single frequency and thus every feature in the sensitivity curve has a strong effect on the S/N (and therefore horizon distance) for circular sources. We also see that, once the merger time of a source is shorter than the LISA mission length (shown by the black dotted line), the horizon distance sharply decreases. This is because, if a source merges before the LISA mission concludes, it has less time to accumulate signal and thus has a lower S/N and horizon distance.

One can use Figure 2 to estimate the horizon distance for any circular source of interest. To illustrate this, we add solid black lines to indicate the typical chirp masses of some possible stellar-mass gravitational-wave sources, along with white dotted lines to show the distances to nearby galaxies as well as the nearest ground-based gravitational-wave detection (GW170817, Abbott et al. 2017). For example, we see that a circular NSNS with an orbital frequency greater than 1 mHz is detectable in the Magellanic Clouds.

4.3. The Role of Eccentricity [⟨⟩](#)

The role of eccentricity is important to consider in the detection of gravitational waves with LISA and other space-based detectors, as sources can still have significant eccentricity during their inspiral phase. A high eccentricity has two major effects on the S/N of a gravitational-wave source in LISA, and we can investigate these effects using LEGWORK.




Figure 3. An illustration of the effect of eccentricity on the detectability of a LISA source. The three sets of points are colored by their eccentricity, and each individual point corresponds to a harmonic frequency, where its height above the curve gives its S/N. We annotate each set of points with its total S/N and overlay the LISA sensitivity curve. The dotted vertical lines indicate the frequency at which the majority of the gravitational-wave signal is concentrated. </>

Let us consider three hypothetical systems that are identical apart from their eccentricities, e_i :

$$m_1 = m_2 = 0.6 M_\odot, f_{\text{orb}} = 1.5 \text{ mHz}, d = 15 \text{ kpc}, \\ e_i = \{0.0, 0.6, 0.9\},$$

Using LEGWORK to calculate their S/Ns in a 4 yr LISA mission, ρ_i , we find

$$\rho_i = \{31.7, 50.2, 38.8\}.$$

We see two effects on the S/N here. First, increasing the eccentricity from essentially circular to $e = 0.6$ results in a higher S/N ($\rho = 31.7 \rightarrow \rho = 50.2$). This is because an eccentric binary has enhanced energy emission via gravitational waves (Peters & Mathews 1963). This means that an eccentric binary will not only inspiral faster than an otherwise identical circular binary, but also will always produce a stronger gravitational-wave strain. We discuss the enhancement factor and its exact dependence on eccentricity in more detail in Section 3 (see specifically Equation (6)).

The second effect is more intriguing. We see that increasing the eccentricity from $e = 0.6$ to $e = 0.9$ results in a relative decrease in S/N ($\rho = 50.2 \rightarrow \rho = 38.8$). The reason for this is that eccentric binaries emit gravitational waves at many harmonic frequencies (unlike circular binaries, which emit predominantly twice the orbital frequency). This leads to the gravitational-wave signal being diluted over many frequencies higher than the orbital frequency, where the higher the eccentricity, the more harmonics are required to capture all of the gravitational luminosity (see Figure 3 of Peters & Mathews 1963). Therefore, if the eccentricity is too high, the majority of the signal may be emitted at a frequency to which LISA is less sensitive.

We can illustrate this point with LEGWORK by calculating the S/N at each individual frequency harmonic using the `SNR` module. We plot this distribution of signal over different frequency harmonics with the LISA sensitivity curve overlaid in Figure 3 using LEGWORK’s `visualisation` module. A point is plotted for each harmonic of each source that has an S/N greater than unity and such that its height above the sensitivity curve corresponds to its S/N.




Figure 4. The merger time for a binary with both a primary and secondary mass of $10 M_\odot$ over different eccentricities and orbital frequencies. The dashed line indicates a merger time of 4 yr, the default LISA mission length. </>

From Figure 3, we can better understand why a source with $e = 0.9$ has a lower S/N than the same source with $e = 0.6$. From the dotted lines, we can note that the signal from the $e = 0.9$ source is concentrated at a frequency of around 40 mHz. The LISA sensitivity at this point is much weaker than the 6 mHz at which the $e = 0.6$ source is concentrated. Therefore, although the strain from a more eccentric binary is stronger, the S/N is lower due to the increased noise in the LISA detector.

Overall, we can therefore conclude that, for LISA sources of this nature, higher eccentricity will produce more detectable binaries only if the orbital frequency is not already at or above the minimum of the LISA sensitivity curve.

Another consideration for more massive binaries is whether the increased eccentricity will cause the binary to merge before the mission ends, which would cause a significant decrease in S/N. We can also use LEGWORK to find how the merger time of a source varies with frequency and eccentricity over a grid of sources.

We plot the results of this calculation in Figure 4. This plot shows that, for most eccentricities, the merger time is largely determined by the orbital frequency. However, for high eccentricities ($e > 0.8$), the eccentricity leads to a significant reduction in the merger time. Additionally, we can see that any binary that is to the right of the dashed line on this plot at the start of a 4 yr LISA mission would merge before the mission ended. Therefore, if increasing a binary’s eccentricity moved it to the right of this line, its S/N would decrease significantly.

4.4. Comparing Gravitational-wave Detectors </>

It may also be useful to consider how changing the specifications of the LISA detector, or using a different detector entirely, could affect the S/N of a particular source. LEGWORK is capable of adjusting the LISA mission specifications or using a different sensitivity curve, and thus we can use it to explore these differences.

As a first step, we can use LEGWORK to plot a series of sensitivity curves in Figure 5. We show the LISA sensitivity curve for the default 4 yr mission length, but also illustrate how the curve changes for shorter mission lengths. At 0.5 and 2 yr, we see a stronger noise level around 3 mHz as a




Figure 5. The strain spectral density of the LISA detector, with different specifications (Robson et al. 2019), and the TianQin detector (Huang et al. 2020). We show the LISA curve for three different mission lengths and once with an approximate response function. [</>](#)

result of the increased Galactic confusion noise. This noise decreases with increasing mission length, because more individual foreground sources can be resolved and thus removed from the confusion noise. We also see that using an approximated response function smooths out the sensitivity curve at higher frequencies. Finally, the TianQin curve is higher than the 4 yr LISA curve until around 5 mHz, beyond which it has a lower noise level.

Although comparing the sensitivity curves would suffice for a stationary and circular source (since it would remain at a single frequency), LEGWORK can also be used to see how the relative S/N between two detectors changes over a range of eccentricities and frequencies.

Using LEGWORK, we can compute the S/N of a grid of sources (spanning a range of frequencies and eccentricities) for both detectors. In Figure 6, we show the ratio of the S/N in LISA to the S/N in TianQin. This plot shows that, for circular binaries, the S/N of the source in LISA is stronger up to an orbital frequency of approximately 2.5 mHz, beyond which the S/N of the source is stronger in TianQin. This transition frequency becomes lower with increasing eccentricity, as one would expect given that eccentric sources emit more at higher harmonics and thus higher frequencies.

4.5. Track S/N of a Binary over Time [</>](#)

As a binary inspirals, its orbital frequency and eccentricity change and this in turn affects the S/N of the binary. For this use case, we will demonstrate how LEGWORK can be used to track the evolution of these parameters and pinpoint the moment at which a binary becomes detectable.

Let us consider a binary with the following initial parameters:

$$m_1 = m_2 = 15M_{\odot}, d = 20 \text{ kpc}, \\ f_{\text{orb},i} = 3 \times 10^{-5} \text{ Hz}, e_i = 0.5,$$

and use LEGWORK’s `evol` module to evolve the system until 100 yr before its merger with 1000 linearly spaced time steps, recording the eccentricity and frequency at each time step.

We plot this evolution of the eccentricity and frequency in the top two panels of Figure 7 as a function of the time before




Figure 6. The ratio of the S/N in LISA (for a 4 yr mission) to the S/N in TianQin. The dashed line indicates the transition at which the S/N is equal in both detectors. We annotate the regions in which either detector has a higher S/N and also annotate the mass and distance of each source in the grid. [</>](#)

the merger. We see that the binary circularizes and increases its orbital frequency as it inspirals, as we would expect.

To take this a step further, we can consider the binary at each time step to be a separate source with the current eccentricity and frequency. It is then trivial to use LEGWORK to calculate the S/N for each of these “sources” and thus attain the S/N evolution, which we plot in the last panel of Figure 7.

We see that the S/N increases monotonically over time and sharply increases as the binary approaches its merger. Around 1 Myr before the merger, the S/N reaches the detection threshold and thus could then be seen by a 4 yr LISA mission.

Note that LEGWORK could also be used in this way to find the S/N of any system in which the orbital evolution is known. Thus, for a triple system or a binary experiencing gas drag (as long as the evolutions of the eccentricity and orbital frequency are known), LEGWORK is entirely capable of calculating the S/N evolution.

4.6. LISA Verification Binaries [</>](#)

Kupfer et al. (2018) present the LISA verification binaries, a collection of known binary systems that have gravitational waves that are strong enough to be detected by LISA. In LEGWORK, we provide easy access to this data through the `VerificationBinaries` class, which is a subclass of `Source`. This means that the class works identically to `Source`, but it has the verification binary data (such as their masses and orbital frequencies) preloaded into the variables.

In addition to the base variables, this class also includes the designation of each binary and the S/N that Kupfer et al. (2018) computed. We note that this S/N differs from the S/N that LEGWORK gives for each binary. This is because Kupfer et al. (2018) run a full, detailed LISA simulation for these binaries, while we follow the orbit-averaged approach (see Section 3.3.3), since the former would be intractable for any large number of sources.

As an example of how you could use this data, we use the `VerificationBinaries` class to plot the sources on the




Figure 7. The evolution of a binary system’s eccentricity (top), orbital frequency (middle), and S/N (bottom). Each panel is annotated with the constant parameters of the system. The S/N is calculated for a 4 yr LISA mission. We use a line in the bottom panel at $S/N = 7$ to highlight the moment at which the source becomes detectable. </>

LISA sensitivity curve with the S/N calculated by Kupfer et al. (2018) for a 4 yr LISA mission and color the binaries by their primary mass. We see that the verification binaries tend to be detected with frequencies between 1–10 mHz and have masses estimated to be between $0.1\text{--}1 M_{\odot}$.

5. Conclusion and Summary

We have presented LEGWORK, a package designed to aid in calculations for stellar-origin binary sources of millihertz gravitational-wave observatories like LISA. We outlined the implementation of orbital evolution due to gravitational-wave emission, gravitational-wave strain, S/N, and visualization modules, and provided a detailed derivation for each of the equations required for each module. Finally, we provided several use-case examples for how LEGWORK can be used to better understand the detectability of compact-object binaries.




Figure 8. The LISA verification binaries from Kupfer et al. (2018) plotted on the LISA sensitivity curve (Robson et al. 2019) for a 4 yr LISA mission. Points are colored by their primary mass. </>

We are grateful to Stas Babak, Floor Broekgaarden, Tom Callister, Will Farr, Yi-Ming Hu, Stephen Justham, Valeria Korol, Mike Lau, Tyson Littenberg, Ilya Mandel, Alberto Sesana, Lieke van Son, the CCA GW group, the BinCosmos group, and the COMPAS group for stimulating discussions that influenced and motivated us to complete this project. We thank the BinCosmos group for testing an early version of the package and providing useful feedback. In particular, we thank Lieke van Son for her innovation in inventing the name LEGWORK! T.W. thanks Floor Broekgaarden for first suggesting that he investigate LISA and the derivation of the S/N calculation. K.B. thanks Shane L. Larson and Kyle Kremer for *several* conversations about the S/N derivation. We additionally thank the anonymous reviewer for providing useful comments, which helped to improve the quality and clarity of this paper.

This project was funded in part by the National Science Foundation under NSF grant No. 2009131, the European Union’s Horizon 2020 research and innovation program from the European Research Council (ERC, grant agreement No. 715063), and by the Netherlands Organization for Scientific Research (NWO) as part of the Vidi research program BinWaves with project number 639.042.728. We further acknowledge the Black Hole Initiative funded by a generous contribution of the John Templeton Foundation and the Gordon and Betty Moore Foundation. The Flatiron Institute is funded by the Simons Foundation.

Software: LEGWORK is written in Python, available from <https://www.python.org>. We make use of the following Python packages: matplotlib (Hunter 2007), NumPy (Harris et al. 2020), Astropy (<http://www.astropy.org>; Astropy Collaboration et al. 2013, 2018), Seaborn (Waskom 2021), SciPy (Virtanen et al. 2020), Numba (Lam et al. 2015), and Schwimmbad (Price-Whelan & Foreman-Mackey 2017). This paper was compiled using showyourwork (Luger et al. 2020).

ORCID iDs

T. Wagg <https://orcid.org/0000-0001-6147-5761>
K. Breivik <https://orcid.org/0000-0001-5228-6598>
S. E. de Mink <https://orcid.org/0000-0001-9336-2825>

References

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017, [PhRvL](#), **119**, 161101
- Andrews, J. J., Breivik, K., Pankow, C., D’Orazio, D. J., & Safarzadeh, M. 2020, [ApJL](#), **892**, L9
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, [AJ](#), **156**, 123
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, [A&A](#), **558**, A33
- Babak, S., Gair, J., Sesana, A., et al. 2017, [PhRvD](#), **95**, 103012
- Babak, S., Hewitson, M., & Petiteau, A. 2021, [arXiv:2108.01167](#)
- Barack, L., & Cutler, C. 2004, [PhRvD](#), **69**, 082005
- Barack, L., & Cutler, C. 2007, [PhRvD](#), **75**, 042003
- Bartolo, N., Caprini, C., Domcke, V., et al. 2016, [JCAP](#), **2016**, 026
- Begelman, M. C., Blandford, R. D., & Rees, M. J. 1980, [Natur](#), **287**, 307
- Belczynski, K., Benacquista, M., & Bulik, T. 2010, [ApJ](#), **725**, 816
- Bellovary, J. M., Cleary, C. E., Munshi, F., et al. 2019, [MNRAS](#), **482**, 2913
- Berti, E., Cardoso, V., & Will, C. M. 2006, [PhRvD](#), **73**, 064030
- Breivik, K., Coughlin, S., Zevin, M., et al. 2020a, [ApJ](#), **898**, 71
- Breivik, K., Mingarelli, C. M. F., & Larson, S. L. 2020b, [ApJ](#), **901**, 4
- Caldwell, R. R., Smith, T. L., & Walker, D. G. E. 2019, [PhRvD](#), **100**, 043513
- Caprini, C., Hindmarsh, M., Huber, S., et al. 2016, [JCAP](#), **2016**, 001
- Chen, W.-C., Liu, D.-D., & Wang, B. 2020, [ApJL](#), **900**, L8
- Christian, P., & Loeb, A. 2017, [MNRAS](#), **469**, 930
- Cornish, N. J., & Larson, S. L. 2003, [PhRvD](#), **67**, 103001
- Fang, X., Thompson, T. A., & Hirata, C. M. 2019, [ApJ](#), **875**, 75
- Finn, L. S., & Thorne, K. S. 2000, [PhRvD](#), **62**, 124021
- Flanagan, É. É., & Hughes, S. A. 1998, [PhRvD](#), **57**, 4535
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, [Natur](#), **585**, 357
- Huang, S.-J., Hu, Y.-M., Korol, V., et al. 2020, [PhRvD](#), **102**, 063021
- Hunter, J. D. 2007, [CSE](#), **9**, 90
- Klein, A., Barausse, E., Sesana, A., et al. 2016, [PhRvD](#), **93**, 024003
- Korol, V., Koop, O., & Rossi, E. M. 2018, [ApJL](#), **866**, L20
- Korol, V., Rossi, E. M., & Barausse, E. 2019, [MNRAS](#), **483**, 5518
- Korol, V., Rossi, E. M., Groot, P. J., et al. 2017, [MNRAS](#), **470**, 1894
- Korol, V., Toonen, S., Klein, A., et al. 2020, [A&A](#), **638**, A153
- Kremer, K., Breivik, K., Larson, S. L., & Kalogera, V. 2017, [ApJ](#), **846**, 95
- Kremer, K., Chatterjee, S., Breivik, K., et al. 2018, [PhRvL](#), **120**, 191103
- Kupfer, T., Korol, V., Shah, S., et al. 2018, [MNRAS](#), **480**, 302
- Lam, S. K., Pitrou, A., Seibert, S., et al. 2015, in Proc. of the Second Workshop on the LLVM Compiler Infrastructure in HPC, LLVM ’15 (New York: Association for Computing Machinery), 7
- Lamberts, A., Blunt, S., Littenberg, T. B., et al. 2019, [MNRAS](#), **490**, 5888
- Lamberts, A., Garrison-Kimmel, S., Hopkins, P. F., et al. 2018, [MNRAS](#), **480**, 2704
- Lau, M. Y. M., Mandel, I., Vigna-Gómez, A., et al. 2020, [MNRAS](#), **492**, 3061
- Liu, J. 2009, [MNRAS](#), **400**, 1850
- Liu, J., & Zhang, Y. 2014, [PASP](#), **126**, 211
- Luger, R., Bedell, M., Foreman-Mackey, D., et al. 2020, [arXiv:2110.06271](#)
- Mandel, I. 2021, [RNAAS](#), **5**, 223
- Moore, C. J., Chua, A. J. K., & Gair, J. R. 2017, [CQGra](#), **34**, 195009
- Moore, C. J., Cole, R. H., & Berry, C. P. L. 2015, [CQGra](#), **32**, 015014
- Nelemans, G., Yungelson, L. R., & Portegies Zwart, S. F. 2001, [A&A](#), **375**, 890
- Nissanke, S., Vallisneri, M., Nelemans, G., & Prince, T. A. 2012, [ApJ](#), **758**, 131
- Peters, P. C. 1964, [PhRv](#), **136**, 1224
- Peters, P. C., & Mathews, J. 1963, [PhRv](#), **131**, 435
- Press, W. H., & Thorne, K. S. 1972, [ARA&A](#), **10**, 335
- Price-Whelan, A. M., & Foreman-Mackey, D. 2017, [JOSS](#), **2**, 357
- Robson, T., Cornish, N. J., & Liu, C. 2019, [CQGra](#), **36**, 105011
- Roebber, E., Buscicchio, R., Vecchio, A., et al. 2020, [ApJL](#), **894**, L15
- Ruiter, A. J., Belczynski, K., Benacquista, M., Larson, S. L., & Williams, G. 2010, [ApJ](#), **717**, 1006
- Sesana, A., Lamberts, A., & Petiteau, A. 2020, [MNRAS](#), **494**, L75
- Shao, Y., & Li, X.-D. 2021, [ApJ](#), **920**, 81
- Thiele, S., Breivik, K., & Sanderson, R. E. 2021, [arXiv:2111.13700](#)
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, [NatMe](#), **17**, 261
- Wagg, T., Breivik, K., & de Mink, S. 2022, [JOSS](#), **7**, 3998
- Wagg, T., Broekgaarden, F. S., de Mink, S. E., et al. 2021, [arXiv:2111.13704](#)
- Waskom, M. 2021, [JOSS](#), **6**, 3021
- Yi, S.-X., Nelemans, G., Brinkerink, C., et al. 2021, [arXiv:2106.13662](#)
- Yu, S., & Jeffery, C. S. 2010, [A&A](#), **521**, A85



Erratum: LEGWORK: A Python Package for Computing the Evolution and Detectability of Stellar-origin Gravitational-wave Sources with Space-based Detectors (2022, ApJS, 260, 52)

T. Wagg^{1,2,3} , K. Breivik⁴ , and S. E. de Mink^{2,3,5}

¹ Department of Astronomy, University of Washington, Seattle, WA, 98195, USA

² Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

³ Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Straße 1, D-85741 Garching, Germany

⁴ Center for Computational Astrophysics, Flatiron Institute, 162 Fifth Avenue, New York, NY, 10010, USA

⁵ Anton Pannekoek Institute for Astronomy and GRAPPA, University of Amsterdam, NL-1090 GE Amsterdam, The Netherlands

Received 2022 August 4; published 2022 September 15

We discovered a typo in our approximation to the general signal-to-noise ratio (S/N) equation, which was carried forward in each S/N approximation equation. Equations (59)–(65) of the published article each have an erroneous factor of 1/4 included. This mistake was not reflected in the code implementation or the online documentation and so will not have affected anyone using the code (e.g., Hamers 2021; Thiele et al. 2021; Wagg et al. 2021; Biscoveanu et al. 2022; Chen et al. 2022). For reference, the functions that implement these approximations are `snr_circ_stationary`, `snr_ecc_stationary` and `snr_circ_evolving` in the `snr`⁶ module and one can see that the typo we describe is not included in the code. The corrected section (replacing part of Section 3.5.5 of the published article) follows below: For a evolving and circular binary, the sum can be removed and so the S/N found as

$$\langle \rho \rangle_{c,e}^2 = \int_{f_{2,i}}^{f_{2,f}} \frac{h_{c,2}^2}{f_2^2 S_n(f_2)} df_2. \quad (59)$$

For a stationary and eccentric binary we can approximate the integral.

$$\langle \rho \rangle_{e,s}^2 = \sum_{n=1}^{\infty} \lim_{\Delta f \rightarrow 0} \int_{f_n}^{f_n + \Delta f_n} \frac{h_{c,n}^2}{f_n^2 S_n(f_n)} df_n, \quad (60)$$

$$= \sum_{n=1}^{\infty} \frac{\Delta f_n \cdot h_{c,n}^2}{f_n^2 S_n(f_n)}, \quad (61)$$

$$= \sum_{n=1}^{\infty} \frac{\dot{f}_n \Delta T \cdot h_{c,n}^2}{f_n^2 S_n(f_n)}, \quad (62)$$

$$= \sum_{n=1}^{\infty} \left(\frac{\dot{f}_n h_{c,n}^2}{f_n^2} \right) \frac{T_{\text{obs}}}{S_n(f_n)}, \quad (63)$$

where we have applied Equation (17) to convert between strains. This gives the following expression

$$\langle \rho \rangle_{e,s}^2 = \sum_{n=1}^{\infty} \frac{h_n^2 T_{\text{obs}}}{S_n(f_n)}. \quad (64)$$




Finally, for a stationary and circular binary the S/N is

$$\langle \rho \rangle_{c,s}^2 = \frac{h_2^2 T_{\text{obs}}}{S_n(f_2)}. \quad (65)$$

⁶ <https://github.com/TeamLEGWORK/LEGWORK/blob/main/legwork/snr.py>



ORCID iDs

T. Wagg  <https://orcid.org/0000-0001-6147-5761>
K. Breivik  <https://orcid.org/0000-0001-5228-6598>
S. E. de Mink  <https://orcid.org/0000-0001-9336-2825>

References

Biscoveanu, S., Kremer, K., & Thrane, E. 2022, arXiv:[2206.15390](#)
Chen, H.-L., Chen, X., & Han, Z. 2022, [ApJ](#), **935**, 9
Hamers, A. S. 2021, [RNAAS](#), **5**, 275

Thiele, S., Breivik, K., & Sanderson, R. E. 2021, arXiv:[2111.13700](#)
Wagg, T., Broekgaarden, F. S., de Mink, S. E., et al. 2021, arXiv:[2111.13704](#)