# **IOP**science

This content has been downloaded from IOPscience. Please scroll down to see the full text.

Download details:

IP Address: 18.118.166.98 This content was downloaded on 09/05/2024 at 16:19

Please note that terms and conditions apply.

You may also like:

Pulsed terahertz tomography S Wang and X-C Zhang

Photonic terahertz technology Alvydas Lisauskas, Torsten Löffler and Hartmut G Roskos

## Concepts and Applications of Nonlinear Terahertz Spectroscopy

## Concepts and Applications of Nonlinear Terahertz Spectroscopy

Thomas Elsaesser, Klaus Reimann and Michael Woerner

Max-Born-Institute, Berlin, Germany

Morgan & Claypool Publishers

Copyright © 2019 Morgan & Claypool Publishers

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the publisher, or as expressly permitted by law or under terms agreed with the appropriate rights organization. Multiple copying is permitted in accordance with the terms of licences issued by the Copyright Licensing Agency, the Copyright Clearance Centre and other reproduction rights organizations.

#### Rights & Permissions

To obtain permission to re-use copyrighted material from Morgan & Claypool Publishers, please contact info@morganclaypool.com.

ISBN978-1-64327-216-0 (ebook)ISBN978-1-64327-213-9 (print)ISBN978-1-64327-214-6 (mobi)

DOI 10.1088/2053-2571/aae931

Version: 20190201

IOP Concise Physics ISSN 2053-2571 (online) ISSN 2054-7307 (print)

A Morgan & Claypool publication as part of IOP Concise Physics Published by Morgan & Claypool Publishers, 1210 Fifth Avenue, Suite 250, San Rafael, CA, 94901, USA

IOP Publishing, Temple Circus, Temple Way, Bristol BS1 6HG, UK

### Contents

Author biographies		vii
Symbols		
1	Introduction	1-1
1.1	Linear optical response	1-2
1.2	Low-energy excitations in condensed matter	1-6
	1.2.1 Electronic excitations	1-7
	1.2.2 Vibrational excitations	1-12
1.3	Charge transport in solids	1-14
	References	1-15
2	Terahertz technology	2-1
2.1	Generation of coherent terahertz radiation	2-1
	2.1.1 Terahertz lasers	2-2
	2.1.2 THz generation by optical frequency conversion	2-6
	2.1.3 Photoconductive switches	2-11
2.2	Phase-resolved detection of terahertz transients	2-12
	2.2.1 Photoconductive antennas	2-14
	2.2.2 Electro-optic sampling	2-14
2.3	Linear terahertz spectroscopy and imaging	2-19
	References	2-23
3	Nonlinear light-matter interactions	3-1
3.1	Nonlinear optical response in the THz range	3-1
	3.1.1 Theoretical framework	3-1
	3.1.2 Approximations	3-5
3.2	Nonlinear currents in condensed matter	3-10
	3.2.1 Shift currents in systems lacking inversion symmetry	3-10
	3.2.2 Nonlinear currents of electrons in a periodic potential	3-15
	3.2.3 Nonlinear currents without real-space transfer of carriers	3-19
3.3	Quantum coherences in coupled multi-level systems	3-20
3.4	Nonperturbative regime of light-matter interaction	3-22
	References	3-26

4	Methods of nonlinear terahertz spectroscopy	4-1
4.1	Concepts and experimental implementation	4-1
	4.1.1 Optical geometries for nonlinear THz experiments	4-1
	4.1.2 Interaction schemes and pulse sequences	4-4
4.2	Two-dimensional spectroscopy	4-11
	4.2.1 Experimental methods of two-dimensional THz spectroscopy	4-13
	4.2.2 Data representation and analysis	4-15
	References	4-19
5	Nonlinear terahertz spectroscopy of condensed matter	5-1
5.1	Dynamics and couplings of low-energy excitations in liquids and solids	5-1
	5.1.1 Condensed-phase molecular systems	5-1
	5.1.2 Solids and nanostructures	5-6
5.2	Field-driven nonlinear response and charge transport in solids	5-17
	5.2.1 Nonlinear THz response of softmodes in a molecular crystal	5-17
	5.2.2 Polaron transport	5-21
	5.2.3 Coherent high-field transport in GaAs on femtosecond time scales	5-24
	5.2.4 THz-induced interband tunneling of carriers	5-29
	5.2.5 Shift current in LiNbO <sub>3</sub>	5-35
	5.2.6 Nonlinear transport in graphene	5-37
5.3	Conclusions and outlook	5-42
	References	5-43

# Author biographies

#### **Thomas Elsaesser**



Thomas Elsaesser is a director at the Max-Born-Institute, Berlin, Germany, and a full professor for experimental physics at Humboldt University, Berlin. He received a Dr rer. nat. degree from the Technical University of Munich in 1986 and worked there as a research associate until 1993. He spent a postdoc period at AT&T Bell Laboratories, Holmdel, in 1990 and joined the newly established Max-Born-Institute in 1993. His research focuses on

ultrafast phenomena in condensed matter, in particular molecular liquids, biomolecules in their aqueous environment, and inorganic solids and nanostructures. Combined in his experimental work are methods of ultrafast spectroscopy and structure research. Thomas is a fellow of the American Physical Society and the Optical Society of America and has received numerous scientific awards.

#### **Klaus Reimann**



Klaus Reimann has worked since 1999 as a scientist at the Max-Born-Institute, Berlin, Germany, in the field of ultrafast mid-infrared and THz spectroscopy. He received a Dr rer. nat. degree from the University (now Technical University) of Dortmund in 1987. Afterwards he joined the Max-Planck-Institut für Festkörperforschung in Stuttgart and worked there on the physics of semiconductors under high pressures. Having received a

five-year Heisenberg-Stipendium of the Deutsche Forschungsgemeinschaft, he spent this time at the Universität Dortmund and at the University of California at Berkeley doing research on nonlinear optics of semiconductors under high pressure before joining the MBI.

### **Michael Woerner**



Michael Woerner is a department head at the Max-Born-Institute, Berlin, Germany, and holds a lecturer qualification (Habilitation) in physics at Humboldt University, Berlin. He received a Dr rer. nat. degree from the Technical University of Munich in 1991 and worked there as a postdoc until 1993. He then joined the Max-Born-Institute in 1993 and spent a postdoc period at Bell Laboratories (Lucent Technologies), Holmdel, in 1997. Michael's

research focuses on ultrafast phenomena in solids and nanostructures with pioneering work in multi-dimensional spectroscopies in the THz frequency range and in femtosecond x-ray diffraction using laser-driven hard x-ray sources.

# Symbols

Α	vector potential
A	magnitude or relevant component of the vector poten-
	tial (the same convention is used for the magnitude of
	all other vectors)
$\hat{a}, \hat{a}^+$	electron annihilation and creation operators
α	absorption coefficient
В	magnetic field
$c = 299\ 792\ 458\ m/s$	speed of light in vacuum
X	electric susceptibility
$\chi^{(2)}, \chi^{(3)}, \dots$	nonlinear susceptibilities
d	dipole moment
δ	Dirac's $\delta$ function
$\Delta T$	(pump-induced) transmission change
$e = 1.602 \times 10^{-19} \text{ C}$	elementary charge
Ε	electric field
Eem	emitted electric field
Ein	incident electric field
E <sub>loc</sub>	local electric field
E <sub>NI</sub>	nonlinear electric field
E	reflected electric field
E <sub>tr</sub>	transmitted electric field
$\mathbf{E}(t)$	electric field as a function of (real) time t
$\mathbf{E}(\nu_t)$	Fourier transform of $\mathbf{E}(t)$ , function of frequency $\nu_t$
$\mathbf{E}(t,\tau)$	electric field as a function of real time t and delay time $\tau$
$\mathbf{E}(\nu_t, \nu_\tau)$	2D Fourier transform of $\mathbf{E}(t, \tau)$ , function of frequen-
-(-1)-1)	cies $\nu_t$ and $\nu_r$
E	energy
Č <sub>F</sub>	Fermi energy
e Ca	bandgap
$\varepsilon$	dielectric function
$\varepsilon_0 = 8.854 \times 10^{-12} \text{ As/Vm}$	permittivity of vacuum
γ	decoherence rate
$h = 6.626 \times 10^{-34} \text{ Js}$	Planck's constant
$= 4.136 \times 10^{-15} \text{ eVs}$	
$\hbar = h/(2\pi) = 1.055 \times 10^{-34} \text{ Js}$	
$= 6.582 \times 10^{-16} \text{ eVs}$	
$\hat{H}$	Hamiltonian
Ι	intensity
i	current density
k	(electron) wavevector
$k_B = 1.381 \times 10^{-23} \text{ J/K}$	Boltzmann's constant
ĸ	imaginary part of the refractive index
l	thickness
т	effective mass of an electron in a crystal
$m_0 = 9.109 \times 10^{-31} \mathrm{kg}$	mass of a free electron
M	magnetization
$\mu_0 = 4\pi \times 10^{-7}$ Vs/Am	permeability of vacuum

n	(real part of the) refractive index
Ne	electron density
ν	frequency
$\tilde{ u}$	wavenumber
$\omega = 2\pi \nu$	angular frequency
р	momentum
P	electric polarization
$\phi, \psi, \Psi$	wavefunctions
q	phonon wavevector
$\bar{q}_e = -e$	electron charge
Q	electric quadrupole polarization
r, R	position
ρ	density matrix
σ	electrical conductivity
$\sigma^{(2)}, \sigma^{(3)},$	nonlinear conductivities
$\sigma_{a}$	absorption cross section
t	(real) time
Т	waiting time
Т	absolute temperature
$T_0$	linear transmission (before interaction with a pump
	pulse)
τ	delay time
$ au_{\mathrm{Drude}}$	Drude scattering time
tr	trace (of a matrix)
$u_{i,\mathbf{k}}(\mathbf{r})$	cell-periodic part of the Bloch function for an electron
	in band <i>i</i> with wavevector <b>k</b>
V	velocity
$v_F$	Fermi velocity
V	volume
$V(\mathbf{r})$	potential